HOKKAIDO UNIVERSITY

Title	Higenamine 4′ -O-β-d-glucoside in the lotus plumule induces glucose uptake of L6 cells through β2-adrenergic receptor
Author(s)	Kato, Eisuke; Inagaki, Yosuke; Kawabata, Jun
Citation	Bioorganic & Medicinal Chemistry, 23(13), 3317-3321 https://doi.org/10.1016/j.bmc.2015.04.054
Issue Date	2015-07-01
Doc URL	https://hdl.handle.net/2115/66355
Rights	©2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	BMC2015_23_3317-3321_Sp_HUSCAP.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Supplementary material

Higenamine 4'-*O*-β-D-glucoside in the lotus plumule induces glucose uptake of L6 cells through β2-adrenergic receptor

Eisuke Kato [a,] *, Yosuke Inagaki [b], and Jun Kawabata [a]

[a] Laboratory of Food Biochemistry, Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

[b] Nippon Supplement, Inc., Applause Tower 18F, 19-19, Chayamachi, Kita-ku, Osaka 530-0013, Japan

*Corresponding author:

Tel/Fax: +81-11-706-2496

E-mail address: eikato@chem.agr.hokudai.ac.jp



Supplementary Fig 1. ^{1}H-NMR spectra of isolated **1** in CD_3OD (270 MHz).



Supplementary Fig 2. ^{1}H-NMR spectra of isolated (black) and synthetic (red) **1** in D_2O (500 MHz).