



HOKKAIDO UNIVERSITY

Title	Non-invasive prediction of the tumor growth rate using advanced diffusion models in head and neck squamous cell carcinoma patients
Author(s)	Fujima, Noriyuki; Sakashita, Tomohiro; Homma, Akihiro et al.
Citation	Oncotarget, 8(20), 33631-33643 https://doi.org/10.18632/oncotarget.16851
Issue Date	2017-05-16
Doc URL	https://hdl.handle.net/2115/66642
Rights(URL)	https://creativecommons.org/licenses/by/3.0/
Type	journal article
File Information	16851-247336-1-SP.pdf, Supplementary Materials



Non-invasive prediction of the tumor growth rate using advanced diffusion models in head and neck squamous cell carcinoma patients

Supplementary Materials

Supplementary Table 1: Details of signal to noise ratio (SNR) in all b-value images

b-value	SNR
0	94.4 ± 32.2
10	87.5 ± 28.7
20	88.3 ± 30.1
30	84.5 ± 27.6
50	82.1 ± 27.1
80	74.5 ± 26.8
100	70.6 ± 27.3
200	75.2 ± 24.7
400	68.3 ± 23.5
800	53.5 ± 20.7
1000	58.2 ± 20.4
2000	45.4 ± 15.2

Data are mean ± standard deviation. SNR: signal to noise ratio (dimensionless).

Supplementary Table 2: The result of five-fold cross validation analysis

	training set (<i>n</i> = 44)			test set (<i>n</i> = 11)	
	Univariate significant parameters	Final output parameters	Correlation coefficient	Correlation coefficient	ICC
1st fold	ADC, D, D _k , DDC, D ₂ , D ₃ , f ₂ , f ₃	D ₂ , D ₃	0.74	0.7	0.69
2nd fold	ADC, D, D _k , DDC, D ₂ , D ₃ , f ₃	D ₂ , D ₃	0.76	0.69	0.66
3rd fold	ADC, D, D _k , DDC, D ₂ , D ₃ , f ₂ , f ₃	DDC, D ₃	0.71	0.63	0.61
4th fold	ADC, D, D _k , DDC, D ₂ , D ₃ , f ₂ , f ₃	D ₂ , D ₃	0.72	0.66	0.64
5th fold	ADC, D, D _k , DDC, D ₂ , D ₃ , f ₃	D ₂ , D ₃	0.77	0.72	0.68
Average			0.74	0.68	0.66

ICC: intraclass correlation coefficient. Abbreviations in 'Univariate significant parameters' are explained in the Table 2 footnote.