



HOKKAIDO UNIVERSITY

Title	Clinical benefit of 1-year certolizumab pegol (CZP) add-on therapy to methotrexate treatment in patients with early rheumatoid arthritis was observed following CZP discontinuation : 2-year results of the C-OPERA study, a phase III randomised trial
Author(s)	Atsumi, Tatsuya; Tanaka, Yoshiya; Yamamoto, Kazuhiko et al.
Citation	Annals of the rheumatic diseases, 76(8), 1348-1356 https://doi.org/10.1136/annrheumdis-2016-210246
Issue Date	2017-08
Doc URL	https://hdl.handle.net/2115/67068
Rights	This article has been accepted for publication in Atsumi T, Tanaka Y, Yamamoto K, et al., Clinical benefit of 1-year certolizumab pegol (CZP) add-on therapy to methotrexate treatment in patients with early rheumatoid arthritis was observed following CZP discontinuation: 2-year results of the C-OPERA study, a phase III randomised trial, Annals of the Rheumatic Diseases 2017;76:1348-1356, following peer review and can also be viewed on the journal's website at http://ard.bmj.com .
Rights(URL)	https://creativecommons.org/licenses/by-nc/4.0/
Type	journal article
File Information	1348.full.pdf





OPEN ACCESS

EXTENDED REPORT

Clinical benefit of 1-year certolizumab pegol (CZP) add-on therapy to methotrexate treatment in patients with early rheumatoid arthritis was observed following CZP discontinuation: 2-year results of the C-OPERA study, a phase III randomised trial

Tatsuya Atsumi,¹ Yoshiya Tanaka,² Kazuhiko Yamamoto,³ Tsutomu Takeuchi,⁴ Hisashi Yamanaka,⁵ Naoki Ishiguro,⁶ Katsumi Eguchi,⁷ Akira Watanabe,⁸ Hideki Origasa,⁹ Shinsuke Yasuda,¹ Yuji Yamanishi,¹⁰ Yasuhiko Kita,¹¹ Tsukasa Matsubara,¹² Masahiro Iwamoto,¹³ Toshiharu Shoji,¹⁴ Osamu Togo,¹⁴ Toshiyuki Okada,¹⁵ Désirée van der Heijde,¹⁶ Nobuyuki Miyasaka,¹⁷ Takao Koike^{1,18}

Handling editor Tore K Kvien

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2016-210246>).

For numbered affiliations see end of article.

Correspondence to

Professor Tatsuya Atsumi, Division of Rheumatology, Endocrinology and Nephrology, Graduate School of Medicine, Hokkaido University, North 15, West 7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan; at3tat@med.hokudai.ac.jp

Received 20 July 2016

Revised 7 December 2016

Accepted 1 January 2017

Published Online First

2 February 2017



CrossMark

To cite: Atsumi T, Tanaka Y, Yamamoto K, et al. *Ann Rheum Dis* 2017;**76**:1348–1356.

ABSTRACT

Objectives To investigate the clinical impact of 1-year certolizumab pegol (CZP) therapy added to the first year of 2-year methotrexate (MTX) therapy, compared with 2-year therapy with MTX alone.

Methods MTX-naïve patients with early rheumatoid arthritis (RA) with poor prognostic factors were eligible to enter Certolizumab-Optimal Prevention of joint damage for Early RA (C-OPERA), a multicentre, randomised, controlled study, which consisted of a 52-week double-blind (DB) period and subsequent 52-week post treatment (PT) period. Patients were randomised to optimised MTX+CZP (n=159) or optimised MTX+placebo (PBO; n=157). Following the DB period, patients entered the PT period, receiving MTX alone (CZP+MTX→MTX; n=108, PBO+MTX→MTX; n=71). Patients who flared could receive rescue treatment with open-label CZP.

Results 34 CZP+MTX→MTX patients and 14 PBO+MTX→MTX patients discontinued during the PT period. From week 52 through week 104, significant inhibition of total modified total Sharp score progression was observed for CZP+MTX versus PBO+MTX (week 104: 84.2% vs 67.5% (p<0.001)). Remission rates decreased after CZP discontinuation; however, higher rates were maintained through week 104 in CZP+MTX→MTX versus PBO+MTX→MTX (41.5% vs 29.3% (p=0.026), 34.6% vs 24.2% (p=0.049) and 41.5% vs 33.1% (p=0.132) at week 104 in SDAI, Boolean and DAS28(erythrocyte sedimentation rate) remission. CZP retreated patients due to flare (n=28) showed rapid clinical improvement. The incidence of overall adverse events was similar between groups.

Conclusions In MTX-naïve patients with early RA with poor prognostic factors, an initial 1 year of add-on CZP to 2-year optimised MTX therapy brings radiographic and clinical benefit through 2 years, even after stopping CZP.

Trial registration number NCT01451203.

INTRODUCTION

Rheumatoid arthritis (RA) is a chronic inflammatory disease characterised by progressive inflammatory synovitis. This results in the destruction of articular cartilage and marginal bone, which is generally thought to be irreversible.¹ Recent studies have demonstrated that the early treatment of patients with antirheumatic drugs is associated with a reduction in inflammation, greater inhibition of structural damage and better long-term outcomes.^{2–3} Furthermore, early aggressive treatment with biological disease-modifying antirheumatic drugs (bDMARDs), such as antitumour necrosis factors (TNFs), was reported to be highly effective at reducing disease progression.⁴ The effect of treatment discontinuation/tapering following successful inhibition of disease progression as a result of using bDMARDs early in the course of the disease has yet to be fully investigated; however, there is the possibility that the positive disease trajectory may be maintained following treatment cessation.

Certolizumab pegol (CZP) is a humanised anti-TNF antibody fragment conjugated to polyethylene glycol, approved for the treatment of inflammatory diseases, including RA. The efficacy and safety of CZP in combination with methotrexate (MTX) during the early stages of RA was assessed in the Certolizumab-Optimal Prevention of joint damage for Early RA (C-OPERA) study. This study consisted of two periods: a 52-week double-blind (DB) period during which patients received either CZP or placebo (PBO) together with MTX, and a subsequent 52-week post-PBO/CZP treatment (PT) period in which patients received MTX therapy without CZP or PBO. Results from the DB period, which showed significant inhibition of structural damage and a reduction in the severity of RA symptoms following treatment with CZP+MTX compared with PBO+MTX, have been reported.⁴ Here, we report the

2-year overall results including the PT period, which investigated whether the clinical benefits of initial 1-year CZP+MTX therapy were sustained through a subsequent 1-year period where patients received MTX alone.

METHODS

for the DB period were previously re provided in the online supplementary

1203) was a multicentre, DB, ised, parallel-group study conducted in ly described.⁴ Full details of the study e online supplementary materials.

e performed every 8 weeks during the analysis of the PT period was change in total Sharp score (mTSS) from baseline pared the progression of joint damage TX→MTX and PBO+MTX→MTX t damage progression was assessed using and 104; mTSS was evaluated by two n accordance with previously reported l analyses comparing clinical efficacy included disease activity score (DAS)28 on rate (ESR)), simple disease activity joint count (SJC), tender joint count ment Questionnaire Disability Index

on in the
ntion of
matoid
mab pegol;

(HAQ-DI), physician's and patient's global assessments of disease activity (PtGADA), patient's assessment of arthritis pain, ESR and C-reactive protein (CRP) levels. Clinical remission was defined as achieving SDAI ≤ 3.3 , DAS28(ESR) < 2.6 or ≤ 1 on all four of the following criteria (Boolean remission): the number of TJC (in 28 joints), number of SJC (in 28 joints), CRP (mg/dL) and PtGADA (100 mm visual analogue scale (VAS) data converted to cm).

Safety assessments

All safety events during the PT period were recorded as adverse events (AEs) or serious AEs (SAEs). Laboratory tests (haematological, blood chemistry, urinalysis), chest radiographs and ECG were also evaluated.


Statistical analyses

Full details of the statistical analyses can be found in the online supplementary materials. In brief, the full analysis set (FAS; defined as all patients who received ≥ 1 dose of study drug and provided any efficacy data thereafter) was used for all efficacy measurements. Missing data were imputed using linear extrapolation for mTSS and last observation carried forward (LOCF) for all other efficacy variables. Change from baseline in mTSS at weeks 52 and 104 was analysed using an analysis of covariance (ANCOVA) model. Fisher's exact test was used to compare rates of mTSS non-progression (mTSS change from baseline ≤ 0.5) and clinical remission at weeks 52 and 104, between the PBO and CZP groups.

RESULTS

Patient characteristics and disposition

Of the 316 patients who were randomised and received at least one dose of study drug (FAS population), 179 patients entered



Clinical and epidemiological research

the PT period and 131 patients completed the study (figure 1). The proportion of patients completing the PT period (from the patients who entered the PT period) was 68.5% and 80.3% in the CZP+MTX→MTX group and the PBO+MTX→MTX group, respectively (figure 1).

Patient baseline demographics and disease characteristics at period entry (week 52) are shown in supplementary table S1. At DB baseline, and disease characteristics were similar both groups, disease activity at baseline was slightly lower than in the total population baseline through week 52 following P+MTX or PBO+MTX.

Disease progression in the total

At week 52 the change from baseline in the CZP+MTX→MTX group compared with the PBO+MTX→MTX group (0.36 ± 2.70 vs 1.58 ± 4.86 , $p=0.002$), and the rate of radiographic non-progression in the CZP+MTX→MTX group compared with the PBO+MTX→MTX group (82.9% vs 70.7%; exact test).⁴ During the PT period, changes from baseline (mean±SD) for HAQ at week 8 vs 3.01±9.66 ($p=0.001$), erosion at week 8 vs 4.43±4.40 ($p=0.003$) and joint space

narrowing score (0.36 ± 4.27 vs 1.58 ± 7.17 ($p=0.002$)) were lower for the CZP+MTX→MTX group compared with the PBO+MTX→MTX group using linear extrapolation for missing data imputation (figure 2A). A sensitivity analysis using an LOCF imputation method (figure 2A) confirmed the results of the primary analysis (linear extrapolation). At week 104, the proportion of patients with radiographic non-progression (ie, mTSS change from baseline ≤ 0.5) was higher for the CZP+MTX→MTX group compared with the PBO+MTX→MTX group (84.2% vs 67.5%, $p<0.001$). Furthermore, the proportion of patients with rapid radiographic progression (RRP: mTSS yearly change from baseline ≥ 5) at week 104 was lower for the CZP+MTX→MTX group compared with the PBO+MTX→MTX group (3.2% vs 9.6%, $p=0.022$). Subgroup analyses revealed that high baseline mTSS, CRP or TNF was associated with poor week 104 radiographic outcomes in the PBO+MTX→MTX group. The CZP+MTX→MTX group also showed higher inhibition of radiographic progression in these populations (see online supplementary table S3). Consistent with radiographic findings, the proportion of patients with HAQ remission (HAQ ≤ 0.5) at week 104 was numerically higher in the CZP+MTX→MTX group than the PBO+MTX→MTX group (73.0% vs 63.7%, $p=0.09$). In addition, the proportion of the patients who achieved HAQ remission at week 104 was higher in patients who showed non-radiographic progression at week 104 than those who did not (76.7% vs

Radiographic and patient characteristics


	CZP+MTX→MTX			PBO+MTX→MTX		
	Total patients n=159	Patients entering PT period n=108		Total patients n=157	Patients entering PT period n=71	
	DB baseline (Week 0)	DB baseline (Week 0)	PT baseline (Week 52)	DB baseline (Week 0)	DB baseline (Week 0)	PT baseline (Week 52)
Age, mean (SD)	49.4±10.6	48.8±11.2	—	49.0±10.3	48.6±10.8	—
Female, n (%)	129 (81.1)	85 (78.7)	—	127 (80.9)	58 (81.7)	—
Weight, mean (SD)	57.4±11.3	57.0±11.5	—	57.4±10.6	57.4±10.3	—
Weight change from baseline, mean (SD)	22.4±3.9	22.2±3.7	—	22.5±3.7	22.4±3.7	—
Weight change from baseline at week 52, mean (SD)	4.0±2.9	4.4±3.1	—	4.3±2.8	4.4±3.1	—
Weight change from baseline at week 104, mean (SD)	159 (100.0)	71 (100.0)	—	157 (100.0)	108 (100.0)	—
Weight change from baseline at week 104, mean (SD)	153 (96.2)	104 (96.3)	—	146 (93.0)	68 (95.8)	—
Weight change from baseline at week 104, mean (SD)	79 (49.7)	41 (47.2)	—	80 (51.0)	34 (47.9)	—
Weight change from baseline at week 104, mean (SD)	8.4±6.1	7.5±5.8	0.5±1.1	8.9±6.5	7.3±6.1	0.6±1.6
Weight change from baseline at week 104, mean (SD)	8.3±5.3	7.6±4.6	0.3±0.7	8.4±5.3	7.0±4.2	0.4±1.4
Weight change from baseline at week 104, mean (SD)	38.4±25.3	36.3±23.7	12.8±9.9	43.7±28.2	36.5±22.2	15.5±14.3
Weight change from baseline at week 104, mean (SD)	1.29±1.82	1.12±1.51	0.06±0.13	1.52±1.91	1.03±1.39	0.17±0.37
Weight change from baseline at week 104, mean (SD)	130.4±135.4	125.3±135.4	47.7±25.7	185.4±214.9	167.3±204.3	52.5±31.1
Weight change from baseline at week 104, mean (SD)	5.4±1.1	5.2±1.1	1.9±0.8	5.5±1.2	5.1±1.0	2.2±0.7
Weight change from baseline at week 104, mean (SD)	28.7±12.5	27.0±11.2	2.4±2.6	30.0±13.6	24.6±11.3	2.7±3.1
Weight change from baseline at week 104, mean (SD)	1.01±0.64	1.04±0.63	0.14±0.26	1.05±0.69	0.79±0.57	0.07±0.14
Weight change from baseline at week 104, mean (SD)	4.1±7.4	3.8±7.4	3.7±7.4	5.5±15.0	3.2±6.2	3.4±6.3
Weight change from baseline at week 104, mean (SD)	1.9±4.0	1.6±3.9	1.6±3.7	2.5±7.8	1.6±3.3	1.8±3.2
Weight change from baseline at week 104, mean (SD)	2.1±4.6	2.2±4.8	2.2±4.8	2.9±8.3	1.5±4.0	1.6±4.1
Weight change from baseline at week 104, mean (SD)	11.4±3.1	11.3±3.2	10.9±4.1	11.5±2.8	11.5±3.1	11.1±3.7

Weight change from baseline at week 104, mean (SD) was indicated. Data in DB baseline columns represent average during weeks 0–104, whereas data in PT baseline columns represent average during

radiographic symptoms.

Weight change from baseline at week 104, mean (SD) was indicated. Data in DB baseline columns represent average during weeks 0–104, whereas data in PT baseline columns represent average during

Weight change from baseline at week 104, mean (SD) was indicated. Data in DB baseline columns represent average during weeks 0–104, whereas data in PT baseline columns represent average during



baseline in modified total Sharp score (mTSS) at week 104. (B) Cumulative probability plot of mTSS change from baseline in mTSS was analysed using an analysis of actual scores were converted to rank scores, using the treatment group as a factor and baseline rank score as a factor. n-progression (mTSS change from baseline ≤ 0.5) was compared using Fisher's exact test. CZP, certolizumab pegol; LOCF, last observation carried forward; MTX, methotrexate; PBO, placebo.

P+MTX→MTX, and 70.8% vs 49.0% X→MTX).

total population


s achieving SDAI, Boolean and DAS28 calculated throughout both the DB and at the end of the DB period (week 52), were significantly higher in the CZP+MTX→MTX group compared with the PBO+MTX→MTX

al remission observed for the CZP+MTX→MTX group decreased during the first 16 weeks of the PT period and then stabilised from week 68 (week 16 to week 104). The remission rates of the PT period were similar to those observed during the DB period. On the end of DB to the PT period. At the end of the PT period, the rates of clinical remission were higher in the CZP+MTX→MTX group (41.5% vs 29.3% and 4.2% (p=0.049) and 41.5% vs 33.1%

(p=0.132) at week 104 in SDAI, Boolean and DAS28(ESR), respectively). The proportion of patients with low disease activity (DAS28(ESR) ≤ 3.2) was also higher in the CZP+MTX→MTX group compared with the PBO+MTX→MTX group throughout the PT period (see online supplementary figure S1).


Impact of CZP discontinuation in the CZP+MTX→MTX group

The impact of CZP discontinuation was assessed on patients who entered the PT period (PT population) from the CZP+MTX group (n=108). Of these, 74 patients (68.5%) completed the 1-year PT period with MTX therapy (figure 4A). Rates of radiographic non-progression during the PT period (94.4% (102/108); figure 4B) were similar to the rates observed during the DB period (91.7% (99/108)), as were mean changes in mTSS ($\pm$ SD) during the first and second 52-week periods (DB period (weeks 0–52): $-0.03 (\pm 0.80)$; PT period (weeks 52–104): $0.06 (\pm 0.76)$). Conversely, clinical remission rates in the PT population of the CZP+MTX→MTX group showed decreases in SDAI, Boolean and DAS28(ESR) definitions of



C-OPERA was designed to assess the clinical benefit of CZP treatment concomitant with MTX as first-line therapy for early RA, particularly for patients who were considered to require aggressive treatment. Patients who had poor prognostic factors, including a high titre of anticyclic citrullinated peptide antibody in addition to either rheumatoid factor positivity or bone erosions, were eligible to enter the study. C-OPERA was a study composed of two periods. The results from the first year of the study demonstrated the clinical benefit of adding CZP to MTX therapy (DAS28(ESR) remission and radiographic non-progression was achieved in more than 50% and 80% of patients, respectively), suggesting that the introduction of CZP at a very early stage led to substantial therapeutic effects, despite poor prognosis.⁴ In this report, we assessed whether the clinical benefit of initial 1-year CZP+MTX treatment was observed after stopping CZP and continuing with MTX therapy for 1 year.

Figure 4 (A) Retention rate after discontinuation of CZP (Kaplan-Meier plot), (B) modified total Sharp score non-progression rate during and after 52-week CZP therapy for patients who entered the PT period from the CZP mission at DAS28 period and who were CZP, double blind; retention rate; treatment.



the PT period reported here, was that the rate of joint destruction during the initial 1 year of treatment with CZP+MTX was significantly lower than with PBO+MTX→MTX. Radiographic progression, as assessed by the modified total Sharp score, remained lower in the CZP group compared with the PBO+MTX→MTX group. This finding is consistent with the results of the mTSS change from baseline. To ensure the validity of the results, the analysis was repeated using LOCF imputation, and the results were consistent. The rate of progression observed in the CZP group during the PT period was significantly lower than during the DB period in the same population. This suggests that joint destruction may be prevented or delayed by the use of CZP in patients who responded to treatment with CZP+MTX. These data, along with the lower rate of RRP in the CZP group compared with the PBO+MTX→MTX group (see online supplementary table S3), suggest that treatment with CZP+MTX benefits those patients with radiographic non-

progression compared with those with radiographic progression, suggesting that radiographic progression associates with functional disability even during the 2 years of the study. These results suggest that early treatment with CZP during the initial stages of the disease, when rapid joint damage may take place,¹³ could prevent long-term progression of joint damage and functional disability as previously suggested in the 'window of opportunity' concept.^{8 14-16}

In addition to joint damage prevention, the rates of clinical remission throughout the PT period remained higher in the CZP+MTX→MTX group compared with the PBO+MTX→MTX group. After CZP discontinuation, approximately 25% of the patients flared; however, they showed rapid response to CZP retreatment, with recovery to preflare disease activity levels. Although joint destruction was consistently prevented in the PT population following CZP withdrawal, clinical remission was sometimes lost. Discrepancies in clinical and radiographic efficacy have been reported for adalimumab (ADA);¹⁷ similar differences in the clinical and radiographic efficacies of CZP that continue following treatment discontinuation could be responsible for the results observed here. A decrease in remission rate was mainly observed during the first 16 weeks after CZP

Table 2 Summary of treatment-emergent adverse events (TEAE)

CZP+MTX→MTX			PBO+MTX→MTX		
Week 0–52 CZP+MTX n=159	Week 52–104 MTX n=108	Week 0–104 CZP+MTX→MTX n=159	Week 0–52 PBO+MTX n=157	Week 52–104 MTX n=71	Week 0–104 PBO+MTX→MTX n=157
136.2	87.7	223.6	116.0	63.4	179.4
n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
153 (96.2)	85 (78.7)	154 (96.9)	148 (94.3)	57 (80.3)	150 (95.5)
542.0	286.1	442.4	548.2	250.7	444.3
13 (8.2)	4 (3.7)	17 (10.7)	14 (8.9)	4 (5.6)	18 (11.5)
11.0	6.8	9.4	12.9	6.3	10.6
0	0	0	0	0	0
n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
97 (61.0)	45 (41.7)	114 (71.7)	87 (55.4)	30 (42.3)	93 (59.2)
5 (3.1)	0	5 (3.1)	7 (4.5)	1 (1.4)	8 (5.1)
7 (4.4)	1 (0.9)	8 (5.0)	8 (5.1)	2 (2.8)	10 (6.4)
0	0	0	0	0	0
5 (3.1)	2 (1.9)	7 (4.4)	1 (0.6)	0	1 (0.6)
1 (0.6)‡	1 (0.9)‡	2 (1.3)‡	0	0	0
68 (42.8)	12 (11.1)	73 (45.9)	69 (43.9)	9 (12.7)	73 (46.5)

ent-years.

at least one TEAE within System Organ Class/Preferred Term.

rms: alanine aminotransferase increased, aspartate aminotransferase increased, γ -glutamyltransferase increased, hepatic function abnormal, hepatitis, hyperbilirubinaemia, liver disorder, liver function test abnormal; MedDRA V.14.1.
umab pegol; MTX, methotrexate; PBO, placebo.

culate that patients who still produce
e flared as the concentration of CZP

uggest that initial aggressive treatment
be a potential treatment option at the
specially for patients who have a poor
nts achieve their treatment targets,
ould be withdrawn. This treatment
ial to prevent irreversible joint damage,
Es and be a more cost-effective way to
g term. This approach is supported by
e Optimal Protocol for Methotrexate
ination Therapy in Early Rheumatoid
y demonstrated minimal loss of clinical
ant radiographic progression after ADA
ith early RA who initiated combination
hile, The High Induction Therapy with
HIT HARD) study showed better radio-
significant difference in disease activity
n the ADA+MTX→MTX group com-
. ¹⁸ Differences in the results of these
C-OPERA suggest that the condition of
ance' regimen ¹² may be important.
w needed to identify the appropriate
required to achieve continued disease
P withdrawal, and to identify patient
d particularly benefit from first-line
oreover, additional analyses are also
ether this approach has significant clin-
s failing to respond to MTX therapy
s the approach currently recommended
es. ¹⁹

similar rates of SAEs for both the CZP
+MTX→MTX groups over the 2 years

of the C-OPERA study, indicating that there are no major safety
concerns when adding CZP to optimised MTX therapy. Incidences
of AEs and SAEs during the PT period were lower compared with
the initial DB period in both groups. One reason for this may be
'survival bias', where patients discontinued the study because of an
intolerance to the study drugs (CZP and/or MTX) in the first
period, resulting in a lower AE rate in the second. ²⁰

This study has several limitations. In clinical practice, only
patients failing to respond to MTX would receive CZP therapy,
and so it is not known how this approach compares with initial
CZP therapy. No patients received CZP for a full 2 years or
were treated with a reduced dose of CZP, so it was not possible
to compare these treatment regimens with CZP discontinuation.
CZP withdrawal had not been optimised; therefore, there is the
potential for further investigation regarding the appropriate treat-
ment targets and the timing of CZP withdrawal in different
patient populations. There were differences between the study
design of C-OPERA and current RA treatment recommendations.
For example, in clinical practice, treatment recommendations for
patients with poor prognostic factors include using additional
DMARDs in addition to MTX, ^{7 21} which was prohibited in
C-OPERA. C-OPERA was not designed as an intercontinental
global study; thus, it is not known whether these results are gen-
eralisable to ethnicities other than Japanese. In particular, the
MTX dose of 16 mg is low compared with similar RA studies
from the European Union and USA (15–17 mg/week). ^{22–26}
However, when considering differences in patient body weight
and MTX metabolism, ²⁷ a lower dose of MTX in this study may
correspond to the doses used in those previous studies. Finally, as
non-responding patients were eligible to receive rescue treatment
from 24 weeks onwards, we cannot exclude the possibility that a
proportion of the 70 PBO+MTX patients switching to rescue
therapy in the first year may have achieved clinical response if
treated for a longer period of time. ⁴

Overall, these results suggest that patients with early RA would benefit from the addition of CZP to MTX therapy during the early stages of disease, particularly with respect to the prevention of joint destruction. Although this aggressive therapeutic strategy would not be recommended for all patients, it may be a potential option for those patients with a high risk of joint destruction. How to identify these patients requires further investigation.

School of Medicine, Sapporo, Japan
Department of Internal Medicine, University of Occupational and Environmental Health, Fukuoka, Japan
Department of Internal Medicine, Graduate School of Medicine, Keio University School of Medicine, Tokyo, Japan

Department of Internal Medicine, Keio University School of Medicine, Tokyo, Japan
Department of Internal Medicine, Nagoya, Japan
Department of Internal Medicine, Sasebo Chuo Hospital, Sasebo, Nagasaki, Japan

Department of Internal Medicine, The University of Toyama, Toyama, Japan
Department of Internal Medicine, Hiroshima, Japan
Department of Internal Medicine, Kato, Japan
Department of Internal Medicine, Jichi Medical University, Maebashi, Japan

Department of Internal Medicine, Leiden University Medical Centre, Leiden, The Netherlands

Tokyo Medical and Dental University, Tokyo, Japan
Department of Internal Medicine, Sapporo, Hokkaido, Japan

This article has been corrected since it published Online First on August 14, 2017. See the full text on the journal website for corrections.

The authors acknowledge Lilia Marinova MD, PhD, UCB Biopharmaceuticals, and Simon Foulcer, PhD, and Danielle Kavanagh, Cambridge, UK, for medical writing and editing this manuscript for publication, based on the draft provided by the authors. UCB Pharma reviewed only for scientific and legal aspects.

The authors involved in the C-OPERA study, reviewed and approved the manuscript and approved the final draft.

UCB Pharma funded this study and manuscript.

The authors taken part in speakers' bureaus for Astellas, Mitsubishi-Tanabe. KY has received consultancy fees from Mitsubishi-Tanabe, Pfizer, Roche and UCB Pharma; and Abbott, Eisai, Mitsubishi-Tanabe, Pfizer, Santen

and received consultancy fees from AstraZeneca, Asahi Kasei, Novartis; research grants from Abbott, Astellas, BMS, Janssen, Mitsubishi-Tanabe, Nippon Shinyaku, Otsuka, Takeda and Teijin; and has taken part in speakers' bureaus for Eisai, Janssen, Mitsubishi-Tanabe, Pfizer and received consultancy fees from Abbott, Astellas, Mitsubishi-Tanabe, Pfizer, Takeda and UCB Pharma; and UCB Pharma. NI has received research grants from Abbott, Chugai, Eisai, Janssen, Kaken

and has taken part in speakers' bureaus for Abbott, Janssen, Kaken, Mitsubishi-Tanabe, Otsuka, Pfizer, has received research grants from Astellas, Abbvie, Mitsubishi-Tanabe, MSD; has received consultancy fees from Kasei, Astellas, AstraZeneca, Chugai, K, Janssen, Mitsubishi-Tanabe, MSD, Pfizer, and has taken part in speakers' bureaus for Astellas, AstraZeneca, Chugai, Daiichi-Sankyo, Eisai, Eli Lilly, Mitsubishi-Tanabe, MSD, Pfizer, Quintiles, Takeda and UCB Pharma. AW has received

research grants from Daiichi-Sankyo, Dainippon-Sumitomo, Kyorin, Meiji Seika; Shionogi, Taiho, Taisho and Toyama Chemical; and has taken part in speakers' bureaus for Daiichi-Sankyo, Dainippon-Sumitomo, GSK, Mitsubishi-Tanabe, MSD, Pfizer, Shionogi and Taisho-Toyama. HO has received consultancy fees from Astellas and UCB Pharma. SY has received research grant from BMS and taken part in speakers' bureaus for Abbvie, Astellas, Chugai, Eisai, Pfizer, Mitsubishi-Tanabe and Takeda. YS has no competing interests to disclose. YK has received speakers' bureau from Astellas, Chugai and Ono. TM has received speaker honoraria from Pfizer Japan, Janssen Pharmaceutical Co. and Astellas Pharma; and research grants from Quintiles Transnational Japan K.K., Janssen Pharmaceutical Co., Takeda Chemical Industries, Daiichi Sankyo Co., Astellas Pharma, Eli Lilly Japan K.K., MSD Co., Nippon Kayaku Co., Parexel International, Pfizer Japan and Bristol-Myers Squibb. MI has received payment for lectures from Astellas, Chugai, Ono and Tanabe-Mitsubishi; has received research grants from Pfizer and a royalty fee from Chugai. TS is an employee of UCB Pharma; TO is an employee of Astellas. DvdH has received consultancy fees from Abbvie, Amgen, AstraZeneca, Augurex, BMS, Boehringer Ingelheim, Celgene, Centocor, Chugai, Covagen, Daiichi-Sankyo, Eli-Lilly, Galapagos, GSK, Janssen Biologics, Merck, Novartis, Novo-Nordisk, Otsuka, Pfizer, Roche, Sanofi-Aventis, Schering-Plough, UCB Pharma and Vertex; and is the Director of Imaging Rheumatology bv. NM has received research grants from Abbott, Astellas, Chugai, Eisai, Mitsubishi-Tanabe, Pfizer and Takeda. TK has received consultancy fees from Abbvie, Astellas, BMS, Chugai, Daiichi-Sankyo, Eisai, Mitsubishi-Tanabe, Pfizer, Santen, Taisho-Toyama, Takeda, Teijin and UCB Pharma, and has taken part in speakers' bureaus for Abbott, Astellas, BMS, Chugai, Daiichi-Sankyo, Eisai, Mitsubishi-Tanabe, Pfizer, Santen, Taisho-Toyama, Takeda, Teijin and UCB Pharma.

Ethics approval This study was conducted after review and approval by the institutional review board designated by each study site after consideration of the ethical, scientific and medical justification for the conduct of the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- McInnes IB, Schett G. The pathogenesis of rheumatoid arthritis. *N Engl J Med* 2011;365:2205–19.
- Finckh A, Liang MH, van Herckenrode CM, et al. Long-term impact of early treatment on radiographic progression in rheumatoid arthritis: A meta-analysis. *Arthritis Rheum* 2006;55:864–72.
- Lard LR, Visser H, Speyer I, et al. Early versus delayed treatment in patients with recent-onset rheumatoid arthritis: comparison of two cohorts who received different treatment strategies. *Am J Med* 2001;111:446–51.
- Atsumi T, Yamamoto K, Takeuchi T, et al. The first double-blind, randomised, parallel-group certolizumab pegol study in methotrexate-naïve early rheumatoid arthritis patients with poor prognostic factors, C-OPERA, shows inhibition of radiographic progression. *Ann Rheum Dis* 2016;75:75–83.
- van der Heijde D. How to read radiographs according to the Sharp/van der Heijde method. *J Rheumatol* 2000;27:261–3.
- van der Heijde DM, van Riel PL, Nuver-Zwart IH, et al. Effects of hydroxychloroquine and sulphasalazine on progression of joint damage in rheumatoid arthritis. *Lancet* 1989;1:1036–8.
- Smolen JS, Landewé R, Breedveld FC, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2013 update. *Ann Rheum Dis* 2014;73:492–509.
- Smolen JS, Emery P, Fleischmann R, et al. Adjustment of therapy in rheumatoid arthritis on the basis of achievement of stable low disease activity with adalimumab plus methotrexate or methotrexate alone: the randomised controlled OPTIMA trial. *Lancet* 2014;383:321–32.
- Lukas C, van der Heijde D, Fatenajad S, et al. Repair of erosions occurs almost exclusively in damaged joints without swelling. *Ann Rheum Dis* 2010;69:851–5.
- Schett G, Gravalles E. Bone erosion in rheumatoid arthritis: mechanisms, diagnosis and treatment. *Nat Rev Rheumatol* 2012;8:656–64.
- Monti S, Montecucco C, Bugatti S, et al. Rheumatoid arthritis treatment: the earlier the better to prevent joint damage. *RMD Open* 2015;1(Suppl 1):e000057.
- van Vollenhoven RF, Nagy G, Tak PP. Early start and stop of biologics: has the time come? *BMC Med* 2014;12:25.
- Ochi T, Iwase R, Yonemasu K, et al. Natural course of joint destruction and fluctuation of serum C1q levels in patients with rheumatoid arthritis. *Arthritis Rheum* 1988;31:37–43.

Clinical and epidemiological research

- 14 Kyburz D, Gabay C, Michel BA, *et al.* The long-term impact of early treatment of rheumatoid arthritis on radiographic progression: a population-based cohort study. *Rheumatology (Oxford)* 2011;50:1106–10.
- 15 Raza K, Saber TP, Kvien TK, *et al.* Timing the therapeutic window of opportunity in early rheumatoid arthritis: proposal for definitions of disease duration in clinical trials. *Ann Rheum Dis* 2012;71:1921–3.
- 16 Smolen JS, Aletaha D. Rheumatoid arthritis therapy reappraisal: strategies, *s. Nat Rev Rheumatol* 2015;11:276–89.
J, *et al.* Adalimumab reduces hand bone loss independent of clinical response: subanalysis of the PREMIER *isord* 2011;12:54.
et al. Induction therapy with adalimumab plus followed by methotrexate monotherapy up to week 48 alone for DMARD-naïve patients with early rheumatoid igator-initiated study. *Ann Rheum Dis* 2013;72:844–50.
rmester GR, *et al.* Treating rheumatoid arthritis to recommendations of an international task force. *Ann*
owicz M, et al. Survival bias associated with n drug effectiveness evaluation: a comparison of 05;162:1016–23.
SL Jr, *et al.* 2015 American College of Rheumatology of Rheumatoid Arthritis. *Arthritis Rheumatol*
- 22 Bathon JM, Martin RW, Fleischmann RM, *et al.* A comparison of etanercept and methotrexate in patients with early rheumatoid arthritis. *N Engl J Med* 2000;343:1586–93.
- 23 Breedveld FC, Weisman MH, Kavanaugh AF, *et al.* The PREMIER study: a multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;54: 26–37.
- 24 Kavanaugh A, Fleischmann RM, Emery P, *et al.* Clinical, functional and radiographic consequences of achieving stable low disease activity and remission with adalimumab plus methotrexate or methotrexate alone in early rheumatoid arthritis: 26-week results from the randomised, controlled OPTIMA study. *Ann Rheum Dis* 2013;72:64–71.
- 25 St Clair EW, van der Heijde DM, Smolen JS, *et al.* Combination of infliximab and methotrexate therapy for early rheumatoid arthritis: a randomized, controlled trial. *Arthritis Rheum* 2004;50:3432–43.
- 26 Takeuchi T, Yamanaka H, Ishiguro N, *et al.* Adalimumab, a human anti-TNF monoclonal antibody, outcome study for the prevention of joint damage in Japanese patients with early rheumatoid arthritis: the HOPEFUL 1 study. *Ann Rheum Dis* 2014;73:536–43.
- 27 Takahashi C, Kaneko Y, Okano Y, *et al.* THU0118 Methotrexate Polyglutamates in Erythrocytes Correlates with Clinical Response in Japanese Patients with Rheumatoid Arthritis. *Ann Rheum Dis* 2014;73(Suppl 2):218–19.