HOKKAIDO UNIVERSITY

Title	Impact of mutations in DNA gyrase genes on quinolone resistance in Campylobacter jejuni
Author(s)	Changkwanyeun, Ruchirada; Yamaguchi, Tomoyuki; Kongsoi, Siriporn et al.
Citation	Drug Testing and Analysis, 8(10), 1071-1076 https://doi.org/10.1002/dta.1937
Issue Date	2016-10
Doc URL	https://hdl.handle.net/2115/67280
Rights	This is the peer reviewed version of the following article: R. Changkwanyeun, T. Yamaguchi, S. Kongsoi, K. Changkaew, K. Yokoyama, H. Kim, O. Suthienkul, M. Usui, Y. Tamura, C. Nakajima, Y. Suzuki. Impact of mutations in DNA gyrase genes on quinolone resistance in Campylobacter jejuni. Drug Test. Anal. 2016, 8, 1071, which has been published in final form at http://dx.doi.org/10.1002/dta.1937. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	journal article
File Information	Supplementary Figure.pdf, Supplementary Figure

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP




FIG. S1. Inhibitory activity of (A) SIT, (B) MXF, (C) OXO, and (D) NAL. Relaxed pBR322 (0.3 mg) was incubated with 1 μM of GyrA and GyrB in the presence of the indicated quinolone concentration. The reactions were stopped and the DNA products analysed by electrophoresis on 1% agarose gel.

399