




Title	Genetically distinct genogroup IV norovirus strains identified in wastewater
Author(s)	Kitajima, Masaaki; Rachmadi, Andri T.; Iker, Brandon C. et al.
Citation	Archives of virology, 161(12), 3521-3525 https://doi.org/10.1007/s00705-016-3036-z
Issue Date	2016-12
Doc URL	https://hdl.handle.net/2115/67731
Rights	The final publication is available at Springer via http://dx.doi.org/10.1007/s00705-016-3036-z .
Type	journal article
File Information	inal_Revision_GIV NoV InaRoger WWTP Manuscript.pdf



Genetically Distinct Genogroup IV Norovirus Strains Identified in Wastewater

Masaaki Kitajima^{a,*}, Andri T. Rachmadi^a, Brandon C. Iker^{b,†}, Eiji Haramoto^c, Charles P. Gerba^b

^a *Division of Environmental Engineering, Faculty of Engineering, Hokkaido University, Japan*

^b *Department of Soil, Water and Environmental Science, The University of Arizona, 1117 E. Lowell St., Tucson, Arizona 85721, USA*

^c *Interdisciplinary Center for River Basin Environment, University of Yamanashi, 4-3-11 Takeda, Kofu, Yamanashi 400-8511, Japan*

[†]Present affiliation

Amway, 7575 Fulton Street East, Ada, MI 49355-0001, USA

***Corresponding author:** Masaaki Kitajima

Mailing address: Division of Environmental Engineering, Faculty of Engineering, Hokkaido University, North 13, West 8, Kita-ku, Sapporo, Hokkaido 060-8628, Japan

Tel: +81(11)-706-7162 **Fax:** +81(11)-706-7162 **E-mail:** mkitajima@eng.hokudai.ac.jp

Word count **Abstract:** 100 words **Main text:** 1,138 words

Running title: GIV norovirus in wastewater in Arizona

Key words: Norovirus; wastewater treatment plant; seminested PCR; genetic diversity; Arizona

ABSTRACT

We investigated the prevalence and genetic diversity of genogroup IV norovirus (GIV NoV) strains in wastewater in Arizona, United States over a 13-month period. Among 50 wastewater samples tested, GIV NoVs were identified in 13 (26%) of the samples. A total of 47 different GIV NoV strains were identified, which were classified into two genetically distinct clusters: GIV.1 human cluster and a unique genetic cluster closely related to strains previously identified in Japanese wastewater. The results provide additional evidence of the considerable genetic diversity among GIV NoV strains through the analysis of wastewater containing virus strains shed from all populations.

TEXT

Noroviruses (NoVs) are the most significant pathogens associated with water- and food-borne outbreaks of nonbacterial acute gastroenteritis in humans worldwide [1]. They are members of the family *Caliciviridae* and possess a positive-sense, polyadenylated, single-stranded RNA genome with three open reading frames (ORFs) [1]. NoVs show considerable genetic diversity and are currently proposed to be divided into genogroups I–VI (GI–GVI), of which GI, GII, and GIV infect humans [2]. GII strains account for the majority of reported outbreaks of acute gastroenteritis due to NoVs worldwide, and GI strains cause a majority of the remaining cases [3]. GIV or “Alphatron-like” NoVs were first identified in stool samples from sporadic cases of gastroenteritis in the Netherlands [4], and they have been found occasionally in fecal specimens from gastroenteritis patients [5–10]. Interestingly, several studies have documented that GIV NoVs were also identified in carnivores, including domestic dogs and cats [11–14]; nevertheless, they are genetically distinct from the human GIV NoVs (genotype GIV.1) and classified as a separate genotype, GIV.2.

Recent studies also demonstrated the dissemination of GIV NoVs in municipal wastewater and river water in European and Asian countries [10, 15–22], suggesting that not only GI and GII but also GIV strains circulate worldwide and contaminate environmental waters. More importantly, Kitajima et al. (2011) identified several unique GIV NoV strains belonging to a novel genetic cluster in Japanese wastewater samples

collected in 2006, which demonstrated considerable genetic diversity among GIV NoV strains [20].

In North America, no report is available on the occurrence of GIV NoVs in water with limited clinical studies reporting the detection of GIV NoVs in feces of gastroenteritis patients [5]. On the basis of this background, we investigated the prevalence and genetic diversity of GIV NoVs in wastewater in Arizona, the United States, over a 13-month period. GIV NoV genomes in wastewater were detected with a seminested reverse transcription (RT)-PCR assay specific for GIV [19], and the strains were further characterized based on partial capsid gene sequences.

Between July 2011 and July 2012, a total of 50 wastewater grab samples were collected monthly from two WWTP (Plants A and B, utilizing activated sludge and trickling filter, respectively) located in southern Arizona, which included 13 influent and 13 effluent samples from Plant A and 12 influent and 12 effluent samples from Plant B. Viruses in the wastewater samples (100 mL influent and 1,000 mL effluent) were concentrated using an electronegative filter (cat. no. HAWP-090-00; Merck Millipore, Billerica, MA) and a Centriprep YM-50 device (Merck Millipore) to obtain a final volume of approximately 650 μ L as described previously [23].

Viral RNA was extracted from the concentrated wastewater sample spiked with murine norovirus (MNV, S7-PP3 strain; kindly provided by Dr. Y. Tohya, Nihon University, Kanagawa, Japan) process control using the ZR Viral DNA/RNA Kit (Zymo Research, Irvine, CA), according to the manufacturer's protocol. The RT reaction was performed using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA). The seminested PCR assay using COG4F, G4SKF, and G4SKR primers was then

performed to amplify a 340-bp region of the GIV NoV partial capsid gene, as previously described [19]. The second PCR products were separated by electrophoresis on a 2 % agarose gel and visualized under a UV lamp after ethidium bromide staining. All of the second PCR products with expected size were excised from the gel and cloned into the pCR4-TOPO vector (Invitrogen, Carlsbad, CA). At least eight colonies (clones) per sample were selected, and both strands of direct colony PCR products were sequenced using a BigDye Cycle Sequencing Kit version 3.1 and a 3730xl Genetic Analyzer (Applied Biosystems). Nucleotide sequences were assembled using the program SequencherTM version 5.0.1 (Gene Codes Corporation, Ann Arbor, MI, USA) and aligned with Clustal W version 2.1 (<http://clustalw.ddbj.nig.ac.jp/top-e.html>). The distances were calculated by Kimura's two-parameter method [24], and the phylogenetic tree from a bootstrap analysis with 1,000 replicates was generated by the neighbor-joining method. The nucleotide sequences determined in the present study have been deposited in GenBank under accession numbers LC150829–LC150875.

Among 50 wastewater samples tested, GIV NoVs were identified in a total of 13 (26%) samples with the seminested RT-PCR (Table 1). Positive rate for Plant A samples (54% for influent and 15% for effluent) was higher than Plant B samples (17% for influent and 8% for effluent). Use of MNV as an internal control, which was quantified with RT-qPCR [25], showed no substantial inhibition in the molecular detection process in any of the wastewater samples tested in this study (mean recovery efficiency of greater than 75%; specific recovery efficiency data was reported in our previous study analyzing 48 of the 50 samples [23]).

Based on nucleotide sequencing analysis, a total of 47 GIV NoV sequences (1 to 6

different sequences per sample) were identified and classified into two genetically distinct clusters: GIV.1 human cluster and a unique genetic cluster (GIV.new) closely related to strains previously identified in Japanese wastewater reported by Kitajima et al. (2011) (Figure 1). Interestingly, the strains belonging to the GIV.new cluster were identified in 11 of 12 GIV NoV-positive samples, whereas GIV.1 strains were identified in only two samples collected in April 2012. Influent samples collected from Plant A in April 2012 contained both GIV.1 and GIV.new strains (Table 1 and Figure 1), whereas other GIV NoV-positive samples contained either GIV.1 or GIV.new strains. The strains originated from the same sample in each genetic cluster shared high nucleotide sequence similarity (>98.2% identity); therefore, only representative strains from each sample are shown in Figure 1. The representative GIV.1 and GIV.new strains identified in the present study (2012/Apr/Plant A/Influent [LC150855] and 2011/July/Plant A/Influent [LC150829], respectively) exhibited highest nucleotide identities of 99.6% to Lake Macquarie (JQ613567) and 96.1% to Wastewater/JPN (GIV strain identified in wastewater in Japan, AB565789), respectively, on the nucleotide sequence database (Table 2). The representative GIV.new strain (2011/July/Plant A/Influent [LC150829]) showed nucleotide identities of only 81.9% to Fort Lauderdale (GIV.1) and 77.9% to Dog/ITA (GIV.2) (Table 2), demonstrating that this strain is genetically distinct from both of previously described GIV genotypes.

The GIV NoV strains belonging to the unique genetic cluster (GIV.new strains) were closely related to the strains identified in Japanese wastewater samples collected in 2006 [20], but related virus strain has not been identified from human or animal stool specimens. Our results demonstrate that diverse GIV NoVs belonging to two genetically distinct clusters were circulating in Arizona, United States. Since wastewater contains viruses shed

125 from all populations regardless of symptoms, virus strains and their genetic information
126 obtained from wastewater should reflect, more precisely, the circulation of genetic variants.
127 In the present study, we report the identification of genetically distinct GIV NoV strains in
128 wastewater, which highlights the need for further clinical and environmental studies on GIV
129 NoVs toward a better understanding of their prevalence, molecular epidemiology, and
130 genetic evolution.

131

132

ACKNOWLEDGMENTS

The authors would like to thank Kelly Reynolds at the University of Arizona Mel and Enid Zuckerman College of Public Health for her laboratory contributions and two anonymous wastewater treatment plants in southern Arizona for providing wastewater samples.

This study was partly supported by National Science Foundation (NSF) Water and Environmental Technology (WET) Center and Water Resources Research Center (WRRC), the University of Arizona.

We also wish to acknowledge the support of the Japan Society for the Promotion of Science (JSPS) to Masaaki Kitajima, under JSPS Postdoctoral Fellowships for Research Abroad. Andri T. Rachmadi was a recipient of 2012 Fulbright Master of Science and Technology Scholarship.

Compliance with ethical standards

Conflict of interest

The authors declare that they have no conflict of interest.

REFERENCES

1. Green KY (2007) Caliciviridae: The noroviruses. In: Knipe D, Howley P (eds) *Fields Virol.*, 5th ed. Lippincott Williams and Wilkins, Philadelphia, pp 949–979
2. Kroneman A, Vega E, Vennema H, et al (2013) Proposal for a unified norovirus nomenclature and genotyping. *Arch Virol* 158:2059–68. doi: 10.1007/s00705-013-1708-5
3. Siebenga JJ, Duizer E, Koopmans MPG (2010) Norovirus epidemiology. In: Hansman GS, Jiang X, Green KY (eds) *Caliciviruses Mol. Cell. Virol.* Caister Academic Press, Norfolk, UK, pp 1–24
4. Vinje J, Koopmans MPG (2000) Simultaneous detection and genotyping of “Norwalk-like viruses” by oligonucleotide array in a reverse line blot hybridization format. *J Clin Microbiol* 38:2595–2601.
5. Fankhauser RL, Monroe SS, Noel JS, et al (2000) Epidemiologic and molecular trends of “Norwalk-like viruses” associated with outbreaks of gastroenteritis in the United States. *J Infect Dis* 186:1–7.
6. Iritani N, Seto Y, Kubo H, et al (2002) Prevalence of “Norwalk-like virus” infections in outbreaks of acute nonbacterial gastroenteritis observed during the 1999–2000 season in Osaka city, Japan. *J Med Virol* 138:131–138. doi: 10.1002/jmv.2121
7. Lindell AT, Grillner L, Svensson L, Wirgart BZ (2005) Molecular epidemiology of norovirus infections in Stockholm, Sweden, during the years 2000 to 2003 : Association of the GGIIb genetic cluster with infection in children. *J Clin Microbiol* 43:1086–1092. doi: 10.1128/JCM.43.3.1086
8. Eden J-S, Lim KL, White PA (2012) Complete genome of the human norovirus GIV.1 strain Lake Macquarie virus. *J Virol* 86:10251–2. doi: 10.1128/JVI.01604-12
9. Ao Y-Y, Yu J-M, Li L-L, et al (2014) Detection of human norovirus GIV.1 in China: A case report. *J Clin Virol* 61:8–11. doi: 10.1016/j.jcv.2014.08.002
10. Muscillo M, Fratini M, Graffeo R, et al (2013) GIV noroviruses in wastewaters and in stool specimens from hospitalized patients. *Food Environ Virol* 5:194–202. doi: 10.1007/s12560-013-9121-5
11. Martella V, Campolo M, Lorusso E, et al (2007) Norovirus in captive lion cub (*Panthera leo*). *Emerg Infect Dis* 13:1071–1073.
12. Martella V, Lorusso E, Decaro N, et al (2008) Detection and molecular characterization of a canine norovirus. *Emerg Infect Dis* 14:1306–8. doi: 10.3201/eid1408.080062
13. Ntafis V, Xylouri E, Radogna A, et al (2010) Outbreak of canine norovirus infection in young dogs. *J Clin Microbiol* 48:2605–8. doi: 10.1128/JCM.02528-09

- 186 14. Pinto P, Wang Q, Chen N, et al (2012) Discovery and genomic characterization of
187 noroviruses from a gastroenteritis outbreak in domestic cats in the US. PLoS One
188 7:e32739. doi: 10.1371/journal.pone.0032739
- 189 15. van den Berg H, Lodder W, van der Poel W, et al (2005) Genetic diversity of noroviruses
190 in raw and treated sewage water. Res Microbiol 156:532–40. doi:
191 10.1016/j.resmic.2005.01.008
- 192 16. La Rosa G, Pourshaban M, Iaconelli M, Muscillo M (2008) Detection of genogroup IV
193 noroviruses in environmental and clinical samples and partial sequencing through rapid
194 amplification of cDNA ends. Arch Virol 153:2077–2083. doi:
195 10.1007/s00705-008-0241-4
- 196 17. Kitajima M, Haramoto E, Phanuwat C, et al (2009) Detection of genogroup IV
197 norovirus in wastewater and river water in Japan. Lett Appl Microbiol 49:655–658. doi:
198 10.1111/j.1472-765X.2009.02718.x
- 199 18. La Rosa G, Iaconelli M, Pourshaban M, et al (2010) Molecular detection and genetic
200 diversity of norovirus genogroup IV: a yearlong monitoring of sewage throughout Italy.
201 Arch Virol 155:589–93. doi: 10.1007/s00705-010-0619-y
- 202 19. Kitajima M, Oka T, Haramoto E, et al (2010) Seasonal distribution and genetic diversity
203 of genogroups I, II, and IV noroviruses in the Tamagawa River, Japan. Environ Sci
204 Technol 44:7116–7122. doi: 10.1021/es100346a
- 205 20. Kitajima M, Oka T, Haramoto E, et al (2011) Genetic diversity of genogroup IV
206 noroviruses in wastewater in Japan. Lett Appl Microbiol 52:181–184. doi:
207 10.1111/j.1472-765X.2010.02980.x
- 208 21. Han T-H, Kim S-C, Kim S-T, et al (2014) Detection of norovirus genogroup IV,
209 klassevirus, and pepper mild mottle virus in sewage samples in South Korea. Arch Virol
210 159:457–63. doi: 10.1007/s00705-013-1848-7
- 211 22. Teixeira DM, Hernandez JM, Silva LD, et al (2015) Occurrence of norovirus GIV in
212 environmental water samples from Belém city, Amazon Region, Brazil. Food Environ
213 Virol 101–104. doi: 10.1007/s12560-015-9220-6
- 214 23. Kitajima M, Iker BC, Pepper IL, Gerba CP (2014) Relative abundance and treatment
215 reduction of viruses during wastewater treatment processes — Identification of potential
216 viral indicators. Sci Total Environ 488-489:290–296. doi:
217 10.1016/j.scitotenv.2014.04.087
- 218 24. Kimura M (1980) A simple method for estimating evolutionary rates of base
219 substitutions through comparative studies of nucleotide sequences. J Mol Evol 16:111–
220 120.
- 221 25. Kitajima M, Oka T, Takagi H, et al (2010) Development and application of a broadly

222 reactive real-time reverse transcription-PCR assay for detection of murine noroviruses. J
223 Virol Methods 169:269–273. doi: 10.1016/j.jviromet.2010.07.018
224

TABLE

Table 1. Detection of GIV NoVs in wastewater in Arizona, USA.

Year/Month	Plant A		Plant B	
	Influent	Effluent	Influent	Effluent
2011/July	GIV.new	— ^a	NA ^b	NA
Aug	—	—	—	—
Sept	GIV.new	—	—	—
Oct	—	—	—	—
Nov	—	—	—	—
Dec	—	GIV.new	—	—
2012/Jan	GIV.new	—	—	—
Feb	GIV.new	GIV.new	—	—
Mar	—	GIV.new	—	GIV.new
Apr	GIV.1 + GIV.new	—	GIV.1	—
May	—	—	—	—
June	GIV.new	—	GIV.new	—
July	GIV.new	—	—	—
% positive	54% (7/13)	23% (3/13)	17% (2/12)	8% (1/12)

^a —, negative.

^b NA, not available.

1 **Table 2.** Nucleotide sequence similarity between the representative GIV NoV strains identified in the present study and previously described GIV
2 NoV strains belonging to each genetic cluster, based on BLAST nucleotide search results^a.


Cluster - Representative Strain	GIV.1			GIV.2		GIV.new
	Alpatron	Fort Lauderdale	Lake Macquarie	Dog/ITA	Cat/US	Wastewater/JPN
GIV.1 - 2012/Apr/Plant A/Influent	94.3% (266/282)	97.5% (274/281)	99.6% (281/282)	84.2% (235/279)	81.4% (228/280)	81.7% (227/278)
GIV.new - 2011/July/Plant A/Influent	80.1% (218/272)	81.9% (221/270)	80.9% (220/272)	77.9% (218/280)	76.8% (209/272)	96.1% (271/282)

3 ^a GenBank accession numbers for GIV NoV strains: 2012/Apr/Plant A/Influent, [LC150855]; 2011/July/Plant A/Influent [LC150829], Alpatron,
4 AF195847; Fort Lauderdale, AF414426; Lake Macquarie, JQ613567; Dog/Italy, EU224456; Cat/US, JF781268; Wastewater/JPN, AB565789.

5

FIGURE LEGEND

Figure. Phylogenetic tree for GIV NoV strains using 282 nucleotides of the partial capsid gene sequences. The tree was generated by the neighbor-joining method with representative strains derived from wastewater and reference strains. The scale represents nucleotide substitutions per site. Strains shown in ***italic bold*** are representative GIV NoV strains identified in the present study, representing the year and month of sample collection, WWTP (Plant A or B), sample type (influent or effluent), and GenBank accession number.



Figure