



HOKKAIDO UNIVERSITY

Title	A Long Noncoding RNA, lncRNA-Amhr2, Plays a Role in Amhr2 Gene Activation in Mouse Ovarian Granulosa Cells
Author(s)	Kimura, Atsushi P.; Yoneda, Ryoma; Kurihara, Misuzu et al.
Citation	Endocrinology, 158(11), 4105-4121 https://doi.org/10.1210/en.2017-00619
Issue Date	2017-11-14
Doc URL	https://hdl.handle.net/2115/67922
Type	journal article
File Information	KimuraAP-rev-final HUSCAP.pdf



A Long Noncoding RNA, *lncRNA-Amhr2*, Plays a Role in *Amhr2* Gene Activation in Mouse Ovarian Granulosa Cells

Atsushi P. Kimura,^{1,2} Ryoma Yoneda,² Misuzu Kurihara,² Shota Mayama,² and Shin Matsubara²

¹Department of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

²Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan

Present address for R.Y.: Research Center for Genomic Medicine, Saitama Medical University, Hidaka 350-1241, Japan

Present address for M.K.: Division of Nuclear Dynamics, Okazaki Institute for Integrative Biosciences, Okazaki 444-8787, Japan

Present address for S.Mat.: Bioorganic Research Institute, Suntory Foundation for Life Sciences, 8-1-1 Seikadai, Seikacho, Sorakugun, Kyoto 619-0284, Japan

Short title: *Amhr2* gene activation by a lncRNA

Keywords: Anti-Müllerian hormone type II receptor, transgenic mouse, long noncoding RNA, granulosa cell, reporter gene, Tet-on

Corresponding author and person to whom reprint request should be addressed: Atsushi P. Kimura, PhD, Department of Biological Sciences, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan, akimura@sci.hokudai.ac.jp

Grants: This work was supported by Grants-in-Aid for Young Scientists 19770048 and 21770068 from

Japan Society for the Promotion of Science.

Disclosure summary: The authors have nothing to disclose.

Précis: A 500-bp promoter was insufficient to mouse *Amhr2* gene activation, and a novel long noncoding RNA, *lncRNA-Amhr2*, contributes to the activation by enhancing promoter activity in granulosa cells.

Abbreviations: AMH, anti-Müllerian hormone; *Amhr2*, Anti-Müllerian hormone type II receptor; ANOVA, analysis of variance; BCA, bichinchoninic acid; CNS, conserved noncoding sequence; DMEM, Dulbecco Modified Eagle medium; Dox, doxycycline; EGFP, enhanced green fluorescent protein; ER, estrogen receptor; FBS, fetal bovine serum; *lncRNA*, long noncoding RNA; PCR, polymerase chain reaction; POP, prolyl oligopeptidase; RACE, rapid amplification of cDNA ends; RT-PCR, reverse transcription-polymerase chain reaction; qRT-PCR, quantitative reverse transcription-polymerase chain reaction; S.D., standard deviation; TSS, transcriptional start site; TTS, transcriptional termination site

Abstract

Anti-Müllerian hormone (AMH) is critical to the regression of Müllerian ducts during mammalian male differentiation and targets ovarian granulosa cells and testicular Sertoli and Leydig cells of adults. Specific effects of AMH are exerted via its receptor, AMH type II receptor (*Amhr2*), but the mechanism by which the *Amhr2* gene is specifically activated is not fully understood. To see whether a proximal promoter was sufficient to *Amhr2* gene activation, we generated transgenic mice that bore the enhanced green fluorescent protein (EGFP) gene driven by a 500-bp mouse *Amhr2* gene promoter. None of the established 10 lines, however, showed appropriate EGFP expression, indicating that the 500-bp promoter was insufficient to *Amhr2* gene activation. As a novel regulatory element, we found a long noncoding RNA, *lncRNA-Amhr2*, transcribed from upstream of the *Amhr2* gene in ovarian granulosa cells and testicular Sertoli cells. In primary granulosa cells, knock down of *lncRNA-Amhr2* resulted in a decrease of *Amhr2* mRNA level, and transient reporter gene assay showed that *lncRNA-Amhr2* activation increased *Amhr2* promoter activity. The activity was correlated with *lncRNA-Amhr2* transcription in stably transfected OV3121 cells that were derived from mouse granulosa cells. Moreover, by the Tet-on system, the induction of *lncRNA-Amhr2* transcription dramatically increased *Amhr2* promoter activity in OV3121 cells. These results indicate that *lncRNA-Amhr2* plays a role in *Amhr2* gene activation in ovarian granulosa cells by enhancing promoter activity, providing a novel insight into the *Amhr2* gene regulation underlying the AMH signaling in the female reproductive system.

Introduction

Anti-Müllerian hormone (AMH) is a glycoprotein which induces the regression of Müllerian ducts during male differentiation in embryos (1, 2) and controls the folliculogenesis and Leydig cell proliferation in the adult ovary and testis (3, 4). Its actions are mediated by type I and type II receptors that form a heterodimer, leading to activation of the Smad signaling pathway (5). Since the type I receptor is an activin-like kinase which is expressed in various tissues (5), it is the AMH type II receptor (*Amhr2*) that determines the specificity of AMH actions. In mammals, the *Amhr2* gene is highly expressed in embryonic and postnatal gonads, especially in ovarian granulosa cells and testicular Sertoli cells (6-9), although Leydig cells also express it (10, 11). Such specific activation of the *Amhr2* gene is essential to sex differentiation and adult reproductive functions (12).

In the adult ovary, AMH is secreted from granulosa cells of preantral follicles, and exhibits inhibitory effects on the entry of primordial follicles into the growing phase and on the sensitivity of antral follicles to follicle stimulating hormone (4, 13). When the AMH signaling is disturbed, the ovulation ceases at a younger age than usual (14). Moreover, in human, AMH is tightly associated with some ovarian dysfunction such as polycystic ovarian syndrome and premature ovarian insufficiency (4, 13), and the defect is caused by genetic mutations in the gene encoding AMH or its receptor (4) or by the aberrant regulation of *AMH* and *AMHR2* without genetic mutations (15). These findings, combined with the fact that the *AMH* gene is simply controlled by its promoter (16-18), and that *Amhr2* is a major determinant of AMH actions, clearly underscore the importance of the regulatory mechanism of the *Amhr2* gene expression. Nevertheless, how the *Amhr2* gene is controlled is not fully understood.

The regulatory mechanism of the *AMHR2* gene was first reported in human. The binding of a transcription factor, SF-1, to the *AMHR2* proximal promoter and a synergistic effect of the β -catenin signaling pathway were shown to be important for gene activation (19, 20). A recent study showed that bone morphogenetic protein 15 could also enhance *AMHR2* promoter activity in granulosa cells (21).

In a rat Leydig cell line, a 1.6-kb *Amhr2* promoter exhibited high activity, and two SF-1 binding sites in the proximal region were critical (22). In mouse, a transcription factor, WT-1, binds to the *Amhr2* proximal promoter and plays a crucial role in gene activation during development (23). The chicken *AMHR2* gene was recently cloned, and found to be positively and negatively regulated by the *DMRT1* gene and estrogen, respectively (24). While all of these studies focused on the proximal promoter, it is not known whether the promoter is really sufficient to specific activation of the *Amhr2* gene and whether any other regulatory elements exist.

It has become clear that many genes require regulatory elements other than proximal promoters, such as distal enhancers and long noncoding RNAs (lncRNAs), to be fully and specifically activated (25-27). Especially, lncRNAs have recently received much attention due to their variability and abundance (28). lncRNAs are defined as RNAs that are longer than 200 nucleotides and function as RNA molecules without being translated (28). In many cases, lncRNA transcripts are important for their functions, but at some loci, noncoding transcription *per se* is effective independently of the resulting transcripts (29, 30). An increasing number of studies have shown that lncRNAs are key regulators in many biological fields, such as immunology, development, and neurobiology, by functioning through gene regulation at various levels (i.e. transcription, RNA stability, RNA degradation, and translation) (31-35). In the field of endocrinology, lncRNAs are known to participate in the hormonal control (36-41) and regulate the genes encoding hormones and hormone receptors (30, 42-45). Therefore, more lncRNAs are presumed to be associated with hormonal control or regulation of hormone/hormone receptor gene expression.

In this study, we assessed whether a proximal promoter was sufficient to *Amhr2* gene activation by generating transgenic mice, and found that a 500-bp promoter was insufficient to specific gene activation. We then identified a novel lncRNA, *lncRNA-Amhr2*, which was transcribed in correlation with *Amhr2*. By knock down and a series of experiments using reporter genes, *lncRNA-Amhr2* was found to enhance *Amhr2* promoter activity in ovarian granulosa cells. The results provide a novel insight into the *Amhr2* gene regulation.

Materials and Methods

Animals

C57/B6 and BDF1 (C57/B6 × DBA2 F1) mice were maintained at 25°C with a photoperiod of 14 hours light and 10 hours dark. Eight tissues (brain, heart, liver, kidney, ovary, spleen, testis, and uterus) were collected from male and/or female mice. Food and water were freely accessed. Experimental procedures used in this study were approved by the Institutional Animal Use and Care Committee at Hokkaido University.

Plasmid constructs

0.5k-EGFP: A 500-bp sequence of the mouse *Amhr2* promoter was amplified by polymerase chain reaction (PCR) with mouse genome DNA and KOD polymerase (Toyobo, Osaka, Japan) using a primer pair of 5'-CTACACAGAGAAACCCTGTC-3' and 5'-CCTCAGCCAAGGCTTCCTA-3'. The product was subcloned into a pBluescript II vector (Stratagene, La Jolla, CA, USA) at the *EcoRV* site, and the resulting plasmid was checked by DNA sequencing method. This plasmid was digested with *HindIII* and *EcoRI*, and a 0.5-kb fragment was blunted and inserted into a pEGFP-1 vector (Clontech, Palo Alto, CA, USA) at the blunted *BamHI* site.

sh-lncAmhr2: A double strand DNA was generated by annealing 5'-GATCCGCAGAAAGCATGTCCATATGAGTGTGCTGTCCTCATATGGACATGCTTTCTGCTT TTTTA-3' and 5'-AGCTTAAAAAAGCAGAAAGCATGTCCATATGAGGACAGCACACTCATATGGACATGCTT TCTGCG-3' oligonucleotides. The DNA was inserted into a pBasi-mU6 Neo DNA vector (Takara, Kusatsu, Japan) at *BamHI* and *HindIII* sites. The construct was checked by DNA sequencing method.

0.5k-Luc: A 500-bp *Amhr2* promoter sequence was obtained as above and subcloned into a

pGL3-Basic vector (Promega, Madison, WI, USA) at the blunted *HindIII* site.

6.1k-Luc: 0.5k-Luc was digested with *BglII*, and a 607-bp fragment, which was obtained by digestion of a B6N mouse BAC clone, B6Ng01-240N22, (RIKEN Bioresource Center) with *BglII*, was inserted. The resulting vector was named 1.0k-Luc. A 6.0-kb fragment was obtained by digestion of B6Ng01-240N22 with *EcoRI* and subcloned into a pBluescript II vector at the *EcoRI* site. The plasmid was digested with *EcoRI* and *NheI*, and a 5126-bp fragment was inserted into 1.0k-Luc at the *EcoRI-NheI* site.

6.1k-toER-Luc: A 1.8-kb cDNA sequence of torazame estrogen receptor (ER) was subcloned into a pcDNA3.1 vector (Invitrogen, Carlsbad, CA, USA) (46). The plasmid was digested with *EcoRI*, and the 1.8-kb ER fragment was purified, blunted, and inserted into 6.1k-Luc which was digested with *AflIII* and *Tth111I* and blunted.

6.1k-ΔlncPr-Luc: 6.1k-Luc was digested with *EcoRV* and *Tth111I*, blunted, and self-ligated.

6.1k-POPpr-Luc: A 914-bp mouse prolyl oligopeptidase (POP) gene promoter was subcloned into a pEGFP-1 vector (Clontech) (47, 48). The plasmid was digested with *BamHI* and *EcoRI*, and the POP promoter was purified, blunted, and inserted into 6.1k-Luc which was digested with *EcoRV* and *Tth111I* and blunted.

RevPOPpr-Luc and POPpr-Luc: The blunted 914-bp POP promoter prepared as above was inserted into pGL3-Basic which was digested with *SmaI*.

RevPOPpr-0.5k-Luc and POPpr-0.5k-Luc: The blunted 914-bp POP promoter prepared as above was inserted into 0.5k-Luc which was digested with *SmaI*.

6.1k-POPpr-toER-Luc: The blunted 914-bp POP promoter prepared as above was inserted into pcDNA3.1 containing toER cDNA which was digested with *BamHI* and blunted. The resulting plasmid was digested with *NotI* and *KpnI*, and a 2.7-kb fragment was purified, blunted, and inserted into 6.1k-Luc which was digested with *EcoRV* and *AflIII* and blunted.

6.1k-EGFP: A 1130-bp *Amhr2* promoter was amplified by PCR with mouse genome DNA and KOD FX Neo (Toyobo) using a primer pair of 5'-CTCTCCATCCCAAGTTGGTT-3' and

5'-CCTCAGCCAAGGCTTCCTA-3', and digested with *KpnI*. The resulting 0.8-kb fragment was subcloned into a pEGFP-1 vector (Clontech) at the *KpnI-SmaI* site, and the plasmid was named 0.8k-EGFP. 6.1k-Luc was digested with *KpnI*, and a 5.3-kb fragment was purified and inserted into 0.8k-EGFP at the *KpnI* site.

6.1k-toER-EGFP: 6.1k-toER-Luc was digested with *KpnI*, and a 5.3-kb fragment was purified and inserted into 0.8k-EGFP at the *KpnI* site.

6.1k-POPpr-EGFP: 6.1k-POPpr-Luc was digested with *KpnI*, and a 5.3-kb fragment was purified and inserted into 0.8k-EGFP at the *KpnI* site.

6.1k-TO-Luc: A 459-bp hCMV*1 promoter sequence, which was responsive to doxycycline (Dox), was obtained by digestion of the pPBhCMV*1-cHA-pA plasmid (49) with *SacII* and *AccI*. The hCMV*1 promoter was blunted and inserted into 6.1k-Luc at the blunted *Tth111I* site.

Transgenic mouse generation

Transgenic mice were generated as previously described (48). The 0.5k-EGFP plasmid was digested with *XhoI* and *AflII* and purified to exclude the vector sequence, and injected into fertilized eggs. 12 founders were obtained by dot blot analysis of tail DNAs with an EGFP fragment (772 bp) as a probe. The probe was prepared by digestion of pEGFP-1 vector with *EcoRI* and *NotI*. All founders were mated with wild type mice and 10 lines were established. Copy numbers of the established lines were determined by Southern blot analysis with genome DNAs digested with *DraI* and *PvuII* using an *EcoRI-NotI* EGFP fragment as a probe. A 422-bp probe, which was amplified from the *Amhr2* gene body by genome PCR using primers of 5'-ACATCTTCTCCAGGCTGGCA-3' and 5'-CCAAGTGGATGGGATGTTG-3', was used as a control of two copies. The procedure of Southern blot was previously described (50).

Reverse transcription-PCR (RT-PCR) and quantitative RT-PCR (qRT-PCR)

RNA isolation, DNase treatment, and reverse transcription followed by PCR or real time PCR

were performed as previously described (50-53). Reverse transcription was performed with a specific primer for *lncRNA-Amhr2* in Figure 2A and with the oligo(dT) primer in the other experiments. In real time PCR, detection of *lncRNA-Amhr2* was done with KOD SYBR qPCR Mix (Toyobo), and for the other transcripts, Power SYBR Green Master Mix (Life Technologies, Foster City, CA, USA) was used. Primer sequences for these experiments are listed in Table 1.

Cell sorting

The tunica albuginea was removed from testes of adult #326 mice and placed in Dulbecco Modified Eagle medium (DMEM) containing 0.5 mg/ml collagenase. The seminiferous tubules were cut into small pieces and incubated at 33°C in water bath for 30 min with occasional agitation. The cells were dispersed by gentle pipetting, and filtrated through a cell strainer tube (falcon 352235) to remove seminiferous tubules. The cells were separated by EGFP intensity, and the EGFP-positive cells were sorted by JSAN cell sorter (Bay Bioscience, Kobe, Japan). OV3121 cells transfected with 6.1k-EGFP were also sorted into EGFP-positive and negative population.

Isolation and culture of primary Sertoli and granulosa cells

Primary Sertoli cells were isolated from 8-12 days old male mice and cultured on collagen-coated dishes as previously described (54). Briefly, testes were decapsulated and treated with 0.1% collagenase at 32°C twice for 15-20 min, and the sedimented tubules were washed with phosphate buffered saline containing 1 mM EDTA three times. The tubules were cut into small pieces, and germ cells and Sertoli cells were dispersed by pipetting them. The dispersed cells were washed with DMEM/F12 supplemented with 10% fetal bovine serum (FBS) three times, and spread onto 24-well dishes that were coated with collagen. After cultured for 1 day at 32°C, the cells were treated with 10 mM Tris-HCl (pH 7.4) for 3 min to eliminate germ cells, and the attached Sertoli cells were cultured for several days.

Primary granulosa cells were isolated from immature female mice that were administrated with

hormones and cultured in fibronectin-coated dishes as previously described (51). Briefly, 3-4 weeks old female mice were administrated with pregnant mare serum gonadotropin and human chorionic gonadotropin to induce follicular growth. Five hours after the injection of human chorionic gonadotropin, ovaries were collected, and follicles were punctured by needles in DMEM/F12 with 10% FBS. The cells were washed with the medium three times and spread onto 24-well dishes that were coated with fibronectin. They were cultured at 37°C for several days.

Fractionation into cytoplasmic and nuclear fractions

Fractionation of primary granulosa cells into cytoplasmic and nuclear fractions followed by RT-PCR analysis was performed as previously described (52).

Rapid amplification of cDNA ends (RACE)

5'RACE and 3' RACE were performed with total RNAs from ovarian granulosa cells and the testis as previously described (52, 53). The first PCR product was diluted to 1:1000 and used for the second PCR. Ten subclones for each RACE product were checked by DNA sequencing method. Primers used for RACE analyses are listed in Table 2.

Knock down of *lncRNA-Amhr2*

sh-lncAmhr2 or pBAsi-mU6 Neo DNA plasmids were transfected into primary granulosa cells using GeneJuice transfection reagent (Merck, Darmstadt, Germany) according to the manufacturer's instruction. Two days later, 1 mg/ml G418 was added to the medium and the cells were cultured for another 4 days. Then, total RNAs were purified with ISOGEN II (Nippongene, Tokyo, Japan), and RNA levels were assessed by qRT-PCR.

Reporter gene assay by transient transfection

Constructs containing the luciferase gene were transfected into primary granulosa or Sertoli cells

with a pRL-CMV vector (Promega) using GeneJuice (Merck) or Genefectine (Genetrone Biotech, Seoul, South Korea) transfection reagents. Two days later, the cells were lysed and firefly luciferase activity was measured as previously described (51). Transfection efficiency was normalized by measuring *Renilla* luciferase activity that was derived from pRL-CMV.

Culture and establishment of OV3121 cells stably transfected with pEGFP plasmids

OV3121 cells derived from mouse granulosa cells were obtained from Japanese Collection of Research Bioresources cell bank (National Institutes of Biomedical Innovation, Health and Nutrition), and cultured in DMEM supplemented with 10% FBS, 100 units/ml penicillin, 100 µg/ml streptomycin, and 292 µg/ml L-glutamine. 0.5k-EGFP was linearized by digestion with *Xho*I, and 6.1k-EGFP, 6.1k-toER-EGFP, and 6.1k-POPpr-EGFP were digested with *Mlu*I. The linearized constructs were transfected into OV3121 cells using GeneJuice transfection reagent, and two days later, selection was started by adding 0.5 mg/ml G418 to the medium. Because the pEGFP-1 vector contained the neomycin-resistant gene, transfection of these constructs generated neomycin-resistant cells. The cells were cultured for another 10-14 days, and resulting survived cells were used for experiments.

Establishment of stably transfected OV3121 cells for Tet-on system and Dox treatment

The 6.1k-TO-Luc plasmid was linearized by digestion with *Mlu*I, and transfected into OV3121 cells with pPBCAGrtTA-IN and pPyCAG-PBase plasmids (49) using GeneJuice transfection reagent. pPBCAGrtTA-IN was a PiggyBac-based construct, and by transposase activity from pPyCAG-PBase, the rtTA gene would be efficiently integrated into the genome of OV3121 cells. Two days after transfection, 0.5 mg/ml G418 was added to the medium, and the cells were cultured for another 12 days. For Dox treatment, the cells were spread on 35-mm dishes, and 1 µg/ml Dox was added to the medium on the next day. Five and twenty-four hours after the addition of Dox, the cells were collected with ISOGEN II for qRT-PCR and with Passive Lysis Buffer (Promega) for luciferase assay. qRT-PCR was done as above, and luciferase activity was measured with LARII (Promega) using Lumat LB9507

(Berthold, Bad Wildbad, Germany). Protein concentration of the cell lysate was measured with a Pierce Bicinchoninic Acid (BCA) Protein Assay Reagent kit (Thermo Scientific, Waltham, MA) according to the manufacturer's instruction. Luciferase activity was normalized to the concentration.

Statistical analysis

The data were presented as means $\pm$ standard deviation (S.D.). Statistical comparison was made by Student *t*-test or one-way analysis of variance (ANOVA) followed by Tukey-Kramer test. $P < 0.05$ was considered statistically significant.

Results

A 500-bp promoter is insufficient to *Amhr2* gene activation

We obtained a 500-bp proximal promoter sequence by genomic PCR, based on the transcriptional start site (TSS) determined in rat and aligned to the mouse *Amhr2* locus (22, 23). This sequence contained the binding sites for SF1 and WT1 that were reported to be critical to *Amhr2/AMHR2* gene activation (23). We successfully established 10 transgenic lines bearing the EGFP gene driven by the 500-bp promoter, and the copy number was determined to be 1-25 (Table 3). We first checked whether green fluorescence was observed in the ovary and testis for all the lines, but most lines showed no fluorescence and only the testis from line #326 was green (Table 3, Figure 1A). To check the EGFP mRNA expression in adult tissues, we picked representative three lines, #321, #326, and #348, and performed RT-PCR with eight adult tissues. We detected no EGFP signals in any tissues of line #321 and faint bands in brain, heart, and spleen of line #348, while in line #326, a specific signal was observed in the testis (Figure 1B). These suggested that the transgene was integrated into an active chromatin region in the testis of #326. Thus, we checked embryos of #326, but did not observe any green fluorescence in gonads or urogenital regions at E12.5-15.5 (Table 3).

We next pursued the possibility that EGFP was specifically activated in somatic cells in the testis of #326. To determine which types of cells expressed EGFP, we dispersed the #326 testis, collected somatic and germ cells together, and sorted them by green fluorescence using a cell sorter. We investigated the expression of marker gene for each type of testicular cells: *Hsd3b* for Leydig cells, *Ddx4/Vasa* for germ cells, and *Sgp2* for Sertoli cells (55). qRT-PCR showed that EGFP-positive cells included both somatic cells and germ cells and were enriched in germ cells, which were inconsistent with the *Amhr2* expression enriched in somatic cells (Figure 1C). Thus, the data demonstrated that in #326, the 500-bp promoter did not reproduce the *Amhr2* expression pattern.

Collectively, the proximal promoter was insufficient to *Amhr2* gene activation, which suggested the presence of other regulatory elements.

Identification and characterization of *lncRNA-Amhr2*

In the process of searching for novel regulatory elements, we found noncoding transcription occurred in an upstream region of the *Amhr2* gene, and named the transcript *lncRNA-Amhr2*. RT-PCR using 8 tissues showed that *lncRNA-Amhr2* was specifically detected in the ovary and testis as with the *Amhr2* gene (Figure 2A). To check whether the cell type expressing *lncRNA-Amhr2* was coincided with *Amhr2*, we isolated and cultured primary granulosa and Sertoli cells and performed RT-PCR. The purity of these cell fractions was confirmed by qRT-PCR for marker genes in the testis and ovary (Supplemental Figure 1). The lncRNA signal was detected in both types of cells (Figure 2B), showing co-expression of *lncRNA-Amhr2* and *Amhr2*. We then examined the subcellular localization of *lncRNA-Amhr2* by RT-PCR with nuclear and cytoplasmic fractions from granulosa cells, and the signal was observed in the nucleus but not in the cytoplasm (Figure 2C). These data suggested the role of *lncRNA-Amhr2* in *Amhr2* gene activation, and we decided to determine the full length of this lncRNA.

The direction of the noncoding transcription was found to be opposite to *Amhr2* by RT-PCR with specific primers for reverse transcription (data not shown), so we attempted to determine the full

length in that direction. RACE analyses were performed with total RNAs from primary granulosa cells and the testis, and we identified tissue-specific TSSs by 5'RACE. In granulosa cells, two nucleotides were identified as TSSs by examining 10 subclones (Figure 3A and 3B). In the testis, we obtained two bands after the second PCR, and each contained one TSS (Figure 3A and 3C). By 3'RACE, we identified 10 transcriptional termination sites (TTSs) within a GA-rich region, and a guanine was determined to be the major one because it was TTS in more subclones than any other sites (Figure 3D, bold red font). Collectively, we discovered a novel lncRNA, namely *lncRNA-Amhr2*, which was co-expressed with *Amhr2*, and identified two variants in ovarian granulosa cells and two in the testis (Figure 3A). The length of the variants was 1655-2099 nucleotides, and the longest one in the testis contained one intron, while the other variants were a single exon gene. Although many lncRNAs were not evolutionarily conserved (33), *lncRNA-Amhr2* contained conserved noncoding sequences (CNSs), when we compared the mouse *Amhr2* locus with the human genome using the mVista program (<http://genome.lbl.gov/vista/index.shtml>) (Figure 3A). Nucleotide sequence data for these four *lncRNA-Amhr2* variants were deposited to DDBJ/EMBL/GenBank databases (LC274966, LC274967, LC274968, and LC274969).

Knock down of *lncRNA-Amhr2* decrease *Amhr2* expression in primary granulosa cells

The co-expression of *lncRNA-Amhr2* and *Amhr2* prompted us to explore the possibility of a novel mechanism for *Amhr2* gene activation by *lncRNA-Amhr2*. Thus, we attempted to knock down *lncRNA-Amhr2* in primary granulosa and Sertoli cells. However, the knock down did not work in Sertoli cells, and we could only obtain the result with granulosa cells. By transfection with the sh-*lncAmhr2* construct and selection with G418 in primary granulosa cells, the *lncRNA-Amhr2* level was decreased by approximately 60% compared to control (Figure 4A). The endogenous *Amhr2* expression was significantly decreased to approximately 50% of control, while *Nr5a1/Sf1* and *Actb* mRNA levels were not changed by this knock down (Figure 4B-D). This indicated that *lncRNA-Amhr2* specifically contributed to *Amhr2* gene activation in granulosa cells.

***lncRNA-Amhr2* activation increases *Amhr2* promoter activity in primary granulosa cells**

To see whether *lncRNA-Amhr2* activated the *Amhr2* gene by enhancing its promoter activity, we generated five constructs (Figure 5A). We obtained a 6.1-kb upstream sequence of the *Amhr2* gene and connected to the luciferase gene (6.1k-Luc). Then, we replaced most of the *lncRNA-Amhr2* sequence with torazame ER cDNA which was similar in length (6.1k-toER-Luc) (46), and the *lncRNA-Amhr2* promoter sequence with the POP promoter (6.1k-POPpr-Luc). The replacement with torazame ER would result in transcription of a sequence unrelated to *lncRNA-Amhr2*, and the POP promoter was known to show strong promoter activity in various types of cells and expected to drive *lncRNA-Amhr2* transcription at a higher level than other constructs (47, 48, 51). As controls, we also prepared the constructs that contained the 0.5-kb *Amhr2* promoter (0.5k-Luc) and lacked a *lncRNA-Amhr2* promoter sequence (6.1k-ΔlncPr-Luc).

We transiently transfected primary granulosa cells with these constructs as well as a control vector bearing no promoter (pGL3-Basic), and measured the luciferase activity. 0.5k-Luc showed significantly higher activity than 6.1k-Luc (Figure 5A), which was consistent with previous studies reporting that a shorter sequence often possessed higher promoter activity in transient assay (56-59). The replacement of *lncRNA* with torazame ER and the deletion of *lncRNA* promoter slightly reduced the activity but the difference was not statistically significant (Figure 5A, 6.1k-Luc vs 6.1k-toER-Luc and 6.1k-ΔlncPr-Luc). In contrast, the replacement of *lncRNA* promoter with POP promoter caused a significant increase in *Amhr2* promoter activity (Figure 5A, 6.1k-Luc vs 6.1k-POPpr-Luc). Notably, the POP sequence in the opposite direction exhibited neither promoter activity (Figure 5B) nor effects on the 0.5-kb *Amhr2* promoter (Figure 5C). These excluded the possibility that the reporter gene expression was increased by bidirectional promoter activity or enhancer activity of the POP promoter. In addition, we replaced both *lncRNA* sequence and *lncRNA* promoter with torazame ER and POP promoter (6.1k-POPpr-toER-Luc), and found that the activation of torazame ER transcription did not significantly increase *Amhr2* promoter activity (Figure 5D). These indicated that *lncRNA-Amhr2*

activation increased *Amhr2* promoter activity in granulosa cells.

In Sertoli cells, we observed significantly higher activity in 0.5k-Luc, similarly to granulosa cells (Figure 6). However, no significant difference was detected among the other constructs, even if 6.1k-POPpr-Luc showed higher activity (Figure 6). Therefore, we conducted further experiments only with granulosa cells.

***lncRNA-Amhr2* transcription is correlated with *Amhr2* promoter activity in stably transfected OV3121 cells**

In the primary culture system, we could only perform transient transfection. We next investigated whether the *lncRNA-Amhr2* transcription also contributed to activation of the *Amhr2* promoter when the construct was integrated into genome DNA. For this purpose, we used OV3121 cells that were derived from mouse granulosa cell tumor and expected to maintain some properties as granulosa cells (60). Here we used the EGFP gene as a reporter and prepared four constructs: 0.5k-EGFP (the construct used for generating transgenic mice), 6.1k-EGFP, 6.1k-toER-EGFP, and 6.1k-POPpr-EGFP (Figure 7A and B). We transfected the constructs into OV3121 cells, established stable cells by the selection with G418, and checked expression of *lncRNA-Amhr2* and EGFP mRNA by qRT-PCR. *lncRNA-Amhr2* transcripts were detected at a significantly higher level in the cells with 6.1k-POPpr-EGFP than those with 6.1k-EGFP (Figure 7A). This showed that the POP promoter actually drove the *lncRNA-Amhr2* transcription more strongly than the original promoter. The other two constructs resulted in no *lncRNA-Amhr2* transcripts as expected (Figure 7A). The EGFP mRNA level was correlated with the *lncRNA-Amhr2* transcription. The cells with 6.1k-POPpr-EGFP showed a significantly higher level of EGFP expression than those with the other three constructs (Figure 7B). 6.1k-EGFP showed slightly higher EGFP expression than 0.5k-EGFP and 6.1k-toER-EGFP (Figure 7B), even in cells showing heterogeneous EGFP expression as observed by a fluorescent microscope (Figure 7C).

To further confirm the relationship between *lncRNA* and EGFP, we concentrated EGFP-positive

cells using a cell sorter, and performed RT-PCR. As a result, *lncRNA-Amhr2* transcripts were detected in EGFP-positive cells at a much higher level than in EGFP-negative cells (Figure 7D). This clearly indicated that the *lncRNA-Amhr2* transcription was well correlated with *Amhr2* promoter activity when the construct was integrated into genome DNA, and supported that *lncRNA-Amhr2* contributed to *Amhr2* activation.

Induction of *lncRNA-Amhr2* transcription increases *Amhr2* promoter activity

We finally attempted to clarify whether the *lncRNA-Amhr2* transcription could enhance *Amhr2* promoter activity, using the Tet-on system. We inserted the hCMV*1 promoter that was responsive to Dox into a restriction site at approximately 100-bp downstream to TSSs of *lncRNA-Amhr2* ovary variants in 6.1k-Luc (Figure 8A). This construct was expected to induce the transcription of most sequence of *lncRNA-Amhr2* in response to Dox. We transfected OV3121 cells with this construct and selected the cells that were integrated with the transgene into genome DNA. We added Dox to the selected cells, and 5 hours and 24 hours later, the *lncRNA-Amhr2* transcription was examined. By qRT-PCR, the *lncRNA-Amhr2* level was about 30-folds higher in the cells treated with Dox than in the control cells at both time points (Figure 8B). This indicated that the induction of *lncRNA-Amhr2* transcription was successful, so we checked the luciferase gene expression by qRT-PCR and luciferase activity assay. The luciferase mRNA level was 5.4-folds and 4.3-folds higher at 5 hours and 24 hours after the addition of Dox than in the control cells (Figure 8C), and luciferase activity was 6.9-folds and 12.5-folds higher with Dox than the control (Figure 8D). These data indicated that the induction of *lncRNA-Amhr2* transcription enhanced *Amhr2* promoter activity and increased the luciferase gene expression. Therefore, we concluded that *lncRNA-Amhr2* controlled the *Amhr2* gene through enhancing its promoter activity in ovarian granulosa cells.

Discussion

431

432 We first assessed *in vivo* activity of an *Amhr2* proximal promoter by generating transgenic mice.
433 The EGFP fluorescence was not observed and the RT-PCR signal was not detected in the ovary and
434 testis of transgenic lines except for the testis of #326 (Figure 1 and Table 3). This suggests that the
435 transgene was integrated into genome regions where the chromatin was tightly packed in the nine lines
436 except for #326, and that the 500-bp promoter does not contain any elements to open the chromatin. In
437 addition, the majority of green cells from the #326 testis were found to be germ cells rather than
438 somatic cells, and embryonic gonads were not green in this line. This further suggests that the
439 transgene does not contain any elements to restrict the expression of a linked reporter gene to testicular
440 somatic cells even if it is inserted into a genomic region that was specifically activated in the testis. We
441 could still obtain some green Sertoli and Leydig cells and the promoter might be partly responsible for
442 this activation, but obviously, the 500-bp sequence could not specify the expression to somatic cells in
443 the testis. Collectively, our present data clearly show that the 500-bp promoter does not reproduce the
444 *Amhr2* expression *in vivo*, in other words, the proximal promoter is insufficient to *Amhr2* gene
445 activation. Given that reporter genes were expressed in mostly identical patterns to *Amhr2* when
446 integrated into exon 5 of the native *Amhr2* gene (61, 62), other regulatory elements could be present at
447 the locus, and we found *lncRNA-Amhr2* as an element responsible for *Amhr2* activation at least in
448 granulosa cells.

449 Here we provided the following evidence supporting the regulation of *Amhr2* by *lncRNA-Amhr2*.
450 First, knock down of *lncRNA-Amhr2* led to a decrease in the *Amhr2* mRNA level in primary granulosa
451 cells (Figure 4). This, together with the data showing that the activation of torazame ER transcription
452 did not increase *Amhr2* promoter activity (Figure 5), also revealed that it was the *lncRNA-Amhr2*
453 transcript but not noncoding transcription *per se* that was required for gene activation. Second,
454 transient reporter gene assay showed that *lncRNA-Amhr2* activation by POP promoter significantly
455 increased *Amhr2* promoter activity in primary granulosa cells (Figure 5). Third, higher *lncRNA-Amhr2*
456 transcription increased *Amhr2* promoter activity even when the transgene was integrated into the

genome DNA of granulosa cell-derived OV3121 cells (Figure 7). Fourth, in those stable OV3121 cells, the EGFP reporter gene was activated only in the cell population expressing *lncRNA-Amhr2* at high levels. Last, the induction of *lncRNA-Amhr2* transcription by the Tet-on system dramatically increased activity of the linked *Amhr2* promoter in stably transfected OV3121 cells (Figure 8). All of these results indicate that *lncRNA-Amhr2* plays an important role in *Amhr2* gene activation in ovarian granulosa cells, and therefore, this lncRNA is a novel regulatory element of the mouse *Amhr2* gene.

In contrast to granulosa cells, the knock down experiment did not work in Sertoli cells, similarly to some nuclear lncRNAs that were not efficiently knocked down with short hairpin RNAs (63, 64). Additionally, the usage of POP promoter for *lncRNA-Amhr2* transcription did not result in a significant increase of *Amhr2* promoter activity in primary Sertoli cells. However, these do not necessarily mean that *lncRNA-Amhr2* has nothing to do with *Amhr2* regulation in this cell lineage. We observed still higher *Amhr2* promoter activity in the cells with 6.1k-POPpr-Luc than 6.1k-Luc, 6.1k-toER-Luc, and 6.1k-ΔlncPr-Luc (Figure 6), suggesting a positive role of *lncRNA-Amhr2* in Sertoli cells. Alternatively, it may be possible that an intron-containing variant (Testis variant 2 in Figure 3A) activates the *Amhr2* gene by a different mechanism in Sertoli cells. Further analyses will be required for revealing the function of *lncRNA-Amhr2* in Sertoli cells.

lncRNAs control their target genes on the same chromosome in *cis* or on another chromosome in *trans*. Our present data strongly suggest that *lncRNA-Amhr2* acts on the *Amhr2* gene only in *cis*, which is consistent with many lncRNAs controlling linked protein-coding genes (26). In transgenic mice, the exogenous *Amhr2* promoter was not activated in the ovary of all 10 lines, despite the presence of endogenous *lncRNA-Amhr2* transcripts in granulosa cells. If *lncRNA-Amhr2* acted in *trans*, we could have observed the EGFP activation in their ovaries. In addition, our preliminary data showed that *Amhr2* expression was not influenced by overexpression of *lncRNA-Amhr2* in some types of cultured cells (our unpublished data). Thus, *lncRNA-Amhr2* is likely to act only in *cis*. *Cis*-acting lncRNAs generally need to interact with the chromatin region of their target gene loci (33, 65), and the endogenous *lncRNA-Amhr2* might not be able to reach genome regions integrated with the transgenes.

In general, lncRNAs are categorized by their localization and functional mechanisms, and *lncRNA-Amhr2* belongs to those localized in the nucleus and involved in gene activation in *cis*. There are some categories for such lncRNAs: enhancer RNAs, activating RNAs, and promoter associated noncoding RNAs. Enhancer RNAs are transcribed bidirectionally from enhancers and mediate the chromatin looping to allow the enhancers to physically contact with their target promoters (66, 67). Activating RNAs bind to mediators and promote the target gene transcription (26, 68), and promoter associated noncoding RNAs are located just upstream to their target gene promoters to control epigenetic modifications (69). *lncRNA-Amhr2* is similar to activating RNAs, in that it is transcribed from a neighboring region of its target, activates the target gene in *cis*, and the transcript is required. Thus, *lncRNA-Amhr2* may activate the *Amhr2* gene by a similar mechanism to activating RNAs.

Our present study revealed that a novel lncRNA, *lncRNA-Amhr2*, contributes to *Amhr2* gene activation in mouse ovarian granulosa cells. AMH is an important hormone for the regulation of folliculogenesis in the ovary, and the *Amhr2* protein is a pivotal molecule to mediate the signaling. Given that the down- and up-regulation of this receptor gene results in abnormal folliculogenesis (14, 70, 71), *lncRNA-Amhr2* is highly likely to be a key molecule controlling the AMH signaling probably via interacting to the *Amhr2* promoter in *cis* and the consequent up-regulation of the *Amhr2* gene. Notably, one of the transcription factors for *Amhr2* activation, WT-1, binds to the proximal promoter of the *Amhr2* gene, and has RNA-binding activity in embryonic stem cells (72, 73). In combination, current results support the view that *lncRNA-Amhr2* interacts with transcription factors binding to the *Amhr2* promoter to activate this gene. Further investigation of the molecular mechanism of the *Amhr2* activation and the physiological functions in the regulation of endocrine system by *lncRNA-Amhr2* is underway.

In conclusion, we demonstrated that a proximal promoter was insufficient to *Amhr2* gene activation *in vivo* and identified a novel lncRNA, *lncRNA-Amhr2*, which plays a role in *Amhr2* gene activation in ovarian granulosa cells. The study reveals a novel mechanism of *Amhr2* gene activation, and provides a new insight into the regulation of AMH signaling.

Acknowledgements

We thank Dr. Yoshinao Katsu for kindly giving us a plasmid containing the torazame ER cDNA sequence. We also thank Dr. Kazuhiro Murakami for kindly giving us plasmids, pPBhCMV*1-CHA-pA, pPBCAGrtTA-IN, and pPyCAG-PBase.

References

1. Josso N. Anti-müllerian hormone. *Baillieres Clin Endocrinol Metab* 1991; 5:635-654.
2. Kimura AP. Anti-Müllerian Hormone. In: Takei Y, Ando H, Tsutsui K, eds. *Handbook of Hormones*. 1st ed. Academic Press; 2016:301-303.
3. Behringer RR, Finegold MJ, Cate RL. Müllerian-inhibiting substance function during mammalian sexual development. *Cell* 1994; 79:415-425.
4. Dewailly D, Andersen CY, Balen A, Broekmans F, Dilaver N, Fanchin R, Griesinger G, Kelsey TW, La Marca A, Lambalk C, Mason H, Nelson SM, Visser JA, Wallace WH, Anderson RA. The physiology and clinical utility of anti-Mullerian hormone in women. *Hum Reprod Update* 2014; 20:370-385
5. Josso N, Clemente N. Transduction pathway of anti-Müllerian hormone, a sex-specific member of the TGF-beta family. *Trends Endocrinol Metab* 2003; 14:91-97.
6. Baarends WM, Hoogerbrugge JW, Post M, Visser JA, De Rooij DG, Parvinen M, Themmen AP, Grootegoed JA. Anti-müllerian hormone and anti-müllerian hormone type II receptor messenger ribonucleic acid expression during postnatal testis development and in the adult testis of the rat. *Endocrinology* 1995; 136:5614-5622.

- 535 7. Baarends WM, Uilenbroek JT, Kramer P, Hoogerbrugge JW, van Leeuwen EC, Themmen AP,
536 Grootegoed JA. Anti-müllerian hormone and anti-müllerian hormone type II receptor messenger
537 ribonucleic acid expression in rat ovaries during postnatal development, the estrous cycle, and
538 gonadotropin-induced follicle growth. *Endocrinology* 1995; 136:4951-4962.
- 539 8. Baarends WM, van Helmond MJ, Post M, van der Schoot PJ, Hoogerbrugge JW, de Winter JP,
540 Uilenbroek JT, Karels B, Wilming LG, Meijers JH. A novel member of the transmembrane
541 serine/threonine kinase receptor family is specifically expressed in the gonads and in
542 mesenchymal cells adjacent to the müllerian duct. *Development* 1994; 120:189-197.
- 543 9. di Clemente N, Wilson C, Faure E, Boussin L, Carmillo P, Tizard R, Picard JY, Vigier B, Josso N,
544 Cate R. Cloning, expression, and alternative splicing of the receptor for anti-Müllerian hormone.
545 *Mol Endocrinol* 1994; 8:1006-1020.
- 546 10. Lee MM, Seah CC, Masiakos PT, Sottas CM, Preffer FI, Donahoe PK, Maclaughlin DT, Hardy
547 MP. Müllerian-inhibiting substance type II receptor expression and function in purified rat Leydig
548 cells. *Endocrinology* 1999; 140:2819-2827.
- 549 11. Racine C, Rey R, Forest MG, Louis F, Ferré A, Huhtaniemi I, Josso N, di Clemente N. Receptors
550 for anti-müllerian hormone on Leydig cells are responsible for its effects on steroidogenesis and
551 cell differentiation. *Proc Natl Acad Sci U S A* 1998; 95:594-599.
- 552 12. Mishina Y, Rey R, Finegold MJ, Matzuk MM, Josso N, Cate RL, Behringer RR. Genetic analysis
553 of the Müllerian-inhibiting substance signal transduction pathway in mammalian sexual
554 differentiation. *Genes Dev* 1996; 10:2577-2587.
- 555 13. Broekmans FJ, Visser JA, Laven JS, Broer SL, Themmen AP, Fauser BC. Anti-Müllerian
556 hormone and ovarian dysfunction. *Trends Endocrinol Metab* 2008; 19:340-347
- 557 14. Durlinger AL, Kramer P, Karels B, de Jong FH, Uilenbroek JT, Grootegoed JA, Themmen AP.
558 Control of primordial follicle recruitment by anti-Müllerian hormone in the mouse ovary.
559 *Endocrinology* 1999; 140:5789-5796.
- 560 15. Dewailly D, Robin G, Peigne M, Decanter C, Pigny P, Catteau-Jonard S. Interactions between

- 561 androgens, FSH, anti-Müllerian hormone and estradiol during folliculogenesis in the human
562 normal and polycystic ovary. Hum Reprod Update 2016; 22:709-724
- 563 16. Peschon JJ, Behringer RR, Cate RL, Harwood KA, Idzerda RL, Brinster RL, Palmiter RD.
564 Directed expression of an oncogene to Sertoli cells in transgenic mice using mullerian inhibiting
565 substance regulatory sequences. Mol Endocrinol 1992; 6:1403-1411
- 566 17. Beau C, Vivian N, Münsterberg A, Dresser DW, Lovell-Badge R, Guerrier D. In vivo analysis of
567 the regulation of the anti-Müllerian hormone, as a marker of Sertoli cell differentiation during
568 testicular development, reveals a multi-step process. Mol Reprod Dev 2001; 59:256-264
- 569 18. Lécureuil C, Fontaine I, Crepieux P, Guillou F. Sertoli and granulosa cell-specific Cre
570 recombinase activity in transgenic mice. Genesis 2002; 33:114-118
- 571 19. de Santa Barbara P, Moniot B, Poulat F, Boizet B, Berta P. Steroidogenic factor-1 regulates
572 transcription of the human anti-müllerian hormone receptor. J Biol Chem 1998;
573 273:29654-29660.
- 574 20. Hossain A, Saunders GF. Synergistic cooperation between the beta-catenin signaling pathway and
575 steroidogenic factor 1 in the activation of the Mullerian inhibiting substance type II receptor. J
576 Biol Chem 2003; 278:26511-26516.
- 577 21. Pierre A, Estienne A, Racine C, Picard JY, Fanchin R, Lahoz B, Alabart JL, Folch J, Jarrier P,
578 Fabre S, Monniaux D, di Clemente N. The Bone Morphogenetic Protein 15 Up-Regulates the
579 Anti-Müllerian Hormone Receptor Expression in Granulosa Cells. J Clin Endocrinol Metab 2016;
580 101:2602-2611.
- 581 22. Teixeira J, Kehas DJ, Antun R, Donahoe PK. Transcriptional regulation of the rat Müllerian
582 inhibiting substance type II receptor in rodent Leydig cells. Proc Natl Acad Sci U S A 1999;
583 96:13831-13838.
- 584 23. Klattig J, Sierig R, Kruspe D, Besenbeck B, Englert C. Wilms' tumor protein Wt1 is an activator
585 of the anti-Müllerian hormone receptor gene Amhr2. Mol Cell Biol 2007; 27:4355-4364.
- 586 24. Cutting AD, Ayers K, Davidson N, Oshlack A, Doran T, Sinclair AH, Tizard M, Smith CA.

587 Identification, expression, and regulation of anti-Müllerian hormone type-II receptor in the
588 embryonic chicken gonad. *Biol Reprod* 2014; 90:106.

589 25. Orom UA, Shiekhattar R. Noncoding RNAs and enhancers: complications of a long-distance
590 relationship. *Trends Genet* 2011; 27:433-439.

591 26. Ørom UA, Shiekhattar R. Long noncoding RNAs usher in a new era in the biology of enhancers.
592 *Cell* 2013; 154:1190-1193.

593 27. Vernimmen D, Bickmore WA. The Hierarchy of Transcriptional Activation: From Enhancer to
594 Promoter. *Trends Genet* 2015; 31:696-708.

595 28. Rinn JL, Chang HY. Genome regulation by long noncoding RNAs. *Annu Rev Biochem* 2012;
596 81:145-166.

597 29. Latos PA, Pauler FM, Koerner MV, Şenergin HB, Hudson QJ, Stocsits RR, Allhoff W, Stricker
598 SH, Klement RM, Warczok KE, Aumayr K, Pasierbek P, Barlow DP. Airn transcriptional overlap,
599 but not its lncRNA products, induces imprinted *Igf2r* silencing. *Science* 2012; 338:1469-1472

600 30. Yoo EJ, Cooke NE, Liebhaber SA. An RNA-independent linkage of noncoding transcription to
601 long-range enhancer function. *Mol Cell Biol* 2012; 32:2020-2029.

602 31. Hirose T, Mishima Y, Tomari Y. Elements and machinery of non-coding RNAs: toward their
603 taxonomy. *EMBO Rep* 2014; 15:489-507.

604 32. Taylor DH, Chu ET, Spektor R, Soloway PD. Long non-coding RNA regulation of reproduction
605 and development. *Mol Reprod Dev* 2015; 82:932-956.

606 33. Chen LL. Linking Long Noncoding RNA Localization and Function. *Trends Biochem Sci* 2016;
607 41:761-772.

608 34. Li W, Notani D, Rosenfeld MG. Enhancers as non-coding RNA transcription units: recent
609 insights and future perspectives. *Nat Rev Genet* 2016; 17:207-223.

610 35. Ulitsky I. Evolution to the rescue: using comparative genomics to understand long non-coding
611 RNAs. *Nat Rev Genet* 2016; 17:601-614.

612 36. Lanz RB, McKenna NJ, Onate SA, Albrecht U, Wong J, Tsai SY, Tsai MJ, O'Malley BW. A

steroid receptor coactivator, SRA, functions as an RNA and is present in an SRC-1 complex. *Cell* 1999; 97:17-27.

37. Lanz RB, Razani B, Goldberg AD, O'Malley BW. Distinct RNA motifs are important for coactivation of steroid hormone receptors by steroid receptor RNA activator (SRA). *Proc Natl Acad Sci U S A* 2002; 99:16081-16086.
38. Takayama K, Horie-Inoue K, Katayama S, Suzuki T, Tsutsumi S, Ikeda K, Urano T, Fujimura T, Takagi K, Takahashi S, Homma Y, Ouchi Y, Aburatani H, Hayashizaki Y, Inoue S. Androgen-responsive long noncoding RNA CTBP1-AS promotes prostate cancer. *EMBO J* 2013; 32:1665-1680.
39. Yang L, Lin C, Jin C, Yang JC, Tanasa B, Li W, Merkurjev D, Ohgi KA, Meng D, Zhang J, Evans CP, Rosenfeld MG. lncRNA-dependent mechanisms of androgen-receptor-regulated gene activation programs. *Nature* 2013; 500:598-602.
40. Vicent GP, Nacht AS, Zaurin R, Font-Mateu J, Soronellas D, Le Dily F, Reyes D, Beato M. Unliganded progesterone receptor-mediated targeting of an RNA-containing repressive complex silences a subset of hormone-inducible genes. *Genes Dev* 2013; 27:1179-1197.
41. Qiu J, Ye L, Ding J, Feng W, Zhang Y, Lv T, Wang J, Hua K. Effects of oestrogen on long noncoding RNA expression in oestrogen receptor alpha-positive ovarian cancer cells. *J Steroid Biochem Mol Biol* 2014; 141:60-70.
42. Ho Y, Elefant F, Liebhaber SA, Cooke NE. Locus control region transcription plays an active role in long-range gene activation. *Mol Cell* 2006; 23:365-375.
43. Huang PP, Brusman LE, Iyer AK, Webster NJ, Mellon PL. A Novel Gonadotropin-Releasing Hormone 1 (Gnrh1) Enhancer-Derived Noncoding RNA Regulates Gnrh1 Gene Expression in GnRH Neuronal Cell Models. *PLoS One* 2016; 11:e0158597.
44. Pnueli L, Rudnizky S, Yosefzon Y, Melamed P. RNA transcribed from a distal enhancer is required for activating the chromatin at the promoter of the gonadotropin α -subunit gene. *Proc Natl Acad Sci U S A* 2015; 112:4369-4374.

- 639 45. Zhou Q, Chen J, Feng J, Wang J. Long noncoding RNA PVT1 modulates thyroid cancer cell
640 proliferation by recruiting EZH2 and regulating thyroid-stimulating hormone receptor (TSHR).
641 Tumour Biol 2016; 37:3105-3113.
- 642 46. Katsu Y, Kohno S, Narita H, Urushitani H, Yamane K, Hara A, Clauss TM, Walsh MT,
643 Miyagawa S, Guillette LJ, Iguchi T. Cloning and functional characterization of Chondrichthyes,
644 cloudy catshark, Scyliorhinus torazame and whale shark, Rhincodon typus estrogen receptors.
645 Gen Comp Endocrinol 2010; 168:496-504.
- 646 47. Kimura A, Yoshida I, Takagi N, Takahashi T. Structure and localization of the mouse prolyl
647 oligopeptidase gene. J Biol Chem 1999; 274:24047-24053.
- 648 48. Matsubara S, Maruyama Y, Kimura AP. A 914-bp promoter is sufficient to reproduce the
649 endogenous prolyl oligopeptidase gene localization in the mouse placenta if not subject to
650 position effect. Gene 2013; 524:114-123.
- 651 49. Murakami K, Günesdogan U, Zyllicz JJ, Tang WW, Sengupta R, Kobayashi T, Kim S, Butler R,
652 Dietmann S, Surani MA. NANOG alone induces germ cells in primed epiblast in vitro by
653 activation of enhancers. Nature 2016; 529:403-407.
- 654 50. Yoneda R, Satoh Y, Yoshida I, Kawamura S, Kotani T, Kimura AP. A genomic region
655 transcribed into a long noncoding RNA interacts with the Prss42/Tessp-2 promoter in
656 spermatocytes during mouse spermatogenesis, and its flanking sequences can function as
657 enhancers. Mol Reprod Dev 2016; 83:541-557.
- 658 51. Matsubara S, Takahashi T, Kimura AP. Epigenetic patterns at the mouse prolyl oligopeptidase
659 gene locus suggest the CpG island in the gene body to be a novel regulator for gene expression.
660 Gene 2010; 465:17-29.
- 661 52. Matsubara S, Kurihara M, Kimura AP. A long non-coding RNA transcribed from conserved
662 non-coding sequences contributes to the mouse prolyl oligopeptidase gene activation. J Biochem
663 2014; 155:243-256.
- 664 53. Kurihara M, Shiraishi A, Satake H, Kimura AP. A conserved noncoding sequence can function as

665 a spermatocyte-specific enhancer and a bidirectional promoter for a ubiquitously expressed gene
666 and a testis-specific long noncoding RNA. *J Mol Biol* 2014; 426:3069-3093.

667 54. Yoneda R, Kimura AP. A testis-specific serine protease, Prss41/Tessp-1, is necessary for the
668 progression of meiosis during murine in vitro spermatogenesis. *Biochem Biophys Res Commun*
669 2013; 441:120-125.

670 55. Yoneda R, Takahashi T, Matsui H, Takano N, Hasebe Y, Ogiwara K, Kimura AP. Three
671 testis-specific paralogous serine proteases play different roles in murine spermatogenesis and are
672 involved in germ cell survival during meiosis. *Biol Reprod* 2013; 88:118.

673 56. Thotakura SR, Karthikeyan N, Smith T, Liu K, George A. Cloning and characterization of rat
674 dentin matrix protein 1 (DMP1) gene and its 5'-upstream region. *J Biol Chem* 2000;
675 275:10272-10277

676 57. Solberg N, Machon O, Krauss S. Characterization and functional analysis of the 5'-flanking
677 promoter region of the mouse Tcf3 gene. *Mol Cell Biochem* 2012; 360:289-299

678 58. Wang XL, Wang K, Zhao S, Wu Y, Gao H, Zeng SM. Oocyte-secreted growth differentiation
679 factor 9 inhibits BCL-2-interacting mediator of cell death-extra long expression in porcine
680 cumulus cell. *Biol Reprod* 2013; 89:56

681 59. Ding X, Hu J, Zhang H, Xu Y. Genetic Variants in the STMN1 Transcriptional Regulatory
682 Region Affect Promoter Activity and Fear Behavior in English Springer Spaniels. *PLoS One*
683 2016; 11:e0158756

684 60. Yanagihara K, Nii M, Tsumuraya M, Numoto M, Seito T, Seyama T. A radiation-induced murine
685 ovarian granulosa cell tumor line: introduction of v-ras gene potentiates a high metastatic ability.
686 *Jpn J Cancer Res* 1995; 86:347-356.

687 61. Jamin SP, Arango NA, Mishina Y, Hanks MC, Behringer RR. Requirement of Bmpr1a for
688 Müllerian duct regression during male sexual development. *Nat Genet* 2002; 32:408-410

689 62. Arango NA, Kobayashi A, Wang Y, Jamin SP, Lee HH, Orvis GD, Behringer RR. A
690 mesenchymal perspective of Müllerian duct differentiation and regression in Amhr2-lacZ mice.

691 Mol Reprod Dev 2008; 75:1154-1162

692 63. Gutschner T, Baas M, Diederichs S. Noncoding RNA gene silencing through genomic integration
693 of RNA destabilizing elements using zinc finger nucleases. *Genome Res* 2011; 21:1944-1954

694 64. Huarte M. The emerging role of lncRNAs in cancer. *Nat Med* 2015; 21:1253-1261

695 65. Roberts TC, Morris KV, Weinberg MS. Perspectives on the mechanism of transcriptional
696 regulation by long non-coding RNAs. *Epigenetics* 2014; 9:13-20

697 66. Kim TK, Hemberg M, Gray JM, Costa AM, Bear DM, Wu J, Harmin DA, Laptewicz M,
698 Barbara-Haley K, Kuersten S, Markenscoff-Papadimitriou E, Kuhl D, Bito H, Worley PF,
699 Kreiman G, Greenberg ME. Widespread transcription at neuronal activity-regulated enhancers.
700 *Nature* 2010; 465:182-187

701 67. Li W, Notani D, Ma Q, Tanasa B, Nunez E, Chen AY, Merkurjev D, Zhang J, Ohgi K, Song X,
702 Oh S, Kim HS, Glass CK, Rosenfeld MG. Functional roles of enhancer RNAs for
703 oestrogen-dependent transcriptional activation. *Nature* 2013; 498:516-520

704 68. Lai F, Orom UA, Cesaroni M, Beringer M, Taatjes DJ, Blobel GA, Shiekhattar R. Activating
705 RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature* 2013;
706 494:497-501

707 69. Tomikawa J, Shimokawa H, Uesaka M, Yamamoto N, Mori Y, Tsukamura H, Maeda K,
708 Imamura T. Single-stranded noncoding RNAs mediate local epigenetic alterations at gene
709 promoters in rat cell lines. *J Biol Chem* 2011; 286:34788-34799

710 70. Durlinger AL, Gruijters MJ, Kramer P, Karels B, Kumar TR, Matzuk MM, Rose UM, de Jong FH,
711 Uilenbroek JT, Grootegoed JA, Themmen AP. Anti-Müllerian hormone attenuates the effects of
712 FSH on follicle development in the mouse ovary. *Endocrinology* 2001; 142:4891-4899

713 71. Durlinger AL, Gruijters MJ, Kramer P, Karels B, Ingraham HA, Nachtigal MW, Uilenbroek JT,
714 Grootegoed JA, Themmen AP. Anti-Müllerian hormone inhibits initiation of primordial follicle
715 growth in the mouse ovary. *Endocrinology* 2002; 143:1076-1084

716 72. Caricasole A, Duarte A, Larsson SH, Hastie ND, Little M, Holmes G, Todorov I, Ward A. RNA

binding by the Wilms tumor suppressor zinc finger proteins. Proc Natl Acad Sci U S A 1996;
93:7562-7566

73. Bharathavikru R, Dudnakova T, Aitken S, Slight J, Artibani M, Hohenstein P, Tollervey D, Hastie N. Transcription factor Wilms' tumor 1 regulates developmental RNAs through 3' UTR interaction. Genes Dev 2017; 31:347-352

Figure legends

Figure 1. A 500-bp *Amhr2* promoter is insufficient to specific gene activation in granulosa and Sertoli cells. Transgenic mice were generated with the 0.5k-EGFP construct in which the EGFP gene was driven by a 500-bp *Amhr2* promoter. (A) Fluorescent microscopic images of testes from a wild type mouse (WT) and a #326 mouse. (B) RT-PCR analysis of the EGFP reporter gene in three transgenic mouse lines. Eight tissues were collected from the indicated transgenic mouse lines, #326, #321, and #348, and total RNAs were purified from them. cDNAs were synthesized with (RT+) or without reverse transcriptase (RT-) using the oligo(dT) primer, and PCR reactions were performed with primer pairs for the indicated genes. The cycle numbers were 30 for *Amhr2* and EGFP and 24 for *Gapdh*. Testis-specific amplification was observed in the #326 line. (C) qRT-PCR analysis of the EGFP gene in cell fractions before (Total) and after sorting (EGFP+). Testicular cells were isolated from testes of #326 mice, and green cells were collected by a cell sorter. Total RNAs were purified from the cells before and after the sorting, and cDNAs were synthesized using the olig(dT) primer. Quantitative PCR was performed to detect marker genes for Leydig cells (*Hsd3b*), germ cells (*Ddx4/Vasa*), and Sertoli cells (*Sgp2*) using the 7300 real-time PCR system (Applied Biosystems). The PCR was also performed for a housekeeping gene, *Gapdh*, and the data were normalized to the *Gapdh* level. The value in the cells before sorting (Total) was set to 1.0, and relative levels of the marker genes after sorting are shown. The EGFP-positive fraction was enriched in germ cells. The data are presented as mean $\pm$ S.D.

from four independent experiments with two sets of sorted cells, and the statistical significance was analyzed by Student *t*-test. **P* < 0.05, ***P* < 0.01.

Figure 2. Expression of *lncRNA-Amhr2*. (A) Tissue-specificity of *lncRNA-Amhr2* was investigated by RT-PCR analysis. Eight tissues were isolated from wild type mice, and RT-PCR was performed by using a specific primer for reverse transcription of *lncRNA-Amhr2* and oligo(dT) for *Gapdh*. We used a primer pair that could amplify all four variants of *lncRNA-Amhr2*. The signal was specifically observed in the ovary and testis. *Gapdh* was amplified as an internal control. The cycle numbers were 40 for *lncRNA-Amhr2* and 27 for *Gapdh*. (B) RT-PCR with primary cells. Sertoli cells were isolated from testes of 8-12 days old male mice, and cultured on collagen-coated dishes for several days. Primary granulosa cells were isolated from ovaries of immature female mice after administration with hormones, and cultured on fibronectin-coated dishes for several days. Total RNAs were purified from these cells, and RT-PCR analysis was performed as in Figure 1B. *Amhr2* was detected as a positive control. The cycle number was 35. (C) Subcellular distribution of *lncRNA-Amhr2* was investigated by RT-PCR. Primary granulosa cells were cultured and fractionated into cytoplasmic and nuclear fractions. Total RNA was purified from each fraction (Cytoplasmic and Nuclear) as well as granulosa cells without fractionation (Total), and RT-PCR was performed as above. A primer pair to amplify a part of exon 6 of the *Gapdh* gene was used as a control. The *lncRNA-Amhr2* signal was exclusively detected in the nuclear fraction. The cycle numbers were 35 for *lncRNA-Amhr2* and 30 for *Gapdh*.

Figure 3. Cloning of *lncRNA-Amhr2* and comparison with human genome. (A) Schematic drawing of the mouse *Amhr2* gene locus. At the top, the *Amhr2* gene is depicted by boxes representing exons, and translated and untranslated regions are indicated as black and white boxes. Below the gene structure, a 7-kb upstream region was enlarged. The 7-kb upstream sequence of the mouse *Amhr2* gene was compared with the corresponding 7-kb sequence at the human *AMHR2* locus, using mVista program (<http://genome.lbl.gov/vista/index.shtml>). This program detected conserved sequences presented by

pink or white painting. Y-axis indicates the percentage of identical nucleotides between the two species. At the bottom, positions of four *lncRNA-Amhr2* variants are shown with their length. Two variants were isolated from the testis and the other two was from ovarian granulosa cells. All variants have poly(A) tails. Testis variant 2 contains two exons. Some of the CNSs, marked with pink, overlapped with the region transcribed into *lncRNA-Amhr2*. (B) TSSs of *lncRNA-Amhr2* ovary variants. A specific band was obtained by 5'RACE, and ten subclones were checked. Two TSSs were identified as indicated by bold red fonts and bent arrows with the numbers of subclones. (C) TSSs of *lncRNA-Amhr2* testis variants. By 5'RACE, two specific bands were obtained and ten subclones were sequenced for each band. All of the ten subclones contained the same nucleotide as a TSS in both cases, and the TSSs are indicated by bold red fonts and bent arrows. (D) TTSs of *lncRNA-Amhr2*. By 3'RACE, a specific band was obtained in both the ovary and testis, and ten subclones were sequenced for each tissue. The TTS position of each subclone from ovary (lower, green fonts) and testis (upper, blue fonts) was indicated by bent arrows with numbers of subclones. A major TTS is shown by a bold red font.

Figure 4. Knock down of *lncRNA-Amhr2* in primary granulosa cells. Primary granulosa cells were cultured as in Figure 2B and transfected with a vector containing shRNA for *lncRNA-Amhr2* (Knock down) or a control vector (Control). The cells were selected by adding G418 to the culture medium for several days. Total RNAs were purified from the cells, and qRT-PCR was performed for expression of *lncRNA-Amhr2* (A), *Amhr2* (B), *Nr5a1/Sf1* (C), and *Actb* (D). The expression level was normalized to *Gapdh*, and the value in the control sample was set to 1.0. *lncRNA-Amhr2* was successfully knocked down, and *Amhr2* expression was significantly decreased. The data are presented as mean $\pm$ S.D. from four independent experiments, and the statistical significance was analyzed by Student *t*-test. **P* < 0.05, ***P* < 0.01.

Figure 5. *lncRNA-Amhr2* activation increase *Amhr2* promoter activity in primary granulosa cells by

transient reporter gene assay. (A) Reporter gene assay in primary granulosa cells. Primary granulosa cells were cultured and transfected with constructs indicated at left. Two days after the transfection, cell lysates were collected and luciferase activity was measured. Relative luciferase activity is shown in the graph at right. The data are presented as mean $\pm$ S.D. from three or five independent experiments. Luciferase activity was significantly higher in 6.1k-POPpr-Luc than other constructs containing the 6.1-kb sequence. (B) Unidirectional activity of the POP promoter. The POP promoter was directly connected to the luciferase gene in both directions, and the constructs were transfected into primary granulosa cells. Luciferase activity was measured and shown as above. The data are presented as mean $\pm$ S.D. from four independent experiments. The POP promoter in the forward direction exhibited much higher activity. (C) The effect of the POP promoter on *Amhr2* promoter activity. The POP promoter was connected to upstream of the 0.5-kb *Amhr2* promoter in 0.5k-Luc, and luciferase activity was assessed in primary granulosa cells as above. The data are presented as mean $\pm$ S.D. from four independent experiments. The POP promoter in the reverse direction had no significant effect on *Amhr2* promoter activity. (D) The effect of the torazame ER sequence driven by the POP promoter on *Amhr2* promoter activity. *lncRNA-Amhr2* and its promoter sequence were replaced with torazame ER and the POP promoter. Luciferase assay was performed in primary granulosa cells as above. The data are presented as mean $\pm$ S.D. from four independent experiments. Torazame ER transcription by the POP promoter did not significantly increase *Amhr2* promoter activity. For all the data, the statistical significance was analyzed by one way ANOVA followed by Tukey-Kramer test. $**P < 0.01$.


Figure 6. Reporter gene assay in primary Sertoli cells. Primary Sertoli cells were cultured, and transfection and luciferase activity assay were done as in Figure 5A. The data are presented as mean $\pm$ S.D. from five or six independent experiments. The usage of POP promoter had no effect on *Amhr2* promoter. The statistical significance was analyzed by one way ANOVA followed by Tukey-Kramer test. $**P < 0.01$.

Figure 7. *lncRNA-Amhr2* transcription is correlated with *Amhr2* promoter activity in stably transfected OV3121 cells. (A) qRT-PCR analysis of *lncRNA-Amhr2* in stably transfected OV3121 cells. OV3121 cells were transfected with the indicated constructs, and the cells were selected with G418. Total RNAs were purified from the cells, and qRT-PCR was performed. Relative *lncRNA-Amhr2* levels were calculated by normalization to the *Gapdh* level and shown as a graph. The value of the cells with the 6.1k-EGFP construct was set to 1.0. The data are presented as mean $\pm$ S.D. from four independent qRT-PCR reactions by two transfection experiments. The *lncRNA-Amhr2* transcription was successfully enhanced by a POP promoter. Student *t*-test was performed to analyze the statistical significance. $**P < 0.01$. (B) qRT-PCR analysis of the EGFP reporter gene in stably transfected OV3121 cells. Stably transfected OV3121 cells were established as above, and EGFP mRNA levels were examined and presented as above. The data are presented as mean $\pm$ S.D. from four independent qRT-PCR reactions by two transfection experiments. EGFP expression was the highest when *lncRNA-Amhr2* was driven by a POP promoter. The statistical significance was analyzed by one way ANOVA followed by Tukey-Kramer test. $**P < 0.01$, $*P < 0.05$. (C) The photograph of OV3121 cells stably transfected with the 6.1k-EGFP construct. Bright field (left) and fluorescent images (right) are shown. Some cells were green, but others were not. (D) RT-PCR analysis of the EGFP gene in stably transfected OV3121 cells sorted by a cell sorter. OV3121 cells that were stably transfected with 6.1k-EGFP were sorted into green cells (EGFP+) and non-fluorescent cells (EGFP-) by a cell sorter. Total RNA was purified from each fraction, and RT-PCR was performed to detect *lncRNA-Amhr2* transcription. The EGFP expression was correlated with *lncRNA-Amhr2* transcription. The cycle numbers were 30 for *lncRNA-Amhr2* and 25 for *Gapdh*.


Figure 8. Induction of *lncRNA-Amhr2* transcription increased *Amhr2* promoter activity in the Tet-on system. (A) A schematic drawing of the 6.1k-TO-Luc construct for establishing the Tet-on inducible system. A 6.1-kb upstream region of the *Amhr2* gene was connected to the luciferase gene, and the

hCMV*1 promoter was inserted to induce the transcription of most sequence of *lncRNA-Amhr2*. (B-D)
OV3121 cells were transfected with the construct and selected with G418. The established stable cells
were treated with (Dox+) or without Dox (Dox-) for 5 or 24 hours, and total RNAs and protein
extracts were collected. Using the total RNAs, qRT-PCR was performed to measure *lncRNA-Amhr2*
(B) and luciferase mRNA levels (C). *Gapdh* was used as an internal control. The protein extracts were
used for luciferase activity assay and BCA assay. The luciferase enzymatic activity was normalized to
the value by BCA assay, and relative activity is shown as a graph (D). *lncRNA-Amhr2* transcription
was successfully induced by Dox, and luciferase expression was dramatically increased. The data are
presented as mean $\pm$ S.D. from three (B, C) or six independent experiments (D). Student *t*-test was
performed to analyze the statistical significance. **P* < 0.05, ***P* < 0.01, ****P* < 0.001.


(A)




(C)




(B)




(A)



(B)



(C)






Figure 4




Figure 5







Figure 6







(A)



(B)



(C)



(D)




Table 1. Oligonucleotide primers used for RT-PCR and qRT-PCR


Designation	Forward	Reverse
RT-PCR followed by electrophoresis		
Amhr2	5'-GCTCCTGGAAGATTGCTGGG-3'	5'-ATGTCGGCCACACGCATGTC-3'
lncRNA-Amhr2	5'-ACAGGGCTGGAGACAGCAT-3'	5'-CCACCTCAGCCAAGTAACTG-3'
EGFP	5'-ACGTAAACGGCCACAAGTTC-3'	5'-TTGAAGTTCACCTTGATGCC-3'
Gapdh	5'-CATGACCACAGTCCATGCCATC-3'	5'-TAGCCCAAGATGCCCTTCAGTG-3'
RT-PCR with a specific primer for RT		
Primer for RT		5'-CCACCTCAGCCAAGTAACTG-3'
lncRNA-Amhr2	5'-AAAGACACTTGTCCCTGGCA-3'	5'-AAGCAGTCTTCCTGCCTCTT-3'
qRT-PCR		
Amhr2	5'-TGCTAGGCCTACAGCATGAC-3'	5'-TCGCTCGTGTACTGAGTCAA-3'
lncRNA-Amhr2	5'-GGCTTGTGTTGCCATAGAGA-3'	5'-AAGCAGTCTTCCTGCCTCTT-3'
Nr5a1/Sf1	5'-GCCACCCTACAGCTATCCAG-3'	5'-CTGGTCCTCCTCTGGCTCTA-3'
Actb	5'-GGTCATCACTATTGGCAACG-3'	5'-ACGGATGTCAACGTCACACT-3'
EGFP	5'-AGCAAAGACCCCAACGAGAA-3'	5'-GGCGGCGGTCACGAA-3'
Luciferase	5'-GGGACGAAGACGAACACTTC-3'	5'-GGTGTGGAGCAAGATGGAT-3'
Gapdh	5'-CATGGCCTTCCGTGTTCTTA-3'	5'-CCTGCTTCACCACCTTCTTGA-3'

Table 2. Oligonucleotide primers used for RACE analyses

Designation	Forward	Reverse
5'RACE with ovarian granulosa cells		
Reverse transcription		5'-CCGTGCTTGGGTTCTGGGTT-3'
Gene specific primer for the first PCR		5'-ACTTATCTGATCTGGAGGCT-3'
Gene specific primer for the second PCR		5'-ATGCTGTCTCCAGCCCTGT-3'
5'RACE with the testis		
Reverse transcription		5'-ATGCTGTCTCCAGCCCTGT-3'
Gene specific primer for the first PCR		5'-AGAGACAGGAGCAGGAACCA-3'
Gene specific primer for the second PCR		5'-TGCCAGGGACAAGTGTCTTT-3'
3'RACE		
Gene specific primer for the first PCR	5'-CATCCTGGGCCTTGTCTCAA-3'	
Gene specific primer for the second PCR	5'-CAGTTACTTGGCTGAGGTGG-3'	

Table 3. Transgene expression and copy number of each transgenic line

Line	Green ovary	Green testis	Green gonad in embryo	Copy number
#318	No	No		1
#321	No	No		10
#326	No	Yes	No	3
#337	No	No		4
#338	No	No		2
#348	No	No		4
#356	No	No		2
#373	No	No		3
#376	No	No		4
#379	No	No		25



Supplemental Figure 1. qRT-PCR analysis of marker genes for the mouse ovary and testis in primary Sertoli and granulosa cells. Primary Sertoli cells were collected from 8-12 days old male mice, and primary granulosa cells were from immature female mice that were administrated with hormones. The cells were cultured for several days, and total RNAs were purified. cDNAs were synthesized with the oligo(dT) primer, and real time PCR was performed to detect the marker gene expression for each cell type. In the testis, *Hsd3b*, *Ddx4/Vasa*, and *Sgp2* were used as markers for Leydig cells, germ cells, and Sertoli cells, and in the ovary, *Cyp17a1*, *Gdf9*, and *Foxl2* were used for theca cells, oocytes, and granulosa cells.