



HOKKAIDO UNIVERSITY

Title	Proceedings of the 38th Sapporo Symposium on Partial Differential Equations
Author(s)	Giga, Y.; Jimbo, S.; Ozawa, T. et al.
Citation	Hokkaido University technical report series in mathematics, 159, i, 1-v, 76
Issue Date	2013-08-21
DOI	https://doi.org/10.14943/81521
Doc URL	https://hdl.handle.net/2115/68083
Type	departmental bulletin paper
File Information	tech159.pdf



Proceedings of the 38th Sapporo Symposium on Partial Differential Equations

Edited by
Y. Giga, S. Jimbo, T. Ozawa, K. Tsutaya, Y. Tonegawa
H. Kubo, T. Sakajo, and H. Takaoka

Series #159. August, 2013

HOKKAIDO UNIVERSITY
TECHNICAL REPORT SERIES IN MATHEMATICS
<http://eprints3.math.sci.hokudai.ac.jp/view/type/techreport.html>

- #137 H. Hida, T. Ito, H. Katsurada, K. Kitagawa (transcribed by T. Suda), Y. Taguchi, A. Murase and A. Yamagami. K. Arai, T. Hiraoka, K. Itakura, T. Kasio, H. Kawamura, I. Kimura, S. Mochizuki, M. Murata and T. Okazaki, 整数論札幌夏の学校, 201 pages. 2008.
- #138 J. Inoguchi, いろいろな幾何と曲線の時間発展, 66 pages. 2008.
- #139 M. Hayashi, I. Saito and S. Miyajima, 第 17 回関数空間セミナー, 91 pages. 2009.
- #140 T. Suda, Y. Umeta, K. Kasai, M. Kasedou, T. Yamanoi and K. Yoshida, 第 5 回数学総合若手研究集会, 252 pages. 2009.
- #141 T. Ozawa, Y. Giga, T. Sakajo, S. Jimbo, H. Takaoka, K. Tsutaya, Y. Tonegawa, G. Nakamura 第 34 回偏微分方程式論札幌シンポジウム, 67 pages. 2009.
- #142 K. Kasai, H. Kuroda, T. Nagai, K. Nishi, S. Tsujie and T. Yamaguchi, 第 6 回数学総合若手研究集会, 267 pages. 2010.
- #143 M. Hayashi, T. Nakazi, M. Yamada and R. Yoneda, 第 18 回関数空間セミナー, 80 pages. 2010.
- #144 Liang Chen, Doctoral thesis “On differential geometry of surfaces in anti de Sitter 3-space”, 79 pages. 2010.
- #145 T. Funaki, Y. Giga, M.-H. Giga, H. Ishii, R. V. Kohn, P. Rybka, T. Sakajo, P. E. Souganidis, Y. Tonegawa, and E. Yokoyama, Proceedings of minisemester on evolution of interfaces, Sapporo 2010, 279 pages. 2010.
- #146 T. Ozawa, Y. Giga, T. Sakajo, H. Takaoka, K. Tsutaya, Y. Tonegawa, and G. Nakamura, Proceedings of the 35th Sapporo Symposium on Partial Differential Equations, 67 pages. 2010.
- #147 M. Hayashi, T. Nakazi, M. Yamada and R. Yoneda, 第 19 回関数空間セミナー, 111 pages. 2011.
- #148 T. Fukunaga, N. Nakashima, A. Sekisaka, T. Sugai, K. Takasao and K. Umeta, 第 7 回数学総合若手研究集会, 280 pages. 2011.
- #149 M. Kasedou, Doctoral thesis “Differential geometry of spacelike submanifolds in de Sitter space”, 69 pages. 2011.
- #150 T. Ozawa, Y. Giga, T. Sakajo, S. Jimbo, H. Takaoka, K. Tsutaya, Y. Tonegawa and G. Nakamura, Proceedings of the 36th Sapporo Symposium on Partial Differential Equations, 63 pages. 2011.
- #151 K. Takasao, T. Ito, T. Sugai, D. Suyama, N. Nakashima, N. Miyagawa and A. Yano, 第 8 回数学総合若手研究集会, 286 pages. 2012.
- #152 M. Hayashi, T. Nakazi and M. Yamada, 第 20 回関数空間セミナー, 89 pages. 2012.
- #153 Y. Giga, S. Jimbo, G. Nakamura, T. Ozawa, T. Sakajo, H. Takaoka, Y. Tonegawa and K. Tsutaya, Proceedings of the 37th Sapporo Symposium on Partial Differential Equations, 81 pages. 2012.
- #154 N. Hu, Doctoral thesis “Affine geometry of space curves and homogeneous surfaces”, 69 pages. 2012.
- #155 2013 代数幾何学シンポジウム, 127 pages. 2013.
- #156 M. Hayashi, S. Miyajima, T. Nakazi, I. Saito and M. Yamada, 第 21 回関数空間セミナー, 90 pages. 2013.
- #157 D. Suyama, T. Ito, M. Kuroda, Y. goto, N. Teranishi, S. Futakuchi, T. Fuda and N. Miyagawa, 第 9 回数学総合若手研究集会, 344 pages. 2013.
- #158 Y. Giga, S. Jimbo, H. Terao, K. Yamaguchi, Proceedings of the 6th Pacific RIM Conference on Mathematics 2013, 154 pages. 2013.

Proceedings of the 38th Sapporo Symposium on Partial Differential Equations

Edited by

Y. Giga, S. Jimbo, T. Ozawa, K. Tsutaya, Y. Tonegawa
H. Kubo, T. Sakajo, and H. Takaoka

Sapporo, 2013

Partially supported by Grant-in-Aid for Scientific Research, the Japan Society
for the Promotion of Science.

日本学術振興会科学研究費補助金	(基盤研究 S 課題番号 21224001)
日本学術振興会科学研究費補助金	(基盤研究 B 課題番号 24340024)
日本学術振興会科学研究費補助金	(基盤研究 B 課題番号 25287022)
日本学術振興会科学研究費補助金	(挑戦的萌芽 課題番号 23654057)

PREFACE

This volume is intended as the proceedings of Sapporo Symposium on Partial Differential Equations, held on August 21 through August 23 in 2013 at Faculty of Science, Hokkaido University.

Sapporo Symposium on PDE has been held annually to present the latest developments on PDE with a broad spectrum of interests not limited to the methods of a particular school. Professor Taira Shirota started the symposium more than 35 years ago. Professor Kôji Kubota and late Professor Rentaro Agemi made a large contribution to its organization for many years.

We always thank their significant contribution to the progress of the Sapporo Symposium on PDE.

Y. Giga, S. Jimbo, T. Ozawa, K. Tsutaya, Y. Tonegawa
H. Kubo, T. Sakajo, and H. Takaoka

CONTENTS

Program

- S. Nakamura (The University of Tokyo)
Application of phase space analysis to the scattering theory for continuous and discrete Schrödinger equations
- J. Byeon (Korea Advanced Institute of Science and Technology)
Phase transition solutions to an Allen-Cahn model equation
- T. Ishiwata (Shibaura Institute of Technology)
Structure-preserving finite difference scheme for some sphere-valued partial differential equations
- S. Ishida (Tokyo University of Science)
Boundedness in degenerate Keller-Segel systems of parabolic-parabolic type
- S. Yoshikawa (Ehime University)
An error estimate of conservative finite difference scheme for the Boussinesq type equations
- T. Sideris (University of California, Santa Barbara)
Existence of solutions for isotropic elastodynamics
- S. Omata (Kanazawa University)
Mathematical modeling and numerical treatment of adhesion, exfoliation and collision
- K. Shirakawa (Chiba University)
Phase field systems of grain boundaries with solidification effects
- T. Kato (Nagoya University)
A cancellation property and the well-posedness for the periodic KdV equations
- K. Abe (The University of Tokyo)
Resolvent estimates for the Stokes equations in spaces of bounded functions
- K. Wakasa (Hokkaido University)
Global existence for semilinear wave equations with the blow-up term in high dimensions
- G. Orlandi (The University of Verona)
Time-like minimal surfaces in Minkowski space
- T. Ogawa (Tohoku University)
 L^1 maximal regularity and local existence of a solution to the compressible Navier-Stokes-Poisson system in a critical Besov space

The 38th Sapporo Symposium on Partial Differential Equations (第38回偏微分方程式論札幌シンポジウム)

組織委員: 久保英夫, 高岡秀夫

Organizers: H. Kubo, H. Takaoka

プログラム委員: 儀我美一, 神保秀一, 小澤徹, 津田谷公利, 利根川吉廣
久保英夫, 坂上貴之, 高岡秀夫

Program Committee: Y. Giga, S. Jimbo, T. Ozawa, K. Tsutaya, Y. Tonegawa
H. Kubo, T. Sakajo, and H. Takaoka

Period (期間) August 21, 2013 - August 23, 2013

Venue (場所) Room 203, Faculty of Science Building #5, Hokkaido University
北海道大学 理学部 5 号館大講義室 (203 号室)

URL <http://www.math.sci.hokudai.ac.jp/sympo/sapporo/program130821.html>

August 21, 2013 (Wednesday)

09:30-09:40 Opening Session

09:40-10:40 中村周 (東京大学) Shu Nakamura (The University of Tokyo)
Application of phase space analysis to the scattering theory for continuous and discrete Schrödinger equations

10:40-11:10 *

11:10-12:10 Jaeyoung Byeon (Korea Advanced Institute of Science and Technology)
Phase transition solutions to an Allen-Cahn model equation

14:00-14:30 *

14:30-15:00 石渡哲哉 (芝浦工業大学) Tetsuya Ishiwata (Shibaura Institute of Technology)
Structure-preserving finite difference scheme for some sphere-valued partial differential equations

15:20-15:50 石田祥子 (東京理科大学) Sachiko Ishida (Tokyo University of Science)
Boundedness in degenerate Keller-Segel systems of parabolic-parabolic type

16:00-16:30 吉川周二 (愛媛大学) Shuji Yoshikawa (Ehime University)
An error estimate of conservative finite difference scheme for the Boussinesq type equations

16:30-17:00 *

August 22, 2013 (Thursday)

09:30-10:30	Thomas Sideris (University of California, Santa Barbara) Existence of solutions for isotropic elastodynamics
10:30-11:00	*
11:00-12:00	小俣正朗 (金沢大学) Seiro Omata (Kanazawa University) Mathematical modeling and numerical treatment of adhesion, exfoliation and collision
14:00-14:30	*
14:30-15:00	白川健 (千葉大学) Ken Shirakawa (Chiba University) Phase field systems of grain boundaries with solidification effects
15:10-15:40	加藤孝盛 (名古屋大学) Takamori Kato (Nagoya University) A cancellation property and the well-posedness for the periodic KdV equations
15:40-16:00	*
16:00-16:30	阿部健 (東京大学) Ken Abe (The University of Tokyo) Resolvent estimates for the Stokes equations in spaces of bounded functions
16:40-17:10	若狭恭平 (北海道大学) Kyohei Wakasa (Hokkaido University) Global existence for semilinear wave equations with the blow-up term in high dimensions
17:10-17:30	*
18:00-20:00	Reception at Enreiso (懇親会, エンレイソウ)

August 23, 2013 (Friday)

09:30-10:30	Giandomenico Orlandi (The University of Verona) Time-like minimal surfaces in Minkowski space
10:30-11:00	*
11:00-12:00	小川卓克 (東北大学) Takayoshi Ogawa (Tohoku University) L^1 maximal regularity and local existence of a solution to the compressible Navier-Stokes-Poisson system in a critical Besov space
12:00-12:30	*

* Free discussion with speakers in the tea room

Application of phase space analysis to the scattering theory for continuous and discrete Schrödinger equations¹

Shu Nakamura²

Abstract

We discuss several applications of phase space analysis (or microlocal analysis) to Schrödinger equations. When we study scattering theory using microlocal analytic methods, it is often useful to utilize the Hörmander pseudodifferential operator theory in the Fourier space. This calculus is sometimes called *the scattering calculus* (following Melrose), but it goes back at least to classical works by Kitada and others.

We review application to the long-range scattering theory for Schrödinger equations on $\mathbb{R}^d$, and discuss its generalizations to discrete Schrödinger equations. “Microlocal analysis on $\mathbb{Z}^d$ ” sounds rather odd, but actually we can consider the problems microlocally on $T^*\mathbb{T}^d$ in a natural manner, where $\mathbb{T}^d = \mathbb{R}^d/(2\pi\mathbb{Z})^d$ denotes the d -dimensional torus, which is the Fourier space for $\mathbb{Z}^d$. T^*M denotes the cotangent bundle of M .

We also discuss the microlocal properties of the scattering matrices. For Schrödinger equations with short-range smooth potentials, we can show the wave operators, the scattering operator, and the scattering matrices are pseudodifferential operators in the Fourier space. Moreover, the asymptotic expansion of the symbol of the scattering matrices are shown to correspond to the classical Born expansion. We then discuss generalization of these results to discrete Schrödinger equations.

These results are still mostly in progress, and some results are preliminary.

1 Schrödinger operators on $\mathbb{R}^d$

We consider Schrödinger operator:

$$H = -\frac{1}{2}\Delta + V(x) \quad \text{on } \mathcal{H} = L^2(\mathbb{R}^d)$$

with the space dimension $d \geq 1$. During the talk, we always suppose V is a real-valued smooth function, and for any multi-index $\alpha \in \mathbb{Z}_+^d$, it satisfies

$$|\partial_x^\alpha V(x)| \leq C_\alpha \langle x \rangle^{-\mu-|\alpha|}, \quad x \in \mathbb{R}^d, \quad (1)$$

where $\mu > 0$ is a given decay rate of V , $C_\alpha > 0$, and $\langle x \rangle = (1 + |x|^2)^{1/2}$. Then it is well-known that H is self-adjoint with $D(H) = H^2(\mathbb{R}^d)$, the Sobolev space of order 2. The spectrum is bounded below, the essential spectrum is $\mathbb{R}_+$, and it is absolutely continuous possibly except for negative discrete eigenvalues. The solution to the time-dependent Schrödinger equation:

$$\frac{\partial}{\partial t} u(t) = -iHu(t), \quad u(0) = u_0 \in L^2(\mathbb{R}^d),$$

is given by $u(t) = e^{-itH}u_0$, where $u \in C(\mathbb{R}, L^2(\mathbb{R}^d))$.

¹Abstract for the 38th Sapporo Symposium on Partial Differential Equations, August 21–23, 2013. (Prepared on July 26, 2013)

²Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1, Komaba, Meguro, Tokyo, Japan 153-8914.

The Hamilton function of the corresponding classical mechanics is given by

$$p(x, \xi) = \frac{1}{2}|\xi|^2 + V(x), \quad (x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d,$$

and the classical flow is defined by the Hamilton vector field:

$$H_p = \sum_{j=1}^d \left(\frac{\partial p}{\partial \xi_j} \frac{\partial}{\partial x_j} - \frac{\partial p}{\partial x_j} \frac{\partial}{\partial \xi_j} \right) = \sum_{j=1}^d \left(\xi_j \frac{\partial}{\partial x_j} - \frac{\partial V}{\partial x_j} \frac{\partial}{\partial \xi_j} \right) \quad \text{on } \mathbb{R}^d \times \mathbb{R}^d.$$

Note that this flow is equivalent to the Newton equation:

$$\dot{x}(t) = \xi(t), \quad \dot{\xi}(t) = -\nabla V(x(t)).$$

Various properties of the Schrödinger evolution group e^{-itH} follow from the properties of the classical flow, as we will see in the context of the scattering theory later. In particular, we note the Hamilton operator H is considered as a *quantization* of $p(x, \xi)$ in the following sense. For a smooth function $a(x, \xi)$ on $\mathbb{R}^d \times \mathbb{R}^d$, the *Weyl quantization* $a(x, D_x)$ is defined by

$$a(x, D_x)u(x) = (2\pi)^{-d} \iint e^{i(x-y)\cdot\xi} a\left(\frac{x+y}{2}, \xi\right) u(y) dy d\xi, \quad u \in \mathcal{S}(\mathbb{R}^d).$$

Then $H = p(x, D_x)$, and we may also consider the time evolution e^{-itH} as a quantization of the Hamilton flow, though the justification of this observation is not obvious in general.

2 Schrödinger operators on $\mathbb{Z}^d$

In solid state physics, the discrete Schrödinger equation, sometimes called the Anderson tight binding model, is widely used. The discrete Schrödinger operator is defined by

$$\tilde{H}u[n] = -\frac{1}{2}\Delta u[n] + \tilde{V}[n]u[n], \quad n \in \mathbb{Z}^d, \quad u = u[\cdot] \in \tilde{\mathcal{H}} = \ell^2(\mathbb{Z}^d),$$

where the discrete Laplacian is defined (for example) by

$$\Delta u[n] = \sum_{|n-m|=1} u[m],$$

and $\tilde{V}[\cdot] : \mathbb{Z}^d \rightarrow \mathbb{R}$ is a potential function. If $\tilde{V}$ is bounded, then $\tilde{H}$ is a bounded self-adjoint operator on $\ell^2(\mathbb{Z}^d)$. Thus the solution to the time-dependent Schrödinger equation is also given by $e^{-it\tilde{H}}u_0$, $u_0 \in \ell^2(\mathbb{Z}^d)$.

Since the configuration space of the quantum particle is the lattice $\mathbb{Z}^d$, it is not obvious (at first) what the corresponding classical mechanics is. We suppose $\tilde{V}[\cdot]$ satisfies the following conditions: Let $\tilde{\partial}_j$ be the difference operator:

$$\tilde{\partial}_j u[n] = u[n] - u[n - e_j], \quad u \in \ell^2(\mathbb{Z}^d), \quad j = 1, \dots, d,$$

where $\{e_j\}_{j=1}^d$ is the standard basis of $\mathbb{R}^d$. Then we assume for any $\alpha \in \mathbb{Z}_d^d$,

$$|\tilde{\partial}^\alpha \tilde{V}[n]| \leq C_\alpha \langle n \rangle^{-\mu-|\alpha|}, \quad n \in \mathbb{Z}^d. \quad (2)$$

Then $\tilde{V}[n]$ is extended to a smooth potential function $V(x)$ which satisfies the conditions (1) in Section 1. We set

$$\tilde{p}(x, \xi) = - \sum_{j=1}^d \cos(\xi_j) + V(x), \quad (x, \xi) \in \mathbb{R}^d \times \mathbb{T}^d \cong T^*\mathbb{T}^d,$$

where $\mathbb{T}^d = \mathbb{R}^d / (2\pi\mathbb{Z})^d$. Then we may consider $\tilde{p}(x, \xi)$ as the classical Hamilton function corresponding to $\tilde{H}$. $\tilde{p}$ generates a Hamilton flow on $T^*\mathbb{T}^d$:

$$H_{\tilde{p}} = \sum_{j=1}^d \left(\frac{\partial \tilde{p}}{\partial \xi_j} \frac{\partial}{\partial x_j} - \frac{\partial \tilde{p}}{\partial x_j} \frac{\partial}{\partial \xi_j} \right) = \sum_{j=1}^d \left(\sin(\xi_j) \frac{\partial}{\partial x_j} - \frac{\partial V}{\partial x_j} \frac{\partial}{\partial \xi_j} \right) \quad \text{on } \mathbb{R}^d \times \mathbb{T}^d.$$

We note, by the Fourier inversion formula,

$$\tilde{H}u[n] = (2\pi)^{-d} \int_{\mathbb{T}^d} \sum_m e^{i(n-m)\cdot\xi} \tilde{p}(n, \xi) u[m] d\xi,$$

and this expression is different from the standard Weyl quantization. However, we can show

$$F\tilde{H}F^* \equiv \tilde{p}(-D_\xi, \xi)$$

modulo smoothing operators on $\mathbb{T}^d$, where $F: \ell^2(\mathbb{Z}^d) \rightarrow L^2(\mathbb{T}^d)$ is the discrete Fourier transform:

$$Fu(\xi) = (2\pi)^{-d/2} \sum_{n \in \mathbb{Z}^d} e^{-in \cdot \xi} u[n], \quad u \in \ell^2(\mathbb{Z}^d), \quad \xi \in \mathbb{T}^d,$$

and the right hand side is defined as a pseudodifferential operator on $\mathbb{T}^d$ with the symbol $\tilde{p}$. We will consider $\tilde{H}$ as a quantization of $\tilde{p}$ in this sense.

3 Short-range scattering theory on $\mathbb{R}^d$

Here we review the well-known formulation of the short-range scattering theory for Schrödinger equations on $\mathbb{R}^d$ (see, e.g., [5] Volume 3, [6]). We suppose V is short-range, i.e., the assumption (1) with $\mu > 1$. Then the wave operators:

$$W_\pm u = \lim_{t \rightarrow \pm\infty} e^{itH} e^{-itH_0} u, \quad u \in L^2(\mathbb{R}^d),$$

exist, where $H_0 = -\frac{1}{2}\Delta$. Moreover, it is proved that $W_\pm$ are asymptotically complete, i.e., $\text{Ran } W_\pm = \mathcal{H}_c(H)$, the continuous subspace of H .

If we write $u = W_\pm u_\pm \in \mathcal{H}_c(H)$, then by the definition of $W_\pm$, we learn

$$u(t) = e^{-itH} u \sim e^{-itH_0} u_\pm \quad \text{as } t \rightarrow \pm\infty.$$

Namely, $u(t)$ asymptotically converges to the free evolutions $e^{-itH_0} u_\pm$ as $t \rightarrow \pm\infty$. $u_\pm$ is called the *scattering data*.

This result corresponds to the fact that if a classical particle $(x(t), \xi(t))$ is not trapped, then there are (classical mechanical) scattering data $(x_\pm, \xi_\pm) \in \mathbb{R}^d \times \mathbb{R}^d$ such that

$$x(t) \sim x_\pm + t\xi_\pm, \quad \xi(t) \sim \xi_\pm \quad \text{as } t \rightarrow \pm\infty.$$

This correspondence plays crucial role in the following discussions.

4 Short-range scattering theory on $\mathbb{Z}^d$

The scattering theory is constructed for the Schrödinger operators on $\mathbb{Z}^d$ similarly. (See, e.g., Isozaki-Korotyaev [4], Boutet de Monvel-Sahbani [2]). Here we also suppose $\tilde{V}$ is short-range, i.e., the assumption (2) with $\mu > 1$. Then the wave operators:

$$\tilde{W}_\pm u = \lim_{t \rightarrow \pm\infty} e^{it\tilde{H}} e^{-it\tilde{H}_0} u, \quad u \in \ell^2(\mathbb{Z}^d),$$

exist, where $\tilde{H}_0 = -\frac{1}{2}\Delta$. Moreover, it is also proved that $\tilde{W}_\pm$ are asymptotically complete, i.e., $\text{Ran } \tilde{W}_\pm = \tilde{\mathcal{H}}_c(\tilde{H})$, the continuous subspace of $\tilde{H}$. Again, if we write $u = \tilde{W}_\pm u_\pm \in \tilde{\mathcal{H}}_c(\tilde{H})$, we learn

$$u(t) = e^{-it\tilde{H}} u \sim e^{-it\tilde{H}_0} u_\pm \quad \text{as } t \rightarrow \pm\infty.$$

The corresponding classical mechanical scattering is not as straightforward. We denote

$$\tilde{p}_0(\xi) = -\sum_{j=1}^d \cos(\xi_j) \quad \text{on } \mathbb{T}^d,$$

and we set the velocity function by

$$v_j(\xi) = \frac{\partial \tilde{p}_0}{\partial \xi_j}(\xi) = \sin(\xi_j), \quad j = 1, \dots, d, \quad \xi \in \mathbb{T}^d.$$

Then the classical free evolution is give by :

$$\exp tH_{\tilde{p}_0}(x, \xi) = x + tv(\xi), \quad t \in \mathbb{R}, \quad (x, \xi) \in \mathbb{R}^d \times \mathbb{T}^d.$$

Thus, if a classical particle i.e., a trajectory along $H_{\tilde{p}}$: $(x(t), \xi(t)) \in \mathbb{R}^d \times \mathbb{T}^d$ is not trapped, then there exist scattering data $(x_\pm, \xi_\pm)$ such that

$$x(t) \sim x_\pm + tv(\xi_\pm), \quad \xi(t) \sim \xi_\pm \quad \text{as } t \rightarrow \pm\infty.$$

We note that the “particle” moves in a *virtual configuration space* $\mathbb{R}^d$, not in the original configuration space $\mathbb{Z}^d$.

5 Long-range scattering theory on $\mathbb{R}^d$

If the potential is long-range type, i.e., if V satisfies the assumption (1) with $0 < \mu \leq 1$, then the above argument does not hold. In this case we need to modify the free evolution to approximate the solution as $t \rightarrow \pm\infty$ (see, e.g., [5] Section X.9).

If μ is modestly long-range, i.e., if $1/2 < \mu \leq 1$, then we can employ the Dollard modifier:

$$\Phi_D(t, \xi) = \int_0^t p(s\xi, \xi) ds = \frac{t}{2} |\xi|^2 + \int_0^t V(s\xi) ds, \quad t \in \mathbb{R}, \quad \xi \in \mathbb{R}^d.$$

We set the Fourier multiplier

$$U_D(t) = \exp(-i\Phi_D(t, D_x)), \quad t \in \mathbb{R},$$

as the *modified free evolution*. Then the modified wave operators:

$$W_{\pm}^D u = \lim_{t \rightarrow \pm\infty} e^{itH} U_D(t) u, \quad u \in L^2(\mathbb{R}^d)$$

exist, and are asymptotically complete: $\text{Ran } W_{\pm}^D = \mathcal{H}_c(H)$.

More generally, we construct solutions to the momentum space Hamilton-Jacobi equation

$$\frac{\partial \Phi}{\partial t}(t, \xi) = p(\partial_{\xi} \Phi(t, \xi), \xi), \quad \pm t \geq 0, \quad \xi \in \mathbb{R}^d,$$

with the initial condition: $\Phi(0, \xi) = \pm R|\xi|$, where $R > 0$ is a sufficiently large constant. Then the modified free evolution is defined similarly: $U(t) = \exp(-i\Phi(t, D_x))$; the modified wave operators exist, and are complete.

We note that the construction of the modified free evolution relies on the classical mechanical argument crucially. The Dollard modifier is a first order approximate solution to the Hamilton-Jacobi equation.

6 Long-range scattering theory on $\mathbb{Z}^d$

The long-range scattering theory for the discrete Schrödinger equations can be constructed similarly using the corresponding classical mechanics on $T^*\mathbb{T}^d$.

If $1/2 < \mu \leq 1$, then we set

$$\tilde{\Phi}_D(t, \xi) = \int_0^t \tilde{p}(sv(\xi), \xi) ds = t\tilde{p}_0(\xi) + \int_0^t V(sv(\xi)) ds, \quad t \in \mathbb{R}, \quad \xi \in \mathbb{T}^d,$$

and

$$\tilde{U}_D(t) = F^* \exp(-i\tilde{\Phi}_D(t, \xi)) F,$$

Then the modified wave operators

$$\tilde{W}_{\pm}^D u = \lim_{t \rightarrow \pm\infty} e^{it\tilde{H}} \tilde{U}_D(t) u, \quad u \in \ell^2(\mathbb{Z}^d),$$

exist and are complete. More generally, we can construct solutions to the Hamilton-Jacobi equation

$$\frac{\partial \tilde{\Phi}}{\partial t}(t, \xi) = \tilde{p}(\partial_{\xi} \tilde{\Phi}(t, \xi), \xi), \quad \pm t \geq 0, \quad \xi \in \mathbb{T}^d,$$

with the initial condition $\tilde{\Phi}(0, \xi) = \pm R\tilde{p}_0(\xi)$. We note the construction of $\tilde{\Phi}(t, \xi)$ is local in the energy: $\lambda = p(x, \xi)$, and we need to avoid the threshold energies $\mathcal{T} = \{-d + 2j \mid j = 0, \dots, d\}$.

7 Scattering matrix for Schrödinger operators on $\mathbb{R}^d$

Here we consider scattering theory from the microlocal point of view. Here we suppose the potential V satisfies the short range condition: $\mu > 1$ in (1). The scattering operator is defined by

$$S = W_+^* W_- : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d),$$

and it is unitary. Moreover, it commutes with the free energy: $SH_0 = H_0S$, and hence

$$\frac{1}{2}|\xi|^2(\mathcal{F}S\mathcal{F}^*)u(\xi) = (\mathcal{F}S\mathcal{F}^*)\frac{1}{2}|\xi|^2u(\xi) \quad \text{for } u(\xi) \in L^2(\mathbb{R}^d).$$

Thus $\mathcal{F}S\mathcal{F}^*$ is reduced to a family of operators on $L^2(\Sigma_\lambda)$, $\lambda > 0$, where

$$\Sigma_\lambda = \{\xi \in \mathbb{R}^d \mid \frac{1}{2}|\xi|^2 = \lambda\}, \quad \lambda > 0,$$

is the energy surface. We denote

$$\mathcal{F}S\mathcal{F}^* = \int^\oplus S(\lambda)d\lambda \quad \text{on } L^2(\mathbb{R}^d) \cong \int^\oplus L^2(\Sigma_\lambda)d\lambda,$$

and $S(\lambda) : L^2(\Sigma_\lambda) \rightarrow L^2(\Sigma_\lambda)$ is called the scattering matrix. The scattering matrix is one of the most fundamental observables in the quantum mechanics.

We set

$$W(t) = e^{itH_0}e^{-itH}, \quad t \in \mathbb{R}.$$

Then it is easy to see

$$\frac{d}{dt}W(t) = -i(e^{itH_0}He^{-itH_0} - H_0)W(t) = -i(e^{itH_0}Ve^{-itH_0})W(t).$$

Using the Weyl quantization, we learn

$$(e^{itH_0}Ve^{-itH_0}) = V(x + tD_x),$$

and hence we may consider $\{W(t)\}$ as an evolution operator with a time-dependent generator $V(x + tD_x)$. Since $V(x + tD_x)$ is bounded, the solution is expressed using the Dyson expansion:

$$\begin{aligned} W(t) &= 1 + \sum_{n=1}^{\infty} (-i)^n \int_0^t \int_0^{t_1} \cdots \int_0^{t_{n-1}} V(x + t_1 D_x) \cdots V(x + t_n D_x) dt_n \cdots dt_1 \\ &= 1 - i \int_0^t V(x + t_1 D_x) dt_1 - \int_0^t \int_0^{t_1} V(x + t_1 D_x) V(x + t_2 D_x) dt_2 dt_1 + \cdots \end{aligned}$$

We may consider this expansion as an asymptotic expansion as pseudodifferential operators in the Fourier space. On the other hand, $W(t)$ converges to $W_\pm^*$ weakly as $t \rightarrow \pm\infty$. Each terms in the Dyson expansion does not converge as $t \rightarrow \pm\infty$ in the standard symbol class, but we can show the wave operators are in fact pseudodifferential operators. Moreover the scattering operator is also a pseudodifferential operator and

$$S = 1 - i \int_{-\infty}^{\infty} V(x + tD_x) dt + (\text{lower order terms}),$$

in some sense. This then implies the scattering matrix is a pseudodifferential operator on Σ_λ with the symbol

$$S(\lambda; x, \xi) = 1 - i \int_{-\infty}^{\infty} V(x + t\xi) dt + (\text{lower order terms}),$$

where $(x, \xi) \in \{(x, \xi) \in \mathbb{R}^d \times \Sigma_\lambda \mid x \perp \xi\} \cong T^*\Sigma_\lambda$, i.e., (x, ξ) is an element of the cotangent bundle of Σ_λ . Using this expression, we can compute the asymptotic distribution of eigenvalues of the scattering matrix, at least formally (see, e.g., Birman-Yafaev [1], Bulgar-Pushnitski [3]).

8 Scattering matrix for Schrödinger operators on $\mathbb{Z}^d$

We can also consider the scattering matrix for discrete Schrödinger operators. For $\lambda \in [-d, d]$, the energy surface is defined by

$$\Lambda_\lambda = \{\xi \in \mathbb{T}^d \mid \tilde{p}_0(\xi) = \lambda\},$$

and it is a regular submanifold of $\mathbb{T}^d$ unless $\lambda \in \mathcal{J}$. The scattering matrix is defined as a unitary operator $\tilde{S}(\lambda)$ on $L^2(\Lambda_\lambda)$ as well as the continuous case. For the wave operators, we compute:

$$\frac{d}{dt}(e^{it\tilde{H}}e^{-it\tilde{H}_0}) = -iV_1(t)(e^{it\tilde{H}}e^{-it\tilde{H}_0})$$

where $FV_1(t)F^*$ is a pseudodifferential operator on $\mathbb{T}^d$ with the symbol:

$$V_1(t; x, \xi) = V(x + tv(\xi)) + (\text{lower order terms}), \quad (x, \xi) \in \mathbb{R}^d \times \mathbb{T}^d \cong T^*\mathbb{T}^d.$$

Then we obtain the Dyson expansion of the evolution:

$$e^{it\tilde{H}}e^{-it\tilde{H}_0} = 1 - i \int_0^t V_1(t_1)dt_1 - \int_0^t \int_0^{t_1} V_1(t_1)V_1(t_2)dt_2dt_1 + \dots$$

Analogously to the previous section, we can show that the scattering matrix $S(\lambda)$ is a pseudodifferential operator on Λ_λ , and the symbol is given by

$$\tilde{S}(\lambda; x, \xi) = 1 - i \int_{-\infty}^{\infty} V(x + tv(\xi))dt + (\text{lower order terms}),$$

where $(x, \xi) \in \{(x, \xi) \in \mathbb{R}^d \times \Lambda_\lambda \mid x \perp v(\xi)\} \cong T^*\Lambda_\lambda$. We can apply this argument to obtain the asymptotic distribution of the eigenvalues of $\tilde{S}(\lambda)$.

References

- [1] Birman, M. Sh., Yafaev, D. R.: Asymptotic behavior of the spectrum of the scattering matrix. J. Sov. Math. **25** (1984), 793–814.
- [2] Boutet de Monvel, A., Sakhbani, J.: On the spectral properties of discrete Schrödinger operators: The multi-dimensional case. Rev. Math. Phys. **11** (1999), 1061–1078.
- [3] Bulger, D., Pushnitski, A.: The spectral density of the scattering matrix for high energies. Preprint 2011. Arxiv:1110.3710.
- [4] Isozaki, H., Korotyaev, E.: Inverse problems, trace formulae for discrete Schrödinger operators. Ann. Inst. H. Poincaré **13** (2012), 751–788.
- [5] Reed, M., Simon, B.: Methods of Modern Mathematical Physics Vols. I–IV. Academic Press, 1972–1979.
- [6] Yafaev, D. R.: Mathematical Scattering Theory, Analytic theory. American Math. Soc. 2010.

Phase transition solutions to an Allen-Cahn model equation

Jaeyoung Byeon
KAIST, Republic of Korea

We consider the following equation

$$(0.1) \quad -\Delta u + A(x)G'(u) = 0 \quad \text{in } \mathbb{R}^n$$

where $A(x) = A(x+i)$ for $x \in \mathbb{R}^n, i \in \mathbb{Z}^n$, $G(u+j) = G(u)$ for $u \in \mathbb{R}, j \in \mathbb{Z}$, $G(u) > 0$ for $u \in \mathbb{R} \setminus \mathbb{Z}$, $G(j) = 0, G''(j) > 0$ for $j \in \mathbb{Z}$ and $G(t) = G(-t)$ for $t \in \mathbb{R}$.

Several authors have studied Allen-Cahn phase transition models in which the spatial phase transition manifests itself as a heteroclinic or homoclinic solution of the corresponding partial differential equation. See e.g. [1]-[4], [16]-[17] and [20]. Similar type of solutions in more general settings arise in extension's of Moser's work [14] on developing an Aubry-Mather theory for PDE's: [5], [6]-[7], [9]-[11], [18]-[19], [21], and [23]. In all of these sources the transition type solutions are *unidirectional* in the sense that they change in a particular direction.

In this talk, I would like to introduce some recent works with Paul Rabinowitz about some *multidirectional* solutions. For autonomous problem, that is, $A = \text{constant}$, there have been many works [22], [12],[13] and references therein which show that the bounded solutions of (0.1) are closely related to the minimal surfaces in $\mathbb{R}^n$.

We believe that there would be a generic condition for existence of *multidirectional* solutions like gap conditions for construction of multi-transition (*unidirectional*) solutions(see [21] and references therein). We do not know how to do this yet for (0.1), but will present a certain type of non-autonomous term A for which such solutions can be found.

To describe our results, let $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ and $A \in C^1(\mathbb{R}^n)$ be a nonnegative function that is 1-periodic in $x_i, 1 \leq i \leq n$. Set $\Omega \equiv$

$\{x \in (0, 1)^n \mid A(x) > 0\}$. We assume $2d^* = |\partial\Omega - \partial[0, 1]^n| > 0$ and $\partial\Omega$ is a smooth manifold. Set $\Omega_d \equiv \{x \in \Omega \mid |x - \partial\Omega| > d\}$. Then for sufficiently small $d \in (0, d^*)$, $\partial\Omega_d$ is diffeomorphic to $\partial\Omega$. Fixing such a small d , let $\varepsilon > 0$ and $A_\varepsilon = 1 + \frac{1}{\varepsilon}A$. Our model equation is

$$(0.2) \quad -\Delta u + A_\varepsilon G'(u) = 0, \quad x \in \mathbb{R}^n.$$

First, we are interested in solutions of (0.2) satisfying $0 < U < 1$ and that are near 1 on A^T and near 0 on B^T , where

$$T \subset \mathbb{Z}^n, \quad A^T = \cup_{i \in T} (i + \Omega), \quad B^T = \cup_{i \in \mathbb{Z}^n \setminus T} (i + \Omega).$$

To describe our results more precisely, for $d \in (0, d^*)$ and Ω_d as above, we define

$$A_T \equiv \cup \{i + \Omega_d \mid i \in T\}$$

and

$$B_T \equiv \cup \{i + \Omega_d \mid i \in \mathbb{Z}^n \setminus T\}.$$

Set

$$L_\varepsilon(u) = \frac{1}{2} |\nabla u|^2 + A_\varepsilon G(u) \quad \text{and} \quad J_\varepsilon(u) = \int_{\mathbb{R}^n} L_\varepsilon(u) \, dx.$$

Choose $0 < b < \frac{1}{2} < a < 1$ and define

$$\tilde{\Gamma}(T) = \{u \in C^2(\mathbb{R}^n, [0, 1]) \mid u \geq a > 1/2 \text{ on } A_T \text{ and } u \leq b < 1/2 \text{ on } B_T\}.$$

Whenever a solution, $u \in \tilde{\Gamma}(T)$, of (0.2) satisfies

$$J_\varepsilon(u + \varphi) - J_\varepsilon(u) = \int_{\text{supp } \varphi} (L_\varepsilon(u + \varphi) - L_\varepsilon(u)) \, dx \geq 0$$

for all $\varphi \in C_0^\infty(\mathbb{R}^n)$ such that $u + \varphi \in \tilde{\Gamma}(T)$, u will be said to be *minimal* in $\tilde{\Gamma}(T)$.

The solutions obtained here will have certain decay properties relative to T . To help describe them, for $x \in \mathbb{R}^n$, define

$$d(x, T) \equiv \text{dist}(x, \partial(T + [0, 1]^n))$$

and for $S \subset \mathbb{R}^n$, let χ_S denote the characteristic function of S . Our first main result [8] is:

Theorem 0.3. *Under the above hypotheses on A and G , there is an $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ and any $T \subset \mathbb{Z}^n$,*

- 1° *there exists a solution $U = U_{\varepsilon, T} \in \tilde{\Gamma}(T)$ of (0.2);*
- 2° *U is minimal in $\tilde{\Gamma}(T)$;*
- 3° *$0 < U_{\varepsilon, T} < 1$;*
- 4° *$U_{\varepsilon, T}$ converges uniformly to 1 on A^T and to 0 on B^T as $\varepsilon \rightarrow 0$;*
- 5° *there exist constants $C, c > 0$, independent of $T \subset \mathbb{Z}^n$ and of $\varepsilon \in (0, \varepsilon_0)$, satisfying*

$$|U_{\varepsilon, T}(x) - \chi_{T+[0,1]^n}(x)| \leq C \exp(-cd(x, T)), \quad x \in \mathbb{R}^n.$$

Second, we are interested in unbounded solutions of (0.2). As an Aubry-Mather theory for PDE's, Moser [14] obtained among other things the following existence result.

Theorem 0.4. *For each $\alpha \in \mathbb{R}^n$, there exists a solution u_α of (0.1) such that for some $C > 0$, $|u_\alpha(x) - \alpha \cdot x| \leq C$, $x \in \mathbb{R}^n$.*

Let $P : \mathbb{Z}^n + \Omega \rightarrow \mathbb{Z}$ be a function such that P is constant on $i + \Omega$ for each $i \in \mathbb{Z}^n$. We say P is Lipschitz continuous if there exists $C > 0$ such that $|P(x) - P(y)| \leq C|x - y|$ for any $x, y \in \mathbb{Z}^n + \Omega$.

As for *multidirectional* (unbounded) solutions generalizing the *unidirectional* solutions obtained by Moser in Theorem 0.4, our second main result is:

Theorem 0.5. *Let A_ε be as above. Suppose P be Lipschitz continuous on S . Then there is an ε_0 depending on the Lipschitz constant of P such that for each $\varepsilon \in (0, \varepsilon_0)$, there is a solution, $U = U_{\varepsilon, P}$ of (0.2) with $U \rightarrow P$ uniformly on $\mathbb{Z}^n + \Omega$ as $\varepsilon \rightarrow 0$.*

References

- [1] FRANCESCA ALESSIO, LOUIS JEANJEAN AND PIERO MONTECCHIARI, Stationary layered solutions in $\mathbb{R}^2$ for a class of non autonomous Allen-Cahn equations, Calc. Var. Partial Differential Equations, **11** (2000), 177–202

- [2] FRANCESCA ALESSIO, LOUIS JEANJEAN AND PIERO MONTECCHIARI, Existence of infinitely many stationary layered solutions in $\mathbb{R}^2$ for a class of periodic Allen-Cahn equations, *Comm. Partial Differential Equations*, **27** (2002), 1537–1574
- [3] FRANCESCA ALESSIO AND PIERO MONTECCHIARI, Entire solutions in $\mathbb{R}^2$ for a class of Allen-Cahn equations, *ESAIM Control Optim. Calc. Var.*, **11** (2005), 633–672
- [4] FRANCESCA ALESSIO AND PIERO MONTECCHIARI, Multiplicity of entire solutions for a class of almost periodic Allen-Cahn type equations, *Adv. Nonlinear Stud.*, **5** (2005), 515–549
- [5] VICTOR BANGERT, On minimal laminations of the torus, *Ann. Inst. Poincare Anal. Non Lineaire*, **6** (1989), 95–138
- [6] UGO BESSI, Many solutions of elliptic problems on $\mathbb{R}^n$ of irrational slope, *Comm. Partial Differential Equations*, **30** (2005), 1773–1804
- [7] UGO BESSI, Slope-changing solutions of elliptic problems on $\mathbb{R}^n$, *Nonlinear Anal.*, **68** (2008), 3923–3947
- [8] JAEYOUNG, PAUL H. RABINOWITZ, On a phase transition model. *Calc. Var. Partial Differential Equations*, **47** (2013), 1–23
- [9] LUIS A. CAFFARELLI AND RAFAEL DE LA LLAVE, Planelike minimizers in periodic media. *Comm. Pure Appl. Math.*, **54** (2001), 1403–1441
- [10] RAFAEL DE LA LLAVE AND ENRICO VALDINOCI, Multiplicity results for interfaces of Ginzburg-Landau Allen-Cahn equations in periodic media, *Adv. Math.*, **215** (2007), 379–426
- [11] RAFAEL DE LA LLAVE AND ENRICO VALDINOCI, A generalization of Aubry-Mather theory to partial differential equations and pseudo-differential equations, *Ann. Inst. H. Poincare Anal. Non Lineaire*, **26** (2009), 1309–1344
- [12] MANUEL DEL PINO, MICHAEL KOWALCZYK, JUNCHENG WEI, On De Giorgi’s conjecture in dimension $N \geq 9$. *Ann. of Math.* **174** (2011), 1485–1569
- [13] MANUEL DEL PINO, MICHAEL KOWALCZYK, JUNCHENG WEI, Entire solutions of the Allen-Cahn equation and complete embedded minimal surfaces of finite total curvature in $\mathbb{R}^3$, *J. Differential Geom.*, **93** (2013), 67–131
- [14] JURGEN MOSER, Minimal solutions of a variational problems on a torus, *Ann. Inst. Poincare Anal. Non Lineaire*, **3** (1986), 229–272
- [15] MATTEO NOVAGA AND ENRICO VALDINOCI, Bump solutions for the mesoscopic Allen-Cahn equation in periodic media. *Calc. Var. Partial Differential Equations*, **40** (2011), no. 1-2, 37–49

- [16] PAUL H. RABINOWITZ AND ED STREDULINSKY, Mixed states for an Allen-Cahn type equation, *Comm. Pure Appl. Math.*, **56** (2003), 1078–1134
- [17] PAUL H. RABINOWITZ AND ED STREDULINSKY, Mixed states for an Allen-Cahn type equation. II, *Calc. Var. Partial Differential Equations*, **21** (2004), 157–207
- [18] PAUL. H. RABINOWITZ AND ED STREDULINSKY, On some results of Moser and of Bangert, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **21** (2004), 673–688
- [19] PAUL H. RABINOWITZ AND ED STREDULINSKY, On some results of Moser and of Bangert. II, *Adv. Nonlinear Stud.*, **4** (2004), 377–396
- [20] PAUL H. RABINOWITZ AND ED STREDULINSKY, On a class of infinite transition solutions for an Allen-Cahn model equation, *Discrete Contin. Dyn. Syst.*, **21** (2008), 319–332
- [21] PAUL H. RABINOWITZ AND ED STREDULINSKY, Extensions of Moser-Bangert Theory: Locally Minimal Solutions, *Progress in Nonlinear Differential Equations and Their Applications*, 81, Birkhauser, Boston, (2011)
- [22] OVIDIU SAVIN, Regularity of flat level sets in phase transitions, *Ann. of Math.*, **169**(2009), 41–78
- [23] ENRICO VALDINOCI, Plane-like minimizers in periodic media: jet flows and Ginzburg-Landau-type functionals, *J. Reine Angew. Math.*, **574** (2004), 147–185

Structure-preserving finite difference scheme for some sphere-valued partial differential equations *

Tetsuya Ishiwata(Shibaura Institute of Technology)

Keywords: (*Finite difference method, structure-preserving scheme, vector-valued PDE, energy structure, sphere-valued solution*)

1 Target PDE and aim

In this talk, we treat some typical sphere-valued partial differential equations, for examples, Heisenberg equation and Landau-Lifshitz equation. These equations describe the evolution of spin fields in continuum ferromagnetism and have the following properties:

1. length preserving,
2. energy conservation or dissipation property.

We propose a finite difference scheme for these equations which inherits the above properties and show some theoretical results on the scheme. And we also demonstrate numerical examples in order to show the effectiveness of our scheme.

2 Landau-Lifshitz equation

Landau-Lifshitz equation is the following vector-valued partial differential equation:

$$\frac{\partial u}{\partial t} = \mu u \times \Delta u - \lambda u \times (u \times \Delta u). \quad (1)$$

Here, $\lambda \geq 0$, $\mu \in \mathbb{R}$ are constants, $\Omega \subseteq \mathbb{R}^N$ is the material and $u = u(x, t) = (u_1(x, t), u_2(x, t), u_3(x, t)) : \Omega \times (0, \infty) \rightarrow \mathbb{R}^3$ describes a spin field. In case $\lambda = 0$, the equation (1) is often called the Heisenberg equation.

By taking an inner product of (1) with $u(x, t)$, we obtain $|u(x, t)| = |u_0(x)|$ for any $t > 0$. Thus we usually consider the case $|u_0| = |u(x, t)| = 1$ as the spin model.

*Collaborators : A. Fuwa (Mizuho Information & Research Institute), K. Kumazaki (Tomakomai National College of Technology) and M. Tsutsumi (Waseda University)

We also obtain the following energy equality:

$$E(u(t)) = E(u_0) - 2\lambda \int_0^t \|u(\cdot, s) \times \Delta u(\cdot, s)\|_{L^2(\Omega)}^2 ds. \quad (2)$$

Here, $E(u(t)) := \|\nabla u(\cdot, t)\|_{L^2(\Omega)}^2$.

In this talk, for simplicity, we only treat one dimensional case with periodic boundary condition. We remark that we can easily extend our results for higher dimensional case and other suitable boundary conditions.

3 Proposed scheme

Let $\Delta t > 0, \Delta x := L/N$ ($N \in \mathbb{N}$) be time and spatial mesh sizes, where $L > 0$ is the spatial period. We denote by $U_n^j = (U_{1,n}^j, U_{2,n}^j, U_{3,n}^j)$ an approximate vector of $u(x, t)$ at $x = x_n := n\Delta x, t = t_j := j\Delta t$. Our proposed scheme is the following:

$$\begin{aligned} \frac{U_n^{j+1} - U_n^j}{\Delta t} &= \mu U_n^{(j+1,j)} \times \tilde{\Delta} U_n^{(j+1,j)} - \lambda U_n^{(j+1,j)} \times \left(U_n^{(j+1,j)} \times \tilde{\Delta} U_n^{(j+1,j)} \right), \\ U_n^0 &= u_0(x_n), \\ U_{N-1}^j &= U_{-1}^j, \quad U_N^j = U_0^j. \end{aligned}$$

Here, $U_n^{(j+1,j)} = (U_n^{j+1} + U_n^j)/2$, $\tilde{\Delta} := D^+ D^-$ is a standard discrete Laplacian, where D^+ (resp. D^-) is a forward (resp. backward) difference operator on space, that is,

$$D^+ X_n = \frac{X_{n+1} - X_n}{\Delta x}, \quad D^- X_n = \frac{X_n - X_{n-1}}{\Delta x}.$$

For a finite difference solution of the above scheme, we have the following properties on a length of the solution vector and an energy structure.

Theorem 1. *Any finite difference solution of the proposed scheme satisfies the following:*

- (1) *The solution keeps its length, that is, $|U_n^{j+1}| = |U_n^j|$ for all j and n .*
- (2) *The following energy equality is satisfied:*

$$E_h(U^j) = E_h(U^0) - 2\lambda \sum_{i=0}^{j-1} \|U^{(i+1,i)} \times \tilde{\Delta}_h U^{(i+1,i)}\|_2^2 \Delta t.$$

Here $E_h(U^j) := \|D^+ U^j\|_2^2$, where $\|v\|_2 = \left(\sum_{n=0}^{N-1} |v_n|^2 \Delta x \right)^{1/2}$.

Note that the above energy equality is a discrete version of that of original problem. That is, the proposed scheme inherits length-preserving property and the energy structure from the original problem.

We also have the following error estimate.

Theorem 2. *Let $T > 0$. Assume that $u_0(x), u(x, t)$ are sufficiently smooth for $x \in [0, L]$ and $t \in (0, T)$. Then, there exist constants $C_c := C_c(\lambda, \mu, u_0, L, T)$ and $C_\gamma := C_\gamma(\lambda, \mu, L, K)$. For any $\Delta t < C_c$ and $\Delta x > 0$ with $\frac{\Delta t}{\Delta x^2} < C_\gamma$, we have*

$$\|U_n^j - u(x_n, t_j)\|_{h^1} = O(\Delta t^2 + \Delta x^2).$$

Here, $\|X\|_{h^1} = \left(\|X\|_2^2 + \|D^+ X\|_2^2 \right)^{1/2}$.

4 Numerical results

In this section we show two numerical examples for the cases $\lambda = 0$ (Heisenberg case) and $\lambda > 0$ (Landau-Lifshitz case) to show the effectiveness of the proposed scheme. Here we use exact solutions and compare them.

First we see the Heisenberg case ($\lambda = 0$). We have $\max ||U_n^j| - 1| \sim 10^{-15}$, that is, we verify the length-preserving property numerically. Figure 1 shows the behavior of the energy. We see that the numerical energy keeps constant value. Thus, we conclude that the numerical energy is conserved. Figure 2 shows the behaviors of (a) U_1 and (b) u_1 of the exact solution. The both behaviors coincide very much.




Fig. 1: Time evolution of the energy ($\lambda = 0$).




Fig. 2: Time evolutions of u_1 of (a) numerical solution and (b) exact solution in the case $\lambda = 0$.

Next we see the Landau-Lifshitz case ($\lambda = 0.1$). We also have $\max ||U_n^j| - 1| \sim 10^{-15}$, that is, we verify the length-preserving property numerically. Figure 3 shows the behaviors of the energy: The curved line describes the energy of numerical solution and “+” marks shows the energy of the exact solution. We see that the behavior of numerical energy is almost same as that of the exact solution. Figure 4 shows the behaviors of (a) U_1 and (b) u_1 of the exact solution. We also conclude that behaviors of numerical solution and the exact solution coincide very much.

From these numerical results via a test problem, we conclude that the proposed scheme is effective.




Fig. 3: Time evolution of energy for $\lambda = 0.1$: numerical solution (curved line) and exact solution (“+” mark).




Fig. 4: Time evolutions of u_1 of (a) numerical solution and (b) exact solution in the case $\lambda = 0.1$.

参考文献

- [1] D. Furihata and T. Matsuo, Discrete variational derivative method, Chapman & Hall/CRC Numerical Analysis and Scientific Computing, CRC Press, Boca Raton, FL, 2011.
- [2] Finite difference scheme for the Ericksen-Leslie equation, A. Fuwa and T. Ishiwata, Proceedings of Czech Japanese Seminar in Applied Mathematics 2010, in COE Lecture Note, Vol. 36, Faculty of Mathematics, Kyushu University Fukuoka, 2012, ISSN 1881-4042, pp. 26–35.
- [3] Finite difference schemes for Landau-Lifshitz equation, A. Fuwa, T. Ishiwata and M. Tsutsumi, Proceedings of Czech–Japanese Seminar in Applied Mathematics 2006, COE Lecture Note Vol. 6, Kyushu University(2007), pp.107-113.
- [4] Finite difference scheme for the Landau-Lifshitz equation, A. Fuwa, T. Ishiwata and M. Tsutsumi, Japan Journal of Industrial and Applied Mathematics Volume 29, Issue 1 (2012), Page 83-110.
- [5] Local well posedness of the Cauchy problem for the Landau-Lifshitz equations, Atsushi Fuwa and Masayoshi Tsutsumi, Differential Integral Equations **18**(2005), no.4, pp.379-404.
- [6] Structure-preserving finite difference scheme for vortex filament motion, T. Ishiwata and K. Kumazaki, Proceedings of ALGORITMY 2012, 230–238.
- [7] On the theory of dispersion of magnetic permeability in ferromagnetic bodies, Landau D., Lifshitz E. M., Phys. Z. Sowj. **8**(1935), pp.101-114.

- [8] Finite Element Approximations of the Ericksen-Leslie Model for Nematic Liquid Crystal Flow, Roland Becker, Xiaobing Feng and Andreas Prohl, SIAM J. Numer. Anal. **Vol. 46**, Issue 4, pp. 1704-1731 (2008).
- [9] Convergence of an implicit finite element method for the Landau-Lifshitz-Gilbert equation, Sören Bartels and Andreas Prohl, SIAM J. NUMER. ANAL. **Vol. 44**(2006), pp.1405-1419.
- [10] Numerical Methods for the Landau-Lifshitz equation, Weinan E, Xiao-Ping Wang, SIAM J. NUMER. ANAL. **Vol. 38**, No. 5, pp.1647-1665.

Boundedness in degenerate Keller-Segel systems of parabolic-parabolic type

Sachiko Ishida

Department of Mathematics
Tokyo University of Science

1. Introduction

We consider the following fully-parabolic Keller-Segel system on the whole space:

$$(KS)_0 \quad \begin{cases} \frac{\partial u}{\partial t} = \nabla \cdot (\nabla u^m - u^{q-1} \nabla v), & x \in \mathbb{R}^N, \ t > 0, \\ \frac{\partial v}{\partial t} = \Delta v - v + u, & x \in \mathbb{R}^N, \ t > 0, \\ u(x, 0) = u_0(x), \ v(x, 0) = v_0(x), & x \in \mathbb{R}^N, \end{cases}$$

where $N \in \mathbb{N}$, $m \geq 1$, $q \geq 2$ and the initial data u_0, v_0 are non-negative functions which has some regularity. The system with $m = 1$ and $q = 2$ is introduced by Keller and Segel in 1970 ([5]). These types of systems describe the chemitactic process of slime molds. Here $u = u(x, t)$ represents the cell density, $v = v(x, t)$ denotes the chemotaxis concentration at place $x \in \mathbb{R}^N$, time $t > 0$.

In the recent study, Sugiyama-Kunii [6] and Ishida-Yokota [1, 2, 3] consider $(KS)_0$ and reach the results of global solvability. However, they did not have the boundedness of the solution. Indeed, in [6], [3] the upper estimate for $\|u\|_{L^\infty(0,T;L^\infty(\mathbb{R}^N))}$ grows up as $T \rightarrow \infty$ (i.e., $\|u\|_{L^\infty(0,T;L^\infty(\mathbb{R}^N))} \leq C_1 T$). The reason of this growing is the following:

- (I) When $q < m + \frac{2}{N}$, it follows that $\|u\|_{L^\infty(0,T;L^\infty(\mathbb{R}^N))} \leq C(\|u\|_{L^\infty(0,T;L^{r^*}(\mathbb{R}^N))})$ for large $r_* > 1$ and the upper estimate of $\|u\|_{L^\infty(0,T;L^{r^*}(\mathbb{R}^N))}$ is depending on T .
- (II) When $q \geq m + \frac{2}{N}$, since they used the maximal Sobolev regularity of the second equation, it follows that $\|u\|_{L^\infty(0,T;L^\infty(\mathbb{R}^N))} \leq C_1 T$ even if u has the L^p -boundedness ($p \in [1, \infty)$).

Note that the approaches (I) and (II) slightly differ from one another because they use the condition $q < m + \frac{2}{N}$ or $q \geq m + \frac{2}{N}$.

On the other hand, in the bounded domain case, Tao-Winkler [7] give a successful result, that is, the global existence and the global-in-time boundedness of solutions to $(KS)_0$, i.e., $\|u\|_{L^\infty(0,\infty;L^\infty(\Omega))} \leq C_2$ where C_2 is not depending on time variable. They show the L^∞ -boundedness from the L^{r^*} -boundedness of u for large $r_* \geq 1$ without using the condition between m and q . Unfortunately, they assume u solves $(KS)_0$ classically, and moreover, this boundedness is depending on $|\Omega|$.

Our goal is the global-in-time L^∞ -boundedness of the solution to $(KS)_0$ on the whole space without using the condition between m and q . Note that it is difficult to apply Tao-Winkler's results for the following reasons; (1st reason) $(KS)_0$ does not have a classical

solution because of the degenerate diffusion; (2nd reason) their boundedness is depending on $|\Omega|$, while our system is on the whole space ($|\mathbb{R}^N| = \infty$).

For the above first reason, it might well have started with the following non-degenerate problem:

$$(KS)_\delta \quad \begin{cases} \frac{\partial u}{\partial t} = \nabla \cdot (\nabla(u + \delta)^m - (u + \delta^{\frac{m-1}{q-2}})^{q-2} u \nabla v), & x \in \mathbb{R}^N, t > 0, \\ \frac{\partial v}{\partial t} = \Delta v - v + u, & x \in \mathbb{R}^N, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \mathbb{R}^N, \end{cases}$$

where $N \in \mathbb{N}$, $m \geq 1$, $q \geq 2$ and $\delta \in (0, 1)$. The non-negative initial data u_0, v_0 belong to $C_0^\infty(\mathbb{R}^N)$. This type of systems is considered in [6, 1, 2, 3] as the approximated problem.

The main purpose in this work is to give a proof of the global-in-time L^∞ -boundedness of the solution to $(KS)_\delta$ on the whole space without using the condition between m and q under assuming the L^p -boundedness of u .

Section 2 gives the main theorem and its proof which are related to L^∞ -boundedness on $(KS)_\delta$. In Section 3 we state L^∞ -boundedness on $(KS)_0$ when $q \geq m + \frac{2}{N}$.

2. Boundedness in $(KS)_\delta$

We discuss the boundedness of the solution to $(KS)_\delta$. Let us begin with giving the definition of solutions to $(KS)_\delta$.

Definition 2.1. Let $T > 0$. A pair (u, v) of non-negative functions defined on $\mathbb{R}^N \times (0, T)$ is called a *solution* to $(KS)_\delta$ on $[0, T)$ if for $r > N + 1$

- (a) $u \in W^{1,p}(0, T; L^p(\mathbb{R}^N)) \cap L^p(0, T; W^{2,p}(\mathbb{R}^N))$ for $p = \begin{cases} 2, r & N = 1, \\ \frac{N}{N-1}, r & N \geq 2, \end{cases}$
- (b) $v \in W^{1,r}(0, T; L^r(\mathbb{R}^N)) \cap L^r(0, T; W^{2,r}(\mathbb{R}^N))$,
- (c) (u, v) satisfies $(KS)_\delta$ almost everywhere on $\mathbb{R}^N \times [0, T)$.

In particular, if we can take $T = \infty$ and

$$\|u\|_{L^\infty(0, \infty; L^\infty(\mathbb{R}^N))} < \infty, \quad \|v\|_{L^\infty(0, \infty; W^{1, \infty}(\mathbb{R}^N))} < \infty,$$

then (u, v) is called a *global bounded solution* to $(KS)_\delta$.

Now we state the main theorem.

Theorem 2.1. Let $m \geq 1$, $q \geq 2$, $n \in \mathbb{N}$. Assume that the initial data $u_0, v_0 \in C_0^\infty(\mathbb{R}^N)$ are non-negative and that the solution (u, v) satisfies $\|u\|_{L^\infty(0, \infty; L^{p_0}(\mathbb{R}^N))} < M_0$ for some large p_0 , where $M > 0$ is a constant which does not depend on δ . Then the solution is globally bounded:

$$\|u\|_{L^\infty(0, \infty; L^\infty(\mathbb{R}^N))} < M.$$

Remark 2.1. It seems that the L^{p_0} -boundedness is a big assumption. Fortunately, in the case where $q \geq m + \frac{2}{N}$ we have already proven it under the small initial data (see [3]).

Proof of Theorem 2.1. We divide the proof into the following five steps:

Step 1: Standard energy inequality derived from first equation.

Step 2: Energy inequality via the L^p -boundedness of ∇v .

Step 3: Energy inequality without ∇u .

Step 4: Creation of the ODI of $Z_r(t) := \|u(t)\|_{L^r}^r + \|u_0\|_{L^1}^r + 1$.

Step 5: Iteration for $Z_r(t)$ and conclusion of boundedness.

【Step 1: Standard energy inequality】 Multiplying the first equation in $(KS)_\delta$ by u^{r-1} and integrating it over $\mathbb{R}^N$, we see from $a^k + b^k \leq (a+b)^k \leq 2^k(a^k + b^k)$ for $a, b, k > 0$ and the Young inequality that

$$\begin{aligned}
(2.1) \quad & \frac{1}{r} \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r \\
&= - \int_{\mathbb{R}^N} \nabla(u + \delta)^m \cdot \nabla u^{r-1} dx + \int_{\mathbb{R}^N} (u + \delta^{\frac{m-1}{q-2}})^{q-2} u \nabla v \cdot \nabla u^{r-1} dx \\
&\leq - \frac{m(r-1)}{2} \left\{ \int_{\mathbb{R}^N} u^{r+m-3} |\nabla u|^2 dx + \delta^{m-1} \int_{\mathbb{R}^N} u^{r-2} |\nabla u|^2 dx \right\} \\
&\quad + 2^{2(q-2)} \frac{(r-1)}{m} \left\{ \int_{\mathbb{R}^N} u^{r+2q-m-3} |\nabla v|^2 dx + \delta^{m-1} \int_{\mathbb{R}^N} u^r |\nabla v|^2 dx \right\}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
(2.2) \quad & \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r \\
&+ \frac{2mr(r-1)}{(r+m-1)^2} \left\| \nabla u^{\frac{r+m-1}{2}}(t) \right\|_{L^2(\mathbb{R}^N)}^2 + \delta^{m-1} \frac{2m(r-1)}{r} \left\| \nabla u^{\frac{r}{2}}(t) \right\|_{L^2(\mathbb{R}^N)}^2 \\
&\leq 2^{2(q-2)} \frac{r(r-1)}{m} \left\{ \int_{\mathbb{R}^N} u^{r+2q-m-3} |\nabla v|^2 dx + \delta^{m-1} \int_{\mathbb{R}^N} u^r |\nabla v|^2 dx \right\}.
\end{aligned}$$

【Step 2: Energy inequality via the L^p -boundedness of ∇v 】 We estimate the right-hand side of the above estimate, in particular, the first term. First we adjust the parameter γ , θ and s as like in [7, Appendix]. Let take $\gamma \in (2, \infty)$ (if $N = 1, 2$), $\gamma \in (2, \frac{2N}{N-2})$ (if $N \geq 3$) and put $\theta = \frac{\gamma}{2} \cdot \frac{r+m-1}{r-m-1}$, $\theta' = (1 - \frac{1}{\theta})^{-1}$. Take $p_1 \geq 1$ such that

$$(2.3) \quad 2(q-1)p_1 \leq p_0$$

and pick $s \in (1, 2)$. Let start deriving estimates. First, because of $\frac{1}{\theta} + \frac{1}{p_1} + \frac{p_1 - \theta'}{p_1 \theta'} = 1$, we have from the Young inequality that

$$\begin{aligned}
(2.4) \quad & \int_{\mathbb{R}^N} u^{r+2q-m-3} |\nabla v|^2 dx \\
&\leq \left(\int_{\mathbb{R}^N} u^{(r-m-1)\theta} dx \right)^{\frac{1}{\theta}} \left(\int_{\mathbb{R}^N} u^{2(q-1)p_1} dx \right)^{\frac{1}{p_1}} \left(\int_{\mathbb{R}^N} |\nabla v|^{2 \cdot \frac{p_1 \theta'}{p_1 - \theta'}} dx \right)^{\frac{p_1 - \theta'}{p_1 \theta'}}.
\end{aligned}$$

Now, from the fact that u has the mass conservation law $\|u(t)\|_{L^1(\mathbb{R}^N)} = \|u_0\|_{L^1(\mathbb{R}^N)}$ ($\forall t > 0$) and the assumption of L^p -boundedness we know

$$\|u\|_{L^\infty(0,\infty;L^p(\mathbb{R}^N))} \leq K_1 \quad (1 \leq p \leq p_0),$$

so it follows from (2.3) that

$$(2.5) \quad \left(\int_{\mathbb{R}^N} u^{2(q-1)p_1}(t) dx \right)^{\frac{1}{p_1}} \leq K_1^{2(q-1)}.$$

Let put $A := \frac{p_1\theta'}{p_1-\theta'}$. Noting that $\frac{p_1-\theta'}{\theta'p_1} \in (0, 1)$ entails $2 < \frac{2p_1\theta'}{p_1-2\theta'} < \infty$, we have

$$\left(\int_{\mathbb{R}^N} |\nabla v|^{2A} dx \right)^{\frac{1}{A}} \leq \|\nabla v(t)\|_{L^\infty(\mathbb{R}^N)}^{\frac{2(A-1)}{A}} \|\nabla v(t)\|_{L^2(\mathbb{R}^N)}^{\frac{2}{A}}.$$

Since $0 < \frac{2(A-1)}{A}, \frac{2}{A} < 2$, it follows that

$$\left(\int_{\mathbb{R}^N} |\nabla v|^{2 \cdot \frac{p_1\theta'}{p_1-\theta'}} dx \right)^{\frac{p_1-\theta'}{p_1\theta'}} \leq \max\{1, \|\nabla v(t)\|_{L^\infty}^2\} \max\{1, \|\nabla v(t)\|_{L^2}^2\}.$$

Here, the L^p - L^q estimate for the heat semigroup and the condition $p_0 > N$ derive $\|\nabla v(t)\|_{L^r} \leq K_2$ ($r \in [1, \infty]$) (see [7, Lemma 1.2], [4, Lemma 2.3], for instance), and so we see that

$$(2.6) \quad \left(\int_{\mathbb{R}^N} |\nabla v|^{2 \cdot \frac{p_1\theta'}{p_1-\theta'}} dx \right)^{\frac{p_1-\theta'}{p_1\theta'}} \leq (\max\{1, K_2^2\})^2.$$

Next, by the choice of γ , we can use the Gagliardo-Nirenberg inequality and find $c_1 > 0$ such that

$$(2.7) \quad \begin{aligned} \left(\int_{\mathbb{R}^N} u^{(r-m-1)\theta} dx \right)^{\frac{1}{\theta}} &= \|u^{\frac{r+m-1}{2}}(t)\|_{L^\gamma(\mathbb{R}^N)}^{\frac{\gamma}{\theta}} \\ &\leq c_1 \|\nabla u^{\frac{r+m-1}{2}}(t)\|_{L^2(\mathbb{R}^N)}^{\frac{\alpha\gamma}{\theta}} \|u^{\frac{r+m-1}{2}}(t)\|_{L^s(\mathbb{R}^N)}^{\frac{(1-\alpha)\gamma}{\theta}}, \end{aligned}$$

where

$$\alpha = \left(\frac{1}{s} - \frac{1}{\gamma} \right) \left(\frac{1}{N} + \frac{1}{s} - \frac{1}{2} \right)^{-1} \in (0, 1).$$

Hence connecting (2.4)–(2.7) gives

$$(2.8) \quad \int_{\mathbb{R}^N} u^{r+2q-m-3} |\nabla v|^2 dx \leq c_2 \|\nabla u^{\frac{r+m-1}{2}}(t)\|_{L^2(\mathbb{R}^N)}^{\frac{\alpha\gamma}{\theta}} \|u^{\frac{r+m-1}{2}}(t)\|_{L^s(\mathbb{R}^N)}^{\frac{(1-\alpha)\gamma}{\theta}},$$

where $c_2 = c_1 K_1^{2(q-1)} \max\{1, K_2^4\}$. Thanks to

$$\frac{\alpha\gamma}{\theta} = \alpha\gamma \cdot \frac{2(r-m-1)}{\gamma(r+m-1)} < 2\alpha < 2$$

because of $m \geq 1$, the Young inequality gives

$$\begin{aligned}
(2.9) \quad & 2^{2(q-2)} \frac{r(r-1)}{m} \int_{\mathbb{R}^N} u^{r+2q-m-3} |\nabla v|^2 dx \\
& \leq \frac{mr(r-1)}{(r+m-1)^2} \left\| \nabla u^{\frac{r+m-1}{2}}(t) \right\|_{L^2(\mathbb{R}^N)}^2 \\
& \quad + r(r-1) \left(\frac{2^{2(q-2)} c_2}{m} \left(\frac{m}{(r+m-1)^2} \right)^{\frac{-\alpha\gamma}{2\theta}} \right)^{\frac{2\theta}{2\theta-\alpha\gamma}} \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2(1-\alpha)\gamma}{s(2\theta-\alpha\gamma)}}.
\end{aligned}$$

From the same argument as above we infer that

$$\begin{aligned}
(2.10) \quad & \frac{r(r-1)}{m} \int_{\mathbb{R}^N} u^r |\nabla v|^2 dx \\
& \leq \frac{m(r-1)}{r} \delta^{m-1} \left\| \nabla u^{\frac{r}{2}} \right\|_{L^2}^2 \\
& \quad + r(r-1) \delta^{m-1} \left(\frac{2^{2q-3} c_3}{m} \left(\frac{m}{r^2} \right)^{-\frac{\alpha\gamma}{2\eta}} \right)^{\frac{2\eta}{2\eta-\alpha\gamma}} \left(\int_{\mathbb{R}^N} u^{\frac{sr}{2}} dx \right)^{\frac{2\gamma(1-\alpha)}{s(2\eta-\alpha\gamma)}},
\end{aligned}$$

where $\eta = \frac{\gamma r}{2(r-2)}$, $c_3 = c_3(K_1, K_2, N) > 0$ is a constant. For large r we find $c_4, c_5 > 0$ which are independent of r such that

$$\left(\frac{2^{2(q-2)} c_2}{m} \left(\frac{m}{(r+m-1)^2} \right)^{\frac{-\alpha\gamma}{2\theta}} \right)^{\frac{2\theta}{2\theta-\alpha\gamma}} \leq c_4, \quad \left(\frac{2^{2q-3} c_3}{m} \left(\frac{m}{r^2} \right)^{-\frac{\alpha\gamma}{2\eta}} \right)^{\frac{2\eta}{2\eta-\alpha\gamma}} \leq c_5$$

because $\theta \rightarrow \frac{\gamma}{2}$ and then from (2.2), (2.9), (2.10) we have

$$\begin{aligned}
(2.11) \quad & \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r + \frac{mr(r-1)}{(r+m-1)^2} \left\| \nabla u^{\frac{r+m-1}{2}} \right\|_{L^2(\mathbb{R}^N)}^2 \\
& \leq r(r-1) c_4 \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2(1-\alpha)\gamma}{s(2\theta-\alpha\gamma)}} + r(r-1) \delta^{m-1} c_5 \left(\int_{\mathbb{R}^N} u^{\frac{sr}{2}} dx \right)^{\frac{2\gamma(1-\alpha)}{s(2\eta-\alpha\gamma)}}.
\end{aligned}$$

[Step 3: Energy inequality without ∇u] Next we consider the second and third terms on the left-hand side of (2.11). From the Gagliardo-Nirenberg inequality and the Young inequality we have

$$\int_{\mathbb{R}^N} u^r dx \leq \left(\left\| \nabla u^{\frac{r+m-1}{2}}(t) \right\|_{L^2(\mathbb{R}^N)} + \left\| u^{\frac{r+m-1}{2}}(t) \right\|_{L^s(\mathbb{R}^N)} \right)^{\frac{2r}{r+m-1}}$$

hence we see that

$$(2.12) \quad \frac{1}{2} \left(\int_{\mathbb{R}^N} u^r dx \right)^{\frac{r+m-1}{r}} - \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2}{s}} \leq \left\| \nabla u^{\frac{r+m-1}{2}}(t) \right\|_{L^2(\mathbb{R}^N)}^2.$$

Plugging (2.12) into (2.11) yields

$$\begin{aligned}
& \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r + \frac{mr(r-1)}{(r+m-1)^2} \left(\frac{1}{2} \left(\int_{\mathbb{R}^N} u^r dx \right)^{\frac{r+m-1}{r}} - \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2}{s}} \right) \\
& \leq r(r-1) c_4 \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2(1-\alpha)\gamma}{s(2\theta-\alpha\gamma)}} + r(r-1) \delta^{m-1} c_5 \left(\int_{\mathbb{R}^N} u^{\frac{sr}{2}} dx \right)^{\frac{2\gamma(1-\alpha)}{s(2\eta-\alpha\gamma)}},
\end{aligned}$$

then we have

$$\begin{aligned}
& \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r + \frac{mr(r-1)}{2(r+m-1)^2} \left(\int_{\mathbb{R}^N} u^r dx \right)^{\frac{r+m-1}{r}} \\
& \leq r(r-1)c_4 \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2(1-\alpha)\gamma}{s(2\theta-\alpha\gamma)}} + \frac{mr(r-1)}{(r+m-1)^2} \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2}{s}} \\
& \quad + r(r-1)\delta^{m-1}c_5 \left(\int_{\mathbb{R}^N} u^{\frac{sr}{2}} dx \right)^{\frac{2\gamma(1-\alpha)}{s(2\eta-\alpha\gamma)}}.
\end{aligned}$$

To simplify this estimate let take $r > r_0$ with r_0 satisfies $\frac{1}{2} \leq \frac{r_0-1}{r_0} \leq 1$, $\frac{1}{2} \leq \frac{r_0(r_0-1)}{(r_0+m-1)^2} \leq 1$. Since

$$\frac{2(1-\alpha)\gamma}{s(2\theta-\alpha\gamma)} \leq \frac{2}{s}, \quad \frac{2(1-\beta)\gamma}{s(2\eta-\beta\gamma)} \leq \frac{2}{s}$$

from $\gamma > 2$ and $\theta, \eta \geq \frac{\gamma}{2}$ and $\delta^{m-1} < 1$, it follows from the Young inequality that

$$\begin{aligned}
(2.13) \quad & \frac{d}{dt} \|u(t)\|_{L^r(\mathbb{R}^N)}^r + \frac{m}{4} \left(\int_{\mathbb{R}^N} u^r dx \right)^{\frac{r+m-1}{r}} \\
& \leq r^2 c_6 \left\{ 1 + \left(\int_{\mathbb{R}^N} u^{\frac{s(r+m-1)}{2}} dx \right)^{\frac{2}{s}} + \left(\int_{\mathbb{R}^N} u^{\frac{sr}{2}} dx \right)^{\frac{2}{s}} \right\},
\end{aligned}$$

where $c_6 = \max\{c_4 + c_5, c_4 + m, c_5\} > 0$ is a constant.

【Step 4: Creation of the ODI of $Z_r(t)$ 】 Here we define

$$Y_r(t) := \int_{\mathbb{R}^N} u^r dx.$$

Then (2.13) is rewritten as

$$\frac{d}{dt} Y_r(t) + \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} \leq c_6 r^2 \left\{ 1 + (Y_{\frac{s(r+m-1)}{2}}(t))^{\frac{2}{s}} + (Y_{\frac{sr}{2}}(t))^{\frac{2}{s}} \right\}.$$

Let us take r such that $1 < \frac{sr}{2}$. Since

$$\begin{aligned}
(Y_{\frac{sr}{2}}(t))^{\frac{2}{s}} &= \|u\|_{L^{\frac{sr}{2}}}^r \leq \|u_0\|_{L^1}^r + \|u(t)\|_{L^{\frac{s(r+m-1)}{2}}}^r \\
&= \|u_0\|_{L^1}^r + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s} \cdot \frac{r}{(r+m-1)}} \\
&\leq \|u_0\|_{L^1}^r + 1 + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}},
\end{aligned}$$

it follows that

$$\begin{aligned}
(2.14) \quad & \frac{d}{dt} Y_r(t) + \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} \\
& \leq c_6 r^2 \left\{ 1 + (Y_{\frac{s(r+m-1)}{2}}(t))^{\frac{2}{s}} + \left(\|u_0\|_{L^1}^r + 1 + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right)^{\frac{2}{s}} \right\} \\
& \leq c_6 r^2 \left(2 + \|u_0\|_{L^1}^r + 2 \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right).
\end{aligned}$$

Next, (2.14) added $\|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1$ gives

$$(2.15) \quad \begin{aligned} \frac{d}{dt} Y_r(t) + \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1 \\ \leq c_6 r^2 \left(2 + \|u_0\|_{L^1}^r + 2 \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right) + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1. \end{aligned}$$

Now we create the unit term of iteration. First we clearly have

$$(2.16) \quad \frac{d}{dt} Y_r(t) = \frac{d}{dt} \left\{ Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1 \right\}.$$

Next we have for $r > m + 1$,

$$(2.17) \quad \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1 \geq c_7 \left\{ Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1 \right\}^{\frac{r+m-1}{r}},$$

where $c_7 = \frac{1}{2} \min\{\frac{m}{4}, 1\}$ because

$$\begin{aligned} c_7 \left\{ Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1 \right\}^{\frac{r+m-1}{r}} &\leq c_7 2^{\frac{r+m-1}{r}-1} \left\{ Y_r(t)^{\frac{r+m-1}{r}} + \|u_0\|_{L^1(\mathbb{R}^N)}^{r \cdot \frac{r+m-1}{r}} + 1^{\frac{r+m-1}{r}} \right\} \\ &\leq \frac{1}{2} \cdot 2^{\frac{m-1}{r}} \left\{ \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1 \right\} \\ &\leq \frac{m}{4} (Y_r(t))^{\frac{r+m-1}{r}} + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1. \end{aligned}$$

Finally it follows that

$$(2.18) \quad \begin{aligned} c_6 r^2 \left(2 + \|u_0\|_{L^1}^r + 2 \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right) + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1 \\ \leq c_6 r^2 \left(2 + 1 + \|u_0\|_{L^1}^{r+m-1} + 2 \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right) + \|u_0\|_{L^1(\mathbb{R}^N)}^{r+m-1} + 1 \\ \leq 4c_6 r^2 \left(1 + \|u_0\|_{L^1}^{r+m-1} + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right) \\ \leq 4c_6 r^2 \left(1 + \|u_0\|_{L^1}^{r+m-1 \cdot \frac{s}{2}} + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right). \end{aligned}$$

Hence (2.15) combined with (2.16)–(2.18) gives that

$$(2.19) \quad \begin{aligned} \frac{d}{dt} \left\{ Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1 \right\} + c_7 \left\{ Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1 \right\}^{\frac{r+m-1}{r}} \\ \leq 4c_6 r^2 \left(1 + \|u_0\|_{L^1}^{r+m-1 \cdot \frac{s}{2}} + \left(Y_{\frac{s(r+m-1)}{2}} \right)^{\frac{2}{s}} \right). \end{aligned}$$

Let $r = p_k$ where p_k is given by

$$p_k := \frac{2}{s} p_{k-1} - m + 1 \text{ with } p_0 = \max\{N, r_0, r_1, \frac{2}{s}, m + 1\}.$$

Note that $p_{k-1} = \frac{s(p_k+m-1)}{2}$, $p_{k-1} < p_k$ when $p_0 > \frac{s(m-1)}{2-s}$ and that

$$p_k = \left(\frac{2}{s}\right)^k p_0 + \left(\left(\frac{2}{s}\right)^{k-1} + \cdots + \frac{2}{s} + 1\right)(-m + 1)$$

implies

$$p_0 \left(\frac{2}{s}\right)^k < p_k < \left(p_0 + \frac{s}{2-s}\right) \left(\frac{2}{s}\right)^k.$$

Let us define

$$Z_r(t) = Y_r(t) + \|u_0\|_{L^1(\mathbb{R}^N)}^r + 1.$$

Then we have

$$\frac{d}{dt} Z_{p_k}(t) + c_7 (Z_{p_k}(t))^{\frac{p_k+m-1}{p_k}} \leq c_8 \left(\frac{2}{s}\right)^k \sup_{t>0} (Z_{p_{k-1}}(t))^{\frac{2}{s}},$$

where $c_8 = 4c_6(p_0 + \frac{s}{2-s})$.

【Step 5: Iteration and conclusion】 From the Gronwall type inequality, the above ODI gives

$$(2.20) \quad Z_{p_k}(t) \leq \max \left\{ Z_{p_k}(0), \left[\frac{c_{10}}{c_9} \left(\frac{2}{s}\right)^k \sup_{t>0} (Z_{p_{k-1}}(t))^{\frac{2}{s}} \right]^{\frac{p_k}{p_k+m-1}} \right\}.$$

Noting that

$$Z_{p_k}(0)^{\frac{1}{p_k}} = \left(\int_{\mathbb{R}^N} u_0^{p_k} dx + \|u_0\|_{L^1(\mathbb{R}^N)}^{p_k} + 1 \right)^{\frac{1}{p_k}} \rightarrow \|u_0\|_{L^\infty(\mathbb{R}^N)} + \|u_0\|_{L^1(\mathbb{R}^N)} + 1 =: \mathcal{R}$$

as $k \rightarrow \infty$ we find that it is enough to consider the following two cases by taking a sub-sequence of $\{k\}$ if necessary:

$$(i): \mathcal{R} < \left[\frac{c_{10}}{c_9} \left(\frac{2}{s}\right)^k \sup_{t>0} (Z_{p_{k-1}}(t))^{\frac{2}{s}} \right]^{\frac{1}{p_k+m-1}} \text{ for any } k \gg 1.$$

In this case (2.20) gives

$$\sup_{t>0} Z_{p_k}(t) \leq \left[C \left(\frac{2}{s}\right)^k \sup_{t>0} (Z_{p_{k-1}}(t))^{\frac{2}{s}} \right]^{\frac{p_k}{p_k+m-1}},$$

where $C = \max\{1, \frac{c_{10}}{c_9}\}$. Because $C, \frac{2}{s} > 1$ and $Z_k > 1$, $\frac{p_k}{p_k+m-1} < 1$ for any $k \geq 1$ it follows that

$$\sup_{t>0} Z_{p_k}(t) \leq C \left(\frac{2}{s}\right)^k \sup_{t>0} (Z_{p_{k-1}}(t))^{\frac{2}{s}}.$$

By Moser's iteration technique, we have

$$\sup_{t>0} Z_{p_k}(t) \leq C^{1+\frac{2}{s}+(\frac{2}{s})^2+\dots+(\frac{2}{s})^{k-2}} \left(\frac{2}{s}\right)^{1 \cdot k + \frac{2}{s} \cdot (k-1) + (\frac{2}{s})^2 \cdot (k-2) + \dots + (\frac{2}{s})^{k-1} \cdot 1} \sup_{t>0} (Z_{p_1}(t))^{\left(\frac{2}{s}\right)^k}.$$

Noting that $(p_0 + \frac{s}{2-s})(\frac{s}{2})^k \leq \frac{1}{p_k} \leq p_0^{-1}(\frac{s}{2})^k$, we have

$$\lim_{k \rightarrow \infty} \frac{1}{p_k} \sum_{n=1}^{k-1} \left(\frac{2}{s}\right)^{n-1} \leq \frac{1}{p_0} \lim_{k \rightarrow \infty} \sum_{n=1}^{k-1} \left(\frac{s}{2}\right)^{n+1} = \frac{s}{2(2-s)p_0},$$

$$\lim_{k \rightarrow \infty} \frac{1}{p_k} \left\{ 1 \cdot k + \frac{2}{s} \cdot (k-1) + \left(\frac{2}{s}\right)^2 \cdot (k-2) + \dots + \left(\frac{2}{s}\right)^{k-1} \cdot 1 \right\} = \frac{2s}{(2-s)^2 p_0},$$

and

$$\lim_{k \rightarrow \infty} \frac{1}{p_k} \left(\frac{2}{s} \right)^{k-1} \leq \frac{s}{2p_0},$$

then we obtain

$$\|u(t)\|_{L^\infty(\mathbb{R}^N)} = \lim_{k \rightarrow \infty} \sup_{t > 0} Z_{p_k}(t)^{\frac{1}{p_k}} - \|u_0\|_{L^1(\mathbb{R}^N)} - 1 \leq M,$$

where $M > 0$ is a constant which depends on $\sup_{t > 0} \|u(t)\|_{L^{p_0}(\mathbb{R}^N)}$, $\|u_0\|_{L^1(\mathbb{R}^N)}$, m , N but not on time variable.

$$(ii): \left[\frac{c_{10}}{c_9} \left(\frac{2}{s} \right)^k \sup_{t > 0} (Z_{p_{k-1}}(t))^{\frac{2}{s}} \right]^{\frac{1}{p_k + m - 1}} < \mathcal{R} \text{ for any } k \gg 1.$$

This case implies

$$\sup_{t > 0} Z_{p_k}^{\frac{1}{p_k}}(t) \leq Z_{p_k}^{\frac{1}{p_k}}(0)$$

and we obtain

$$\sup_{t > 0} \|u(t)\|_{L^\infty(\mathbb{R}^N)} \leq \|u_0\|_{L^\infty(\mathbb{R}^N)}.$$

The L^∞ -boundedness of u can be obtained from the above two cases, i.e., there exist a positive constant M which is not depending on time such that

$$\|u(t)\|_{L^\infty} \leq M.$$

Moreover, the $W^{1,\infty}$ -boundedness of v is given by L^p - L^q estimate of heat semigroup. $\square$

3. Boundedness in $(KS)_0$ in the super-critical case

In this section we state the L^∞ -boundedness in $(KS)_0$. As mentioned in Remark 2.1, we already had the L^p -boundedness on $(KS)_\delta$ as the approximate problem in super-critical case (i.e., $q \geq m + \frac{2}{N}$). Hence, this boundedness and Theorem 2.1 drive the global-in-time boundedness to the degenerate system $(KS)_0$:

Theorem 3.1. *Assume that m and q satisfy*

$$q \geq m + \frac{2}{N}.$$

Assume that the initial data satisfies

$$u_0, v_0 \geq 0, \quad u_0, v_0 \in L^1 \cap L^\infty(\mathbb{R}^N), \\ v_0 \in B_{r+q-1, r+q-1}^{2-\frac{2}{r+q-1}} \cap B_{r+1, r+1}^{2-\frac{2}{r+1}}(\mathbb{R}^N), \quad \Delta v_0 \in L^{q_c+q-1} \cap L^{q_0+q-1} \cap L^\infty(\mathbb{R}^N)$$

with some $r > 1$ and $q_c := \frac{N}{2}(q - m)$, $q_0 := \frac{N}{2}$ and satisfies following smallness:

$$\|u_0\|_{L^{q_c}} \leq \delta_u, \quad \|v_0\|_{B_{q_c+q-1, q_c+q-1}^{2-2/(q_c+q-1)}} \leq \delta_v \text{ when } q \geq m + 1 \text{ } (N \geq 3) \text{ or } N = 2, \\ \|u_0\|_{L^{q_0}} \leq \delta_u, \quad \|v_0\|_{B_{q_0+q-1, q_0+q-1}^{2-2/(q_0+q-1)}} \leq \delta_v \text{ when } q < m + 1 \text{ } (N \geq 3),$$

where $\delta_u = \delta_u(m, q, N)$, $\delta_v = \delta_v(m, q, N)$ are positive constants. Then $(\text{KS})_0$ has a non-negative global-in-time solution (u, v) such that

$$\begin{aligned} u &\in L^\infty(0, \infty; L^p(\mathbb{R}^N)) \quad (\forall p \in [1, \infty]), \quad u^m \in L^2(0, T; H^1(\mathbb{R}^N)) \quad (\forall T > 0), \\ v &\in L^\infty(0, \infty; H^1(\mathbb{R}^N)), \\ (u, v) &\text{ satisfies } (\text{KS})_0 \text{ in the sense of distributions} \end{aligned}$$

with

$$\|u\|_{L^\infty(0, \infty; L^p(\mathbb{R}^N))} \leq M,$$

where $M > 0$ is independent of time variable.

Future work: In the present work we derived the globally L^∞ -boundedness on $(\text{KS})_\delta$ without using the condition of m and q (Theorem 2.1), and moreover, can obtain it on $(\text{KS})_0$ only in the case where $q \geq m + \frac{2}{N}$ because [3] allows the L^p -boundedness of u in this case (Theorem 3.1). So, our next interest is suitable for finding L^p -boundedness of u with the condition $q < m + \frac{2}{N}$.

References

- [1] S. Ishida, T. Yokota, *Global existence of weak solutions to quasilinear degenerate Keller-Segel systems of parabolic-parabolic type*, J. Differential Equations **252** (2012), 1421–1440.
- [2] S. Ishida, T. Yokota, *Global existence of weak solutions to quasilinear degenerate Keller-Segel systems of parabolic-parabolic type with small data*, J. Differential Equations **252** (2012), 2469–2491.
- [3] S. Ishida, T. Yokota, *Remarks on the global existence of weak solutions to quasilinear degenerate Keller-Segel systems*, AIMS Proceedings, to appear.
- [4] S. Ishida, T. Yokota, *Blow-up in finite or infinite time for quasilinear degenerate Keller-Segel systems of parabolic-parabolic type*, preprint.
- [5] E. F. Keller, L. A. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol. **26** (1970), 399–415.
- [6] Y. Sugiyama, H. Kunii, *Global existence and decay properties for a degenerate Keller-Segel model with a power factor in drift term*, J. Differential Equations **227** (2006), 333–364.
- [7] Y. Tao, M. Winkler, *Boundedness in a quasilinear parabolic-parabolic Keller-Segel system with subcritical sensitivity*, J. Differential Equations **252** (2012), 692–715.

An error estimate of conservative finite difference scheme for the Boussinesq type equations

Shuji Yoshikawa*

1 Introduction

We consider the Boussinesq type equation

$$\partial_t^2 u + \gamma \partial_x^4 u - \alpha_1 \partial_x^2 u = \partial_x^2 f(u), \quad (x, t) \in (0, L) \times (0, T), \quad (1.1)$$

with the periodic boundary condition $u(x, t) = u(x + L, t)$. We assume that constants γ and α_1 are positive and that f is a polynomial and a *definite energy type* such that for some constant C and a primitive F of f

$$F(u) \geq -C \quad (u \in \mathbb{R}). \quad (1.2)$$

Examples of f are given by

$$f(u) = \alpha_\rho u^\rho \quad (\rho \text{ is a odd number, } \alpha_\rho > 0), \quad (1.3)$$

$$f(u) = -\alpha_3 u^3 + \alpha_5 u^5 \quad (\alpha_3, \alpha_5 > 0) \quad (1.4)$$

etc. The equation (1.1) with (1.4) was derived by Falk et al. as the model representing the phase transition on shape memory alloys (we refer to [2]). Originally, the equation with the opposite sign in the fourth derivative $\gamma < 0$ was derived by Boussinesq [1] as the model of water waves. In [4] Furihata proposed the discrete variational derivative method (DVDM) which is the method to derive the structure preserving finite difference scheme for evolution equations. More concretely, the finite difference scheme derived by DVDM preserves the property such as energy conservation law or dissipative law. In [4] the conservative finite difference scheme for the Cahn-Hilliard equation was derived by DVDM and the error estimate between an exact solution and a solution of the scheme was given. For more precise informations and recent progress of DVDM, we refer to the monograph by Furihata and Matsuo [5]. In [7] Matsuo proposed several numerical schemes for the good Boussinesq equation by DVDM, on the other hand, an error estimate has not been gave in [7].

In this study we give an error estimate between the solution for (1.1) and the solution for one of numerical schemes derived in [7]. Moreover our motivation of this study is to give the error estimate as general as possible in order to apply to other equations. This study is a joint work with Kyosuke Ichikawa based on [6]

2 Preliminary

We denote by ∂_t and ∂_x partial differential operators with respect to variables t and x , respectively. We split space interval $[0, L]$ into K -th parts and time interval $[0, T]$ into N -th parts, and hence we see that $L = K\Delta x$ and $T = N\Delta t$. For $k = 0, 1, \dots, K$ and $n = 0, 1, \dots, N$ we write $u_k^{(n)} = u(k\Delta x, n\Delta t)$ for the solution u of the original equation. Let $U_k^{(n)}$ an approximate solution for difference scheme corresponding $u_k^{(n)}$.

*Department of Engineering for Production and Environment, Graduate School of Science and Engineering, Ehime University, E-mail: yoshikawa@ehime-u.ac.jp

We introduce several definitions of difference operators which are the almost same as the notation in Furihata-Matsuo [5]. We define several difference operators δ^+ , δ^- , $\delta^{(1)}$, $\delta^{(2)}$ as an approximation to the differential operators by

$$\begin{aligned}\delta_k^+ u_k^{(n)} &:= \frac{u_{k+1}^{(n)} - u_k^{(n)}}{\Delta x}, & \delta_k^- u_k^{(n)} &:= \frac{u_k^{(n)} - u_{k-1}^{(n)}}{\Delta x}, \\ \delta_k^{(1)} u_k^{(n)} &:= \frac{u_{k+1}^{(n)} - u_{k-1}^{(n)}}{2\Delta x}, & \delta_k^{(2)} u_k^{(n)} &:= \frac{u_{k+1}^{(n)} - 2u_k^{(n)} + u_{k-1}^{(n)}}{\Delta x^2}.\end{aligned}$$

We approximate an integral by the trapezoid formula

$$\sum_{k=0}^K {}''u_k \Delta x := \left(\frac{1}{2} u_0 + \sum_{k=1}^{K-1} u_k + \frac{1}{2} u_K \right) \Delta x.$$

Shift operators s and mean value operators μ are defined by

$$\begin{aligned}s_k^+ u_k^{(n)} &:= u_{k+1}^{(n)}, & s_k^- u_k^{(n)} &:= u_{k-1}^{(n)}, & s_k^{(1)} &:= \frac{s_k^+ + s_k^-}{2}, \\ \mu_k^+ &:= \frac{s_k^+ + 1}{2}, & \mu_k^- &:= \frac{1 + s_k^-}{2}, & \mu_k^{(1)} &:= \frac{\mu_k^+ + \mu_k^-}{2} = \frac{s_k^+ + 2 + s_k^-}{4}.\end{aligned}$$

In the same way as the above definitions, we also define difference, shift and mean value operators with respect to time variable by $\delta_n^\pm$, $\delta_n^{(m)}$, $s_n^\pm$, $s_n^{(1)}$, $\mu_n^\pm$ and $\mu_n^{(1)}$.

From the direct calculation, the formulae corresponding to the fundamental theorem of differentiation-integral hold

$$\sum_{k=0}^K {}''\delta_k^{(1)} u_k^{(n)} \Delta x = \left[\mu_k^{(1)} u_k^{(n)} \right]_{k=0}^K, \quad \sum_{k=0}^K {}''\delta_k^{(2)} u_k^{(n)} \Delta x = \left[\delta_k^{(1)} u_k^{(n)} \right]_{k=0}^K. \quad (2.1)$$

Moreover, we also obtain

$$\begin{aligned}\sum_{k=0}^K {}''\left(\delta_k^{(2)} u_k^{(n)} \right) \tilde{u}_k^{(n)} \Delta x &+ \sum_{k=0}^K {}''\frac{\left(\delta_k^+ u_k^{(n)} \right) \left(\delta_k^+ \tilde{u}_k^{(n)} \right) + \left(\delta_k^- u_k^{(n)} \right) \left(\delta_k^- \tilde{u}_k^{(n)} \right)}{2} \Delta x \\ &= \left[\frac{\left(\delta_k^+ u_k^{(n)} \right) \left(\mu_k^+ \tilde{u}_k^{(n)} \right) + \left(\delta_k^- u_k^{(n)} \right) \left(\mu_k^- \tilde{u}_k^{(n)} \right)}{2} \right]_{k=0}^K,\end{aligned} \quad (2.2)$$

which corresponds to the integration by part

$$\int_0^L u \partial_x^2 v \, dx + \int_0^L \partial_x u \partial_x v \, dx = [u \partial_x v]_{x=0}^L.$$

For simplicity, we denote by $\phi^{(\rho)}$ the following ρ -th homogeneous polynomial:

$$\phi^{(\rho)}(u, v) := \sum_{j=0}^{\rho} u^{\rho-j} v^j \quad (\rho \in \mathbb{N} \cup \{0\}), \quad (2.3)$$

which appears naturally through the procedure of the calculation for the discrete variational derivative.

3 Difference Scheme

In this section we introduce the derivation of the difference scheme for the Boussinesq type equation (1.1) applying DVDM to the equation. For simplicity we only treat the equation with simple nonlinearity (1.3). The Hamiltonian form of the equation (1.1) with (1.3) is rewritten as

$$\begin{cases} \partial_t u = \partial_x^2 v, \\ \partial_t v = -\gamma \partial_x^2 u + \alpha_1 u + \alpha_\rho u^\rho. \end{cases} \quad (3.1)$$

The energy of this system is given as

$$G(u, v) = \frac{1}{2} (\partial_x v)^2 + \frac{1}{2} \gamma (\partial_x u)^2 + \frac{1}{2} \alpha_1 u^2 + \frac{1}{\rho+1} \alpha_\rho u^{\rho+1}. \quad (3.2)$$

By using the energy G we may rewrite the equation (3.1) to

$$\partial_t \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \frac{\delta G}{\delta u} \\ \frac{\delta G}{\delta v} \end{pmatrix}. \quad (3.3)$$

To introduce the discrete variational derivative, we give more precise explanation of the above. The total energy I is defined by

$$I(u, v) = \int_0^L G(u, v) dx. \quad (3.4)$$

Since from the direct calculation

$$\begin{aligned} I(u + \delta u, v) - I(u, v) &= \int_0^L (-\gamma \partial_x^2 u + \alpha_1 u + \alpha_\rho u^\rho) \delta u dx, \\ I(u, v + \delta v) - I(u, v) &= - \int_0^L \partial_x^2 v \delta v dx, \end{aligned}$$

the variational derivatives $\frac{\delta G}{\delta u}$ and $\frac{\delta G}{\delta v}$ with respect to u and v are given as

$$\frac{\delta G}{\delta u} = -\gamma \partial_x^2 u + \alpha_1 u + \alpha_\rho u^\rho, \quad \frac{\delta G}{\delta v} = -\partial_x^2 v.$$

From now on, we start to derive the difference scheme approximating the problem (3.3). Corresponding to the definition (3.2), we define the discrete energy G_d by

$$G_d(U_k, V_k) := \frac{1}{2} \frac{(\delta_k^+ V_k)^2 + (\delta_k^- V_k)^2}{2} + \frac{1}{2} \gamma \frac{(\delta_k^+ U_k)^2 + (\delta_k^- U_k)^2}{2} + \frac{1}{2} \alpha_1 U_k^2 + \frac{1}{\rho+1} \alpha_\rho U_k^{\rho+1}. \quad (3.5)$$

We remark that for the discretizing the energy we need to be able to apply the summation by part (2.2). Corresponding to the periodic boundary condition we assume that

$$U_k^{(n)} = U_{k \bmod K}^{(n)}, \quad V_k^{(n)} = V_{k \bmod K}^{(n)}. \quad (3.6)$$

We define the discrete total energy I_d corresponding (3.4) by

$$I_d(\mathbf{U}, \mathbf{V}) := \sum_{k=0}^K G_d(U_k, V_k) \Delta x.$$

An similar calculation to (3.2) yields

$$I_d(\mathbf{U}, \tilde{\mathbf{V}}) - I_d(\mathbf{U}, \mathbf{V}) = -\frac{1}{2} \sum_{k=0}^K {}'' \left(\tilde{V}_k - V_k \right) \left\{ \delta_k^{(2)} \phi^{(1)}(\tilde{V}_k, V_k) \right\} \Delta x,$$

due to (2.2). From this we define *discrete variational derivative* with respect to $\mathbf{V}$ by

$$\left(\frac{\delta G_d}{\delta(\tilde{\mathbf{V}}; \mathbf{V})} \right)_k := -\frac{1}{2} \delta_k^{(2)} \phi^{(1)}(\tilde{V}_k, V_k). \quad (3.7)$$

Similarly we define the *discrete variational derivative* with respect to $\mathbf{U}$ by

$$\left(\frac{\delta G_d}{\delta(\tilde{\mathbf{U}}; \mathbf{U})} \right)_k := -\frac{1}{2} \gamma \delta_k^{(2)} \phi^{(1)}(\tilde{U}_k, U_k) + \frac{1}{2} \alpha_1 \phi^{(1)}(\tilde{U}_k, U_k) + \frac{1}{\rho + 1} \alpha_\rho \phi^{(\rho)}(\tilde{U}_k, U_k). \quad (3.8)$$

Corresponding to (3.3), the difference scheme of DVDM for (1.1) is given by

$$\delta_n^+ \begin{pmatrix} U_k^{(n)} \\ V_k^{(n)} \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \left(\frac{\delta G_d}{\delta(\mathbf{U}^{(n+1)}; \mathbf{U}^{(n)})} \right)_k \\ \left(\frac{\delta G_d}{\delta(\mathbf{V}^{(n+1)}; \mathbf{V}^{(n)})} \right)_k \end{pmatrix}, \quad k = 1, 2, \dots, K. \quad (3.9)$$

Our scheme is (3.8)-(3.9) with the boundary condition (3.6).

Since the difference scheme (3.9) is nonlinear and implicit, it is not trivial whether the solution for (3.9) exists or not. We can show the existence of the solution for (3.9) by using the fixed point principle under some assumptions for Δx and Δt . If we take

$$\max \left\{ \frac{\Delta t}{\Delta x^2}, \frac{\Delta t}{\Delta x^{\frac{\rho-1}{2}}} \right\} \quad (3.10)$$

sufficiently small, then these assumptions are satisfied.

4 Stability and Error Estimate

The equation (3.3) has the following conservation laws:

$$\frac{d}{dt} I(u(t), v(t)) = 0, \quad \frac{d}{dt} M(u(t)) = 0,$$

where the momentum $M(u(t))$ is defined by

$$M(u(t)) := \int_0^L u(x, t) dx.$$

Correspondingly, we can easily show

$$I_d(\mathbf{U}^{(n)}, \mathbf{V}^{(n)}) = I_d(\mathbf{U}^{(0)}, \mathbf{V}^{(0)}),$$

$$M_d(\mathbf{U}^{(n)}) = M_d(\mathbf{U}^{(0)}) \quad \text{for} \quad M_d(\mathbf{U}^{(n)}) := \sum_{k=0}^K {}'' U_k^{(n)} \Delta x.$$

By using these conservation laws we easily obtain the stability result.

Theorem 4.1 (Stability). *The solution for the difference equation (3.9) satisfies*

$$\max_{0 \leq k \leq K} |U_k^{(n)}| \leq \left| M_d(\mathbf{U}^{(0)}) \right| + \sqrt{\frac{2L}{\gamma} I_d(\mathbf{U}^{(0)}, \mathbf{V}^{(0)})}.$$

We define error terms $e_k^{(n)}, \tilde{e}_k^{(n)}$ between exact solutions $u_k^{(n)}, v_k^{(n)}$ and approximate solutions $U_k^{(n)}, V_k^{(n)}$ by

$$e_k^{(n)} := U_k^{(n)} - u_k^{(n)}, \quad \tilde{e}_k^{(n)} := V_k^{(n)} - v_k^{(n)}.$$

We shall estimate the error terms for the difference scheme (3.9).

Theorem 4.2 (Error Estimate). *We set*

$$C_1 := \max_{0 \leq \ell \leq N} \left\{ \max_{0 \leq k \leq K} |U_k^{(\ell)}|, \max_{0 \leq k \leq K} |u_k^{(\ell)}| \right\} \quad (4.1)$$

Assume that the smooth solution for (3.3) satisfying $u \in \mathcal{C}^4([0, L] \times [0, T])$ exists for the initial data (u_0, v_0) and that $e_k^{(0)} = 0$ and $\tilde{e}_k^{(0)} = 0$. If the time-step Δt is small satisfying

$$\Delta t < \frac{4\sqrt{\gamma}}{\alpha_1 + \alpha_\rho \rho C_1^{\rho-1}}, \quad (4.2)$$

then the solution $(U_k^{(n)}, V_k^{(n)})$ for (3.9) converges to (u, v) in the sense that

$$\|e\|_{L_d^2} + \|\tilde{e}\|_{L_d^2} = O(\Delta x^2 + \Delta t^2), \quad \text{as } \Delta x, \Delta t \rightarrow 0,$$

where

$$\|e\|_{L_d^2} := \left(\sum_{k=0}^K |e_k|^2 \Delta x \right)^{1/2},$$

Remarks 4.3. (i) The above results can be extended for more general polynomial nonlinearity satisfying (1.2) by a slightly modification.

(ii) In the same fashion the above results holds true in the Neumann boundary conditions for (u, v) (see [4]).

5 Computation Example

By using the proposed scheme (3.9) we can simulate a variety of dynamics of solutions for the Boussinesq type equations through a numerical calculation. Particularly, in this section we consider the isothermal Falk model:

$$\partial_t^2 u + \gamma \partial_x^4 u - \alpha_1 \partial_x^2 u = \partial_x^2 \{ \alpha_5 u^5 - \alpha_3 u^3 \}, \quad (x, t) \in \mathbb{R} \times \mathbb{R}^+, \quad (5.1)$$

which corresponds to the Boussinesq type equation (1.1) with nonlinearity (1.4). The equation is the isothermal model of Falk's thermoelastic system (see [2, Chapter 5]). In [3], this equation has a traveling wave solution in the austenite (high-temperature, namely large α_1) phase.

We demonstrate the travelling wave solution through a numerical computation. We first take the physical constant as the same as in [3]: $\gamma = 2$, $\alpha_5 = 6$, $\alpha_3 = 4$, $\alpha_1 = 1/4$, $L = 100$, $T = 50$. Moreover, we assume that the velocity of the travelling wave c is given by $c = -\sqrt{3}/2$ which satisfies the condition to observe the travelling wave solution given in [3]. According to the paper [3],

$$u(z) = u(x - ct - x_0) = \frac{1}{\sqrt{4 + 2\sqrt{2} + 4\sqrt{2} \sinh^2 \left(\frac{x - ct - x_0}{2\sqrt{2}} \right)}} \quad (5.2)$$

is a solution for (5.1), where x_0 is a initial position of the wave. Corresponding to (5.2), we choose the initial values. For the error estimate we need to replace the assumption (4.2) to

$$\Delta t < \frac{4\sqrt{\gamma}}{\alpha_1 + 3\alpha_3 C_1^2 + 5\alpha_5 C_1^4}, \quad (5.3)$$

In our case, we easily check the following $I_d(U_k^{(0)}, V_k^{(0)}) \leq 0.73$ and $|M_d(\mathbf{U}^{(0)})| \leq 3.57$. Then we see that $\max_{0 \leq k \leq K} |U_k^{(n)}| \leq 8.58$. Since the exact solution obviously satisfies $\|u(t)\|_{L^\infty} < 1$, we see that $C_1 \leq 8.58$. To satisfy the assumption (5.3) it is sufficient to take, for example, $\Delta t = \frac{1}{30000}$. If we take $\Delta x = 1/2$, then assumptions related to (3.10) which assure the existence of solution for the scheme are also satisfied for some M . In order to obtain new time-step solutions using our nonlinear scheme, we use Newton's method. Figure 1 shows the profiles of the numerical solutions U .




Figure 1: Numerical solution U

The energy fluctuation obtained by using our scheme is shown in Figure 2. From this we can confirm that the discrete energy values are conserved well.




Figure 2: Discrete conserved quantities I_d and M_d

References

- [1] M.J. Boussinesq, Thèory des ondes et des remous quise propagent..., J. Math. Pure Appl. sect 2 **117**(1872), 55–107.
- [2] M. Brokate and J. Sprekels, *Hysteresis and Phase Transitions*, Springer, New York, 1996.
- [3] F. Falk, E. W. Laedke and K. H. Spatschek, Stability of solitary-wave pulses in shape-memory alloys, Physical Rev. B, **36** (1987), 3031–3041.
- [4] D. Furihata, A stable and conservative finite difference scheme for the Cahn-Hilliard equation, Numer. Math., **87** (2001), 675-699.
- [5] D. Furihata and M. Matsuo, *Discrete Variational Derivative Method*, Numerical Analysis and Scientific Computing series, CRC Press/Taylor & Francis, 2010.
- [6] K. Ichikawa and S. Yoshikawa, An error estimate of conservative finite difference scheme for the Boussinesq type equations, submitted.
- [7] T. Matsuo, New conservative schemes with discrete variational derivatives for nonlinear wave equations, J. Comput. Appl. Math., **203** (2007), 32–56.

Existence of Solutions for Isotropic Elastodynamics

Thomas C. Sideris
University of California, Santa Barbara

The talk will present an overview of global existence results for the initial value problem associated to the motion of isotropic elastic and viscoelastic materials. In the absence of boundaries, the motion of a body is described by a smooth one-parameter family of orientation preserving diffeomorphisms or *deformations* $x(t, \cdot) : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $t \geq 0$, $n = 2, 3$. Under such a family of deformations, each material point $y \in \mathbb{R}^n$ follows the trajectory $t \mapsto x(t, y)$, $t \geq 0$. A homogeneous elastic material can be characterized by a smooth strain energy function

$$W : GL_+(n, \mathbb{R}) \rightarrow \mathbb{R}_+,$$

where $GL_+(n, \mathbb{R})$ is the set of $n \times n$ real matrices with positive determinant. Formally, the equations of motion are obtained by applying the principle of stationary action to the Lagrangian

$$(1) \quad \mathcal{L}[x] = \iint \left[\frac{1}{2} |D_t x|^2 - W(D_y x) \right] dy dt.$$

This yields the system

$$(2) \quad D_t^2 x_i - \sum_{j=1}^n D_j [S_{ij}(D_y x)] = 0, \quad S_{ij}(F) = \frac{\partial W(F)}{\partial F_{ij}}.$$

It is physically reasonable to assume that this system is Galilean invariant and that the equilibrium deformation $x(t, y) = y$ has vanishing energy and stress. We also assume that the material is isotropic. This implies that $W(I) = 0$, $S(I) = 0$, and that $W(F)$ is a function only of the principal invariants of $F^\top F$.

The construction of solutions relies on a combination of energy estimates and decay estimates. The results are perturbative in that at every time t the displacement from equilibrium, $u(t, y) = x(t, y) - y$, is assumed to be “small” in an appropriate energy norm, defined below. Therefore, a key role is played by the linearized problem

$$D_t^2 x^i - \sum_{j,k,\ell=1}^n A_{ijk\ell} D_j D_\ell x^k = 0, \quad A_{ijk\ell} = \frac{\partial^2 W}{\partial F_{ij} \partial F_{k\ell}}(I).$$

We impose the standard Legendre-Hadamard ellipticity condition

$$(3) \quad \sum_{i,j,k,\ell=1}^n A_{ijkl} x^i \xi_j x^k \xi_\ell = D_\varepsilon^2 W(I + \varepsilon x \otimes \xi)|_{\varepsilon=0} > 0,$$

for all $0 \neq x, \xi \in \mathbb{R}^n$.

This can also be justified by physical considerations. In the isotropic case, the linearized equation takes the form

$$(4) \quad Lx = D_t^2 x - c_2^2 \Delta x - (c_1^2 - c_2^2) D(D \cdot x) = 0,$$

with $c_1 > c_2 > 0$, $D = (D_1, \dots, D_n)$.

The energy

$$\mathcal{E}[x](t) = \frac{1}{2} \int_{\mathbb{R}^r} [|D_t x(t)|^2 + c_2^2 |Dx(t)|^2 + (c_1^2 - c_2^2) (D \cdot x(t))^2] dy$$

is conserved for equation (4).

Equation (4) is invariant under the Galilean (but not Lorentz) group, the generators of which contain the collection of vector fields

$$(5) \quad \Gamma = \{D_j, S = tD_t + y \cdot D, \Omega_{ij} = I(y_j D_i - y_i D_j) + e_j \otimes e_i - e_i \otimes e_j\},$$

where $\{e_i\}_{i=1}^n$ is the standard basis for $\mathbb{R}^n$. This implies the commutation properties

$$LS = S(L + 2) \quad \text{and} \quad L\Omega_{ij} = \Omega_{ij}L,$$

for the linear operator L . It follows that the higher order energies $\mathcal{E}_m[x](t) = \sum_{|\alpha| \leq m-1} \mathcal{E}[\Gamma^\alpha x](t)$ are also conserved for solutions of (4).

Solutions of (4) are a superposition of pressure and shear waves, propagating with speeds c_1 and c_2 respectively. This decomposition is approximated by

$$x(t, y) = \mathbb{P}(y)x(t, y) + [I - \mathbb{P}(y)]x(t, y), \quad \text{with} \quad \mathbb{P}(y) = \frac{y}{|y|} \otimes \frac{y}{|y|}.$$

Using scaling and rotational invariance of L , we obtain the uniform bound (see [6])

$$\mathcal{X}[D_j D_\ell x](t) \lesssim \mathcal{E}_2[x](t) + |t| \|Lx(t)\|_{L^2(\mathbb{R}^n)}, \quad j, \ell = 1, \dots, n$$

in which

$$\mathcal{X}[u](t) = \|\langle c_1 t - |y| \rangle \mathbb{P}(y) u(t, y)\|_{L^2(\mathbb{R}^n)} + \|\langle c_2 t - |y| \rangle [I - \mathbb{P}(y)] u(t, y)\|_{L^2(\mathbb{R}^n)}.$$

In the full nonlinear problem, the self-interaction of the two wave families influences the existence of global solutions. For the pressure

waves, this interaction must be limited by imposing an additional *null condition*:

$$(6) \quad D_\varepsilon^3 W(I + \varepsilon \xi \otimes \xi)|_{\varepsilon=0} = 0, \quad \text{for all } \xi \in \mathbb{R}^n.$$

In the isotropic case, the nonlinear interactions of shear waves is inherently null.

Theorem 1 ([4], see also [1]). *Let $n = 3$. Suppose that equation (2) arises from a strain energy function W which satisfies $W(I) = 0$, $S(I) = 0$, the Legendre-Hadamard condition (3), and the null condition (6).*

Choose initial data for (2) such that the displacement from equilibrium $u(t, y) = x(t, y) - y$ satisfies $\mathcal{E}_m[u](0) < \infty$, $m \geq 9$.

Let $\varepsilon > 0$ be sufficiently small. There is a constant $C > 0$, depending on W , such that if

$$\mathcal{E}_{m-2}[u](0) \exp [C \mathcal{E}_m^{1/2}[u](0)] < \varepsilon,$$

then (2) has a unique global solution $x(t, y) = y + u(t, y)$ with

$$u(t, y) \in \cap_{k=0}^m C^k(\mathbb{R}_+, H^{m-k}(\mathbb{R}^3))$$

and $\mathcal{E}_{m-2}[u](t) < \varepsilon$.

Incompressible isotropic materials can be viewed as a limiting case where the speed of the pressure waves is infinite, see [5]. In this case, the linearized problem is the wave equation, and the nonlinear interactions are null. However, with the incompressibility constraint the problem becomes nonlocal, requiring special estimates for the pressure term.

The equations of motion can be obtained by extremizing the Lagrangian (1) constrained to the class of volume preserving deformations. For simplicity, we consider only the Hookean case

$$W(F) = \frac{1}{2} \operatorname{tr} F^\top F,$$

whence the equations are

$$(7) \quad D_t^2 x_i - \Delta_y x_i + \sum_{j=1}^n (D_y x^{-1})_{ji} D_j \lambda = 0, \quad \det D_y x = 1.$$

Here, λ is a Lagrange multiplier. Using the constraint, it follows that

$$\Delta_x \lambda + \sum_{i,j=1}^n \partial_i \partial_j [D_t x_i D_t x_j - \sum_{k=1}^n D_k x_i D_k x_j] = 0,$$

where the derivatives Δ_x and ∂_i are taken in the spatial coordinates x . This shows that λ can be regarded as a nonlocal null form.

Theorem 2 ([7], [3]). *Choose initial data for (7) which is compatible with the incompressibility constraint*

$$\det D_y x(0) = 1, \quad \text{tr}[D_y x(0)^{-1} D_t D_y x(0)] = 0.$$

When $n = 3$, suppose that the displacement from equilibrium $u(t, y) = x(t, y) - y$ satisfies $\mathcal{E}_m[u](0) < \infty$, with $m \geq 8$. Let $\varepsilon > 0$ be sufficiently small. There is a constant $C > 0$ such that if

$$\mathcal{E}_{m-2}[u](0) \exp [C\mathcal{E}_m^{1/2}[u](0)] < \varepsilon,$$

then (7) has a unique global solution $x(t, y) = y + u(t, y)$ with

$$u(t, y) \in \cap_{k=0}^m C^k(\mathbb{R}_+, H^{m-k}(\mathbb{R}^3))$$

and $\mathcal{E}_{m-2}[u](t) < \varepsilon$.

When $n = 2$, suppose that $\mathcal{E}_m[u](0) < \varepsilon$, $m \geq 5$, for ε sufficiently small. Then (7) has a unique local solution $x(t, y) = y + u(t, y)$ with

$$u(t, y) \in \cap_{k=0}^m C^k([0, T_\varepsilon], H^{m-k}(\mathbb{R}^2)),$$

$T_\varepsilon > \exp(C/\varepsilon)$, and $\mathcal{E}_m[u](t) < 2\varepsilon$.

Viscoelastic materials arise through the addition of weak viscous dissipation to the classical elastic case, resulting in a hyperbolic-parabolic system. Obtaining global existence results which are uniform in the magnitude of the dissipation parameter ν requires the adaptation of the hyperbolic estimates to a system without scaling invariance, see [2]. As a model, consider the nonlinear wave equation in 3-D

$$(8) \quad D_t^2 u - \Delta u + \nu \Delta D_t u = \sum_{i,j,k=1}^3 B_{ijk} D_i (D_j u D_k u).$$

Theorem 3. *Choose initial data for (8) with $\mathcal{E}_m[u](0) < \infty$, $m \geq 6$. Define $\delta = \max\{|\sum_{i,j,k=1}^3 B_{ijk} \omega_i \omega_j \omega_k| : \omega \in \mathbb{R}^3, |\omega| = 1\}$. Let ε be sufficiently small. Put $m^* = \lfloor \frac{m+5}{2} \rfloor$. There is a constant $C > 0$ such that if*

$$\mathcal{E}_{m^*}[u](0) \{ \exp [C\mathcal{E}_m^{1/2}[u](0)] + \exp [C\mathcal{E}_m[u](0)\delta^2/\nu] \} < \varepsilon,$$

then (8) has a unique global solution with $u(t, y) \in \cap_{k=0}^m C^k(\mathbb{R}_+, H^{m-k}(\mathbb{R}^3))$ and $\mathcal{E}_{m^}[u](t) < \varepsilon$.*

The number of occurrences of the scaling operator S is restricted to $m^* + 1$ in $\mathcal{E}_m[u](t)$. Also, since (8) is a scalar equation, we use $\Omega_{ij} = y_i D_j - y_j D_i$ instead of the vectorial version in (5). This result says, in particular, if the null condition is satisfied, $\delta = 0$, then (8) has global solutions for small initial data, *independent of the size of ν .*

REFERENCES

- [1] Agemi, R. *Global existence of nonlinear elastic waves*. Invent. Math. 142 (2000), no. 2, 225–250.
- [2] Kessenich, P. *Global Existence with Small Initial Data for Three-Dimensional Incompressible Isotropic Viscoelastic Materials*, available online at arXiv:0903.2824.
- [3] Lei, Z., Sideris, T.C., and Zhou, Y. *Almost global existence for 2-D incompressible isotropic elastodynamics*. To appear in Trans. A.M.S.
- [4] Sideris, T. C. *Nonresonance and global existence of prestressed nonlinear elastic waves*. Ann. of Math. (2) 151 (2000), no. 2, 849–874.
- [5] Sideris, T. C. and Thomases, B. *Global existence for three-dimensional incompressible isotropic elastodynamics via the incompressible limit*. Comm. Pure Appl. Math. 58 (2005), no. 6, 750–788.
- [6] Sideris, T. C. and Thomases, B. *Local energy decay for solutions of multi-dimensional isotropic symmetric hyperbolic systems*. J. Hyperbolic Differ. Equ. 3 (2006), no. 4, 673–690.
- [7] Sideris, T. C. and Thomases, B. *Global existence for 3d incompressible isotropic elastodynamics*. Comm. Pure Appl. Math. 60 (2007), no. 12, 1707–1730.

Mathematical modeling and numerical treatment of adhesion, exfoliation and collision

OMATA, Seiro (Kanazawa University)

1 Scalar problem

In this talk, we would like to treat mathematical modeling and numerical treatment of adhesion, exfoliation and collision. Phenomena of adhesion and exfoliation come from droplet motion on a plane. We describe the surface of droplet by using a graph of scalar function. For this problem, key words are 'volume constraint', 'free boundary' and 'potential energy'. The constraint produces a non-local term in the partial differential equations. Hence, we need to introduce variational method to these problems. Lagrangian will be

$$L(u) = \int_{\Omega} (-\chi_{u>0} u_t^2 + |\nabla u|^2 + Q\chi_{u>0}) dx, \quad (1)$$

where $\chi_{u>0}$ is the characteristic function and Q is a force of adhesion. If the droplet keeps its volume, we can get the following equation:

$$\chi_{u>0} u_{tt} = \Delta u - Q\chi'_{u>0} + \left(\int_{\Omega} [u_{tt}u + |\nabla u|^2] dx \right) \chi_{u>0}. \quad (2)$$

2 Shell bouncing problem

The model equation is derived by calculating the energy stored in the shell. The considered types of energy are stretching energy, bending energy, energy related to the compression of the enclosed gas, potential energy and kinetic energy. We assume that the strip of initial radius r_0 is bent to radius r and stretched by the ratio $\tilde{\mu}$. We adopt the following form of elastic energy

$$E_e(\mathbf{p}) = \frac{1}{24}kh^3 \int_0^{2\pi} (\kappa - \kappa_0)^2 |\mathbf{p}_{\theta}| d\theta + \frac{1}{2}kh \int_0^{2\pi} \left(\frac{|\mathbf{p}_{\theta}|}{|\mathbf{q}_{\theta}|} - 1 \right)^2 |\mathbf{q}_{\theta}| d\theta.$$

Denoting the mass density of the shell in equilibrium by σ , the local mass density of the shell $\mathbf{p}$ becomes $\sigma|\mathbf{q}_{\theta}|/|\mathbf{p}_{\theta}|$ and thus the kinetic energy is given by

$$E_k(\mathbf{p}) = \frac{1}{2}h \int_0^{2\pi} \sigma|\mathbf{q}_{\theta}| |\mathbf{p}_t|^2 \chi_{p^2>0} d\theta.$$

Physically, this corresponds to the assumption of zero reflection and infinite friction between the shell and the obstacle.

When the shell is closed, it is necessary to take into account also the energy related to the compression of the gas present inside the shell. The energy stored due to compression of the enclosed volume V of gas can now be calculated as minus the work done by pressure

forces:

$$E_g(\mathbf{p}) = - \int_{V_0}^V P dV = - \int_{V_0}^V \left\{ P_0 + c_g \left(\frac{1}{V} - \frac{1}{V_0} \right) \right\} dV = -P_0(V - V_0) - c_g \left(1 - \frac{V}{V_0} + \ln \frac{V}{V_0} \right). \quad (3)$$

The constant c_g is the product of the total mass of the gas and the square of the sound of speed. The equilibrium volume V_0 is known and the volume of $\mathbf{p}(t)$ is given by

$$V = \frac{1}{2} \int_0^{2\pi} (\mathbf{p} \cdot A\mathbf{p}_\theta) \chi_{p^2 > 0} d\theta.$$

In real situations, the impact of a shell is influenced by the action of gravity. Therefore, we introduce also the gravity potential of the shell

$$E_p(\mathbf{p}) = gh \int_0^{2\pi} \sigma |\mathbf{q}_\theta| p^2 \chi_{p^2 > 0} d\theta.$$

Employing the obtained elastic, kinetic and gas energies, the action integral is written as

$$I(\mathbf{p}) = \int_0^T (E_e(\mathbf{p}) + E_g(\mathbf{p}) + E_p(\mathbf{p}) - E_k(\mathbf{p})) dt. \quad (4)$$

Applying Hamilton's principle, the governing equation for the free part of the shell is obtained from

$$\frac{d}{d\epsilon} I(\mathbf{p} + \epsilon \phi) \Big|_{\epsilon=0} = 0 \quad \forall \phi \in [C_0^\infty((0, T) \times (0, 2\pi) \cap \{p^2 > 0\})]^2, \quad (5)$$

where I is the action functional defined in (4).

In the sequel, we shall use the notation $\rho(\theta) = |\mathbf{p}_\theta(\theta)|/|\mathbf{q}_\theta(\theta)|$ for the local relative density with respect to the equilibrium state.

Taking into account the influence of the obstacle on the computed energy, one may expect that the following equation expresses, in a rough approximation, the deformation of the whole shell:

$$\begin{aligned} \sigma \chi_{p^2 > 0} \mathbf{p}_{tt} = & \left\{ -\frac{1}{12} k h^2 \rho (\kappa_{ss} + \frac{1}{2} \kappa^3) \chi_{p^2 > 0} + \frac{1}{24} k h^2 \kappa_0^2 \rho \kappa - k \rho (\rho - 1) \kappa \chi_{p^2 > 0} \right\} \boldsymbol{\nu} \\ & + \frac{\rho}{h} \left(P_0 + c_g \left(\frac{1}{V} - \frac{1}{V_0} \right) \right) \chi_{p^2 > 0} \boldsymbol{\nu} + k \rho \rho_s \chi_{p^2 > 0} \boldsymbol{\tau} + g \sigma \chi_{p^2 > 0} \mathbf{e}_2. \end{aligned} \quad (6)$$

In our talk, we will explain how to treat the above equation numerically and will show the numerical result.

References

- [1] E. Ginder, K. Svadlenka: *A variational approach to a constrained hyperbolic free boundary problem*, Nonlinear Analysis, Theory Methods Appl. 71(12), 2009, pp. 1527–1537.

- [2] H. Imai, K. Kikuchi, N. Nakane, S. Omata, T. Tachikawa: *A numerical approach to the asymptotic behavior of solutions of a one-dimensional free boundary problem of hyperbolic type*, Japan J. Indust. Appl. Math. 18 (1), 2001, pp. 43–58.
- [3] K. Ito, M. Kazama, H. Nakagawa, K. Svadlenka: *Numerical solution of a volume-constrained free boundary problem by the discrete Morse flow method*, Gakuto International Series 29, 2008, pp. 383–398.
- [4] K. Kikuchi, S. Omata: *A free boundary problem for a one dimensional hyperbolic equation*, Adv. Math. Sci. Appl. 9(2), 1999, pp. 775–786.
- [5] S. Omata: *A numerical treatment of film motion with free boundary*, Adv. Math. Sci. Appl. 14(1), 2004, pp. 129–137.
- [6] K. Rektorys, “*On application of direct variational methods to the solution of parabolic boundary value problems of arbitrary order in the space variables*”, Czech. Math. J. **21** (1971), 318–339.
- [7] K. Svadlenka, S. Omata: *Construction of solutions to heat-type problems with volume constraint via the discrete Morse flow*, Funkcialaj Ekvacioj 50, 2007, pp. 261–285.
- [8] K. Svadlenka, S. Omata: *Mathematical modelling of surface vibration with volume constraint and its analysis*, Nonlinear Analysis 69, 2008, pp. 3202–3212.
- [9] K. Svadlenka, S. Omata: *Mathematical analysis of a constrained parabolic free boundary problem describing droplet motion on a surface*, Indiana Univ. Math. J. 58(5), 2009, pp. 2073–2102.
- [10] A. Tachikawa, “*A variational approach to constructing weak solutions of semilinear hyperbolic systems*”, Adv. Math. Sci. Appl. 4 (1994), 93–103.
- [11] T. Yamazaki, S. Omata, K. Svadlenka, K. Ohara: *Construction of approximate solution to a hyperbolic free boundary problem with volume constraint and its numerical computation*, Adv. Math. Sci. Appl. 16(1), 2006, pp. 57–67.
- [12] H. Yoshiuchi, S. Omata, K. Svadlenka, K. Ohara: *Numerical solution of film vibration with obstacle*, Adv. Math. Sci. Appl. 16(1), 2006, pp. 33–43.

Phase field systems of grain boundaries with solidification effects

KEN SHIRAKAWA*

Introduction

This study is based on a recent collaborations with Prof. Salvador Moll[†], Dr. Hiroshi Watanabe[‡] and Prof. N. Yamazaki^{††} (cf. [10, 11, 12, 15]), which are communicated and supported by Prof. J. M. Mazón^{‡‡}.

Let $0 < T < \infty$ be a fixed constant, let $1 < N \in \mathbb{N}$ be a fixed number, and let $\Omega \subset \mathbb{R}^N$ be a bounded domain with a smooth boundary $\partial\Omega$. Besides, we denote by $\nu_{\partial\Omega}$ the unit outer normal vector on $\partial\Omega$, and we set $Q := (0, T) \times \Omega$ and $\Sigma := (0, T) \times \partial\Omega$.

Let $\nu > 0$ be a fixed small constant. In this paper, a coupled system of parabolic initial-boundary value problems is considered. This system is denoted by $(S)_\nu$, and formally described as follows.

$$(S)_\nu: \begin{cases} w_t - \Delta w + p(w, \eta) + \nu\beta'(w)|\nabla\theta|^2 = 0 & \text{in } Q, \\ \nabla w \cdot \nu_{\partial\Omega} = 0 & \text{on } \Sigma, \\ w(0, x) = w_0(x), & x \in \Omega; \\ \eta_t - \Delta \eta + q(w, \eta) + \alpha'(\eta)|\nabla\theta| = 0 & \text{in } Q, \\ \nabla \eta \cdot \nu_{\partial\Omega} = 0 & \text{on } \Sigma, \\ \eta(0, x) = \eta_0(x), & x \in \Omega; \\ \alpha_0(w, \eta) \theta_t - \operatorname{div} \left(\alpha(\eta) \frac{D\theta}{|D\theta|} + \nu\beta(w)\nabla\theta \right) = 0 & \text{in } Q, \\ \left(\alpha(\eta) \frac{D\theta}{|D\theta|} + \nu\beta(w)\nabla\theta \right) \cdot \nu_{\partial\Omega} = 0 & \text{on } \Sigma, \\ \theta(0, x) = \theta_0(x), & x \in \Omega. \end{cases}$$

The system $(S)_\nu$ is based on a mathematical model of planar grain boundary motions with solidification effects, proposed by Kobayashi [9] and Warren-Kobayashi-Lobkovski-Carter [14]. In accordance with the modelling method as in [9, 14], the system $(S)_\nu$ is

* Department of Mathematics, Faculty of Education, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba, 263-8522, Japan; sirakawa@faculty.chiba-u.jp.

† Departament d'Anàlisi Matemàtica, Universitat de València, Spain; ‡ Department of General Education, Salesian Polytechnic, Japan; †† Department of Mathematics, Faculty of Engineering, Kanagawa University, Japan; ‡‡ Departament d'Anàlisi Matemàtica, Universitat de València, Spain.

This study is supported by Grant-in-Aid for Encouragement of Young Scientists (B) (No. 24740099) JSPS.

derived as a gradient system of a governing free-energy, defined as:

$$\begin{aligned} [w, \eta, \theta] &\in [H^1(\Omega) \cap L^\infty(\Omega)] \times [H^1(\Omega) \cap L^\infty(\Omega)] \times H^1(\Omega) \\ \mapsto \mathcal{F}_\nu(w, \eta, \theta) &:= \frac{1}{2} \int_\Omega |\nabla w|^2 dx + \frac{1}{2} \int_\Omega |\nabla \eta|^2 dx + \int_\Omega f(w, \eta) dx \\ &\quad + \int_\Omega \alpha(\eta) |\nabla \theta|^2 dx + \nu \int_\Omega \beta(w) |\nabla \theta|^2 dx. \end{aligned}$$

In the context, the unknown $w = w(t, x)$ is an order parameter to indicate the solidification order of grain. Unknowns $\eta = \eta(t, x)$ and $\theta = \theta(t, x)$ are order parameters to reproduce the crystalline orientation in Q , by using a vector field:

$$(t, x) \in Q \mapsto \eta(t, x) [\cos \theta(t, x), \sin \theta(t, x)] \in \mathbb{R}^2.$$

Then, the components η and θ are supposed to indicate the orientation order and the orientation angle of the grain, respectively. In particular, w and η are supposed to satisfy range constraint properties $0 \leq w, \eta \leq 1$ in Q , and the cases when $[w, \eta] = [1, 1]$ and $[w, \eta] = [0, 0]$ are supposed to reproduce two stable states of grains: *the solidified-oriented state*; *the liquefied-disoriented state*, respectively.

The major theme of this study is to ensure qualitative properties for the system $(S)_\nu$ from the theoretical viewpoint in mathematics. As part of this theme, we here focus on an approach based on the time-discretization, and we conclude the existence theorem of the system $(S)_\nu$. Furthermore, we will mention about the problems in future with the application possibilities of similar approaches by time-discretizations.

1 Statement of the Main Theorem

First of all, let us confirm the assumptions for the given functions $f = f(w, \eta)$, $p = p(w, \eta)$, $q = q(w, \eta)$, $\alpha_0 = \alpha_0(w, \eta)$, $\alpha = \alpha(\eta)$, $\beta = \beta(w)$, η_0 and θ_0 as in the system $(S)_\nu$.

(A1) $f \in C^2([0, 1]^2)$ and $p, q \in C^1([0, 1]^2)$ such that

$$\left\{ \begin{array}{l} \bullet f(w, \eta) \geq 0 \text{ and } \nabla f(w, \eta) = [p(w, \eta), q(w, \eta)] \text{ for all } [w, \eta] \in [0, 1]^2, \\ \bullet p(0, \eta) \leq 0 \text{ and } p(1, \eta) \geq 0 \text{ for all } \eta \in [0, 1], \\ \bullet q(w, 0) \leq 0 \text{ and } q(w, 1) \geq 0 \text{ for all } w \in [0, 1]. \end{array} \right.$$

(A2) $\alpha_0 \in C^1([0, 1]^2)$ and $\alpha, \beta \in C^2[0, 1]$ such that:

$$\left\{ \begin{array}{l} \bullet \alpha_0(w, \eta) > 0 \text{ for all } [w, \eta] \in [0, 1]^2, \\ \bullet \alpha'(0) = 0, \alpha(\eta) > 0 \text{ and } \alpha''(\eta) \geq 0 \text{ for all } \eta \in [0, 1], \\ \bullet \beta'(0) = 0, \beta(w) > 0 \text{ and } \beta''(w) \geq 0 \text{ for all } w \in [0, 1], \end{array} \right.$$

where α' and β' are the differentials of α and β , respectively, and α'' and β'' are the second differentials. Note that α and β turn out to be convex functions on $\mathbb{R}$, and

$$\delta_* := \min_{[\tilde{w}, \tilde{\eta}] \in [0, 1]^2} \{\alpha_0(\tilde{w}, \tilde{\eta}), \alpha(\tilde{\eta}), \beta(\tilde{w})\} > 0.$$

(A3) The pair (triplet) of initial data $[w_0, \eta_0, \theta_0]$ belongs to a class $D_* \subset H^1(\Omega)^2 \times H^1(\Omega)$, defined as:

$$D_* := \{ [\tilde{w}, \tilde{\eta}, \tilde{\theta}] \in H^1(\Omega)^3 \mid 0 \leq \tilde{w}, \tilde{\eta} \leq 1 \text{ a.e. in } \Omega, \tilde{\theta} \in L^\infty(\Omega) \}.$$

In addition, for the convenience of descriptions, we prepare the following notations.

Notation 1.1 For any $\tilde{v} = [\tilde{w}, \tilde{\eta}] \in L^\infty(\Omega)^2$, let $\Phi_\nu(\tilde{v}; \cdot) = \Phi_\nu(\tilde{w}, \tilde{\eta}; \cdot)$ be a proper l.s.c. and convex function on $L^2(\Omega)$, defined as:

$$\begin{aligned} \vartheta \in L^2(\Omega) &\mapsto \Phi_\nu(\tilde{v}; \vartheta) = \Phi_\nu(\tilde{w}, \tilde{\eta}; \vartheta) \\ &:= \begin{cases} \int_\Omega \alpha(\tilde{\eta}) |\nabla \vartheta| dx + \nu \int_\Omega \beta(\tilde{w}) |\nabla \vartheta|^2 dx, & \text{if } \vartheta \in H^1(\Omega), \\ \infty, & \text{otherwise,} \end{cases} \end{aligned}$$

and let $\partial\Phi_\nu(\tilde{v}; \cdot) = \partial\Phi_\nu(\tilde{w}, \tilde{\eta}; \cdot)$ be the L^2 -subdifferential of $\Phi_\nu(\tilde{v}; \cdot) = \Phi_\nu(\tilde{w}, \tilde{\eta}; \cdot)$.

Now, the Main Theorem is stated as follows.

Main Theorem. *(Solvability of the system $(S)_\nu$) Under the assumptions (A1)-(A3), the system $(S)_\nu$ admits at least one solution $[w, \eta, \theta]$, in the sense of the following items.*

(S1) $w \in W^{1,2}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^1(\Omega))$, $0 \leq w \leq 1$ a.e. in Q ;
 $\eta \in W^{1,2}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^1(\Omega)) \cap L^2(0, T; H^2(\Omega))$, $0 \leq \eta \leq 1$ a.e. in Q ;
 $\theta \in W^{1,2}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^1(\Omega)) \cap L^\infty(Q)$, $|\theta|_{L^\infty(Q)} \leq |\theta_0|_{L^\infty(\Omega)}$.

(S2) w solves the following variational identity of parabolic type:

$$\begin{aligned} \int_\Omega (w_t(t) + p(w(t), \eta(t)) + \nu \beta'(w(t)) |\nabla \theta(t)|^2) \varphi dx + \int_\Omega \nabla w(t) \cdot \nabla \varphi dx &= 0, \\ \text{for any } \varphi \in H^1(\Omega) \cap L^\infty(\Omega) \text{ and a.e. } t \in (0, T). \end{aligned} \quad (1.1)$$

(S3) η solves the following variational identity of parabolic type:

$$\begin{aligned} \int_\Omega (\eta_t(t) + q(w(t), \eta(t)) + \alpha'(\eta(t)) |\nabla \theta(t)|) \psi dx + \int_\Omega \nabla \eta(t) \cdot \nabla \psi dx &= 0, \\ \text{for any } \psi \in H^1(\Omega) \text{ and a.e. } t \in (0, T). \end{aligned} \quad (1.2)$$

(S4) θ solves the following variational inequality of parabolic type:

$$\begin{aligned} \int_\Omega \alpha_0(w(t), \eta(t)) \theta_t(t) (\theta(t) - \omega) dx + \Phi_\nu(w(t), \eta(t); \theta(t)) &\leq \Phi_\nu(w(t), \eta(t); \omega), \\ \text{for any } \omega \in H^1(\Omega) \text{ and a.e. } t \in (0, T). \end{aligned} \quad (1.3)$$

(S5) $[w(0), \eta(0), \theta(0)] = [w_0, \eta_0, \theta_0]$ in $L^2(\Omega)^3$.

Remark 1.1 For the solution $[w, \eta, \theta]$ to the system $(S)_\nu$, let us put $v := [w, \eta]$. Then, we note that the variational identities (1.1)-(1.2) can be reduced to the following form of a single evolution equation:

$$\begin{aligned} v_t(t) + Fv(t) - v(t) + (\nabla f)(v(t)) + [\nu\beta'(w(t))|\nabla\theta(t)|^2, \alpha'(\eta(t))|\nabla\theta(t)|] &= 0 \\ \text{in } [H^1(\Omega) \cap L^\infty(\Omega)]^* \times H^1(\Omega)^*, \quad \text{a.e. } t \in (0, T), \end{aligned}$$

where $F : H^1(\Omega) \longrightarrow H^1(\Omega)^*$ is the duality mapping between $H^1(\Omega)$ and $H^1(\Omega)^*$, and hence $F\tilde{v} := [F\tilde{w}, F\tilde{\eta}]$ for any $\tilde{v} = [\tilde{w}, \tilde{\eta}]$.

Also, we note that the variational inequality (1.3) is reformulated to the following form of an evolution equation:

$$\alpha_0(v(t))\theta_t(t) + \partial\Phi_\nu(v(t); \theta(t)) \ni 0 \quad \text{in } L^2(\Omega), \quad \text{a.e. } t \in (0, T),$$

governed by the subdifferentials $\partial\Phi_\nu(v(t); \cdot)$ of unknown-dependent convex functions $\Phi_\nu(v(t); \cdot)$ for $t \in [0, T]$.

2 Solution method for the system $(S)_\nu$

As mentioned in Introduction, the mathematical approach to the system $(S)_\nu$ is based on the time-discretization. To this end, we need to prepare a class $\{(RS)_\varepsilon^\nu \mid 0 < \varepsilon < 1\}$ of relaxation systems, denoted by $(RS)_\varepsilon^\nu$, for the original system $(S)_\nu$.

Let us fix a constant $N_* > 1 + N/2$, and let us fix any $\tilde{\theta}_0 \in H^{N_*}(\Omega)$. Also, for any $0 < \varepsilon < 1$ and any $\tilde{v} = [\tilde{w}, \tilde{\eta}] \in H^1(\Omega)^2$, let us define a relaxed convex function $\Psi_\varepsilon^\nu(\tilde{v}; \cdot) = \Psi_\varepsilon^\nu(\tilde{w}, \tilde{\eta}; \cdot)$, by putting:

$$\begin{aligned} \vartheta \in L^2(\Omega) &\mapsto \Psi_\varepsilon^\nu(\tilde{v}; \vartheta) = \Psi_\varepsilon^\nu(\tilde{w}, \tilde{\eta}; \vartheta) \\ &:= \begin{cases} \Phi_\nu(\tilde{v}; \vartheta) + \frac{\varepsilon}{2}|\vartheta|_{H^{N_*}(\Omega)}^2, & \text{if } \vartheta \in H^{N_*}(\Omega), \\ \infty, & \text{otherwise,} \end{cases} \end{aligned}$$

and let us denote by $\partial\Psi_\varepsilon^\nu(\tilde{v}; \cdot) = \partial\Psi_\varepsilon^\nu(\tilde{w}, \tilde{\eta}; \cdot)$ its subdifferential in $L^2(\Omega)$.

By using the above notations, the relaxation system $(RS)_\varepsilon^\nu$, for each $0 < \varepsilon < 1$, is prescribed as follows.

$(RS)_\varepsilon^\nu$: to find $[v, \theta] \in C([0, T]; L^2(\Omega))^3$ with $v = [w, \eta] \in C([0, T]; L^2(\Omega))^2$, such that

$$\begin{cases} v_t(t) - \Delta_N v(t) + (\nabla f)(v(t)) + [\nu\beta'(w(t))|\nabla\theta(t)|^2, \alpha'(\eta(t))|\nabla\theta(t)|] = 0 \\ \quad \text{in } L^2(\Omega)^2, \quad \text{a.e. } t \in (0, T), \\ v(0) = [w(0), \eta(0)] = v_0 := [w_0, \eta_0] \text{ in } L^2(\Omega)^2; \end{cases} \quad (2.1)$$

$$\begin{cases} \alpha_0(v(t))\theta_t(t) + \partial\Psi_\varepsilon^\nu(v(t); \theta(t)) \ni 0 \text{ in } L^2(\Omega), \quad \text{a.e. } t \in (0, T), \\ \theta(0) = \tilde{\theta}_0 \text{ in } L^2(\Omega), \end{cases} \quad (2.2)$$

where Δ_N is the operator of the Laplacian subject to the Neumann-zero boundary condition, i.e.

$$\Delta_N : z \in \{ \tilde{z} \in H^2(\Omega) \mid \nabla \tilde{z} \cdot \nu_{\partial\Omega} = 0 \text{ a.e. on } \partial\Omega \} \mapsto \Delta z \in L^2(\Omega).$$

On that basis, for any $0 < \varepsilon < 1$, we call a pair (triplet) of functions $[v, \theta] = [w, \eta, \theta]$ a solution to $(\text{RS})_\varepsilon^\nu$ if and only if $v = [w, \eta] \in W^{1,2}(0, T; L^2(\Omega))^2 \cap L^\infty(0, T; H^1(\Omega))^2 \cap L^2(0, T; H^2(\Omega))^2$, $\theta \in W^{1,2}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^{N^*}(\Omega))$, and the components $v = [w, \eta]$ and θ fulfill the Cauchy problems (2.1) and (2.2), respectively. Note that the embedding $H^{N^*}(\Omega) \subset W^{1,\infty}(\Omega)$ enables us to treat $\beta'(w(t))|\nabla\theta(t)|^2$ and $\alpha'(\eta(t))|\nabla\theta(t)|$ as in (2.1), as L^∞ -perturbations.

Now, we prepare an approximation index $h \in (0, 1)$ of the time-step, and denote by $(\text{AP})_h^\varepsilon$ the time-discretization system for $(\text{RS})_\varepsilon^\nu$, formulated as follows.

$(\text{AP})_h^\varepsilon$:

$$\frac{v_{h,i} - v_{h,i-1}}{h} - \Delta_N v_{h,i} + (\nabla f)(v_{h,i}) + [\nu \beta'(w_{h,i}) |\nabla \theta_{h,i-1}|^2, \alpha'(\eta_{h,i}) |\nabla \theta_{h,i-1}|] = 0 \quad \text{in } L^2(\Omega)^2, \quad (2.3)$$

$$\alpha_0(v_{h,i}) \frac{\theta_{h,i} - \theta_{h,i-1}}{h} + \partial \Psi_\varepsilon^\nu(\alpha(v_{h,i}); \theta_{h,i}) \ni 0 \quad \text{in } L^2(\Omega), \quad (2.4)$$

for $i = 1, 2, 3, \dots$, subject to:

$$[v_{h,0}, \theta_{h,0}] := [w_{h,0}, \eta_{h,0}, \theta_{h,0}] = [v_0, \tilde{\theta}_0] \quad \text{in } L^2(\Omega)^3. \quad (2.5)$$

Here, for any $0 < h < 1$, we call a pair (triplet) of sequences $[\{v_{h,i}\}, \{\theta_{h,i}\}] = [\{w_{h,i}\}, \{\eta_{h,i}\}, \{\theta_{h,i}\}] \subset L^2(\Omega)^3$ a solution to $(\text{AP})_h^{(\nu)}$, or simply an approximating solution, if and only if $\{v_{h,i} \mid i \in \mathbb{N}\} \subset H^2(\Omega)^2$, $\{\theta_{h,i} \mid i \in \mathbb{N}\} \subset H^{N^*}(\Omega)$, and for any $i \in \mathbb{N}$, the components $v_{h,i}$ and $\theta_{h,i}$ fulfill the respective elliptic type problems (2.3) and (2.4) subject to (2.5).

In this paper, the class $\{(\text{AP})_h^\varepsilon \mid 0 < \varepsilon, h < 1\}$ of the time-discretization systems is adopted as that of approximation problems for $(\text{S})_\nu$. With regard to each approximation problem, we can prove the following theorem.

Theorem 2.1 (*Solvability of the approximation problem*) *There exists a small constant $0 < h_* < 1$, and for any $0 < \varepsilon < 1$ and any $0 < h \leq h_*$, the approximation problem $(\text{AP})_h^\varepsilon$ admits a unique solution $[\{v_{h,i}\}, \{\theta_{h,i}\}] = [\{w_{h,i}\}, \{\eta_{h,i}\}, \{\theta_{h,i}\}] \subset L^2(\Omega)^3$, such that:*

$$0 \leq w_{h,i}, \eta_{h,i} \leq 1 \quad \text{a.e. in } \Omega, \quad |\theta_{h,i}|_{L^\infty(\Omega)} \leq |\theta_{h,i-1}|_{L^\infty(\Omega)}, \quad \text{and} \quad (2.6)$$

$$\begin{aligned} \frac{1}{2h} |v_{h,i} - v_{h,i-1}|_{L^2(\Omega)^2}^2 + \frac{1}{h} \left| \sqrt{\alpha_0(v_{h,i})} (\theta_{h,i} - \theta_{h,i-1}) \right|_{L^2(\Omega)}^2 + \mathcal{E}_\varepsilon^\nu(v_{h,i}, \theta_{h,i}) \\ \leq \mathcal{E}_\varepsilon^\nu(v_{h,i-1}, \theta_{h,i-1}), \quad i = 1, 2, 3, \dots, \end{aligned} \quad (2.7)$$

where $\mathcal{E}_\varepsilon^\nu$ is the relaxed free-energy, defined as:

$$[\tilde{v}, \tilde{\theta}] = [\tilde{w}, \tilde{\eta}, \tilde{\theta}] \in H^1(\Omega)^2 \times H^{N^*}(\Omega) \mapsto \mathcal{E}_\varepsilon^\nu(\tilde{v}, \tilde{\theta}) := \mathcal{F}_\nu(\tilde{w}, \tilde{\eta}, \tilde{\theta}) + \frac{\varepsilon}{2} |\tilde{\theta}|_{H^{N^*}(\Omega)}^2.$$

Outline of the proof. Note that (2.3) and (2.4) can be regarded as independent variational problems of elliptic types. Indeed, it is not so difficult to show that the problem (2.3) has a unique (vectorial) unknown $v_{h,i} = [w_{h,i}, \eta_{h,i}]$. Hence, after solving (2.3), we can restrict the unknown in (2.4) only to $\theta_{h,i}$. Consequently, for each step $i \in \mathbb{N}$, these problems can be solved in the order of (2.3) and (2.4) by means of the usual variational

method (e.g. [3]). The property (2.6) can be deduced on the basis of the theory of T-monotonicity (cf. [2, 7]). Furthermore, the inequality (2.7) is obtained by multiplying the both sides of (2.3) and (2.4) by $(v_{h,i} - v_{h,i-1})$ and $(\theta_{h,i} - \theta_{h,i-1})$, respectively, and taking the sum of the results. Incidentally, the smallness $0 < h \leq h_*$ for h will be needed only for the discussions associated with $\{v_{h,i}\}$: the solvability of (2.3); the range constraint property as in (2.6); the derivation of the coefficient $\frac{1}{2h}$ at the head of (2.7). $\blacksquare$

3 Outline of the proof of the Main Theorem

In this paper, we show just the outline of the proof. The detailed arguments will be published in the forthcoming paper (in preparation).

Roughly summarized, the proof is proceeded through several steps, lined up below.

Step 1: preparation of notation. We begin with taking a sequence $\{\tilde{\theta}_0^\varepsilon\} \subset H^{N^*}(\Omega)$ such that:

$$\begin{cases} |\tilde{\theta}_0^\varepsilon|_{L^\infty(\Omega)} \leq |\theta_0|_{L^\infty(\Omega)} & \text{for all } 0 < \varepsilon < 1, \\ \tilde{\theta}_0^\varepsilon \rightarrow \theta_0 & \text{in } H^1(\Omega) \text{ and } \mathcal{E}_\varepsilon^\nu(v_0; \tilde{\theta}_0^\varepsilon) \rightarrow \mathcal{F}_\nu(w_0, \eta_0, \theta_0), \text{ as } \varepsilon \searrow 0. \end{cases} \quad (3.1)$$

To this end, we need to prove the following (Fact 1), in advance.

(Fact 1) The sequence $\{\Psi_\varepsilon^\nu(v_0; \cdot) \mid 0 < \varepsilon < 1\}$ of convex functions converges to the convex function $\Phi_\nu(v_0; \cdot)$ on $L^2(\Omega)$, in the sense of Γ -convergence, as $\varepsilon \searrow 0$.

Subsequently, let $0 < h_* < 1$ be the small constant as in Theorem 2.1, and for any $0 < h \leq h_*$, let $[\{v_{h,i}^\varepsilon\}, \{\theta_{h,i}^\varepsilon\}] = [\{w_{h,i}^\varepsilon\}, \{\eta_{h,i}^\varepsilon\}, \{\theta_{h,i}^\varepsilon\}]$ be the solution to $(AP)_h^\varepsilon$ in the case when $\tilde{\theta}_0 = \tilde{\theta}_0^\varepsilon$. On that basis, we set:

$$\begin{cases} t_{h,i} := ih, & i = 0, 1, 2, 3, \dots, \\ \Delta_{h,i} := [t_{h,i-1}, t_{h,i}), & i = 1, 2, 3, \dots, \end{cases} \quad \text{for any } 0 < h \leq h_*,$$

and we construct sequences:

$$\left\{ \begin{array}{l} \{\bar{v}_h^\varepsilon, \bar{\theta}_h^\varepsilon\} = \{[\bar{w}_h^\varepsilon, \bar{\eta}_h^\varepsilon, \bar{\theta}_h^\varepsilon] \mid 0 < \varepsilon < 1, 0 < h \leq h_*\} \\ \quad \subset L^\infty(0, T; H^2(\Omega))^2 \times L^\infty(0, T; H^{N^*}(\Omega)), \\ \{\underline{v}_h^\varepsilon, \underline{\theta}_h^\varepsilon\} = \{[\underline{w}_h^\varepsilon, \underline{\eta}_h^\varepsilon, \underline{\theta}_h^\varepsilon] \mid 0 < \varepsilon < 1, 0 < h \leq h_*\} \\ \quad \subset L^\infty(0, T; H^1(\Omega))^2 \times L^\infty(0, T; H^{N^*}(\Omega)), \\ \{\widehat{v}_h^\varepsilon, \widehat{\theta}_h^\varepsilon\} = \{[\widehat{w}_h^\varepsilon, \widehat{\eta}_h^\varepsilon, \widehat{\theta}_h^\varepsilon] \mid 0 < \varepsilon < 1, 0 < h \leq h_*\} \\ \quad \subset W^{1,\infty}(0, T; H^1(\Omega))^2 \times W^{1,\infty}(0, T; H^{N^*}(\Omega)), \end{array} \right.$$

by using the following different kinds of time-interpolations:

$$\left\{ \begin{array}{l} [\bar{v}_h^\varepsilon(t), \bar{\theta}_h^\varepsilon(t)] = [\bar{w}_h^\varepsilon(t), \bar{\eta}_h^\varepsilon(t), \bar{\theta}_h^\varepsilon(t)] := [v_{h,i}, \theta_{h,i}] = [w_{h,i}, \eta_{h,i}, \theta_{h,i}] \\ \quad \text{in } H^2(\Omega)^2 \times H^{N^*}(\Omega), \\ [\underline{v}_h^\varepsilon(t), \underline{\theta}_h^\varepsilon(t)] = [\underline{w}_h^\varepsilon(t), \underline{\eta}_h^\varepsilon(t), \underline{\theta}_h^\varepsilon(t)] := [\eta_{h,i-1}, \theta_{h,i-1}] = [w_{h,i-1}, \eta_{h,i-1}, \theta_{h,i-1}] \\ \quad \text{in } H^1(\Omega)^2 \times H^{N^*}(\Omega), \\ [\widehat{v}_h^\varepsilon(t), \widehat{\theta}_h^\varepsilon(t)] = [\widehat{w}_h^\varepsilon(t), \widehat{\eta}_h^\varepsilon(t), \widehat{\theta}_h^\varepsilon(t)] := \frac{t_{h,i}-t}{h} [v_{h,i-1}^\varepsilon, \theta_{h,i-1}^\varepsilon] + \frac{t-t_{h,i-1}}{h} [v_{h,i}^\varepsilon, \theta_{h,i}^\varepsilon] \\ \quad \text{in } H^1(\Omega)^2 \times H^{N^*}(\Omega), \end{array} \right.$$

for all $0 < h \leq h_*$ and all $t \in \Delta_{h,i}$, $i = 1, 2, 3, \dots$.

Step 2: finding the approximation limits as $\varepsilon, h \searrow 0$. Taking into account the assumptions (A1)-(A3), (2.6)-(2.7) and (3.1), we can see that:

$$\left\{ \begin{array}{l} \bullet \{ \bar{v}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } L^\infty(0, T; H^1(\Omega))^2, \\ \bullet \{ \underline{v}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } L^\infty(0, T; H^1(\Omega))^2, \\ \bullet \{ \widehat{v}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } W^{1,2}(0, T; L^2(\Omega))^2 \text{ and} \\ \quad \text{bounded in } L^\infty(0, T; H^1(\Omega))^2, \\ \bullet 0 \leq \bar{w}_h^\varepsilon, \bar{\eta}_h^\varepsilon \leq 1, 0 \leq \underline{w}_h^\varepsilon, \underline{\eta}_h^\varepsilon \leq 1 \text{ and } 0 \leq \widehat{w}_h^\varepsilon, \widehat{\eta}_h^\varepsilon \leq 1, \text{ a.e. in } Q, \\ \quad \text{for all } 0 < \varepsilon < 1, 0 < h \leq h_*; \end{array} \right.$$

$$\left\{ \begin{array}{l} \bullet \{ \bar{\theta}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } L^\infty(0, T; H^1(\Omega)), \\ \bullet \{ \underline{\theta}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } L^\infty(0, T; H^1(\Omega)), \\ \bullet \{ \widehat{\theta}_h^\varepsilon \mid 0 < \varepsilon < 1, 0 < h \leq h_* \} \text{ is bounded in } W^{1,2}(0, T; L^2(\Omega)) \text{ and} \\ \quad \text{bounded in } L^\infty(0, T; H^1(\Omega)), \\ \bullet |\bar{\theta}_h^\varepsilon|_{L^\infty(Q)} \leq |\theta_0|_{L^\infty(\Omega)}, |\underline{\theta}_h^\varepsilon|_{L^\infty(Q)} \leq |\theta_0|_{L^\infty(\Omega)} \text{ and } |\widehat{\theta}_h^\varepsilon|_{L^\infty(Q)} \leq |\theta_0|_{L^\infty(\Omega)}, \\ \quad \text{for all } 0 < \varepsilon < 1 \text{ and } 0 < h \leq h_*. \end{array} \right.$$

Therefore, applying the compactness theory of Aubin's type [13], we find sequences $\{\varepsilon_n \mid n \in \mathbb{N}\} \subset (0, 1)$, $\{h_n \mid n \in \mathbb{N}\} \subset (0, h_*]$, and a pair (triplet) of functions $[v, \theta] = [w, \eta, \theta] \in L^2(0, T; L^2(\Omega))^3$, such that:

$$\varepsilon_n \searrow 0 \text{ and } h_n \searrow 0 \text{ as } n \rightarrow \infty,$$

$$\left\{ \begin{array}{l} v = [w, \eta] \in W^{1,2}(0, T; L^2(\Omega))^2 \cap L^\infty(0, T; H^1(\Omega))^2, \quad 0 \leq w, \eta \leq 1 \text{ a.e. in } Q, \\ \theta \in W^{1,2}(0, T; L^2(\Omega)) \cap L^\infty(0, T; H^1(\Omega)), \quad |\theta|_{L^\infty(Q)} \leq |\theta_0|_{L^\infty(\Omega)}, \end{array} \right. \quad (3.2)$$

$$\left\{ \begin{array}{l} \bar{v}_n = [\bar{w}_n, \bar{\eta}_n] := \bar{v}_{h_n}^{\varepsilon_n} \rightarrow v \text{ and } \underline{v}_n = [\underline{w}_n, \underline{\eta}_n] := \underline{v}_{h_n}^{\varepsilon_n} \rightarrow v, \\ \quad \text{in } L^\infty(0, T; L^2(\Omega))^2, \text{ weakly-* in } L^\infty(0, T; H^1(\Omega))^2, \\ \quad \text{weakly-* in } L^\infty(Q)^2, \text{ and in the pointwise sense a.e. in } Q, \\ \widehat{v}_n = [\widehat{w}_n, \widehat{\eta}_n] := \widehat{v}_{h_n}^{\varepsilon_n} \rightarrow v \text{ in } C([0, T]; L^2(\Omega))^2, \\ \quad \text{weakly in } W^{1,2}(0, T; L^2(\Omega))^2, \text{ weakly-* in } L^\infty(0, T; H^1(\Omega))^2, \\ \quad \text{weakly-* in } L^\infty(Q)^2, \text{ and in the pointwise sense a.e. in } Q, \end{array} \right. \quad \text{as } n \rightarrow \infty, \quad (3.3)$$

and

$$\left\{ \begin{array}{l} \bar{\theta}_n := \bar{\theta}_{h_n}^{\varepsilon_n} \rightarrow \theta \text{ and } \underline{\theta}_n := \underline{\theta}_{h_n}^{\varepsilon_n} \rightarrow \theta \text{ in } L^\infty(0, T; L^2(\Omega)), \\ \quad \text{weakly-* in } L^\infty(0, T; H^1(\Omega)), \text{ weakly-* in } L^\infty(Q), \text{ and} \\ \quad \text{in the pointwise sense a.e. in } Q, \\ \widehat{\theta}_n := \widehat{\theta}_{h_n}^{\varepsilon_n} \rightarrow \theta \text{ in } C([0, T]; L^2(\Omega)), \text{ weakly in } W^{1,2}(0, T; L^2(\Omega)), \\ \quad \text{weakly-* in } L^\infty(0, T; H^1(\Omega)), \text{ weakly-* in } L^\infty(Q), \text{ and} \\ \quad \text{in the pointwise sense a.e. in } Q, \end{array} \right. \quad \text{as } n \rightarrow \infty. \quad (3.4)$$

Step 3: verification of (S1)-(S5). From (3.2)-(3.4), we easily check almost all conditions in (S1) and (S5), except for $\eta \in L^2(0, T; H^2(\Omega))$.

Alternatively, the conditions (S2)-(S4) will be verified on the basis of the following (Fact 2) associated with the Γ -convergence of functionals on $L^2(0, T; L^2(\Omega))$.

(Fact 2) A sequence $\{\hat{\Psi}_n \mid n \in \mathbb{N}\}$ of convex functions, defined as:

$$\vartheta \in L^2(0, T; L^2(\Omega)) \mapsto \hat{\Psi}_n(\vartheta) := \int_0^T \Psi_{\varepsilon_n}^\nu(\bar{v}_n(t); \vartheta(t)) dt, \quad \text{for } n \in \mathbb{N},$$

converges to a convex function $\hat{\Phi}_\nu$, defined as:

$$\vartheta \in L^2(0, T; L^2(\Omega)) \mapsto \hat{\Phi}_\nu(\vartheta) := \int_0^T \Phi_\nu(v(t); \vartheta(t)) dt,$$

on $L^2(0, T; L^2(\Omega))$, in the sense of Γ -convergence, as $n \rightarrow \infty$.

In fact, by this Γ -convergence, we obtain (S4), together with the following strong convergence:

$$\bar{\theta}_n \rightarrow \theta, \quad \underline{\theta}_n \rightarrow \theta \quad \text{and} \quad \hat{\theta}_n \rightarrow \theta \quad \text{in } L^2(0, T; H^1(\Omega)), \quad \text{as } n \rightarrow \infty. \quad (3.5)$$

From (3.3)-(3.5), it further follows that:

$$\begin{cases} \int_0^T \int_\Omega \varphi \beta'(\bar{w}_n(t)) |\nabla \underline{\theta}_n(t)|^2 dx dt \rightarrow \int_0^T \int_\Omega \varphi \beta'(w) |\nabla \theta(t)|^2 dx dt, \\ \text{for any } \varphi \in H^1(\Omega) \cap L^\infty(\Omega), \\ \int_0^T \int_\Omega \psi \alpha'(\bar{\eta}_n(t)) |\nabla \underline{\theta}_n(t)| dx dt \rightarrow \int_0^T \int_\Omega \psi \alpha'(\eta) |\nabla \theta(t)| dx dt, \\ \text{for any } \psi \in H^1(\Omega), \end{cases} \quad \text{as } n \rightarrow \infty. \quad (3.6)$$

Taking into account (3.3)-(3.4) and (3.6), we verify (S2)-(S3).

Now, the remaining condition $\eta \in L^2(0, T; H^2(\Omega))$ is a direct consequence from the standard regularity theory of evolution equations (cf. [2]). $\blacksquare$

4 Problems in future

Finally, we mention about the problems in future.

(Problem 1) Further qualitative properties for (S) $_\nu$

Based on the solvability result, it will be possible to verify further qualitative properties for (S) $_\nu$, such as *smoothing effect*, *energy-dissipation*, *large-time behavior*, and so on. Then, the key-point will be how we derive the pointwise convergence of the energy:

$$\begin{aligned} \mathcal{E}_{\varepsilon_n}^\nu(\bar{v}_n(t), \bar{\theta}_n(t)) &\rightarrow \mathcal{F}_\nu(w(t), \eta(t), \theta(t)) \quad \text{and} \quad \mathcal{E}_{\varepsilon_n}^\nu(\underline{v}_n(t), \underline{\theta}_n(t)) \rightarrow \mathcal{F}_\nu(w(t), \eta(t), \theta(t)), \\ &\text{for a.e. } t \in (0, T), \quad \text{as } n \rightarrow \infty, \end{aligned}$$

to obtain the limiting formula for the energy-inequality (2.7) in Theorem 2.1:

$$\begin{aligned} &\frac{1}{2} \int_s^t |v_t(\tau)|_{L^2(\Omega)^2}^2 d\tau + \int_s^t |\sqrt{\alpha_0(v(\tau))} \theta_t(\tau)|_{L^2(\Omega)}^2 d\tau + \mathcal{F}_\nu(w(t), \eta(t), \theta(t)) \\ &\leq \mathcal{F}_\nu(w(s), \eta(s), \theta(t)), \quad \text{for all } 0 \leq s \leq t \leq T. \end{aligned}$$

(Problem 2) The case when $\nu = 0$

This theme will be concerned with the study of association between the limiting situation of (S) $_\nu$ as $\nu \searrow 0$, and the following simplified system, denoted by (S) $_0$.

(S)₀:

$$\begin{cases} w_t - \Delta w + p(w, \eta) = 0 & \text{in } Q, \\ \eta_t - \Delta \eta + q(w, \eta) + \alpha'(\eta)|D\theta| = 0 & \text{in } Q, \\ \alpha_0(w, \eta)\theta_t - \operatorname{div} \left(\alpha(\eta) \frac{D\theta}{|D\theta|} \right) = 0 & \text{in } Q, \end{cases}$$

subject to suitable initial-boundary conditions. Then, analytic methods established in [1, 10] will act key-roles in mathematical treatments of the nonstandard terms $\alpha'(\eta)|D\theta|$ and $-\operatorname{div}(\alpha(\eta) \frac{D\theta}{|D\theta|})$ as in (S)₀.

(Problem 3) The anisotropic case

We may consider the anisotropic versions for the system (S)_ν (actually, also for (S)₀) by modifying the density of the functionals $\Phi_\nu(v; \cdot)$ for $v = [w, \eta] \in H^1(\Omega)^2 \cap L^\infty(\Omega)^2$. However, since *the Wulff shape*, i.e. the structural unit of the crystal, is forced to rotate with the variation of θ , the modified formula of the functional should be settled in the following form:

$$\begin{aligned} \vartheta \in H^1(\Omega) &\mapsto \int_\Omega \alpha(\eta) \gamma(\theta, \nabla \theta) dx + \nu \int_\Omega \beta(w) \gamma(\theta, \nabla \theta)^2 dx, \\ &\text{for } v = [w, \eta] \in H^1(\Omega)^2 \cap L^\infty(\Omega)^2, \end{aligned}$$

where $\gamma : \mathbb{R} \times \mathbb{R}^N \longrightarrow [0, \infty)$ is a given function to characterize the anisotropy.

(Problem 4) The non-isothermal case

According to the modelling method of [9, 14], we can consider full-versions for the systems (S)_ν and (S)₀ (and also for their anisotropic versions), by coupling with the equation for the temperature $u = u(t, x)$. Then, the equation for u should be described in the form of the heat equation:

$$(u + w)_t - \Delta u = h \quad \text{in } Q,$$

with the given forcing term $h = h(t, x)$ and the initial-boundary condition. Also, the perturbations $p = p(w, \eta)$, $q = q(w, \eta)$ and their potential $f = f(w, \eta)$ should be replaced by more interactive forms $p = p(u, w, \eta)$, $q = q(u, w, \eta)$ and $f = f(u, w, \eta)$, respectively.

Conclusive comments. In the mathematical analysis for the above problems, our strategy will be based on the time-discretization approach as presented here. In addition, we expect that similar kinds of approaches might be effective not only for the problems in future but also for mathematical models treated in the relevant previous studies, such as [4, 5, 6, 8, 10, 11, 12, 15].

References

- [1] L. Ambrosio, N. Fusco and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Science Publications, 2000.

- [2] H. Brézis, *Opérateurs Maximaux Monotones et Semi-groupes de Contractions dans les Espaces de Hilbert*, North-Holland Mathematics Studies **5**, Notas de Matemática (50), North-Holland Publishing and American Elsevier Publishing, 1973.
- [3] I. Ekeland and R. Temam, *Convex analysis and variational problems*, Classics in Applied Mathematics, **28**. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [4] A. Ito, N. Kenmochi and N. Yamazaki, A phase-field model of grain boundary motion, *Appl. Math.*, **53** (2008), no. 5, 433–454.
- [5] A. Ito, N. Kenmochi and N. Yamazaki, Weak solutions of grain boundary motion model with singularity, *Rend. Mat. Appl.* (7), **29** (2009), no. 1, 51–63.
- [6] A. Ito, N. Kenmochi and N. Yamazaki, Global solvability of a model for grain boundary motion with constraint, *Discrete Contin. Dyn. Syst. Ser. S*, **5** (2012), no. 1, 127–146.
- [7] N. Kenmochi, Y. Mizuta and T. Nagai, Projections onto convex sets, convex functions and their subdifferentials, *Bull. Fac. Edu.*, Chiba Univ., **29** (1980), 11–22.
- [8] N. Kenmochi and N. Yamazaki, Large-time behavior of solutions to a phase-field model of grain boundary motion with constraint, *Current advances in nonlinear analysis and related topics*, 389–403, GAKUTO Internat. Ser. Math. Sci. Appl., **32**, Gakkōtoshō, Tokyo, 2010.
- [9] R. Kobayashi, Modelling of grain structure evolution, *Variational Problems and Related Topics*, RIMS Kōkyūroku, **1210** (2001), 68–77.
- [10] S. Moll and K. Shirakawa, Existence of solutions to the Kobayashi-Warren-Carter system, *Calculus of Variations and Partial Differential Equations* (to appear).
- [11] K. Shirakawa and H. Watanabe, Energy-dissipative solution to a one-dimensional phase field model of grain boundary motion, *Discrete Contin. Dyn. Syst. Ser. S*, **7** (2014), no. 1, 139–159. DOI:10.3934/dcdss.2014.7.139
- [12] K. Shirakawa, H. Watanabe and N. Yamazaki, Solvability for one-dimensional phase field system associated with grain boundary motion, *Math. Ann.* (2013), **356**, 301–330. DOI: 10.1007/s00208-012-0849-2
- [13] J. Simon, Compact set in the space $L^p(0, T; B)$, *Ann. Mat. Pura Appl.* (4), **146** (1987), 65–96.
- [14] J. A. Warren, R. Kobayashi, A. E. Lobkovski and W. C. Carter, Extending phase field models of solidification to polycrystalline materials, *Acta Materialia*, **51** (2003), 6035–6058.
- [15] H. Watanabe and K. Shirakawa, Qualitative properties of a one-dimensional phase-field system associated with grain boundary, submitted to *GAKUTO Internat. Ser. Math. Sci. Appl.*.

A CANCELLATION PROPERTY AND THE WELL-POSEDNESS FOR THE PERIODIC KDV EQUATIONS

TAKAMORI KATO

JOINT WORK WITH KOTARO TSUGAWA (NAGOYA UNIVERSITY)

In this paper, we consider the Cauchy problem of the fifth order KdV equation in the periodic setting $\mathbb{T} := \mathbb{R}/2\pi\mathbb{Z}$ as follows:

$$\begin{cases} \partial_t u + a_1 \partial_x^5 u + c_1 \partial_x (\partial_x u)^2 + c_2 \partial_x^2 (u \partial_x u) + c_3 \partial_x (u^3) = 0, & (t, x) \in \mathbb{R} \times \mathbb{T}, \\ u|_{t_0} = u_0 \in H^s(\mathbb{T}) \end{cases} \quad (1)$$

where an unknown function $u = u(t, x)$ is real valued and $a_1, c_1, c_2, c_3 \in \mathbb{R}$ with $a_1 \neq 0$. Such equation and its generalizations

$$\partial_t u + a_0 \partial_x^3 u + a_1 \partial_x^5 u + c_0 \partial_x (u^2) + c_1 \partial_x (\partial_x u)^2 + c_2 \partial_x^2 (u \partial_x u) + c_3 \partial_x (u^3) = 0$$

arise as long-wave approximations to the water wave equations like to the KdV equation

$$\partial_t u + \partial_x^3 u + \partial_x (u^2) = 0 \quad (2)$$

(see Craig, Guyenne and Kalisch [5], Oiver [9] and so on). By the scaling $u(t, x) \mapsto |c_3/10a_0|^{1/2} u(t/a_0, x)$, (1) is rewritten into

$$\partial_t u + \partial_x^5 u + \alpha \partial_x (\partial_x u)^2 + \beta \partial_x^2 (u \partial_x^2 u) + 10\gamma \partial_x (u^3) = 0 \quad (3)$$

where $\alpha, \beta \in \mathbb{R}$ and $\gamma = \text{sgn } c_3/a_0 = -1, 0$ or 1 . In the special case $(\alpha, \beta, \gamma) = (-5, 10, 1)$ or $(5, -10, 1)$, the equations in (3) are in the KdV hierarchy discovered by Lax [8]. In this case, (3) is complete integrable and has an infinite number of conservation laws, which implies that (3) has rich structure. Put

$$E_0(u(t)) := \frac{1}{2\pi} \int_{\mathbb{T}} u(t) \, dx, \quad E_1(u(t)) := \frac{1}{2\pi} \int_{\mathbb{T}} u^2(t) \, dx.$$

E_0 and E_1 are conserved if and only if the following condition holds.

$$2\alpha + \beta = 0. \quad (4)$$

Note that in the case (4), the Hamiltonian

$$H(u(t)) := \frac{1}{2\pi} \int_{\mathbb{T}} (\partial_x^2 u(t))^2 + 2\alpha u(t)(\partial_x u(t))^2 + 5\gamma u^4(t) \, dx$$

is well-defined. In this paper, we only consider the Hamiltonian case, namely

$$\partial_t u + \partial_x^5 u + \alpha \partial_x (\partial_x u)^2 - 2\alpha \partial_x^2 (u \partial_x u) + 10\gamma \partial_x (u^3) = 0. \quad (5)$$

In the periodic case, there are many results on the well-posedness of the KdV equation (2) but no results on that of the fifth order KdV equation (5) when initial data is given in $H^s(\mathbb{T})$. Therefore our aim is to establish the method reflecting the algebraic structure and show the well-posedness of the Cauchy problem for (5) in $H^s(\mathbb{T})$.

The strong singularity in the nonlinear terms having some derivatives makes the problem difficult. This is called a loss of derivatives. It is the most important point in this study how to cancel a loss of three derivatives in the quadratic terms of (5). We can divide problem into two parts. First one is how to cancel the resonant parts with a loss of derivatives (For the definition of the resonant parts, see below). Second one is how to recover three derivatives on the non-resonant parts. To overcome this difficulty, we try to use the Fourier restriction norm method established by Bourgain [3] or the normal form method (differentiation by parts). Since there are the resonant parts with a loss of derivatives, the solution map associated to (5) fails to be C^3 in $H^s(\mathbb{T})$ for any $s \in \mathbb{R}$. As a consequence, we cannot solve the Cauchy problem of (5) by a Picard iteration method in $H^s(\mathbb{T})$ with $s \in \mathbb{R}$. This is because there are no smoothing effects associated to the linear parts in the periodic case unlike $\mathbb{R}$ case. Here we give the definition of the resonant parts and the non-resonant parts. We consider the following integral equation

$$u(t) = e^{-t\partial_x^5} u(0) + \int_0^t e^{-(t-s)\partial_x^5} P^{(N)}(u, \partial_x u, \dots, \partial_x^l u)(s) ds$$

where $P^{(N)}$ is an N th homogeneous polynomial. We apply a change of coordinates: $v(t) = e^{t\partial_x^5} u(t)$. We substitute v into the above integral equation and use the spatial Fourier series expansion to obtain

$$\hat{v}(t, k) = \hat{v}(0, k) + \int_0^t \sum_{k=k_1+k_2+\dots+k_N} e^{s\Phi_N} m^{(N)}(k_1, k_2, \dots, k_N) \prod_{i=1}^N \hat{v}(s, k_i) ds$$

where $m^{(N)}$ is the N th multiplier and Φ_N is the oscillation term defined as

$$\Phi_N = i \left[\left(\sum_{i=1}^N k_i \right)^5 - \sum_{i=1}^N k_i^5 \right].$$

The resonant parts are expressed by

$$\Omega_R^{(N)} := \{ (k_1, k_2, \dots, k_N) \in \mathbb{Z}^N : k = k_{1,2,\dots,N}, \Phi_N = 0 \}$$

and the non-resonant parts are defined as

$$\Omega_{NR}^{(N)} := \{(k_1, k_2, \dots, k_N) \in \mathbb{Z}^N : k = k_{1,2,\dots,N}, \Phi_N \neq 0\}$$

where $k_{1,2,\dots,N} := k_1 + k_2 + \dots + k_N$. Note that the oscillation terms Φ_2 and Φ_3 can be factorized exactly as follows:

$$\begin{aligned}\Phi_2 &= i\frac{5}{2}k_1k_2k_{1,2}(k_1^2 + k_2^2 + k_{1,2}^2), \\ \Phi_3 &= i\frac{5}{2}k_{1,2}k_{2,3}k_{3,1}(k_{1,2}^2 + k_{2,3}^2 + k_{3,1}^2),\end{aligned}$$

which implies that the resonant cases are $k_1k_2k_{1,2} = 0$ and $k_{1,2}k_{2,3}k_{3,1} = 0$ for $N = 2$ and $N = 3$, respectively.

Let us briefly go over recent results on the well-posedness of the periodic KdV equation (2). Bourgain [3] proved the local well-posedness (LWP) of (2) for $s \geq 0$. His idea is to remove the resonant parts by using the conservation laws $E_0(u(t)) = E_0(u_0)$ and modifying the linear parts. Actually, with $v(t) = e^{t\partial_x^3}u(t)$, it follows from (2) that v satisfies

$$\partial_t \hat{v} = - \sum_{k=k_1+k_2} e^{-t\Phi_3} i k_{1,2} \prod_{i=1}^2 \hat{v}(t, k_i)$$

where $\Phi_3 = i(k_{1,2}^3 - k_1^3 - k_2^3) = 3ik_1k_2k_{1,2}$. This implies that the resonant case is $k_1k_2k_{1,2} = 0$. Note that

$$\hat{v}(t, 0) = \hat{u}(t, 0) = \frac{1}{2\pi} \int_{\mathbb{T}} u(t) dx = E_0(u_0).$$

(2) is rewritten into

$$\partial_t u + \partial_x^3 u + 2E_0(u_0)\partial_x u = -\partial_x \left(u - \frac{1}{2\pi} \int_{\mathbb{T}} u dx \right)^2. \quad (6)$$

The nonlinear term of (6) is the non-resonant parts. A loss of one derivative in the non-resonant parts can be recovered by the Fourier restriction norm method. Bourgain's result was improved to $s \geq -1/2$ by Kenig, Ponce and Vega [6] (see also [4]). Later they also the solution map associated to (2) fails to be uniformly continuous when $s < -1/2$. As a consequence, $s = -1/2$ is the critical regularity in this sense. Moreover Babin, Ilyin and Titi [1] proved LWP and the unconditional uniqueness in $C([0, T] : H^s(\mathbb{T}))$ for (2) when $s \geq 0$. The unconditional uniqueness means that uniqueness holds in the whole of $C([0, T] : H^s(\mathbb{T}))$ without using any auxiliary function space. On the other hand, the nonlinear term $\partial_x(u^2)$ cannot be defined in the distribution sense for $s < 0$. Therefore this result is optimal in the

above sense. Their idea is to recover one derivative in (6) by the normal form method (differentiation by parts). In fact, it follows from (6) that $v(t) = e^{t\partial_x^3}u(t)$ satisfies

$$\partial_t \hat{v}(t, k) = - \sum_{\Phi_2 \neq 0} e^{-t\Phi_2} i k_{1,2} \prod_{i=1}^2 \hat{v}(t, k_i). \quad (7)$$

We can differentiate the non-resonant parts (7) by parts to have

$$\partial_t \hat{v}(t, k) = \frac{\partial}{\partial t} \left[\sum_{\Phi_2 \neq 0} e^{-t\Phi_2} \frac{i k_{1,2}}{\Phi_2} \prod_{i=1}^2 \hat{v}(t, k_i) \right] \quad (8)$$

$$- \sum_{\Phi_2 \neq 0} e^{-t\Phi_2} \frac{i k_{1,2}}{\Phi_2} (\partial_t \hat{v}(t, k_1) \hat{v}(t, k_2) + \hat{v}(t, k_1) \partial_t \hat{v}(t, k_2)). \quad (9)$$

Both (8) and (9) have $\Phi_2 (\neq 0)$ in the denominators, and this provides smoothing. As a consequence, we can obtain the nonlinear terms with no loss by the above procedure i.e. the normal form method. Note that at most one derivative can be recovered by using the normal form method once.

(5) has the nonlinear terms with a loss of three derivatives. But at most two derivatives can be recovered by the Fourier restriction norm method. We apply the normal form method three times to recover three derivatives and obtain the following results.

Theorem 1. *Let $s \geq 1$. Then (5) is unconditionally locally well-posed in $H^s(\mathbb{T})$. Namely, for any $u_0 \in H^s(\mathbb{T})$, there exist $T = T(\|u_0\|_{H^s}) > 0$ and the unique solution $u \in C([0, T] : H^s(\mathbb{T}))$ to (5). Moreover the solution map, $H^s(\mathbb{T}) \ni u_0 \mapsto u \in C([0, T] : H^s(\mathbb{T}))$, is continuous.*

Theorem 2. *Let $s \geq 1/2$. (5) is locally well-posed in $X_w^s([0, T]) \cap C([0, T] : H^s(\mathbb{T}))$ in the sense of Theorem 1. X_w^s is the modified $X^{s,b}$ space defined below.*

Remark 1. When $s < 1$, the nonlinear term $\partial_x(\partial_x u)^2$ cannot be defined in the distribution sense. Therefore Theorem 1 is optimal in this sense.

Remark 2. In the case $s \geq 2$, the Hamiltonian of (5) is well-defined. The conserved quantities E_1 and H provide a control of the H^2 norm and allow to prove (5) is unconditionally global well-posed in H^2 .

Remark 3. The results of Theorems 1 and 2 are corresponding to those of Babin, Ilyin and Titi [1] and Kenig, Ponce and Vega [6].

Remark 4. The results of Theorems 1 and 2 are also valid for the following equation and their proofs are similar.

$$\partial_t u + a_0 \partial_x^3 u + \partial_x^5 u + c_0 \partial_x(u^2) + \alpha \partial_x(\partial_x u)^2 - 2\alpha \partial_x^2(u \partial_x u) = 0.$$

The key idea in the proof of the main results is to cancel the resonant parts with a loss of derivatives by using the conserved quantities E_0 and E_1 and modifying the linear parts of (5). In fact, if E_0 and E_1 are conserved, then (5) is rewritten into

$$\begin{aligned} \partial_t u - \partial_x^5 u - 2\alpha E_0(u_0) \partial_x^3 u + (30\gamma - \frac{4}{5}\alpha^2) E_1(u_0) \partial_x u + \frac{4}{5}\alpha^2 E_0^2(u_0) \partial_x u \\ = -30\gamma \left(u^2 - \frac{1}{2\pi} \int_{\mathbb{T}} u^2 dx \right) \partial_x u \end{aligned} \quad (10)$$

$$- \alpha \partial_x(\partial_x u)^2 + 2\alpha \partial_x^2 \left\{ \left(u - \frac{1}{2\pi} \int_{\mathbb{T}} u dx \right) \partial_x u \right\} \quad (11)$$

$$- \frac{4}{5}\alpha^2 \left\{ \frac{1}{2\pi} \int_{\mathbb{T}} u^2 dx - \left(\frac{1}{2\pi} \int_{\mathbb{T}} u dx \right)^2 \right\} \partial_x u. \quad (12)$$

Recall the resonant parts in the case $N = 2, 3$. There are no resonant parts with a loss of derivatives in the nonlinear terms except the resonant parts (12). Therefore we can apply the normal form method to (10) and (11) once. However, to cancel a loss of three derivatives in (11), we need to apply the normal form method three times. So it is necessary that the resonant parts with a loss of derivatives coming from the normal form reduction of (11) are canceled with the resonant parts (12). This is the most interesting point in our study. All nonlinear terms with no loss coming from the normal form reduction can be estimated by only Sobolev's embedding theorem when $s \geq 1$. As a consequence, we obtain the claim of Theorem 1 by a Picard iteration method.

The claim of Theorem 1 can be extended to $s \geq 1/2$ by introducing the function space X_w^s . This is the modified $X^{s,b}$ space such as Bejenaru and Tao [2] and equipped with the norm as follows:

$$\|u\|_{X_w^s} := \|w_{u_0}^s(\tau, k) \hat{u}\|_{l_k^2 L_\tau^2}$$

Here the weight function $w_{u_0}^s$ is defined as follows;

$$w_{u_0}^s(\tau, k) = \begin{cases} \langle k \rangle^s \langle \tau - p_{u_0}(k) \rangle^{1/4}, & \text{when } |k|^4 \leq |\tau - p_{u_0}(k)| \leq |k|^5 \\ \langle k \rangle^s \langle \tau - p_{u_0}(k) \rangle^{1/2}, & \text{otherwise} \end{cases}.$$

where

$$p_{u_0}(k) = k^5 - 2\alpha E_0(u_0)k^3 + (30\gamma - \frac{4}{5}\alpha^2)E_1(u_0)k + \frac{4}{5}\alpha^2 E_0^2(u_0)k.$$

REFERENCES

- [1] A. Babin, A. Ilyin and E. Titi, *On the regularization mechanism for the periodic Korteweg-de Vries equation*, Comm. Pure. Appl. Math. **64** (2011), no. 5, 591–648.
- [2] J. Bejenaru and T. Tao, *Sharp well-posedness and ill-posedness results for a quadratic nonlinear Schrödinger equation*, J. Funct. Anal. **233** (2006), 228–259.
- [3] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation*, Geom. Funct. Anal. **3** (1993), no. 3, 209–262.
- [4] J. Colliander, M. Keel, G. Staffilani, H. Takaoka and T. Tao, *Sharp global well-posedness for KdV and modified KdV on $\mathbf{R}$ and $\mathbf{T}$* , J. Amer. Math. Soc. **16** (2003), no. 3, 705–749.
- [5] W. Craig, P. Guyenne and H. Kalisch, *Hamiltonian long wave expansions for free surfaces and interfaces*, Comm. Pure Anal. Math., **58** (2005), 1587–1641.
- [6] C. E. Kenig, G. Ponce and L. Vega, *A bilinear estimate with applications to the KdV equation*, J. Amer. Math. Soc. **9** (1996), no. 2, 573–603.
- [7] C. E. Kenig, G. Ponce and L. Vega, *On the ill-posedness of some canonical dispersive equations*, Duke Math. J. **106** (2001), no. 3, 617–633.
- [8] P. Lax, *Integrals of nonlinear equations of evolutions and solitary waves*, Comm. Pure Appl. Math., **21** (1968), 467–490.
- [9] P. J. Olver, *Hamiltonian and non-Hamiltonian models for water-waves*, Lecture Notes in Physics, Springer, Berlin, **195** (1995), 273–290.

(Takamori Kato) INSTITUTE FOR ADVANCED RESEARCH, NAGOYA UNIVERSITY CHIKUSA-KU,
NAGOYA, 464-8601, JAPAN

E-mail address, Takamori Kato: `kato.takamori@math.nagoya-u.ac.jp`

Resolvent estimates for the Stokes equations in spaces of bounded functions

Ken Abe (University of Tokyo)*

Abstract

The Stokes equation is well understood in the L^p -setting for a large class of domains including bounded and exterior domains with smooth boundaries provided $1 < p < \infty$. The situation is different for the case $p = \infty$ since in this case the Helmholtz projection does not act as a bounded operator anymore. Nevertheless, it was recently shown by a contradiction argument that the Stokes operator generates an analytic semigroup on L^∞ -type spaces for a large class of domains. In this talk, we present a new approach as well as new a priori estimate to the resolvent Stokes equation. They in particular implies that the Stokes operator generates an analytic semigroup of angle $\pi/2$ on L^∞ -type spaces for a large class of domains. This talk is based on a joint work with Y. Giga and M. Hieber.

1 Introduction

It is well known that the Stokes semigroup is an analytic semigroup on L^p -solenoidal spaces, $p \in (1, \infty)$, for various kind of domains, e.g., bounded and exterior domains with smooth boundaries [8], [5]. However, it was unknown whether or not the Stokes semigroup is an analytic semigroup on L^∞ -type spaces except a half space where explicit solution formulas are available [4], [9], [6]. Recently, an affirmative answer to this problem was given for a large class of domains including bounded and exterior domains based on an a priori L^∞ -estimate for solutions to the non-stationary Stokes equations [1], [2].

In this talk, we present a new approach as well as new a priori estimate for the resolvent Stokes equations. We consider the resolvent Stokes equations in the domain $\Omega \subset \mathbb{R}^n$, $n \geq 2$:

$$\lambda v - \Delta v + \nabla q = f \quad \text{in } \Omega, \tag{1.1}$$

$$\operatorname{div} v = 0 \quad \text{in } \Omega, \tag{1.2}$$

$$v = 0 \quad \text{on } \partial\Omega, \tag{1.3}$$

*kabe@ms.u-tokyo.ac.jp

and establish a bound for

$$M_p(v, q)(x, \lambda) = |\lambda| |v(x)| + |\lambda|^{1/2} |\nabla v(x)| + |\lambda|^{n/2p} \|\nabla^2 v\|_{L^p(\Omega_{x,|\lambda|^{-1/2}})} + |\lambda|^{n/2p} \|\nabla q\|_{L^p(\Omega_{x,|\lambda|^{-1/2}})},$$

and $p > n$ of the form,

$$\sup_{\lambda \in \Sigma_{\vartheta, \delta}} \|M_p(v, q)\|_{L^\infty(\Omega)}(\lambda) \leq C \|f\|_{L^\infty(\Omega)} \quad (1.4)$$

for some constant $C > 0$ independent of f . Here, $\Omega_{x,r}$ denotes the intersection of Ω with an open ball $B_x(r)$ centered at $x \in \Omega$ with radius $r > 0$, and $\Sigma_{\vartheta, \delta}$ denotes the sectorial region in the complex plane given by $\Sigma_{\vartheta, \delta} = \{\lambda \in \mathbb{C} \setminus \{0\} \mid |\arg \lambda| < \vartheta, |\lambda| > \delta\}$ for $\vartheta \in (\pi/2, \pi)$ and $\delta > 0$. The estimate (1.4) in particular implies that the Stokes operator generates an analytic semigroup on L^∞ -type spaces.

Our approach is inspired by the corresponding approach for general elliptic operators. K. Masuda was the first to prove the analyticity of the semigroup associated to general elliptic operators in the whole space [7]. This result was then extended by H. B. Stewart to the case for the Dirichlet problem [10], [11]. By now, this Masuda-Stewart method applies to many other situations. However, it seems that their localization argument does not directly apply to the resolvent Stokes equations (1.1)–(1.3) because of the presence of pressure.

In order to prove the estimate (1.4), we apply *the harmonic-pressure gradient estimate* introduced in [1], [2], i.e.,

$$\sup_{x \in \Omega} d_\Omega(x) |\nabla q(x)| \leq C_\Omega \|W(v)\|_{L^\infty(\partial\Omega)}, \quad (1.5)$$

for $W(v) = -(\nabla v - \nabla^T v) n_\Omega$. When $n = 3$, $W(v)$ is nothing but the tangential component of vorticity, i.e., $-\text{curl } v \times n_\Omega$. The estimate (1.5) plays a key role in transferring results from the elliptic situation to the situation of the Stokes system. Here, n_Ω denotes the unit outward normal vector field on $\partial\Omega$ and d_Ω denotes the distance from $x \in \Omega$ to the boundary $\partial\Omega$. A key observation is that the Neumann data of the pressure q is transformed into the surface divergence, i.e., $\Delta v \cdot n_\Omega = \text{div}_{\partial\Omega} W(v)$ as $\text{div } v = 0$ in Ω . So the estimate (1.5) is reduced to investigating an a priori estimate for solutions of the homogeneous Neumann problem:

$$\Delta q = 0 \text{ in } \Omega, \quad \frac{\partial q}{\partial n_\Omega} = \text{div}_{\partial\Omega} W \text{ on } \partial\Omega.$$

Of course, the estimate (1.5) holds for a half space. It is proved in [1], [2] by a blow-up argument that the estimate (1.5) holds for bounded and exterior domains with C^3 -boundaries.

To state a result, let $C_{0,\sigma}(\Omega)$ be the L^∞ -closure of all smooth solenoidal vector fields with compact support in Ω . Let $L_d^\infty(\Omega)$ be the space of all locally integrable functions f such that $d_\Omega f$ is bounded in Ω . Our typical main result is:

Theorem 1.1. *Let Ω be a bounded or an exterior domain in $\mathbb{R}^n$, $n \geq 2$, with C^3 -boundary. Let $p > n$. For $\vartheta \in (\pi/2, \pi)$ there exists constants δ and C such that the a priori estimate (1.4) holds for all solutions $(v, \nabla q) \in W_{\text{loc}}^{2,p}(\Omega) \times (L_{\text{loc}}^p \cap L_d^\infty)(\Omega)$ for $f \in C_{0,\sigma}(\Omega)$ and $\lambda \in \Sigma_{\vartheta, \delta}$.*

The a priori estimates (1.4) and (1.5) imply that the solution operator $R(\lambda) : f \mapsto v_\lambda$ is invertible on $C_{0,\sigma}$, i.e., there exists a closed operator A such that $R(\lambda) = (\lambda - A)^{-1}$. We call A *the Stokes operator* in $C_{0,\sigma}$. The estimate (1.4) says that the Stokes operator A is a sectorial operator in $C_{0,\sigma}$.

Theorem 1.2. *Let Ω be a bounded or an exterior domain with C^3 -boundary. Then, the Stokes operator A generates a C_0 -analytic semigroup of angle $\pi/2$ on $C_{0,\sigma}(\Omega)$.*

References

- [1] K. Abe, Y. Giga, Analyticity of the Stokes semigroup in spaces of bounded functions, *Acta Math.*, to appear
- [2] K. Abe, Y. Giga, The L^∞ -Stokes semigroup in exterior domains, *J. Evol. Equ.*, to appear
- [3] K. Abe, Y. Giga, M. Hieber, Stokes resolvent estimates in spaces of bounded functions. Hokkaido University Preprint Series in Mathematics, no.1022 (2012)
- [4] W. Desch, M. Hieber, J. Prüss, L^p -theory of the Stokes equation in a half space, *J. Evol. Equ.*, **1** (2001), 115–142.
- [5] Y. Giga, Analyticity of the semigroup generated by the Stokes operator in L_r spaces, *Math. Z.* **178** (1981), 297–329.
- [6] P. Maremonti, G. Starita, Nonstationary Stokes equations in a half-space with continuous initial data, *Zapiski Nauchnykh Seminarov POMI*, **295** (2003) 118–167; translation: *J. Math. Sci. (N.Y.)* **127** (2005), 1886–1914.
- [7] K. Masuda, On the generation of analytic semigroups of higher-order elliptic operators in spaces of continuous functions (in Japanese), *Proc. Katata Symposium on Partial Differential Equations* (1972), 144–149.
- [8] V. A. Solonnikov, Estimates for solutions of nonstationary Navier-Stokes equations, *J. Soviet Math.* **8** (1977), 467–529.
- [9] V. A. Solonnikov, On nonstationary Stokes problem and Navier-Stokes problem in a half-space with initial data nondecreasing at infinity, *J. Math. Sci. (N. Y.)*, **114** (2003), 1726–1740.
- [10] H. B. Stewart, Generation of analytic semigroups by strongly elliptic operators, *Trans. Amer. Math. Soc.*, **199** (1974), 141–162.
- [11] H. B. Stewart, Generation of analytic semigroups by strongly elliptic operators under general boundary conditions, *Trans. Amer. Math. Soc.*, **259** (1980), 299–310.

Global existence for semilinear wave equations with the blow-up term in high dimensions *

Kyouhei Wakasa

Department of Mathematics, Hokkaido University,
Sapporo, Hokkaido 060-0810, Japan.
e-mail : wakasa@math.sci.hokudai.ac.jp

1 General theory for nonlinear wave equations

First we shall outline the general theory on the initial value problem for fully nonlinear wave equations,

$$\begin{cases} u_{tt} - \Delta u = H(u, Du, D_x Du) & \text{in } \mathbf{R}^n \times [0, \infty), \\ u(x, 0) = \varepsilon f(x), \quad u_t(x, 0) = \varepsilon g(x), \end{cases} \quad (1)$$

where $u = u(x, t)$ is a scalar unknown function of space-time variables,

$$\begin{aligned} Du &= (u_{x_0}, u_{x_1}, \dots, u_{x_n}), \quad x_0 = t, \\ D_x Du &= (u_{x_i x_j}, \quad i, j = 0, 1, \dots, n, \quad i + j \geq 1), \end{aligned}$$

$f, g \in C_0^\infty(\mathbf{R}^n)$ and $\varepsilon > 0$ is a “small” parameter. We note that it is impossible to construct a general theory for “large” ε due to blow-up results. For example, see Glassey [4], Levine [7], or Sideris [17]. Let

$$\widehat{\lambda} = (\lambda; (\lambda_i), i = 0, 1, \dots, n; (\lambda_{ij}), i, j = 0, 1, \dots, n, \quad i + j \geq 1).$$

Suppose that the nonlinear term $H = H(\widehat{\lambda})$ is a sufficiently smooth function with

$$H(\widehat{\lambda}) = O(|\widehat{\lambda}|^{1+\alpha})$$

in a neighborhood of $\widehat{\lambda} = 0$, where $\alpha \geq 1$ is an integer. Let us define the lifespan $\widetilde{T}(\varepsilon)$ of classical solutions of (1) by

$$\widetilde{T}(\varepsilon) = \sup\{t > 0 : \exists \text{ classical solution } u(x, t) \text{ of (1) for arbitrarily fixed data, } (f, g).\}.$$

*This is a joint work with Professor Hiroyuki Takamura, Department of Complex and Intelligent Systems Faculty of Systems Information Science, Future University Hakodate 116-2 Kamedanakano-cho, Hakodate, Hokkaido 041-8655, Japan. e-mail : takamura@fun.ac.jp. This talk is presented in The 38th Sapporo Symposium on Partial Differential Equations at Hokkaido University on August 22, 2013.

When $\tilde{T}(\varepsilon) = \infty$, the problem (1) admits a global in time solution, while we only have a local in time solution on $t \in [0, \tilde{T}(\varepsilon))$ when $\tilde{T}(\varepsilon) < \infty$. For local in time solutions, one can measure the long time stability of a zero solution by orders of ε . Because the uniqueness of the solution of (1) may yield that $\lim_{\varepsilon \rightarrow +0} \tilde{T}(\varepsilon) = \infty$. Such an uniqueness theorem can be found in Appendix of John [12] for example.

In Chapter 2 of Li and Chen [9], we have long histories on the estimate for $\tilde{T}(\varepsilon)$. The lower bounds of $\tilde{T}(\varepsilon)$ are summarized in the following table. Let $a = a(\varepsilon)$ satisfy

$$a^2 \varepsilon^2 \log(a + 1) = 1 \quad (2)$$

and c stands for a positive constant independent of ε . Then, due to the fact that it is impossible to obtain an L^2 estimate for u itself by standard energy methods, we have

$\tilde{T}(\varepsilon) \geq$	$\alpha = 1$	$\alpha = 2$	$\alpha \geq 3$
$n = 2$	$ca(\varepsilon)$ in general case, $c\varepsilon^{-1}$ if $\int_{\mathbf{R}^2} g(x)dx = 0$, $c\varepsilon^{-2}$ if $\partial_u^2 H(0) = 0$	$c\varepsilon^{-6}$ in general case, $c\varepsilon^{-18}$ if $\partial_u^3 H(0) = 0$, $\exp(c\varepsilon^{-2})$ if $\partial_u^3 H(0) = \partial_u^4 H(0) = 0$	∞
$n = 3$	$c\varepsilon^{-2}$ in general case, $\exp(c\varepsilon^{-1})$ if $\partial_u^2 H(0) = 0$	∞	∞
$n = 4$	$\exp(c\varepsilon^{-2})$ in general case, ∞ if $\partial_u^2 H(0) = 0$	∞	∞
$n \geq 5$	∞	∞	∞

The result for $n = 1$ is that

$$\tilde{T}(\varepsilon) \geq \begin{cases} \varepsilon^{-\alpha/2} & \text{in general case,} \\ c\varepsilon^{-\alpha(1+\alpha)/(2+\alpha)} & \text{if } \int_{\mathbf{R}} g(x)dx = 0, \\ c\varepsilon^{-\alpha} & \text{if } \partial_u^\beta H(0) = 0 \text{ for } 1 + \alpha \leq \forall \beta \leq 2\alpha. \end{cases} \quad (3)$$

For references on these results, see Li and Chen [9]. We shall skip to refer them here. But we note that two parts in this table are different from the one in Li and Chen [9]. One is the general case in $(n, \alpha) = (4, 1)$. In this part, the lower bound of $\tilde{T}(\varepsilon)$ is $\exp(c\varepsilon^{-1})$ in Li and Chen [9]. But later, it has been improved by Li and Zhou [10]. Another is the case for $\partial_u^3 H(0) = 0$ in $(n, \alpha) = (2, 2)$. This part is due to Katayama [14]. But it is missing in Li and Chen [9]. Its reason is closely related to the sharpness of results in the general theory. The sharpness is achieved by the fact that there is no possibility to improve the lower bound of $\tilde{T}(\varepsilon)$ in sense of order of ε by blow-up results for special equations and special data. It is expressed in the upper bound of $\tilde{T}(\varepsilon)$ with the same order of ε as

the lower bound. On this matter, Li and Chen [9] says that all these lower bounds are known to be sharp except for $(n, \alpha) = (4, 1)$. But before this article, Li [8] says that $(n, \alpha) = (2, 2)$ has also open sharpness while the case for $\partial_u^3 H(0) = 0$ is still missing. Li and Chen [9] might have dropped the open sharpness in $(n, \alpha) = (2, 2)$ by conjecture that $\partial_u^4 H(0) = 0$ is a technical condition. No one disagrees with this observation because the model case of $H = u^4$ has a global solution in two space dimensions, $n = 2$. However, surprisingly, Zhou and Han [23] have obtained this final sharpness in $(n, \alpha) = (2, 2)$ by studying $H = u_t^2 u + u^4$. This puts Katayama's result into the table after 20 years from Li and Chen [9]. We note that Zhou and Han [22] have also obtained the sharpness of the case for $\partial_u^3 H(0) = \partial_u^4 H(0) = 0$ in $(n, \alpha) = (2, 2)$ by studying $H = u_t^3$. This part had been verified by $H = |u_t|^3$ only.

We now turn back to another open sharpness of the general case in $(n, \alpha) = (4, 1)$. It has been obtained by our previous work, Takamura and Wakasa [19], by studying model case of $H = u^2$. This part had been open more than 20 years in the analysis on the critical case for model equations, $u_{tt} - \Delta u = |u|^p$ ($p > 1$). In this way, the general theory and its optimality have been completed.

2 The final problem and related results

After the completion of the general theory, we are interested in “almost” global existence, namely, the case where $\tilde{T}(\varepsilon)$ has an lower bound of the exponential function of ε with a negative power. Such a case appears in $(n, \alpha) = (2, 2), (3, 1), (4, 1)$ in the table of the general theory. It is remarkable that Klainerman [15] and Christodoulou [3] have independently found a special structure on $H = H(Du, D_x Du)$ in $(n, \alpha) = (3, 1)$ which guarantees the global existence. This algebraic condition on nonlinear terms of derivatives of the unknown function is so-called “null condition”. It has been also established by Godin [5] for $H = H(Du)$ and Katayama [13] for $H = H(Du, D_x Du)$ in $(n, \alpha) = (2, 2)$. The null condition has been supposed to be not sufficient for the global existence in $(n, \alpha) = (2, 2)$. Finally Hoshiga [6] and Kubo [16] have independently succeeded to establish “non-positive” condition in this case for $H = H(Du)$. It might be necessary and sufficient condition to the global existence. On the other hand, the situation in $(n, \alpha) = (4, 1)$ is completely different from $(n, \alpha) = (2, 2), (3, 1)$ because H has to include u^2 .

In the sense of the first section, one of the final open problem on the optimality of the general theory for fully nonlinear wave equations can be established by model problem;

$$\begin{cases} u_{tt} - \Delta u = u^2 & \text{in } \mathbf{R}^4 \times [0, \infty), \\ u(x, 0) = \varepsilon f(x), \quad u_t(x, 0) = \varepsilon g(x). \end{cases} \quad (4)$$

We note that this is an extended problem of John [11] to high dimensional case which has the “critical” exponent of Strauss' conjecture [18]. The lifespan $T(\varepsilon)$ of the solution of (4) should have an estimate of the form;

$$\exp(c\varepsilon^{-2}) \leq T(\varepsilon) \leq \exp(C\varepsilon^{-2}). \quad (5)$$

This final problem on the upper bound has been solved by our previous work, Takamura and Wakasa [19]. In its proof, the analysis on $\|u(\cdot, t)\|_{L^2(\mathbf{R}^4)}^2$ is a key because we cannot use any pointwise estimate of the solution due to so-called derivative loss in fundamental solutions in high dimensions. Therefore one may have questions;

- Do we have any possibility to get $T(\varepsilon) = \infty$ if the nonlinear term is not single while it includes u^2 ?
- Do we have any possibility to get a pointwise positivity of the solution for some special nonlinear term ?

For these questions, we get the following partial answers.

Theorem 1 (Takamura and Wakasa [20]) *Even if the right-hand side of the equation in (4) additionally has integral terms of the form;*

$$-\frac{1}{\pi^2} \int_0^t d\tau \int_{|\xi| \leq 1} \frac{(u_t u)(x + (t - \tau)\xi, \tau)}{\sqrt{1 - |\xi|^2}} d\xi - \frac{\varepsilon^2}{2\pi^2} \int_{|\xi| \leq 1} \frac{f(x + t\xi)^2}{\sqrt{1 - |\xi|^2}} d\xi, \quad (6)$$

there is no change on the estimate the lifespan (5).

(6) comes from a removal of the derivative loss factor in the nonlinear term of the equivalent integral equation to (4). This observation already appeared in Agemi, Kubota and Takamura [1] in which a global solution is obtained for the “super-critical” case. The proof of this theorem is established by iteration argument in weighted L^∞ space.

In contrast with (6), we have the following.

Theorem 2 (Takamura and Wakasa [21]) *If the right-hand side of the equation in (4) additionally has integral terms of the form;*

$$-\frac{1}{2\pi^2} \int_0^t d\tau \int_{|\omega|=1} (u_t u)(x + (t - \tau)\omega, \tau) dS_\omega - \frac{\varepsilon}{4\pi^2} \int_{|\omega|=1} (\varepsilon f^2 + \Delta f + 2\omega \cdot \nabla g)(x + t\omega) dS_\omega, \quad (7)$$

then, $T(\varepsilon) = \infty$ holds.

(7) follows from the following fact due to Agemi and Takamura [2]. When $n \geq 3$, a classical solution u of (1) satisfies

$$\begin{aligned} (n - 2)\omega_n u(x, t) = & \varepsilon \int_{|\omega|=1} \{t\omega \cdot \nabla f + (n - 2)f + tg\} (x + t\omega) dS_\omega \\ & + (n - 3) \int_0^t d\tau \int_{|\omega|=1} u_t(x + (t - \tau)\omega, \tau) dS_\omega \\ & + \int_0^t (t - \tau) d\tau \int_{|\omega|=1} H(x + (t - \tau)\omega, \tau) dS_\omega, \end{aligned} \quad (8)$$

where ω_n is an area of the unit sphere in $\mathbf{R}^n$. If we neglect the second term in the right-hand side of (8), we get (7) by replacing g by $2g$ when $n = 4$ and $H = u^2$. The proof of this theorem is also established by iteration argument in weighted L^∞ space. But the key estimate is

$$\left| \int_{|\omega|=1} (t\omega \cdot \nabla f + 2f + 2tg)(x + t\omega) dS_\omega \right| \leq \frac{C_{f,g}}{(1 + t)^2},$$

where $C_{f,g}$ is a positive constant. This is faster than $(1 + t)^{-3/2}$ which is a decay of a solution of the free equation.

References

- [1] R.Agemi, K.Kubota and H.Takamura, *On certain integral equations related to nonlinear wave equations*, Hokkaido. Math. J., **23**(1994), 241-276.
- [2] R.Agemi and H.Takamura, *Remarks on representations of solutions to the wave equations*, Mathematical Research Note 94-004, Institute of Mathematics, University of Tsukuba (June, 1994).
- [3] D.Christodoulou, *Global solutions of nonlinear hyperbolic equations for small initial data*, Comm. Pure Appl. Math., **39**(1986), 267-282.
- [4] R.T.Glassey, *Blow-up theorems for nonlinear wave equations*, Math. Z., **132**(1973), 183-203.
- [5] P.Godin, *Lifespan of solutions of semilinear wave equations in two space dimensions*, Comm. Partial Differential Equations, **18**(1993), 895-916.
- [6] A.Hoshiga, *The existence of the global solutions to semilinear wave equations with a class of cubic nonlinearities in 2-dimensional space*, Hokkaido. Math. J., **37**(2008), 669-688.
- [7] H.A.Levine, *Instability and nonexistence of global solutions to nonlinear wave equations of the form $Pu_{tt} = -Au + \mathcal{F}(u)$* , Trans. Amer. Math. Soc., **192**(1974), 1-21.
- [8] T-T.Li, *Lower bounds of the life-span of small classical solutions for nonlinear wave equations*, Microlocal Analysis and Nonlinear Waves (Minneapolis, MN, 1988-1989), The IMA Volumes in Mathematics and its Applications, vol.30 (M.Beals, R.B.Melrose and J.Rauch ed.), 125-136, Springer-Verlag New York, Inc., 1991.
- [9] T-T.Li and Y.Chen, "Global Classical Solutions for Nonlinear Evolution Equations", Pitman Monographs and Surveys in Pure and Applied Mathematics 45, Longman Scientific & Technical, 1992.
- [10] T-T.Li and Y.Zhou, *A note on the life-span of classical solutions to nonlinear wave equations in four space dimensions*, Indiana Univ. Math. J., **44**(1995), 1207-1248.
- [11] F.John, *Blow-up of solutions of nonlinear wave equations in three space dimensions*, Manuscripta Math., **28**(1979), 235-268.
- [12] F.John, "Nonlinear Wave Equations, Formation of Singularities", Pitcher Lectures in Mathematical Sciences, Lehigh University, AMS, 1990.
- [13] S.Katayama, *Global existence for systems of nonlinear wave equations in two space dimensions. II*, Publ. Res. Inst. Math. Sci., **31**(1995), 645-665.
- [14] S.Katayama, *Lifespan of solutions for two space dimensional wave equations with cubic nonlinearity*, Comm. Partial Differential Equations, **26**(2001), 205-232.

- [15] S.Klainerman, *The null condition and global existence to nonlinear wave equations*, Nonlinear systems of partial differential equations in applied mathematics, Part 1 (Santa Fe, N.M., 1984), 293-326, Lectures in Appl. Math., 23, Amer. Math. Soc., Providence, RI, 1986.
- [16] H.Kubo, *Asymptotic behavior of solutions of to semilinear wave equations with dissipative structure*, Discerte Contin. Dyn. Syst. 2007, Dynamical Systems and Differential Equations. Proceedings of the 6th AIMS International Conference, suppl., 602-613.
- [17] T.C.Sideris, *Formation of singularities in solutions to nonlinear hyperbolic equations*, Arch. Rational Mech. Anal., **86**(1984), 369-381.
- [18] W.A.Strauss, *Nonlinear scattering theory at low energy*, J. Funct. Anal., **41**(1981), 110-133.
- [19] H.Takamura and K.Wakasa, *The sharp upper bound of the lifespan of solutions to critical semilinear wave equations in high simensions*, J. Differential Equations **251** (2011), 1157-1171.
- [20] H.Takamura and K.Wakasa, *Almost global solutions of semilinear wave equations with the critical exponent in high dimensions*, preprint.
- [21] H.Takamura and K.Wakasa, *Global existence for semilinear wave equations with the blow-up term in high dimensions*, in preparation.
- [22] Y.Zhou and W.Han, *Sharpness on the lower bound of the lifespan of solutions to nonlinear wave equations*, Chin. Ann. Math. Ser.B., **32B**(4)(2011), 521-526. (doi: 10.1007/s11401-011-0652-5)
- [23] Y.Zhou and W. Han, *Blow up for some semilinear wave equations in multi-space dimensions*, arXiv:1207.5306 [Math.AP] 30 Jul. 2012.

Timelike minimal surfaces in Minkowski space

Giandomenico Orlandi*

We discuss here some properties of timelike minimal surfaces in flat Minkowski space-time, reviewing in particular some results proved in [3, 2, 4, 10].

Recall a for a smooth lorentzian time-like minimal submanifold, its time-slices satisfy

$$a = (1 - |v|_e^2)\kappa, \quad (0.1)$$

where a , v and κ are respectively the normal acceleration, the normal velocity and the euclidean mean curvature of the time-slice of the submanifold, $|v|_e$ is the euclidean length of v , and where we normalize units so that the speed of light equals one. Equation (0.1) can be considered as a geometric evolution equation, and the lorentzian structure of the ambient space $\mathbb{R}^{1+N}$ reflects in the hyperbolic nature of the equation. Smooth lorentzian timelike minimal submanifolds are thus characterized by the fact that their (spacetime) lorentzian mean curvature is identically zero. A particular case, relevant for its applications in physics, is the one of two-dimensional surfaces of codimension one or two, namely when $N = 2$ or $N = 3$. Under these assumptions, minimal surfaces are called (classical) relativistic strings, of codimension one and two respectively. In case the minimal submanifold is a two-dimensional graph, the evolution equation (0.1) is also known as the Born-Infeld equation, a nonlinear model for electromagnetism (see [7, 6]).

Solutions to (0.1) develop generically singularities of various type in finite time [15, 8].

A general short-time existence result for smooth solutions to (0.1) has been obtained in [12]. If the manifold is supposed sufficiently close to a linear subspace, then global existence results has been proved in [5], [11]. If one is interested in non regular or weak solutions to the lorentzian minimal surface equation, globally defined in time, one may for instance use non smooth parametrizations, or introduce generalized lorentzian surfaces in the spirit of Almgren's varifold theory and Di Perna-Majda generalized Young measures solutions the Euelr and Navier-Stokes equation [4], but in this case one is faced also to the fact that, differently from the euclidean case, the space of such solutions is not compact under uniform convergence, and a satisfactory characterization of its closure is still missing for weak or generalized minimal submanifolds of dimension larger than two.

Remark also that some suggestions in the direction of a suitable definition of weak solution arise from the analysis of the asymptotic limits, as $\varepsilon \rightarrow 0^+$, of hyperbolic Ginzburg-Landau equation, a semilinear wave equation that allows concentration of energy on timelike minimal surfaces, as observed by Neu [13]. In the more simple one-codimensional case, this is the scaled semilinear hyperbolic PDE

$$u_{tt} - \Delta u + \frac{1}{\varepsilon^2} W'(u) = 0, \quad (0.2)$$

*Dipartimento di Informatica, Università di Verona, strada le Grazie 15, 37134 Verona, Italy, email: giandomenico.orlandi@univr.it

where $W(x) = \frac{1}{4}(1 - x^2)^2$ is a standard double-well potential. As showed in [13] by a formal asymptotic expansion argument, equation (0.1) can be approximated by solutions to (0.2). Next, it has been rigorously shown in [9] that solutions to (0.2) with a well-prepared initial datum converge, as $\varepsilon \rightarrow 0^+$, to a smooth lorentzian minimal hypersurface, when the latter exists, this result being generalized by the author also in codimension two. Again, due to the presence of singularities, the validity of this convergence result is restricted to short times. On the other hand, a preliminary analysis of the limit behaviour of the hyperbolic Ginzburg-Landau equations in the varifolds framework has been pursued in [3], without restricting to short times, but under rather strong assumptions on the limiting varifold solution, in the spirit of [1] for the parabolic case.

It is worth noticing that for surfaces of dimension two, i.e. the case of relativistic strings, the problem of characterizing the closure of the space of solutions of (0.1) has a reasonable positive answer. Indeed, in this case one may parametrize the string in the so called orthogonal gauge, so that the parametrization $X(s, t)$ solves the linear wave system, and hence the solution is explicitly given by D'Alembert formula $X(s, t) = (t, \gamma(t, s))$, with $\gamma(t, s) = \frac{1}{2}(a(s - t) + b(s + t))$, where a and b are arc-length parametrized closed curves (i.e. $|a'| = |b'| = 1$).

The closure of two-dimensional minimal surfaces has been completely characterized, at least in a parametric setting, leading to the concept of subrelativistic string (see [15, 13, 6, 2] for a detailed discussion), i.e. a lipschitz parametrization $\gamma(t, s) = \frac{1}{2}(a(s - t) + b(s + t))$ with the relaxed constraint $|a'| \leq 1$, $|b'| \leq 1$ a.e..

In terms of lorentzian varifolds, as introduced in [4], the notion of subrelativistic string corresponds to the notion of a stationary weakly rectifiable varifold, which can be in turn thought as a limit of rectifiable varifolds, and enjoys some useful properties reflected by equation (0.1) such as energy and momentum conservation during the evolution.

In the case of evolution of strings in three dimensional Minkowski space-time we analyze also convexity preserving properties of (0.1), in the spirit of Gage-Hamilton result for curvature flow of planar curves, and Kong-Kefeng-Zhang in the case of the (nonrelativistic) hyperbolic curvature flow, proving such a result for centrally symmetric uniformly convex planar curves initially at rest in [2], together with an analysis of the asymptotic profile near the collapsing time, which corresponds to a round circle.

Differently from the parabolic case, this collapsing singularity is non generic, while the generic singularity expected is a cusp, as proven in [15, 8].

The extinction singularity concerns also the case of non uniformly convex boundary curves (e.g. a square), but in this case the asymptotic profile will not be a circle (for the square, the profile remains a square). The D'Alembert parametrization provides a global weak periodic solution (pulsating kink), which strongly lacks of uniqueness (one can for instance restart the evolution of a pulsating circle after collapsing with a suitable expanding centrally symmetric convex boundary, see [10]).

The special case of minimal immersed cylinders in Minkowski space-time has also been investigated, in particular by Nguyen-Tian in [14], where it is proved that any immersed timelike minimal cylinder in three-dimensional space-time necessarily develops singularities, and it is conjectured that this doesn't hold in dimension higher than four. This conjecture is proved to be true in [10], where also an estimate of the dimension of the singular set of parametrizations of minimal C^1 cylinders in a specific form (not necessarily immersions).

More precisely, we prove in [10] that, generically, in space dimension higher than three, given

a closed immersed curve Γ and a velocity field v with $|v| < 1$ and orthogonal to Γ , there exists a smooth globally immersed timelike minimal surface containing Γ and tangent to $(1, v)$. In the three-dimensional case, roughly speaking, both globally smooth immersed solutions and solutions that develop singularities occur for large sets of initial data (i.e. sets with nonempty interior).

References

- [1] L. Ambrosio, H.M. Soner. A measure-theoretic approach to higher codimension mean curvature flows. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, 25(1-2):27–49, 1998.
- [2] G. Bellettini, J. Hoppe, M. Novaga, G. Orlandi. Closure and convexity properties of closed relativistic strings. *Complex Anal. Oper. Theory*, 4(3):473–496, 2010.
- [3] G. Bellettini, M. Novaga, G. Orlandi. Time-like minimal submanifolds as singular limits of nonlinear wave equations. *Physica D*, 239(6):335–339, 2010.
- [4] G. Bellettini, M. Novaga, G. Orlandi. Lorentzian varifolds and applications to relativistic string theory, *Indiana Univ. Math. J.*, to appear.
- [5] S. Brendle. Hypersurfaces in Minkowski space with vanishing mean curvature. *Comm. Pure Appl. Math.*, 55(10):1249–1279, 2002.
- [6] Y. Brenier. Nonrelativistic strings may be approximated by relativistic strings. *Methods Appl. Anal.*, 2005.
- [7] M. Born, L. Infeld. Foundations of a new field theory. *Proc. Roy. Soc. A*, 144:425–451, 1934.
- [8] J. Eggers, J. Hoppe. Singularity formation for time-like extremal hypersurfaces. *Physics Letters B*, 680:274–278, 2009.
- [9] R.L. Jerrard. Defects in semilinear wave equations and timelike minimal surfaces in Minkowski space, *Preprint*, 2010.
- [10] R.L. Jerrard, M. Novaga, G. Orlandi. On the regularity of timelike extremal surfaces, preprint 2013
- [11] H. Lindblad. A remark on global existence for small initial data of the minimal surface equation in Minkowskian space time. *Proc. Am. Math. Soc.*, 132(4):1095–1102, 2004.
- [12] O. Milbredt. The Cauchy problem for membranes. *Preprint arXiv:0807.3465*, 2008.
- [13] J.C. Neu. Kinks and the minimal surface equation in Minkowski space. *Physica D*, 43(2-3):421–434, 1990.
- [14] L. Nguyen, G. Tian. On smoothness of timelike maximal cylinders in three dimensional vacuum spacetimes. *arXiv:1201.5183*, 2012.
- [15] A. Vilenkin, E. P. S. Shellard. *Cosmic Strings and Other Topological Defects*. Cambridge University Press, 1994.

L^1 MAXIMAL REGULARITY AND LOCAL EXISTENCE OF A SOLUTION TO THE COMPRESSIBLE NAVIER-STOKES-POISSON SYSTEM IN A CRITICAL BESOV SPACE

Takayoshi Ogawa (小川卓克)

Mathematical Institute, Tohoku University
 Sendai 980-8578, JAPAN

1. INTRODUCTION

We consider the Cauchy problem of the compressible Navier-Stokes-Poisson system in $\mathbb{R}^n$ with $n \geq 2$.

$$\begin{cases}
 \partial_t \rho + \operatorname{div}(\rho u) = 0, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^n, \\
 \begin{cases}
 \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla P(\rho) \\
 = \mu \Delta u + (\mu + \lambda) \nabla \operatorname{div} u + \kappa \rho \nabla \psi, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^n, \\
 -\Delta \psi = \rho - \bar{\rho}, & (t, x) \in \mathbb{R}_+ \times \mathbb{R}^n, \\
 u(0, x) = u_0(x), \quad \rho(0, x) = \rho_0(x), & x \in \mathbb{R}^n,
 \end{cases}
 \end{cases} \quad (1.1)$$

where $\rho = \rho(t, x)$, $u = u(t, x)$ and $\psi = \psi(t, x)$ are the unknown fluid density, the velocity vector and the potential force, respectively. $P = P(\rho)$ is the pressure given by ρ , and $u \otimes u$ denotes the tensor product of velocity vector u . μ, λ are Lamé constants satisfying $\mu + 2\lambda > 0$, $\kappa = \pm$ is a coupling constant and $\bar{\rho}$ is a given background density. Without losing generality, we assume that $\bar{\rho} = 1$. The system is strongly relevant to the simplified system of degenerate drift-diffusion equations and the Smoluchowski-Poisson system appeared in a semiconductor device models (cf. [16]).

Introducing the perturbed density by $a(t, x) \equiv \rho(t, x) - 1$ with $a_0(x) \equiv \rho_0(x) - 1$, the problem (1.1) is reduced into the following problem of (a, u) :

$$\begin{cases}
 \partial_t a + u \cdot \nabla a = -(1 + a) \operatorname{div} u, \\
 \partial_t u - \mathcal{L}u + \nabla(-\Delta)^{-1}a = -\frac{a}{1+a} \mathcal{L}u - u \cdot \nabla u - \nabla(Q(a)), \\
 u(0, x) = u_0, \quad a(0, x) = a_0.
 \end{cases} \quad (1.2)$$

Here, we denote the elliptic operator $\mathcal{L}$ by $\mathcal{L} = \mu \Delta + (\lambda + \mu) \nabla \operatorname{div}$ and Q is a smooth function determined by P by

$$Q(a) := - \int_0^t \frac{P'((1+z)^{-1})}{(1+z)^2} dz.$$

Nash [20] considered the local well-posedness of the compressible Navier-Stokes system for smooth data away from a vacuum. Itaya [15] also obtained the existence and uniqueness of the system assuming sufficient smoothness to the data. Matsumura-Nishida [19] proved the existence of global classical solution provided the initial data with high regularity is close to the equilibrium state.

We recall that the compressible Navier-Stokes system (1.1) has a scaling invariance: For $\nu > 0$,

$$\begin{cases} \rho_\nu(t, x) = \rho(\nu^2 t, \nu x), \\ u_\nu(t, x) = \nu u(\nu^2 t, \nu x) \end{cases} \quad (1.3)$$

provided the pressure term has been changed accordingly. Extending the classical idea initiated by Fujita-Kato [11] applied to the incompressible Navier-Stokes system, Danchin [5], [8] considered the local existence and the uniqueness of the solution for the problem in the “*scaling-critical*” homogeneous Besov space. Haspot [14] improved Danchin’s result [8] to the general Besov space and a larger space for the density by introducing an effective velocity. In order to consider the critical solvability, we necessarily introduce the homogeneous Besov spaces. Since the system (1.1) involves the hyperbolic equation for the density equation, it is required to consider the equation in the suitable space, where the supremum of the density has to be controlled. To this end, Danchin introduced the homogeneous Besov space that embedded into $L^\infty(\mathbb{R}^n)$.

Let $\{\phi_j\}_{j \in \mathbb{Z}}$ be the Littlewood-Paley dyadic decomposition of unity satisfying that

$$\sum_{j \in \mathbb{Z}} \hat{\phi}_j(\xi) = 1$$

for all $\xi \neq 0$ and $\text{supp } \hat{\phi}_j \subset \{\xi \in \mathbb{R}^n \mid 2^{j-1} < |\xi| < 2^{j+1}\}$. For $s \in \mathbb{R}$ and $1 \leq p, \sigma \leq \infty$, we define the homogeneous Besov space $\dot{B}_{p,\sigma}^s(\mathbb{R}^n)$ by

$$\dot{B}_{p,\sigma}^s(\mathbb{R}^n) = \{f \in \mathcal{S}'/\mathcal{P}; \|f\|_{\dot{B}_{p,\sigma}^s} < \infty\}$$

with the norm

$$\|f\|_{\dot{B}_{p,\sigma}^s} \equiv \begin{cases} \left(\sum_{j \in \mathbb{Z}} 2^{js\sigma} \|\phi_j * f\|_p^\sigma \right)^{1/\sigma}, & 1 \leq \sigma < \infty, \\ \sup_{j \in \mathbb{Z}} 2^{js} \|\phi_j * f\|_p, & \sigma = \infty, \end{cases} \quad (1.4)$$

where $\mathcal{P}$ denotes polynomials (see Triebel [23] for details).

Since our system involves the Poisson term and it brings our problem disturbing the low frequency in the Fourier spaces. Namely, the inverse operator of the Laplacian gives a stronger restriction for the low frequency part of the solution (cf. Yukawa potential case [2]). To handle with low frequency part, we also introduce another homogeneous Besov space of hybrid type; $\dot{B}_{p,\sigma}^{s_*} \oplus \dot{B}_{p,\sigma}^{s^*}$ by

$$\|f\|_{\dot{B}_{p,\sigma}^{s_*} \oplus \dot{B}_{p,\sigma}^{s^*}} \equiv \left(\sum_{j \leq 0} 2^{\sigma s_* j} \|\phi_j * f\|_p^\sigma + \sum_{j > 0} 2^{\sigma s^* j} \|\phi_j * f\|_p^\sigma \right)^{1/\sigma}$$

for all $1 \leq p, \sigma \leq \infty$ and $s_*, s^* \in \mathbb{R}$. We note that if $s_* < s^*$, then it holds that

$$\dot{B}_{p,\sigma}^{s_*} \oplus \dot{B}_{p,\sigma}^{s^*} = \dot{B}_{p,\sigma}^{s_*} \cap \dot{B}_{p,\sigma}^{s^*}$$

and hereafter we only use this setting. We define the *critical* inhomogeneous space as follows:

$$a \in L^\infty(0, T; \dot{B}_{p,1}^{\frac{N}{p}}), \quad u \in L^\infty(0, T; \dot{B}_{p,1}^{\frac{N}{p}-1}), \quad f \in L_{loc}^1(\mathbb{R}^+; \dot{B}_{p,1}^{\frac{N}{p}-1}).$$

Zheng [24] used the linearized formulation to (1.1) and solve the system by the way of integral equations. The key idea is to consider the Poisson term as the linear term and he

introduced the semi-group

$$\frac{d}{dt} \begin{pmatrix} a \\ u \end{pmatrix} = \begin{pmatrix} 0 & \text{div} \\ -|\nabla| - \nabla(-\Delta)^{-1} & \mathcal{L} \end{pmatrix} \begin{pmatrix} a \\ u \end{pmatrix}. \quad (1.5)$$

Establishing the L^p - L^q type estimate of the semi-group generated by the above operator he constructed a global solution for small data to (1.1) in the critical Besov space $\rho_0 - 1 \in \dot{B}_{2,1}^{\frac{n}{2}-2} \oplus \dot{B}_{p,1}^{\frac{n}{p}}$, $u_0 \in \dot{B}_{2,1}^{\frac{n}{2}-2} \oplus \dot{B}_{p,1}^{\frac{n}{p}-1}$. However the critical case $p = n$ was not treated, since the product formula necessarily required in the case $p = n$ such as

$$\|fg\|_{\dot{B}_{p,1}^{n/p-1}} \leq C \|f\|_{\dot{B}_{p,1}^{n/p-1}} \|g\|_{\dot{B}_{p,1}^{n/p+1}}$$

failes in general.

We now recover the local existence result in the critical hybrid Besov spaces.

Theorem 1.1 ([3]). *Let $n = 3$, $1 < p \leq 3$. $\mu > 0$ with $\mu + 2\lambda > 0$. For any $\rho_0 - 1 \in \dot{B}_{p,1}^{\frac{n}{p}-2}(\mathbb{R}^3) \oplus \dot{B}_{p,1}^{\frac{n}{p}}(\mathbb{R}^3)$, $u_0 \in \dot{B}_{p,1}^{\frac{n}{p}-2}(\mathbb{R}^3) \oplus \dot{B}_{p,1}^{\frac{n}{p}-1}(\mathbb{R}^3)$. Then there exists a weak solution to (1.1) such that for some $T > 0$ with $I = [0, T)$, (ρ, u, ψ) : a solution of (1.1) satisfying*

$$\begin{aligned} \rho - 1 &\in C(I; \dot{B}_{p,1}^{\frac{n}{p}-1} \oplus \dot{B}_{p,1}^{\frac{n}{p}}), \\ u &\in (C(I; \dot{B}_{p,1}^{\frac{n}{p}-2} \oplus \dot{B}_{p,1}^{\frac{n}{p}-1}) \cap L^1(I; \dot{B}_{p,1}^{\frac{n}{p}+1}))^N, \\ \psi &\in C(I; \dot{B}_{p,1}^{\frac{n}{p}+1} \oplus \dot{B}_{p,1}^{\frac{n}{p}+2}). \end{aligned} \quad (1.6)$$

2. KEY ESTIMATES

2.1. The mass conservation equation. To prove the case $p = n = 3$, we employ the following proposition. Let (a, u) solves the following equation.

$$\begin{cases} \partial_t a + u \cdot \nabla a = -(1 + a) \text{div } u, & (t, x) \in I \times \mathbb{R}^n, \\ a(0, x) = a_0(x), & x \in \mathbb{R}^N, \end{cases} \quad (2.1)$$

where $I = [0, T)$.

Proposition 2.1. *Let $a_0 \in \dot{B}_{3,1}^{-1}(\mathbb{R}^3)$, $u \in L^\infty(I; \dot{B}_{3,1}^0(\mathbb{R}^3)) \cap L^1(I; \dot{B}_{3,1}^2(\mathbb{R}^3))$ and $U(t) := \int_0^t \|\nabla u(\tau)\|_{\dot{B}_{3,1}^1} d\tau$. Suppose that $a \in L^\infty(I; \dot{B}_{3,1}^{-1}(\mathbb{R}^3) \cap \dot{B}_{3,1}^1(\mathbb{R}^3))$ solves the equation (2.1). Then there exists a constant $C > 0$ depending on n and p such that the following inequality holds.*

$$\|a\|_{L_t^\infty(I; \dot{B}_{3,1}^{-1})} \leq e^{U(t)} \left[\|a_0\|_{\dot{B}_{3,1}^{-1}} + C \int_0^t e^{-U(\tau)} (1 + \|a\|_{\dot{B}_{3,1}^1}) \|u\|_{\dot{B}_{3,1}^0} d\tau \right], \quad (2.2)$$

for $t \in [0, T]$.

In view of the above estimate (2.2), it is required that the velocity field has to have maximal regularity in L^1 in time variable. This is the key point to show the main theorem.

2.2. Maximal L^1 Regularity. Let u_0 be the initial data, and a and h be given functions. The momentum governed by the following linearized parabolic equation.

$$\begin{cases} \partial_t u - \mu \Delta u - (\lambda + \mu) \nabla(\operatorname{div} u) = F, & (t, x) \in I \times \mathbb{R}^n, \\ u(0, x) = u_0(x), & x \in \mathbb{R}^n. \end{cases} \quad (2.3)$$

It is known that the Cauchy problem for the heat equation (2.3) has maximal regularity in such non-reflexive function spaces. One of the general result can be seen in [21].

Proposition 2.2 ([21]). *Let $1 < \rho, \sigma \leq \infty$ and $I = [0, T)$ be an interval with $T \leq \infty$. For $f \in L^\rho(I; \dot{B}_{1,\rho}^0(\mathbb{R}^n))$ and $u_0 \in \dot{B}_{1,\rho}^{2(1-1/\rho)}(\mathbb{R}^n)$, let u be a solution of the Cauchy problem of the heat equation (2.3). Then we have*

$$\|\partial_t u\|_{L^\rho(I; \dot{B}_{1,\rho}^0)} + \|\nabla^2 u\|_{L^\rho(I; \dot{B}_{1,\rho}^0)} \leq C(\|u_0\|_{\dot{B}_{1,\rho}^{2(1-1/\rho)}} + \|F\|_{L^\rho(I; \dot{B}_{1,\rho}^0)}). \quad (2.4)$$

The above result does not cover the end-point exponent $\rho = 1$. In general, the end-point case $p = 1$ is eliminated in the abstract theory and we need to develop the each cases. Danchin [6] (see also Haspot [14]) obtained maximal regularity in the homogeneous Besov space for the case $\rho = 1$.

Theorem 2.3 ([8], [22]). *Let $1 \leq p \leq \infty$. For $F \in L^1(\mathbb{R}_+; \dot{B}_{p,1}^0(\mathbb{R}^n))$ and $u_0 \in \dot{B}_{p,1}^0(\mathbb{R}^n)$ there exists a unique solution u to (2.3) which satisfies the estimate:*

$$\begin{aligned} \|\partial_t u\|_{L^1(\mathbb{R}_+; \dot{B}_{p,1}^0)} + \|\nabla^2 u\|_{L^1(\mathbb{R}_+; \dot{B}_{p,1}^0)} \\ \leq C \left(\|u_0\|_{\dot{B}_{p,1}^0} + \|F\|_{L^1(\mathbb{R}_+; \dot{B}_{p,1}^0)} \right), \end{aligned} \quad (2.5)$$

where constant C is depending only on n . Moreover the estimate is optimal for the class of initial data. Namely if the data is $u_0 \in L^p(\mathbb{R}^N)$ or $\dot{F}_{p,1}^0(\mathbb{R}^N)$ the above estimate fails.

Remark 2.1. The upper estimate of (2.5) was obtained by Danchin- Mucha [9, Proposition 5] with $1 < p < \infty$. For $p = 1$, Danchin essentially obtained the same estimate even for the variable coefficient case. Giga-Saal considered time L^1 maximal regularity in some space [12].

If we replace $u_0 \in \dot{B}_{p,1}^0(\mathbb{R}^n)$ into $u_0 \in \dot{B}_{p,\sigma}^0(\mathbb{R}^n)$ for $1 < \sigma \leq \infty$, then maximal regularity fails since the lower bound by the initial data and the strict inclusion result for the sub-suffix σ as $\dot{B}_{p,1}^0(\mathbb{R}^n) \subsetneq \dot{B}_{p,\sigma}^0(\mathbb{R}^n)$.

To avoid the difficulty on using the limiting case of the bi-linear estimate in the homogeneous Besov spaces, we employ the following bi-linear estimate to treat the nonlinear term of the equation (2.1).

Lemma 2.4. *For $u \in \dot{B}_{3,1}^0 \cap \dot{B}_{3,1}^2$ and $\tilde{a} \in \dot{B}_{3,1}^{-1} \cap \dot{B}_{3,1}^1$ it holds that*

$$\|\operatorname{div}(\tilde{a}u)\|_{\dot{B}_{3,1}^{-1}} \leq C(\|u\|_{\dot{B}_{3,1}^0} \|\tilde{a}\|_{\dot{B}_{3,1}^1} + \|u\|_{\dot{B}_{3,1}^2} \|\tilde{a}\|_{\dot{B}_{3,1}^{-1}}).$$

REFERENCES

- [1] Abidi, H. and Paicu, M., *Existence globale pour un fluide inhomogène*, Ann. Inst. Fourier (Grenoble) **57** (2007), 883–917.
- [2] Chikami, N., *The blow-up criterion for the compressible Navier-Stokes system with a Yukawa-potential in the critical Besov space*, preprint.
- [3] Chikami, N. and Ogawa, T., *Existence of solutions to compressible Navier-Stokes-Poisson system in a critical Besov space*, in preparation.
- [4] Chen, Q., Miao, C. and Zhang, Z., *Well-posedness in critical spaces for the compressible Navier-Stokes equations with density dependent viscosities*, Rev. Mat. Iberoam. **26** (2010), no. 3, 915–946.
- [5] Danchin, R., *Global existence in critical spaces for compressible Navier-Stokes equations.*, Invent. Math. **141** (2000), 579–614.
- [6] ———, *Local theory in critical spaces for compressible viscous and heat-conductive gases*, Comm. Partial Differential Equations **26** (2001), 1183–1233.
- [7] ———, *On the uniqueness in critical spaces for compressible Navier-Stokes equations*, NoDEA Nonlinear Differential Equations Appl. **12** (2005), 111–128.
- [8] ———, *Well-posedness in critical spaces for barotropic viscous fluids with truly not constant density*, Comm. Partial Differential Equations **32** (2007), 1373–1397.
- [9] Danchin R. and Mucha P.B., *A critical functional framework for the inhomogeneous Navier-Stokes equations in the half-space*, J. Funct. Anal. **256** (2009), 881–927.
- [10] Ducomet, B., *Simplified models of quantum fluids in nuclear physics*, Proceedings of Partial Differential Equations and Applications (Olomouc, 1999), 2001, pp. 323–336.
- [11] Fujita, H. and Kato, T., *On the Navier-Stokes initial value problem. I*, Arch. Rational Mech. Anal. **16** (1964), 269–315.
- [12] Giga, Y. and Saal, J., *L^1 maximal regularity for the Laplacian and applications*, Discrete Contin. Dyn. Syst. **Suppl. 1** (2011), 495–504.
- [13] Hao, C. and Li, H.-L., *Global existence for compressible Navier-Stokes-Poisson equations in three and higher dimensions*, J. Differential Equations **246** (2009), 4791–4812.
- [14] Haspot, B., *Well-posedness in critical spaces for the system of compressible Navier-Stokes in larger spaces*, J. Differential Equations **251** (2011), 2262–2295.
- [15] Itaya, N., *On the Cauchy problem for the system of fundamental equations describing the movement of compressible viscous fluid*, Kōdai Math. Sem. Rep. **23** (1971), 60–120.
- [16] Kobayashi, T. and Ogawa, T., *Fluid mechanical approximation to the degenerate drift-diffusion and chemotaxis equations in barotropic model*, Indiana Univ. Math. J., to appear.
- [17] Kozono, H., Ogawa, T., and Taniuchi, Y., *The critical Sobolev inequalities in Besov spaces and regularity criterion to some semi-linear evolution equations*, Math. Z. **242** (2002), no. 2, 251–278.
- [18] Li, H.-L., Matsumura, A., and Zhang, G., *Optimal decay rate of the compressible Navier-Stokes-Poisson system in $\mathbb{R}^3$* , Arch. Ration. Mech. Anal. **196** (2010), 681–713.
- [19] Matsumura, A. and Nishida, T., *The initial value problem for the equations of motion of viscous and heat-conductive gases*, J. Math. Kyoto Univ. **20** (1980), 67–104.
- [20] Nash, J., *Le problème de Cauchy pour les équations différentielles d’un fluide général*, Bull. Soc. Math. France **90** (1962), 487–497.
- [21] Ogawa, T. and Shimizu, S., *End-point maximal regularity and wellposedness of the two dimensional Keller-Segel system in a critical Besov space*, Math. Z. **56** (2010), 601–628.
- [22] ———, *Remarks on optimality of end-point maximal regularity in L^1 type*, preprint.
- [23] Triebel, H., *Theory of function spaces*, Monographs in Mathematics, vol. 78, Birkhäuser Verlag, Basel, 1983.
- [24] Zheng, X., *Global well-posedness for the compressible Navier-Stokes-Poisson system in the L^p framework*, Nonlinear Anal. **75** (2012), no. 10, 4156–4175.