HOKKAIDO UNIVERSITY

Title	Fock-space representations of the relativistic supersymmetry algebra in the two-dimensional spacetime
Author(s)	Arai, Asao
Citation	Hokkaido University Preprint Series in Mathematics, 123, 2-13
Issue Date	1991-09
DOI	https://doi.org/10.14943/83268
Doc URL	https://hdl.handle.net/2115/68870
Type	departmental bulletin paper
File Information	re123.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Fock-Space Representations of the Relativistic Supersymmetry Algebra in the Two-dimensional Space-time

Asao Arai

Series #123. September 1991

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

96: S. Ninomiya, The Fourier-Sato transformation of pure sheaves, 22 pages. 1990.

97: Y. Okabe, A. Inoue, The theory of KM_2O-Langevin equations and its applications to data analysis (II): Causal analysis (1), 51 pages. 1990.

98: J. Lawrynowicz, S. Koshi and O. Suzuki, Dualities generated by the generalised Hurwitz problem and variation of the Yang-Mills field, 17 pages. 1991.

99: R. Agemi, K. Kubota and H. Takamura, On certain integral equations related to nonlinear wave equations, 52 pages. 1991.

100: S. Izumiya, Geometric singularities for Hamilton-Jacobi equation, 13 pages. 1991.

101: S. Izumiya, Legendrian singularities and first order differential equations, 16 pages. 1991.

102: A. Munemasa, Y. Watatani, Orthogonal pairs of $*$-subalgebras and association schemes, 11 pages. 1991.

103: A. Arai, O. Ogurisu, Meromorphic $N = 2$ Wess-Zumino supersymmetric quantum mechanics, 27 pages. 1991.

104: H. Takamura, Global existence of classical solutions to nonlinear wave equations with spherical symmetry for small data with noncompact support in three space dimensions, 14 pages. 1991.

105: R. Agemi, Blow-up of solutions to nonlinear wave equations in two space dimensions, 11 pages. 1991.

106: T. Nakazi, Extremal problems in H^p, 13 pages. 1991.

107: T. Nakazi, ρ-dilations and hypo-Dirichlet algebras, 15 pages. 1991.

108: A. Arai, An abstract sum formula and its applications to special functions, 25 pages. 1991.

109: Y.-G. Chen, Y. Giga and S. Goto, Analysis toward snow crystal growth, 18 pages. 1991.

110: T. Hibi, M. Wakayama, A q-analogue of Capelli's identity for $GL(2)$, 7 pages. 1991.

111: T. Nishimori, A qualitative theory of similarity pseudogroups and an analogy of Sacksteder's theorem, 13 pages. 1991.

112: K. Matsuda, An analogy of the theorem of Hector and Duminy, 10 pages. 1991.

113: S. Takahashi, On a regularity criterion uo to the boundary for weak solutions of the Navier-Stokes equations, 23 pages. 1991.

114: T. Nakazi, Sum of two inner functions and exposed points in H^1, 18 pages. 1991.

115: A. Arai, De Rham operators, Laplacians, and Dirac operators on topological vector spaces, 27 pages. 1991.

116: T. Nishimori, A note on the classification of non-singular flows with transverse similarity structures, 17 pages. 1991.

117: T. Hibi, A lower bound theorem for Ehrhart polynomials of convex polytopes, 6 pages. 1991.

118: R. Agemi, H. Takamura, The lifespan of classical solutions to nonlinear wave equations in two space dimensions, 30 pages. 1991.

119: S. Altschuler, S. Angenent and Y. Giga, Generalized motion by mean curvature for surfaces of rotation, 15 pages. 1991.

120: T. Nakazi, Invariant subspaces in the bidisc and commutators, 20 pages. 1991.

121: A. Arai, Commutation properties of the partial isometries associated with anticommuting self-adjoint operators, 25 pages. 1991.

122: Y.-G. Chen, Blow-up solutions to a finite difference analogue of $u_t = \Delta u + u^{1+\alpha}$ in N-dimensional balls, 31 pages. 1991.

Fock-Space Representations of the Relativistic Supersymmetry Algebra in the Two-dimensional Space-time

ASAO ARAI
Department of Mathematics, Hokkaido University
Sapporo 060, Japan

Abstract. An operator theoretical analysis is made on the representation of the supersymmetry(SUSY) algebra originating from the relativistic supersymmetry in the two-dimensional space-time with a proper treatment of anticommutatvity and commutativity of operators in the SUSY algebra. It is shown that, given on a Hilbert space $\mathfrak{H}$ two self-adjoint supercharges anticommuting *in the strong sense*, one can construct a representation of the SUSY algebra on $\mathfrak{H}$ such that the supersymmetric Hamiltonian and the momentum operator commute *in the strong sense*. As an application, a general class of representations of the SUSY algebra is constructed on an abstract Boson-Fermion Fock space.

AMS subject classification (1990): 81R05, 81T60, 81Q10,17B70, 47A67

1. Introduction

In this paper we consider, from an operator theoretical point of view, the representation of the supersymmetry (SUSY) algebra

$$\begin{aligned} &Q_1^2 = H + P, \quad Q_2^2 = H - P, \\ &\{Q_1, Q_2\} = 0, \\ &[Q_j, H] = [Q_j, P] = 0, \quad j = 1, 2, \end{aligned} \tag{1.1}$$

where $\{\,\cdot\,,\,\cdot\,\}$ and $[\,\cdot\,,\,\cdot\,]$ denote anticommutator and commutator, respectively. This SUSY algebra originates from the relativistic supersymmetry in the two-dimensional space-time [16,17]. In the context of supersymmetric quantum field theory (SQFT) the quantities $\{Q_1, Q_2\}$, H, and P are called supercharges, a supersymmetric Hamiltonian, and a momentum operator, respectively. It has been shown in [7–9] that supersymmetric Wess-Zumino models on a two-dimensional cylindrical space-time give realizations of (1.1).

A representation of (1.1) requires a realization of Q_1, Q_2, H, and P as self-adjoint operators satisfying (1.1) on a Hilbert space. However, from an operator theoretical point of view, this is not sufficient to characterize a representation of (1.1) completely, because Q_1, Q_2, H, and P are not necessarily bounded in a realization of them as self-adjoint operators on a Hilbert space and hence the anticommutativity of Q_1 and Q_2 and the commutativity of Q_j with H and P may not be uniquely defined ; we need specify the meaning of them to define a representation of (1.1) unambiguously. In the present paper we use the concept of *strong commutativity* and *strong anticommutativity* given in the following definition.

DEFINITION 1.1. *Let A and B be self-adjoint operators on a Hilbert space.*

(i) *We say that A and B strongly commute if*

$$e^{itA}B \subset Be^{itA}$$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

for all $t \in \mathbf{R}$.

(ii) *We say that* A *and* B *strongly anticommute if*

$$e^{itA}B \subset Be^{-itA}$$

for all $t \in \mathbf{R}$.

Remark. Two self-adjoint operators strongly commute if and only if their spectral projections commute (e.g.,[11, §VIII.5],[13, Part 1, §1.1]). The definition of the strong anticommutativity of two self-adjoint operators is taken from [10,15] (cf. also [13]). Definitions (i) and (ii) are symmetric with respect to A and B.

DEFINITION 1.2. *A representation of (1.1) is a set* $\{Q_1, Q_2, H, P\}$ *of self-adjoint operators on a Hilbert space with the following properties:*

(i)

$$H + P \subset Q_1^2, \quad H - P \subset Q_2^2. \tag{1.2}$$

(ii) Q_1 *and* Q_2 *strongly anticommute.*

(iii) *Each* Q_j $(j = 1, 2)$ *strongly commutes with* H *and* P.

Remark. In our definition of representation of (1.1), we replace the first two equations in (1.1) by (1.2), because $H \pm P$ are not self-adjoint in general.

The outline of the present paper is as follows. In Section 2 we show that, for each pair $\{Q_1, Q_2\}$ of strongly anticommuting self-adjoint operators on a Hilbert space $\mathfrak{H}$, there exists a unique representation $\{Q_1, Q_2, H, P\}$ of (1.1) on $\mathfrak{H}$ such that H and P strongly commute (Theorems 2.4 and 2.5). We also show that, in the representation, the spectral condition for $\{H, P\}$ holds (Corollary 2.6). In view of constructive quantum field theory, it is of particular interest to construct representations of (1.1) on Fock spaces. For this purpose, we define in Section 3 a general class of strongly anticommuting self-adjoint operators on the Boson-Fermion Fock space $\mathfrak{F} = \mathfrak{F}(\mathcal{H}, \mathcal{K})$ based on the pair $\{\mathcal{H}, \mathcal{K}\}$ of two complex Hilbert spaces with $\mathcal{H}$ and $\mathcal{K}$ being the one particle space of bosons and of fermions, respectively. We introduce a set $\mathfrak{A} = \mathfrak{A}(\mathcal{H}, \mathcal{K})$ of pairs $\{A, B\}$ of two densely defined closed linear operators from $\mathcal{H}$ to $\mathcal{K}$. For each $\{A, B\} \in \mathfrak{A}$ and $\lambda \in \mathbf{R}$, we define two strongly anticommuting self-adjoint operators $\{Q_A(\lambda), Q_B\}$ on $\mathfrak{F}$. In the last section, we apply the main result (Theorem 2.4) in Section 2 to construct a Fock-space representation of (1.1) for each $\{A, B\} \in \mathfrak{A}$ and $\lambda \in \mathbf{R}$ with $Q_1 = Q_A(\lambda)$ and $Q_2 = Q_B$. In this way we obtain a general class of Fock-space representations. If $\lambda \neq 0$, then the partial breaking of the supersymmetry occurs, i.e., there exists no state $\Psi \in \mathfrak{F}$ such that $Q_A(\lambda)\Psi = 0$, while there exists a state $\Phi \in \mathfrak{F}$ such that $Q_B\Phi = 0$. We also discuss the unitary equivalence between two of these Fock space representations. Concrete realizations of these Fock-space representations (e.g., $\mathcal{H} = \mathcal{K} = L^2(\mathbf{R})$) corresponds to free SQFT's. But the Fock-space representations constructed here can be used as a starting point for constructing other types of Fock-space representations of (1.1) that correspond to SQFT's with interactions. This subject will be discussed in a forthcoming paper.

2. Representation Based on Two Strongly Anticommuting Self-Adjoint Operators

Let Q_1 and Q_2 be strongly anticommuting self-adjoint operators on a Hilbert space. For each $j = 1, 2$, one can define, via the functional calculus for self-adjoint operators, the 'absolute value' $|Q_j|$ of Q_j, which is the nonnegative self-adjoint operator such that $|Q_j|^2 = Q_j^2$. A fundamental property of $\{Q_1, Q_2\}$ is given by the following lemma [10,15].

LEMMA 2.1. *The self-adjoint operators $|Q_1|$ and $|Q_2|$ strongly commute.*

Using this lemma and the functional calculus for self-adjoint operators, we can prove that the operator

$$H = \frac{1}{2}(Q_1^2 + Q_2^2), \tag{2.1}$$

which is equal to $|Q_1|^2 + |Q_2|^2$, is nonnegative and self-adjoint with domain $D(H) = D(Q_1^2) \cap D(Q_2^2)$ [10,15] and that $Q_1^2 - Q_2^2 (= |Q_1|^2 - |Q_2|^2)$ is essentially self-adjoint. The latter fact allows us to define the self-adjoint operator

$$P = \frac{1}{2}\overline{(Q_1^2 - Q_2^2)}, \tag{2.2}$$

the closure of $(Q_1^2 - Q_2^2)/2$.

LEMMA 2.2.

(i) *Each Q_j $(j = 1, 2)$ strongly commutes with H and P.*
(ii) *The operators H and P strongly commute.*

Proof. (i) One can show that Q_1^2 and Q_2^2 strongly commute (see the proof of [10, Theorem 3.1]). The functional calculus for self-adjoint operators shows that Q_2^2 and Q_2 strongly commute. Hence it follows that H and Q_2 strongly commute. By symmetry H strongly commutes also with Q_1. Similarly we can prove that Q_j strongly commutes with P.

(ii) This follows from the functional calculus for the strongly commuting self-adjoint operators $|Q_j|, j = 1, 2$. ∎

LEMMA 2.3. *The operators H and P defined by (2.1) and (2.2), respectively, satisfy (1.2).*

Proof. It is obvious that

$$D(H) \cap D(P) = D(H) = D(Q_1^2) \cap D(Q_2^2) \subset D(Q_j^2), j = 1, 2.$$

For $f \in D(H) \cap D(P)$, we have

$$Hf = \frac{1}{2}(Q_1^2 + Q_2^2)f, \quad Pf = \frac{1}{2}(Q_1^2 - Q_2^2)f.$$

Hence

$$(H + P)f = Q_1^2 f, \quad (H - P)f = Q_2^2 f \tag{2.3}$$

Thus (1.2) holds. ∎

In summary we have obtained the following result.

THEOREM 2.4. *Let Q_1 and Q_2 be strongly anticommuting self-adjoint operators on a Hilbert space $\mathfrak{H}$. Then $\{Q_1, Q_2, H, P\}$ with H and P given by (2.1) and (2.2), respectively, is a representation of the SUSY algebra (1.1) on $\mathfrak{H}$ such that H and P strongly commute.*

We can prove the uniqueness of the representation given in Theorem 2.4.

THEOREM 2.5. *Let $\{Q_1, Q_2, H, P\}$ be a representation of the SUSY algebra (1.1) such that H and P strongly commute. Then H and P are given by (2.1) and (2.2), respectively.*

Proof. It follows from (1.2) that $D(H) \cap D(P) \subset D(Q_1^2) \cap D(Q_2^2)$ and

$$Hf = \frac{1}{2}(Q_1^2 + Q_2^2)f, \quad Pf = \frac{1}{2}(Q_1^2 - Q_2^2)f, \quad f \in D(H) \cap D(P).$$

Since H and P are strongly commuting self-adjoint operators by the assumption, it follows that $D(H) \cap D(P)$ is a common core for H and P. We already know that $Q_1^2 + Q_2^2$ is self-adjoint and $Q_1^2 - Q_2^2$ is essentially self-adjoint. Thus (2.1) and (2.2) follow. ∎

In concluding this section we show that the 'energy-momentum operator' $\{H, P\}$ in the representation given in Theorem 2.4 satisfies the spectral condition in relativistic quantum field theory in the two-dimensional space-time. For any pair $\{A, B\}$ of strongly commuting self-adjoint operators, there exists a unique resolution of the identity $E_{A,B}$ on $\mathbf{R}^2$ such that $E_{A,B}(S_1 \times S_2) = E_A(S_1)E_B(S_2)$ for all Borel sets $S_1, S_2 \subset \mathbf{R}$, where E_A (resp. E_B) denotes the resolution of the identity of A (resp. B). The joint spectrum $\sigma(A, B)$ of the pair $\{A, B\}$ is defined by

$$\sigma(A, B) = \text{supp } E_{A,B},$$

the suppot of $E_{A,B}$(e.g., [13]).

COROLLARY 2.6 (SPECTRAL CONDITION). *Let $\{Q_1, Q_2, H, P\}$ be the representation of (1.1) constructed in Theorem 2.4. Then the joint spectrum of $\{H, P\}$ is included in the forward light cone in $\mathbf{R}^2$:*

$$\sigma(H, P) \subset \{(p_0, p_1) \in \mathbf{R}^2 |\ p_0^2 - p_1^2 \geq 0, p_0 \geq 0\}.$$

Proof. This follows from the strong commutativity of H and P and the fact that

$$H \pm P \geq 0 \text{ on } D(H) \cap D(P)$$

which follows from (2.3) and the nonnegativity of $Q_j^2, j = 1, 2$. ∎

3. A Class of Strongly Anticommuting Self-Adjoint Operators on Boson-Fermion Fock Spaces

In this section, as a preliminary for the construction of Fock-space representations of the SUSY algebra (1.1), we define a class of strongly anticommuting self-adjoint operators on an abstract Boson-Fermion Fock space. This can be done by employing the theory of infinite dimensional Dirac operators developed in [2,4,5](cf. also [3,6]). In [2,4,5] we used the Q-space representation for the Boson Fock space over an abstract (one particle) Hilbert space (e.g,[12,14]). Here we treat the Boson Fock space in its original form.

Let $\mathcal{H}$ and $\mathcal{K}$ be separable complex Hilbert spaces. Let $\mathfrak{F}_{\mathrm{b}}(\mathcal{H})$ and $\mathfrak{F}_{\mathrm{f}}(\mathcal{K})$ be the Boson Fock space over $\mathcal{H}$ and the Fermion Fock space over $\mathcal{K}$, respectively [11, §II.4]. The Hilbert space in which we shall work is the Boson-Fermion Fock space

$$\mathfrak{F} = \mathfrak{F}(\mathcal{H}, \mathcal{K}) = \mathfrak{F}_{\mathrm{b}}(\mathcal{H}) \otimes \mathfrak{F}_{\mathrm{f}}(\mathcal{K}).$$

Let $a(f), f \in \mathcal{H}$, and $b(u), u \in \mathcal{K}$, be the annihilation operators on $\mathfrak{F}_{\mathrm{b}}(\mathcal{H})$ (*antilinear in* f) and on $\mathfrak{F}_{\mathrm{f}}(\mathcal{K})$ (*antilinear in* u), respectively. We denote by Ω_{b} and Ω_{f} the Fock vacuum in $\mathfrak{F}_{\mathrm{b}}(\mathcal{H})$ and in $\mathfrak{F}_{\mathrm{f}}(\mathcal{K})$, respectively. *The Fock vacuum in* $\mathfrak{F}$ is defined by

$$\Omega = \Omega_{\mathrm{b}} \otimes \Omega_{\mathrm{f}} \in \mathfrak{F}.$$

Every densely defined closed linear operator T on $\mathfrak{F}_{\mathrm{b}}(\mathcal{H})$ (resp. $\mathfrak{F}_{\mathrm{f}}(\mathcal{K})$) extends to a closed linear operator on $\mathfrak{F}$ as $T \otimes I$ (resp. $I \otimes T$). For notational simplicity we denote the extension by the same symbol T.

For a linear operator S on a Hilbert space with domain $D(S)$, we denote by $C^\infty(S)$ the set of C^∞-vectors of S:

$$C^\infty(S) = \cap_{n=1}^\infty D(S^n).$$

Let $A : \mathcal{H} \to \mathcal{K}$ be a densely defined closed linear operator and

$$\begin{aligned}\mathfrak{D} = \mathcal{L}\{a(f_1)^* \cdots a(f_n)^* b(u_1)^* \cdots b(u_m)^* \Omega | f_j \in C^\infty(A^*A), j = 1, \cdots, n, \\ u_k \in C^\infty(AA^*), k = 1, \cdots, m, n, m \geq 0\},\end{aligned} \tag{3.1}$$

where $\mathcal{L}\{\cdots\}$ means the algebraic linear span of elements in the set $\{\cdots\}$. The subspace $\mathfrak{D}$ is dense in $\mathfrak{F}$. We define an operator d_A with domain $\mathfrak{D}$ by

$$d_A \Psi = (1 - \delta_{n0}) \sum_{j=1}^{n} a(f_1)^* \cdots \widehat{a(f_j)}^* \cdots a(f_n)^* b(Af_j)^* b(u_1)^* \cdots b(u_m)^* \Omega, \tag{3.2}$$

for $\Psi \in \mathfrak{D}$ of the form

$$\Psi = a(f_1)^* \cdots a(f_n)^* b(u_1)^* \cdots b(u_m)^* \Omega \tag{3.3}$$

and by extending it by linearity to all $\Psi \in \mathfrak{D}$, where $\widehat{a(f_j)}^*$ indicates the omission of $a(f_j)^*$. The operator d_A is well-defined. In fact, for each complete orthonormal system $\{e_n\}_{n=1}^\infty$ of $\mathcal{K}$ with $e_n \in D(A^*)$, we have

$$d_A \Psi = \mathrm{s} - \lim_{N \to \infty} \sum_{n=1}^{N} a(A^* e_n) b(e_n)^* \Psi, \quad \Psi \in \mathfrak{D}. \tag{3.4}$$

LEMMA 3.1.

(i) *The operator* d_A *leaves* $\mathfrak{D}$ *invariant and*

$$d_A^2 = 0. \tag{3.5}$$

(ii) $\mathfrak{D} \subset D(d_A^*)$ *and* d_A^* *leaves* $\mathfrak{D}$ *invariant.*

(iii) d_A *is closable.*

Proof. (i) The invariance of $\mathfrak{D}$ under the action of d_A follows from (3.1) and (3.2). For $\Psi \in \mathfrak{D}$ of the form (3.3), we have

$$d_A^2\Psi = (1-\delta_{(n-1)0})(1-\delta_{n0})\sum_{k\neq j}^{n}\sum_{j=1}^{n} a(f_1)^*\cdots\widehat{a(f_j)^*}\cdots\widehat{a(f_k)^*}\cdots a(f_n)^* \times b(Af_k)^*b(Af_j)^*b(u_1)^*\cdots b(u_m)^*\Omega.$$

Using the anticommutativity of $b(u)^*$ and $b(v)^*, u, v \in \mathcal{K}$, we see that the right hand side is eqaul to $-d_A^2\Psi$. Thus (3.5) follows.

(ii) By (3.4) one can easily check that $\mathfrak{D} \subset D(d_A^*)$ and

$$d_A^*\Psi = (1-\delta_{m0})\sum_{k=1}^{m}(-1)^{k-1}a(A^*u_k)^*a(f_1)^*\cdots a(f_n)^*b(u_1)^*\cdots\widehat{b(u_k)^*}\cdots b(u_m)^*\Omega \quad (3.6)$$

for $\Psi \in \mathfrak{D}$ of the form (3.3). Formula (3.6) shows that d_A^* leaves $\mathfrak{D}$ invariant.

(iii) By part (ii), $D(d_A^*)$ is dense. Hence d_A is closable. ∎

By a limiting argument, we can show that (3.5) with the closure of d_A in place of d_A holds. *We denote the closure of d_A by the same symbol.*

We define *Dirac operators on $\mathcal{F}$* by

$$Q_A = d_A + d_A^*, \quad (3.7)$$

$$\widetilde{Q}_A = i(d_A - d_A^*), \quad (3.8)$$

with $D(Q_A) = D(\widetilde{Q}_A) = D(d_A) \cap D(d_A^*)$.

We denote by $d\Gamma(\cdot)$ (resp. $d\Lambda(\cdot)$) the second quantization on $\mathfrak{F}_b(\mathcal{H})$ (resp. $\mathfrak{F}_f(\mathcal{K})$)[11, §VIII.10]. We introduce the nonnegative self-adjoint operator

$$\Delta_A = d\Gamma(A^*A) + d\Lambda(AA^*) \quad (3.9)$$

on $\mathfrak{F}$.

THEOREM 3.2. *The operators Q_A and $\widetilde{Q}_A$ are self-adjoint and are essentially self-adjoint on every core for Δ_A. Moreover, the following operator equations hold :*

$$\Delta_A = Q_A^2 = \widetilde{Q}_A^2 = d_A^*d_A + d_Ad_A^*. \quad (3.10)$$

$$\widetilde{Q}_A = Q_{iA}. \quad (3.11)$$

Proof. The outline of the proof is as follows. We first show that Q_A and $\widetilde{Q}_A$ are closed symmetric (cf.[2, Lemma 4.4(b)]). Then we prove that (3.10) holds on $\mathfrak{D}$. Noting that $\mathfrak{D}$ is a core for Δ_A (apply [11, Theorem VIII.33 and Corollary]), we use a commutator theorem [12, Theorem X.37] to prove the essential self-adjointness of Q_A and $\widetilde{Q}_A$ on $\mathfrak{D}$. See [4] for details. It is easy to see that (3.11) holds on $\mathfrak{D}$. Since $\mathfrak{D}$ is a common core for Q_{iA} and $\widetilde{Q}_A$, the operator equation (3.11) follows. ∎

The Boson-Fock space $\mathfrak{F}$ has the following orthogonal decompsition:

$$\mathfrak{F} = \mathfrak{F}_+ \oplus \mathfrak{F}_-$$

with

$$\mathfrak{F}_+ = \bigoplus_{m=0}^{\infty} \mathfrak{F}_b(\mathcal{H}) \otimes \wedge^{2m}(\mathcal{K}), \quad \mathfrak{F}_- = \bigoplus_{m=0}^{\infty} \mathfrak{F}_b(\mathcal{H}) \otimes \wedge^{2m+1}(\mathcal{K}),$$

where $\wedge^m(\mathcal{K})$ denotes the m-fold antisymmetric tensor product of $\mathcal{K}$ ($\wedge^0(\mathcal{K}) = \mathbf{C}$). Let $P_\pm$ be the orthogonal projection onto $\mathfrak{F}_\pm$ and

$$N_F = P_+ - P_-.$$

Then one can show [4] that

$$N_F : D(Q_A) \to D(Q_A), \tag{3.12}$$

$$\{Q_A, N_F\} = \{\tilde{Q}_A, N_F\} = 0 \quad \text{on } D(Q_A). \tag{3.13}$$

It follows that the quadruple $\{\mathfrak{F}, \{Q_A, \tilde{Q}_A\}, \Delta_A, N_F\}$ is a supersytmmetric quantum theory [1,3,17].

Using the boundedness of N_F, (3.12), and (3.13), one can easily see that

$$e^{itN_F} Q_A \subset Q_A e^{-itN_F}, \quad e^{itN_F} \tilde{Q}_A \subset \tilde{Q}_A e^{-itN_F}.$$

for all $t \in \mathbf{R}$. Thus we obtain the following result.

PROPOSITION 3.3. *The self-adjoint operator N_F strongly anticommutes with Q_A and $\tilde{Q}_A$.*

Let $\lambda \in \mathbf{R}$ and

$$Q_A(\lambda) = Q_A + \lambda N_F. \tag{3.14}$$

Then, by Proposition 3.3 and [15, Corollary 2.2], $Q_A(\lambda)$ is self-adjoint and

$$Q_A(\lambda)^2 = Q_A^2 + \lambda^2, \tag{3.15}$$

where we have used the fact $N_F^2 = I$.

Let $C(\mathcal{H}, \mathcal{K})$ be the set of closed linear operators from $\mathcal{H}$ to $\mathcal{K}$. We introduce a set of pairs $\{A, B\}$ with $A, B \in C(\mathcal{H}, \mathcal{K})$.

DEFINITION 3.4. *Let $A, B \in C(\mathcal{H}, \mathcal{K})$. We say that $\{A, B\}$ is in the set $\mathfrak{A} = \mathfrak{A}(\mathcal{H}, \mathcal{K})$ if the following conditions are satisfied:*

(i) *The self-adjoint operators A^*A and B^*B strongly commute.*

(ii) *The self-adjoint operators AA^* and BB^* strongly commute.*

(iii)

$$C^\infty(A^*A) \cap C^\infty(B^*B) \subset D(A^*B) \cap D(B^*A),$$
$$C^\infty(AA^*) \cap C^\infty(BB^*) \subset D(AB^*) \cap D(BA^*),$$

and

$$A^*B + B^*A = 0, \quad AB^* + BA^* = 0.$$

The set $\mathfrak{A}$ is not empty. For example, $\{A, iA\} \in \mathfrak{A}$ for all $A \in C(\mathcal{H}, \mathcal{K})$.

The main result in this section is the following.

THEOREM 3.5. *For all* $\{A, B\} \in \mathfrak{A}$ *and* $\lambda \in \mathbf{R}$, $Q_A(\lambda)$ *and* Q_B *strongly anticommute.*

To prove this theorem, we need some preliminaries. For a self-adjoint operator S on a Hilbert space, we define the subspace

$$C_0^\infty(S) = \cup_{-\infty<a<b<\infty} R(E_S((a,b])),$$

where E_S is the spectral measure of S and $R(T)$ denotes the range of operator T. The subspace $C_0^\infty(S)$ is dense in the Hilbert space on which S is defined. Moreover, each $f \in C_0^\infty(S)$ is an entire analytic vector of S, i.e., for all $t > 0$,

$$\sum_{n=0}^{\infty} \frac{\|S^n f\|}{n!} t^n < \infty.$$

LEMMA 3.6.

(i) *Let* S *be a self-adjoint operator on* $\mathcal{H}$ *and*

$$\mathfrak{D}_S(\mathcal{H}) = \mathcal{L}\{a(f_1)^* \cdots a(f_n)^* \Omega_b | f_j \in C_0^\infty(S), j = 1, \cdots, n, n \geq 0\}.$$

Then $d\Gamma(S)$ *leaves* $\mathfrak{D}_S(\mathcal{H})$ *invariant and each* $\Psi \in \mathfrak{D}_S(\mathcal{H})$ *is an entire analytic vector of* $d\Gamma(S)$.

(ii) *Let* T *be a self-adjoint operator on* $\mathcal{K}$ *and*

$$\mathfrak{D}_T(\mathcal{K}) = \mathcal{L}\{b(u_1)^* \cdots b(u_m)^* \Omega_f | u_j \in C_0^\infty(T), j = 1, \cdots, m, m \geq 0\}.$$

Then $d\Lambda(T)$ *leaves* $\mathfrak{D}_T(\mathcal{K})$ *leaves invariant and each* $\Psi \in \mathfrak{D}_T(\mathcal{K})$ *is an entire analytic vector of* $d\Lambda(T)$.

Proof. Let

$$\Psi = a(f_1)^* \cdots a(f_m)^* \Omega_b, \quad f_j \in C_0^\infty(S), j = 1, \cdots, m.$$

Without loss of generality, we can assume that for some $a, b \in \mathbf{R}$, $f_j \in R(E_S((a,b])), j = 1, \cdots, m$. Then $\Psi \in C^\infty(d\Gamma(S))$ and the vector $d\Gamma(S)^n \Psi$ is written as a sum of m^n terms of the form

$$a(S^{j_1} f_1)^* \cdots a(S^{j_m} f_m)^* \Omega_b, \quad 0 \leq j_k \leq n, k = 1, \cdots, m.$$

The standard number operator estimate gives

$$\|a(S^{j_1} f_1)^* \cdots a(S^{j_m} f_m)^* \Omega_b\| \leq \sqrt{(m-1)!} \prod_{k=1}^{m} \|S^{j_k} f_k\|.$$

Using the spectral representation of S, we can show that

$$\|S^k f_j\| \leq c^k \|f_j\|$$

with $c = \max\{|a|, |b|\}$. Hence we obtain

$$||a(S^{j_1} f_1)^* \cdots a(S^{j_m} f_m)^* \Omega_b|| \leq \sqrt{(m-1)!} c^{mn} \prod_{j=1}^{m} ||f_j||.$$

Therefore

$$||d\Gamma(S)^n \Psi|| \leq m^n c^{mn} \sqrt{(m-1)!} \prod_{j=1}^{m} ||f_j||.$$

This estimate implies that Ψ is an entire analytic vector of $d\Gamma(S)$. Similarly we can prove part (ii). ∎

LEMMA 3.7. *Let $A \in C(\mathcal{H}, \mathcal{K})$ and*

$$\mathfrak{D}_0 = \mathcal{L}\{a(f_1)^* \cdots a(f_n)^* b(u_1)^* \cdots b(u_m)^* \Omega | f_j \in C_0^\infty(A^*A), \\ u_k \in C_0^\infty(AA^*), j = 1, \cdots, n, k = 1, \cdots, m, n, m \geq 0\}.$$

Then Q_A leaves $\mathfrak{D}_0$ invariant and each $\Psi \in \mathfrak{D}_0$ is an entire analytic vector of Q_A.

Proof. By (3.2) and (3.6) one easily sees that $\mathfrak{D}_0 \subset C^\infty(Q_A)$. By Theorem 3.2 we have

$$Q_A^{2n} = \Delta_A^n.$$

It follows from (3.9) and Lemma 3.6 that each $\Psi \in \mathfrak{D}_0$ is an entire analytic vector of Δ_A. Hence we have for all $t > 0$

$$\sum_{n=0}^{\infty} \frac{||Q_A^{2n} \Psi||}{n!} t^n < \infty.$$

Since $Q_A \Psi \in \mathfrak{D}_0$, we have also

$$\sum_{n=0}^{\infty} \frac{||Q_A^{2n+1} \Psi||}{n!} t^n < \infty.$$

These estimates imply that Ψ is an entire analytic vector of Q_A. ∎

Proof of Theorem 3.5: Let $\{A, B\} \in \mathfrak{A}$. The strong commutativity of A^*A and B^*B implies that $C_0^\infty(A^*A) \cap C_0^\infty(B^*B)$ is dense in $\mathcal{H}$. Similarly $C_0^\infty(AA^*) \cap C_0^\infty(BB^*)$ is dense in $\mathcal{K}$. Hence the subspace

$$\mathfrak{D}_{A,B} = \mathcal{L}\{a(f_1)^* \cdots a(f_n)^* b(u_1)^* \cdots b(u_m)^* \Omega | f_j \in C_0^\infty(A^*A) \cap C_0^\infty(B^*B), \\ u_k \in C_0^\infty(AA^*) \cap C_0^\infty(BB^*), j = 1, \cdots, n, k = 1, \cdots, m, n, m \geq 0\}$$

is dense in $\mathfrak{F}$. By (3.12), (3.13), and property (iii) in Definition 3.4, we see that

$$\{Q_A(\lambda), Q_B\} = 0 \quad \text{on } \mathfrak{D}_{A,B}.$$

Each $\Psi \in \mathfrak{D}_{A,B} \subset \mathfrak{D}_0$ is an entire analytic vector for Q_B and $Q_A(\lambda)$ leaves $\mathfrak{D}_{A,B}$ invariant. Hence we have for all $t \in \mathbf{R}$

$$e^{itQ_B}Q_A(\lambda)\Psi = \mathrm{s}-\lim_{N\to\infty}\sum_{n=1}^{N}\frac{(it)^n}{n!}Q_B^nQ_A(\lambda)\Psi$$

$$= \mathrm{s}-\lim_{N\to\infty}Q_A(\lambda)\sum_{n=1}^{N}\frac{(-it)^n}{n!}Q_B^n\Psi.$$

Since

$$\mathrm{s}-\lim_{N\to\infty}\sum_{n=1}^{N}\frac{(-it)^n}{n!}Q_B^n\Psi = e^{-itQ_B}\Psi$$

and $Q_A(\lambda)$ is closed, it follows that $e^{-itQ_B}\Psi \in D(Q_A(\lambda))$ and

$$e^{itQ_B}Q_A(\lambda)\Psi = Q_A(\lambda)e^{-itQ_B}\Psi. \tag{3.16}$$

The operator $Q_A(\lambda)$ is essentially self-adjoint on $\mathfrak{D}_{A,B}$, because Q_A is essentially self-adjoint on $\mathfrak{D}_{A,B}$ (the Nelson analytic vector theorem [12, §X.6]) and λN_F is bounded. Hence (3.16) implies that

$$e^{itQ_B}Q_A(\lambda) \subset Q_A(\lambda)e^{-itQ_B}.$$

Thus $Q_A(\lambda)$ and Q_B strongly anticommute. ∎

4. Fock-Space Representations of the SUSY Algebra

We can now apply Theorem 2.4 to construct representations of the SUSY algebra (1.1) on the Boson-Fermion Fock space $\mathfrak{F}$. Let $\{A, B\} \in \mathfrak{A}$ and define

$$H_{A,B}(\lambda) = \frac{1}{2}(Q_A(\lambda)^2 + Q_B^2), \tag{4.1}$$

$$P_{A,B}(\lambda) = \frac{1}{2}\overline{(Q_A(\lambda)^2 - Q_B^2)} \tag{4.2}$$

Theorems 3.5 and 2.4 imply the following result.

THEOREM 4.1. *Let $\{A, B\} \in \mathfrak{A}$ and $\lambda \in \mathbf{R}$. Then $\{Q_A(\lambda), Q_B, H_{A,B}(\lambda), P_{A,B}(\lambda)\}$ is a representation of (1.1) with $Q_1 = Q_A(\lambda), Q_2 = Q_B, H = H_{A,B}(\lambda), P = P_{A,B}(\lambda)$. Moreover we have*

$$H_{A,B}(\lambda) = \frac{1}{2}\{d\Gamma(A^*A + B^*B) + d\Lambda(AA^* + BB^*) + \lambda^2\}, \tag{4.3}$$

$$P_{A,B} = \frac{1}{2}\{d\Gamma(\overline{A^*A - B^*B}) + d\Lambda(\overline{AA^* - BB^*}) + \lambda^2\}. \tag{4.4}$$

Proof. It remains to prove formulae (4.3) and (4.4). They easily follow from (4.1), (4.2), Theorem 3.2 and (3.15). ∎

Let $\lambda \neq 0$. Then formula (4.3) shows that $H_{A,B}(\lambda) \geq \lambda^2/2 > 0$ and hence the supersymmetric Hamiltonian $H_{A,B}(\lambda)$ has no zero-energy states, i.e., the supersymmetry is broken. Similarly fromula (3.15) implies that there exists no vector Ψ such that $Q_A(\lambda)\Psi = 0$. On the other hand, we have $Q_B\Omega = 0$. Thus, in the case $\lambda \neq 0$, the supersymmetry is partially broken in the representation given in Theorem 4.1. One can completely identify the subspace Ker Q_A [4].

We next consider the unitary equivalence of the representations $\{Q_A(\lambda), Q_B, H_{A,B}(\lambda), P_{A,B}(\lambda)\}$, $\{A, B\} \in \mathfrak{A}$.

DEFINITION 4.2. *Let $\{Q_1, Q_2, H, P\}$ and $\{Q_1', Q_2', H', P'\}$ be representations of (1.1) on Hilbert spaces $\mathfrak{H}$ and $\mathfrak{H}'$, respectively. They are said to be unitarily equivalent if there exists a unitary operatror $\Upsilon : \mathfrak{H} \to \mathfrak{H}'$ such that*

$$Q_j' = \Upsilon Q_j \Upsilon^{-1}, j = 1,2,\ H' = \Upsilon H \Upsilon^{-1},\ P' = \Upsilon P \Upsilon^{-1}.$$

The following fact is easily proven.

LEMMA 4.3. *Let $\mathcal{H}', \mathcal{K}'$ be Hilbert spaces and $\{A, B\} \in \mathfrak{A}(\mathcal{H}, \mathcal{K})$. Let $U : \mathcal{H} \to \mathcal{H}'$ and $V : \mathcal{K} \to \mathcal{K}'$ be unitary operators and set*

$$A' = VAU^{-1}, \quad B' = VBU^{-1}. \tag{4.5}$$

Then $\{A', B'\} \in \mathfrak{A}(\mathcal{H}', \mathcal{K}')$.

THEOREM 4.4. *Let $\{A, B\} \in \mathfrak{A}(\mathcal{H}, \mathcal{K})$ and $\{A', B'\} \in \mathfrak{A}(\mathcal{H}', \mathcal{K}')$ be as in Lemma 4.3 and $\lambda \in \mathbf{R}$. Then $\{Q_A(\lambda), Q_B, H_{A,B}(\lambda), P_{A,B}(\lambda)\}$ is unitarily equivalent to $\{Q_{A'}(\lambda), Q_{B'}, H_{A',B'}(\lambda), P_{A',B'}(\lambda)\}$.*

Proof. Let (4.5) be satisfied. It is well-known that the operators

$$\Gamma(U) = \bigoplus_{n=0}^{\infty} \underbrace{U \otimes U \otimes \cdots \otimes U}_{n \text{ factors}} : \mathfrak{F}_{\mathrm{b}}(\mathcal{H}) \to \mathfrak{F}_{\mathrm{b}}(\mathcal{H}')$$

and

$$\Lambda(V) = \bigoplus_{m=0}^{\infty} \underbrace{V \otimes V \otimes \cdots \otimes V}_{m \text{ factors}} : \mathfrak{F}_{\mathrm{f}}(\mathcal{K}) \to \mathfrak{F}_{\mathrm{f}}(\mathcal{K}')$$

are unitary [12, §X.7]. Hence

$$\Upsilon = \Gamma(U) \otimes \Lambda(V) : \mathfrak{F}(\mathcal{H}, \mathcal{K}) \to \mathfrak{F}(\mathcal{H}', \mathcal{K}')$$

is unitary. Direct computations using (4.5) show that

$$\Upsilon Q_A(\lambda) \Upsilon^{-1} = Q_{A'}(\lambda), \quad \Upsilon Q_B \Upsilon^{-1} = Q_{B'},$$

on $\mathfrak{D}_{A',B'}$. Since $\mathfrak{D}_{A',B'}$ is a common core for $Q_{A'}(\lambda)$ and $Q_{B'}$, it follows that $\{Q_A(\lambda), Q_B\}$ is unitarily equivalent to $\{Q_{A'}(\lambda), Q_{B'}\}$ via Υ. In the present case, the unitary equivalence

of $\{Q_A(\lambda), Q_B\}$ and $\{Q_{A'}(\lambda), Q_{B'}\}$ implies that of $\{H_{A,B}(\lambda), P_{A,B}(\lambda)\}$ and $\{H_{A',B'}(\lambda), P_{A',B'}(\lambda)\}$. ∎

To formulate a sufficient condition for two representations given in Thereom 4.1 not to be unitarily equivalent, we intoroduce a concept related to the spectrum of $\{A, B\} \in \mathfrak{A}(\mathcal{H}, \mathcal{K})$. For $\{A, B\} \in \mathfrak{A}(\mathcal{H}, \mathcal{K})$, we have

$$S_{A,B} \equiv \sigma(A^*A + B^*B) = \{\lambda + \mu | \lambda \in \sigma(A^*A), \mu \in \sigma(B^*B)\}^-,$$

where $\{\cdot\}^-$ denotes the closure of the set $\{\cdot\}$. Let

$$\Sigma_{A,B} = \cup_{m,n=0,m+n\geq 1}^{\infty} \{\lambda_1 + \cdots + \lambda_n + \mu_1 + \cdots + \mu_m | \lambda_j \in S_{A,B}, \\ \mu_k \in S_{A^*,B^*}, j = 1, 2 \cdots, n, k = 1, \cdots, m, \}.$$

THEOREM 4.5. *Let* $\{A, B\} \in \mathfrak{A}(\mathcal{H}, \mathcal{K}), \{A', B'\} \in \mathfrak{A}(\mathcal{H}', \mathcal{K}')$. *Suppose that* $\Sigma^-_{A,B} \neq \Sigma^-_{A',B'}$. *Then* $\{Q_A(\lambda), Q_B, H_{A,B}(\lambda), P_{A,B}(\lambda)\}$ *is not unitarily equivalent to* $\{Q_{A'}(\lambda), Q_{B'}, H_{A',B'}(\lambda), P_{A',B'}(\lambda)\}$.

Proof. By the assumption and the fact

$$\sigma(2H_{A,B}(\lambda)) = \{x + y + \lambda^2 | x \in \sigma(d\Gamma(A^*A + B^*B)), y \in \sigma(d\Lambda(AA^* + BB^*))\}^- \\ = \{\lambda^2\} \cup \{\Sigma^-_{A,B} + \lambda^2\},$$

we have $\sigma(2H_{A,B}(\lambda)) \neq \sigma(2H_{A',B'}(\lambda))$, which implies that $H_{A,B}(\lambda)$ cannot be unitarily equivalent to $H_{A',B'}(\lambda)$. Hence the desired result follows. ∎

We give only one example.

EXAMPLE: Let $\mathcal{H} = \mathcal{K} = L^2(\mathbf{R})$ and $\omega(p)$ be a nonnegative measurable function on $\mathbf{R}$ such that

$$|p| \leq \omega(p), \quad p \in \mathbf{R}.$$

Let

$$\nu(p) = (p + \omega(p))^{1/2}$$

and $\theta(p)$ be a real-valued measurable function on $\mathbf{R}$. Define the operators A and B on $L^2(\mathbf{R})$ to be the multiplication operatros by the functions

$$A(p) = i\nu(p)e^{i\theta(p)}$$

and

$$B(p) = \nu(-p)e^{i\theta(p)},$$

respectively. Then we have

$$A^*A = AA^* = p + \omega,$$
$$B^*B = BB^* = -p + \omega,$$
$$A^*B = -i\nu(p)\nu(-p) = -i\sqrt{\omega^2 - p^2},$$
$$B^*A = i\nu(p)\nu(-p) = i\sqrt{\omega^2 - p^2}.$$

These relations imply that $\{A, B\}$ is in $\mathfrak{A}$ under consideration and

$$H_{A,B}(\lambda) = d\Gamma(\omega) + d\Lambda(\omega) + \frac{\lambda^2}{2},$$

$$P_{A,B}(\lambda) = d\Gamma(p) + d\Lambda(p) + \frac{\lambda^2}{2},$$

Note that $H_{A,B}(\lambda)$ and $P_{A,B}(\lambda)$ are independent of θ. If $\omega(p) = \sqrt{p^2 + m^2}$ with a constant $m \geq 0$, then $H_{A,B}(0)$ and $P_{A,B}(0)$ are the Hamiltonian and the momentum operator of a free relativistic SQFT in the two-dimensional space-time. The present example with $\theta = 0$ is related to the $N = 1$ Wess-Zumino model in the two-dimensional space-time (cf.[8,9]).

References

1. Arai, A., Supersymmetry and singular perturbations, *J.Funct.Anal.* **60**, 378-393(1985).
2. Arai, A., Path integral representation of the index of Kähler-Dirac operators on an infinite dimensional manifold, *J.Funct.Anal.***82**, 330-369(1989)
3. Arai, A., Supersymmetric embedding of a model of a quantum harmonic oscillator interacting with infinitely many bosons, *J.Math.Phys.***30**, 512-520(1989).
4. Arai,A., A general class of infinite dimensional Dirac operators and path integral representation of their index, *J.Funct.Anal.* (in press)
5. Arai,A., De Rham operators, Laplacians, and Dirac operators on topological vector spaces, *Hokkaido Univ. Preprint Series in Math.* No.115, 1991.
6. Arai, A., and Mitoma, I., De Rham-Hodge-Kodaira decomposition in ∞-dimensions, *Math.Ann.* (in press).
7. Jaffe,A., Lesniewski,A., and Weitsman, J., The two-dimensional, N=2 Wess-Zumino model on a cylinder, *Commun.Math.Phys.* **114**, 147-165(1988).
8. Jaffe,A., Lesniewski,A., and Weitsman, J., The loop space $S^1 \to \mathbf{R}$ and supersymmetric quantum fields, *Ann.Phys.* **114**, 553-575(1988).
9. Jaffe,A., Lesniewski,A., and Wieczerkowski, C., Supersymmetry and the spectral condition on a cylinder, *Lett.Math.Phys.* **16**, 385-388(1988).
10. Pedersen, S., Anticommuting self-adjoint operators, *J.Funct.Anal.***89**, 428-443(1990).
11. Reed, M., and Simon, B., *Methods of Modern Mathematical Physics Vol.I:Functional Analysis*, Academic Press, New York, 1972.
12. Reed, M., and Simon, B., *Methods of Modern Mathematical Physics Vol.II:Fourier Analysis, Self-adjointness*, Academic Press, New York, 1975.
13. Samoilenko, *Spectral Theory of Families of Self-Adjoint Operators*, Kluwer Academic Publishers, Dordrecht, 1991.
14. Simon, B., *The $P(\phi)_2$ Euclidean (Quantum) Field Theory*, Princeton University Press, Princeton, New Jersey, 1974.
15. Vasilescu, F.-H., Anticommuting self-adjoint operators, *Rev.Roum.Math.Pures et Appl.*28, 77-91(1983).
16. West, P, *Introduction to Supersymmetry and Supergravity* (Extended Second Edition), World Scientific, Singapore, 1990.
17. Witten, E., Supersymmetry and Morse theory, *J.Diff.Geom.*17, 661-692(1982).