HOKKAIDO UNIVERSITY

Title	Canonical modules and Cohen-Macaulay types of partially ordered sets
Author(s)	Hibi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 184, 1-6
Issue Date	1993-03
DOI	https://doi.org/10.14943/83328
Doc URL	https://hdl.handle.net/2115/68930
Type	departmental bulletin paper
File Information	re184.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Canonical modules and Cohen-Macaulay types of partially ordered sets

Takayuki Hibi

Series ♯184. March 1993

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

♯ 156: S. Izumiya, Perestroikas of optical wave fronts and graphlike Legendrian unfoldings, 13 pages. 1992.

♯ 157: A. Arai, Momentum operators with gauge potentials, local quantization of magnetic flux, and representation of canonical commutation relations, 11 pages. 1992.

♯ 158: S. Izumiya, W.L. Marar, The Euler number of a topologically stable singular surface in a 3-manifold, 11 pages. 1992.

♯ 159: T. Hibi, Cohen-Macaulay types of Cohen-Macaulay complexes, 26 pages. 1992.

♯ 160: A. Arai, Properties of the Dirac-Weyl operator with a strongly singular gauge potential, 26 pages. 1992.

♯ 161: A. Arai, Dirac operators in Boson-Fermion Fock spaces and supersymmetric quantum field theory, 30 pages. 1992.

♯ 162: S. Albeverio, K. Iwata, T. Kolsrud, Random parallel transport on surfaces of finite type, and relations to homotopy, 8 pages. 1992.

♯ 163: S. Albeverio, K. Iwata, T. Kolsrud, Moments of random fields over a family of elliptic curves, and modular forms, 9 pages. 1992.

♯ 164: Y. Giga, M. Sato, Neumann problem for singular degenerate parabolic equations, 12 pages. 1992.

♯ 165: J. Wierzbicki, Y. Watatani, Commuting squares and relative entropy for two subfactors, 18 pages. 1992.

♯ 166: Y. Okabe, A new algorithm driven from the view-point of the fluctuation-dissipation theorem in the theory of KM_2O-Langevin equations, 13 pages. 1992.

♯ 167: Y. Okabe, H. Mano and Y. Itoh, Random collision model for interacting populations of two species and its strong law of large numbers, 14 pages. 1992.

♯ 168: A. Inoue, On the equations of stationary precesses with divergent diffusion coefficients, 25 pages. 1992.

♯ 169: T. Ozawa, Remarks on quadratic nonlinear Schrödinger equations, 19 pages. 1992.

♯ 170: T. Fukui, Y. Giga, Motion of a graph by nonsmooth weighted curvature, 11 pages. 1992.

♯ 171: J. Inoue, T. Nakazi, Finite dimensional solution sets of extremal problems in H^1, 10 pages. 1992.

♯ 172: S. Izumiya, A characterization of complete integrability for partial differential equations of first order, 6 pages. 1992.

♯ 173: T. Suwa, Unfoldings of codimension one complex analytic foliation singularities, 49 pages. 1992.

♯ 174: T. Ozawa, Wave propagation in even dimensional spaces, 15 pages. 1992.

♯ 175: S. Izumiya, Systems of Clairaut type, 7 pages. 1992.

♯ 176: A. Hoshiga, The initial value problems for quasi-linear wave equations in two space dimensions with small data, 25 pages. 1992.

♯ 177: K. Sugano, On bicommutators of modules over H-separable extension rings III, 9 pages. 1993.

♯ 178: T. Nakazi, Toeplitz operators and weighted norm inequalities, 17 pages. 1993.

♯ 179: O. Ogurisu, Existence and structure of infinitely degenerate zero-energy ground states of a Wess-Zumino type model in supersymmetric quantum mechanics, 26 pages. 1993.

♯ 180: O. Ogurisu, Ground state of a spin 1/2 charged particle in an even dimensional magnetic field, 9 pages. 1993.

♯ 181: K. Sugano, Note on H-separable Galois extension, 6 pages. 1993.

♯ 182: M. Yamada, Distance formulas of asymptotic Toeplitz and Hankel operators, 13 pages. 1993

♯ 183: G. Ishikawa, T. Ozawa, The genus of a connected compact real algebraic surface in the affine three space, 11 pages. 1993.

Advances in Mathematics, to appear.

Canonical modules and Cohen-Macaulay types of partially ordered sets

Takayuki Hibi

Department of Mathematics
Faculty of Science
Hokkaido University
Kita-ku, Sapporo 060, Japan

Every partially ordered set (or "poset" for short) to be studied is finite. We write $\#(X)$ for the cardinality of a finite set X.

(1.1) Let P be a poset. Then we set $P^ = P \cup \{ 0^ , 1^ \}$ with $0^ < \alpha < 1^$ for every $\alpha \in P$. A chain C of the poset $P^$ is a totally ordered subset of $P^$. From now on, we assume that $P^$ is graded of rank $d+1$ [6, p.99], that is to say, there exists a unique rank function $\rho : P^ \to \{0,1, \ldots ,d+1\}$ such that $\rho(0^) = 0$, $\rho(1^) = d + 1$, and $\rho(\beta) = \rho(\alpha) + 1$ if $\alpha < \beta$ and $\alpha < \gamma < \beta$ for no $\gamma \in P^$.

(1.2) Fix a field k. Let $A = k[x ; x \in P]$ be the polynomial ring over k with the "standard grading," i.e., each $\deg x = 1$, whose variables are the elements of P. We write I for the ideal of A generated by those quadratic monomials xy such that x and y are incomparable in P. The quotient algebra $k[P] = A/I$ is called the *Stanley-Reisner ring* of P over k. Define $\vartheta_i \in k[P]$ by $\vartheta_i = \Sigma_{\rho(x)=i} x$ for each $1 \leq i \leq d$. Then the sequence $\vartheta_1 , \vartheta_2 , \ldots , \vartheta_d$ is a system of parameters for $k[P]$, in other words, (i) $\vartheta_1 , \ldots , \vartheta_d$ are algebraically independent over k and (ii) $k[P]$ is finitely generated as a module over the

subalgebra $k[\Theta] = k[\theta_1,\theta_2,\ldots,\theta_d]$. We say that $k[P]$ is *Cohen-Macaulay* (or P is Cohen-Macaulay over k) if $k[P]$ is a free module over $k[\Theta]$.

For example, if $L = P\hat{}$ is a semimodular lattice then $k[P]$ is Cohen-Macaulay. We refer the reader to, e.g., [2] for basic information about Cohen-Macaulay posets.

(1.3) Suppose that the Stanley-Reisner ring $k[P]$ of a poset P is Cohen-Macaulay. First, the *canonical module* $\Omega(k[P])$ of $k[P]$ is defined to be the graded module

$$\Omega(k[P]) = \mathrm{Hom}_{k[\Theta]}(k[P],k[\Theta])$$

over $k[P]$. Secondly, the *socle* $\mathrm{Soc}(k[P]/(\Theta))$ of $k[P]/(\Theta)$ is

$$\mathrm{Soc}(k[P]/(\Theta)) = \{\, y \in k[P]/(\Theta)\,;\, xy = 0 \text{ for every } x \in P \,\},$$

where (Θ) is the parameter ideal $(\theta_1,\theta_2,\ldots,\theta_d)$ of $k[P]$. The dimension $\dim_k \mathrm{Soc}(k[P]/(\Theta))$ of $\mathrm{Soc}(k[P]/(\Theta))$ as a vector space over k is called the *Cohen-Macaulay type* of $k[P]$.

It is known, e.g., [3, Corollary 6.11] that the Cohen-Macaulay type of $k[P]$ is equal to the minimal number of generators of $\Omega(k[P])$ as a module over $k[P]$. Moreover, in the language of homological algebra, the Cohen-Macaulay type of $k[P]$ is equal to $\dim_k \mathrm{Tor}_{v-d}^A(k[P],k)$ with $v = \#(P)$.

(1.4) Now, if $\alpha < \beta$ in $P\hat{}$ with $\rho(\beta) - \rho(\alpha) = r + 1$, then the $(r-1)$-th reduced homology group $\tilde{H}_{r-1}((\alpha,\beta);k)$ of the open interval $(\alpha,\beta) = \{ x \in P\hat{}\,;\, \alpha < x < \beta \}$ of $P\hat{}$ can be imbedded in $k[P]$. Given a chain $C : 0\hat{} = x_0 < x_1 < \ldots < x_s < x_{s+1} = 1\hat{}$ of $P\hat{}$ with each $\rho(x_{i+1}) - \rho(x_i) = r(i) + 1$, we write $\mathcal{R}(C)$ for the subspace of $k[P]$ spanned by those polynomials

$$f_0 x_1^{2} f_1 x_2^{2} \cdots f_{s-1} x_s^{2} f_s$$

with $f_i \in \tilde{H}_{r(i)-1}((x_i,x_{i+1});k)$ for every $0 \leq i \leq s$. Here we

employ the convention that each monimial of f_i is of the form $\alpha_1\alpha_2\ldots\alpha_{r(i)}$ with $x_i < \alpha_1 < \alpha_2 < \ldots < \alpha_{r(i)} < x_{i+1}$. Moreover, we define $\mathcal{Q}(C)$ to be the subspace of $k[P]/(\theta)$ which is the image of $x_1^{-2}x_2^{-2}\ldots x_s^{-2}\mathcal{R}(C)$ in $k[P]/(\theta)$.

(1.5) Let $\mu = \mu_{P^\wedge}$ be the Möbius function of $P^\wedge$ as defined in, e.g., [6, p.116]. If $C : 0^\wedge = x_0 < x_1 < \ldots < x_s < x_{s+1} = 1^\wedge$ is a chain of $P^\wedge$, then we set

$$\mu(C) = \mu(x_0,x_1)\mu(x_1,x_2)\ldots\mu(x_s,x_{s+1}).$$

Thus, in particular,

$$\dim_k(\mathcal{R}(C)) = |\mu(C)|.$$

We say that a chain $C : 0^\wedge = x_0 < x_1 < \ldots < x_s < x_{s+1} = 1^\wedge$ of $P^\wedge$ is *essential* if $\mu(C) \neq 0$. Let $\mathcal{E}(P^\wedge)$ be the set of essential chains of $P^\wedge$ and $\mathcal{E}^*(P^\wedge)$ ($\subset \mathcal{E}(P^\wedge)$) the set of minimal essential chains (or *fundamental chains* [4]) of $P^\wedge$.

We write $\mathcal{I}^*(k[P])$ for the ideal of $k[P]$ generated by all $\mathcal{R}(C)$ with $C \in \mathcal{E}^*(P^\wedge)$. Moreover, we define $\mathcal{J}^*(k[P])$ to be the ideal of $k[P]$ which is generated by those square-free monomials $x_1x_2\ldots x_s$ such that $x_i < x_{i+1}$ in P for every $1 \leq i \leq s$ and $0^\wedge = x_0 < x_1 < \ldots < x_s < x_{s+1} = 1^\wedge \in \mathcal{E}^*(P^\wedge)$.

We are now in the position to state our main result in the paper.

(1.6) THEOREM. Suppose that the Stanley-Reisner ring $k[P]$ of a poset P is Cohen-Macaulay. Then the ideal $\mathcal{I}^*(k[P])$ is isomorphic to the canonical module $\Omega(k[P])$ of $k[P]$, up to shift in grading, if and only if the Cohen-Macaulay type of $k[P]$ is equal to $\sum_{C\in\mathcal{E}^*(P^\wedge)} |\mu(C)|$.

Proof. Let $\mathcal{I}(k[P])$ be the ideal of $k[P]$ generated by all $\mathcal{R}(C)$ with $C \in \mathcal{E}(P^\wedge)$. The monomorphism ψ in [1, Theorem 1]

enables us to define a monomorphism $\psi : \mathcal{J}(k[P]) \to \Omega(k[P])$ of graded modules over $k[P]$ in the analogous way. Note that the lowest degree of a non-zero homogeneous element of $\Omega(k[P])$ (resp. $\mathcal{J}(k[P])$) is $-d + \min\{\#(C); C \in \mathcal{E}(P^\wedge)\}$ (resp. $d + \min\{\#(C); C \in \mathcal{E}(P^\wedge)\}$), while ψ has degree $-2d$. We write $\mathcal{Q}(k[P])$ for the subspace of $k[P]$ which is spanned by those polynomials $f_0 x_1^{n_1} f_1 x_2^{n_2} \cdots f_{s-1} x_s^{n_s} f_s$ such that $0^\wedge = x_0 < x_1 < \cdots < x_s < x_{s+1} = 1^\wedge \in \mathcal{E}(P^\wedge)$ with $\rho(x_{i+1}) - \rho(x_i) = r(i) + 1$, $f_i \in \tilde{H}^{r(i)-1}((x_i, x_{i+1}); k)$ for every $0 \leq i \leq s$, and each $n_i \geq 2$ is an integer. Since $\mathcal{Q}(k[P]) \subset \mathcal{J}(k[P])$, it follows from [5, Theorem 7.1, p.80] that the above monomorphism $\psi : \mathcal{J}(k[P]) \to \Omega(k[P])$ is an isomorphism and $\mathcal{Q}(k[P]) = \mathcal{J}(k[P])$. Moreover, each subspace $\mathcal{R}(C)$ with $C \in \mathcal{E}^*(P^\wedge)$ is contained in the subspace of $k[P]$ spanned by an arbitrary system of generators of the ideal $\mathcal{J}(k[P])$. Hence $\mathcal{J}^*(k[P]) = \mathcal{J}(k[P])$ if and only if the minimal number of generators of $\Omega(k[P])$, i.e., the Cohen-Macaulay type of $k[P]$, is equal to $\sum_{C \in \mathcal{E}^*(P^\wedge)} |\mu(C)|$ as desired. Q.E.D.

When $\mu(0^\wedge, 1^\wedge) \neq 0$, the above Theorem (1.6) essentially coincides with [1, Theorem 2]. Now, it is shown in [4] that, for a poset P such that $L = P^\wedge$ is a modular lattice, the Cohen-Macaulay type of $k[P]$ is equal to $\sum_{C \in \mathcal{E}^*(P^\wedge)} |\mu(C)|$.

(1.7) COROLLARY. If $L = P^\wedge$ is a modular lattice, then $\mathcal{J}^*(k[P])$ is isomorphic to the canonical module $\Omega(k[P])$ of $k[P]$.

REMARK. Every subspace $\mathcal{Q}(C)$ of $k[P]/(\theta)$ with $C \in \mathcal{E}^*(P^\wedge)$ is contained in $\mathrm{Soc}(k[P]/(\theta))$. However, even though the Cohen-Macaulay type of $k[P]$ is equal to $\sum_{C \in \mathcal{E}^*(P^\wedge)} |\mu(C)|$, the subspace of $\mathrm{Soc}(k[P]/(\theta))$ spanned by all $\mathcal{Q}(C)$ with $C \in \mathcal{E}^*(P^\wedge)$ does not necessarily coincide with $\mathrm{Soc}(k[P]/(\theta))$.

It might be of interest to give a similar result to Hochster's theorem [5, Theorem7.3, p.81]. We refer the reader to, e.g., [5, p.28] for the definition of an orientable pseudo-manifold with boundary. It follows easily that the order complex $\Delta(P)$

(see, e.g., [6, p.120]) of a Cohen-Macaulay poset P possesses the connectivity property [5, Definition 3.15 (c), p.28]. Moreover, if the order complex $\Delta(P)$ of a Cohen-Macaulay poset P over a field k is an orientable pseudo-manifold with boundary, then the poset P satisfies (☆) $|\mu(x,y)| \leq 1$ for every $0\hat{} \leq x \leq y \leq 1\hat{}$ in $P\hat{}$.

(1.8) PROPOSITION. Let P be a Cohen-Macaulay poset over k. Then there exists a homogeneous non-zero divisor $\Theta \in k[P]$ of degree d with $\vartheta^*(k[P])\cdot\Theta = \vartheta(k[P])$ ($= \Omega(k[P])$) if and only if $\Delta(P)$ is an orientable pseudo-manifold with boundary.

Proof. Let $\mathfrak{M}(P\hat{})$ be the set of maximal chains of $P\hat{}$. Given a function $\varepsilon : \mathfrak{M}(P\hat{}) \to k - \{0\}$, for each $F : 0\hat{} = y_0 < y_1 < \ldots < y_d < y_{d+1} = 1\hat{} \in \mathfrak{M}(P\hat{})$, we set $m(F;\varepsilon) = \varepsilon(F) \prod_{1\leq i\leq d} y_i \in k[P]$, and define $\Theta(\varepsilon) \in k[P]$ by $\Theta(\varepsilon) = \sum_{F\in \mathfrak{M}(P\hat{})} m(F;\varepsilon)$, which is a homogeneous non-zero divisor on $k[P]$ of degree d.

Now, let P be a Cohen-Macaulay poset over k and suppose the existence of a homogeneous non-zero divisor $\Theta \in k[P]$ of degree d with $\vartheta^*(k[P])\cdot\Theta = \Omega(k[P])$. Then Θ must be of the form $\Theta(\varepsilon)$ for some ε. Hence, the poset P satisfies the above condition (☆), in particular, $\Delta(P)$ is a pseudo-manifold with boundary. If $F : 0\hat{} < \alpha_1 < \ldots < \alpha_p < \gamma < \beta_1 < \ldots < \beta_q < 1\hat{}$, $F' : 0\hat{} < \alpha_1 < \ldots < \alpha_p < \delta < \beta_1 < \ldots < \beta_q < 1\hat{} \in \mathfrak{M}(P\hat{})$ with $\gamma \neq \delta$, then $\varepsilon(F) = -\varepsilon(F')$ since $C : 0\hat{} < \alpha_1 < \ldots < \alpha_p < \beta_1 < \ldots < \beta_q < 1\hat{} \in \mathfrak{E}(P\hat{})$ and $\alpha_1{}^2 \ldots \alpha_p{}^2 (\gamma - \delta) \beta_1{}^2 \ldots \beta_q{}^2 \in \mathfrak{R}(C)$. Thus $0 \neq \Theta \in H_{d-1}(\Delta(P),\partial\Delta(P);k)$, hence $\Delta(P)$ is orientable.

On the other hand, suppose that the order complex $\Delta(P)$ of a Cohen-Macaulay poset P over k is an orientable pseudo-manifold with boundary. Since $H_{d-1}(\Delta(P),\partial\Delta(P);k) \neq (0)$, there exists $\varepsilon : \mathfrak{M}(P\hat{}) \to k - \{0\}$ such that $\varepsilon(F) = -\varepsilon(F')$ if $\#(F \cap F') = d - 1$. Then $0 \neq x_1x_2 \ldots x_s\Theta(\varepsilon) \in \mathfrak{R}(C)$ for every $C : 0\hat{} = x_0 < x_1 < \ldots < x_s < x_{s+1} = 1\hat{} \in \mathfrak{E}(P\hat{})$. Thus, since $\dim_k \mathfrak{R}(C) = 1$, the subspace $\mathfrak{R}(C)$ is spanned by $x_1x_2 \ldots x_s\Theta$ over k. Hence $\vartheta^*(k[P])\cdot\Theta = \Omega(k[P])$ as required. **Q.E.D.**

REFERENCES

[1] K. Baclawski, Canonical modules of partially ordered sets, J. of Algebra 83 (1983), 1-5.

[2] A. Björner, A. Garsia and R. Stanley, An introduction to Cohen-Macaulay partially orgered sets, in "Ordered Sets" (I. Rival, ed.), Reidel, Dordrecht/Boston, 1982, pp. 583-615.

[3] J. Herzog and E. Kunz, "Der kanonische Modul eines Cohen-Macaulay-Rings," Lect. Notes in Math. No. 238, Springer-Verlag, Berlin/Heidelberg/New York, 1971.

[4] T. Hibi, Cohen-Macaulay types of Cohen-Macaulay complexes, J. Algebra, submitted.

[5] R. Stanley, "Combinatorics and Commutative Algebra," Birkhäuser, Boston/Basel/Stuttgart, 1983.

[6] _____, "Enumerative Combinatorics, Volume I," Wadsworth & Brooks/Cole, Monterey, Calif., 1986.