



HOKKAIDO UNIVERSITY

Title	Multipliers of invariant subspaces in the bidisc
Author(s)	Nakazi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 189, 1-12
Issue Date	1993-03
DOI	https://doi.org/10.14943/83333
Doc URL	https://hdl.handle.net/2115/68935
Type	departmental bulletin paper
File Information	re189.pdf



**Multipliers Of Invariant Subspaces
In The Bidisc**

Takahiko Nakazi

Series #189. March 1993

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 162: S. Albeverio, K. Iwata, T. Kolsrud, Random parallel transport on surfaces of finite type, and relations to homotopy, 8 pages. 1992.
- # 163: S. Albeverio, K. Iwata, T. Kolsrud, Moments of random fields over a family of elliptic curves, and modular forms, 9 pages. 1992.
- # 164: Y. Giga, M. Sato, Neumann problem for singular degenerate parabolic equations, 12 pages. 1992.
- # 165: J. Wierzbicki, Y. Watatani, Commuting squares and relative entropy for two subfactors, 18 pages. 1992.
- # 166: Y. Okabe, A new algorithm driven from the view-point of the fluctuation-dissipation theorem in the theory of KM_2O -Langevin equations, 13 pages. 1992.
- # 167: Y. Okabe, H. Mano and Y. Itoh, Random collision model for interacting populations of two species and its strong law of large numbers, 14 pages. 1992.
- # 168: A. Inoue, On the equations of stationary precesses with divergent diffusion coefficients, 25 pages. 1992.
- # 169: T. Ozawa, Remarks on quadratic nonlinear Schrödinger equations, 19 pages. 1992.
- # 170: T. Fukui, Y. Giga, Motion of a graph by nonsmooth weighted curvature, 11 pages. 1992.
- # 171: J. Inoue, T. Nakazi, Finite dimensional solution sets of extremal problems in H^1 , 10 pages. 1992.
- # 172: S. Izumiya, A characterization of complete integrability for partial differential equations of first order, 6 pages. 1992.
- # 173: T. Suwa, Unfoldings of codimension one complex analytic foliation singularities, 49 pages. 1992.
- # 174: T. Ozawa, Wave propagation in even dimensional spaces, 15 pages. 1992.
- # 175: S. Izumiya, Systems of Clairaut type, 7 pages. 1992.
- # 176: A. Hoshiga, The initial value problems for quasi-linear wave equations in two space dimensions with small data, 25 pages. 1992.
- # 177: K. Sugano, On bicommutators of modules over H-separable extension rings III, 9 pages. 1993.
- # 178: T. Nakazi, Toeplitz operators and weighted norm inequalities, 17 pages. 1993.
- # 179: O. Ogürisu, Existence and structure of infinitely degenerate zero-energy ground states of a Wess-Zumino type model in supersymmetric quantum mechanics, 26 pages. 1993.
- # 180: O. Ogürisu, Ground state of a spin 1/2 charged particle in an even dimensional magnetic field, 9 pages. 1993.
- # 181: K. Sugano, Note on H-separable Galois extension, 6 pages. 1993.
- # 182: M. Yamada, Distance formulas of asymptotic Toeplitz and Hankel operators, 13 pages. 1993.
- # 183: G. Ishikawa, T. Ozawa, The genus of a connected compact real algebraic surface in the affine three space, 11 pages. 1993.
- # 184: T. Hibi, Canonical modules and Cohen-Macaulay types of partially ordered sets, 6 pages. 1993.
- # 185: Y. Giga, K. Yama-uchi, On a lower bound for the extinction time of surfaces moved by mean curvature, 16 pages. 1993.
- # 186: Y. Kurokawa, On functional moduli for first order ordinary differential equations, 9 pages. 1993.
- # 187: O. Ogürisu, Unitary equivalence between a spin 1/2 charged particle in a two-dimensional magnetic field and a spin 1/2 neutral particle with an anomalous magnetic moment in a two-dimensional electric field, 4 pages. 1993.
- # 188: A. Jensen, T. Ozawa, Existence and non-existence results for wave operators for perturbations of the laplacian, 30 pages. 1993.

Multipliers Of Invariant Subspaces In The Bidisc

By

Takahiko Nakazi*

Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060, Japan

*This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education

1980 Mathematics Subject Classification (1985 Revision): Primary 46 J 15, 47 A 15, 47 B 20 ; Secondary 32 A 35.

Keywords and phrases : bidisc, invariant subspace, multipliers

Abstract. For any nonzero invariant subspace M in $H^2(T^2)$, set $M^\times = [\bigcup_{n=0}^{\infty} \bar{z}^n M] \cap [\bigcup_{n=0}^{\infty} \bar{w}^n M]$ then $M^\times$ is also an invariant subspace of $H^2(T^2)$ that contains M . If M is of finite codimension in $H^2(T^2)$ then $M^\times = H^2(T^2)$ and if $M = qH^2(T^2)$ for some inner function q then $M^\times = M$. In this paper invariant subspaces with $M^\times = M$ are studied. If $M = q_1 H^2(T^2) \cap q_2 H^2(T^2)$ and q_1, q_2 are inner functions then $M^\times = M$. However in general this invariant subspace may not be of the form $qH^2(T^2)$ for some inner function q . Put $\mathcal{M}(M) = \{\phi \in L^\infty : \phi M \subseteq H^2(T^2)\}$ then $\mathcal{M}(M)$ is described and $\mathcal{M}(M) = \mathcal{M}(M^\times)$ is shown. This is the set of all multipliers of M in the title. A necessary and sufficient condition for $\mathcal{M}(M) = H^\infty(T^2)$ is given. It is noted that the kernel of a Hankel operator is an invariant subspace M with $M^\times = M$. The argument applies to the polydisc case.

§1. Introduction

Let T^2 be the torus that is the Cartesian product of 2 unit circles in $\mathbb{C}$. The usual Lebesgue spaces, with respect to the Haar measure m of T^2 , are denoted by $L^p = L^p(T^2)$, and the Hardy spaces $H^p = H^p(T^2)$ are spaces of all $f \in L^p(T^2)$ whose Fourier coefficients

$$\hat{f}(j, l) = \int_{T^2} f(z, w) \bar{z}^j \bar{w}^l dm(z, w)$$

are 0 as soon as at least one component of (j, l) is negative where $1 \leq p \leq \infty$. Let U^2 be the unit bidisc that is the Cartesian product of 2 open unit discs in $\mathbb{C}$. Any function f in H^p has an analytic extension to U^2 which is also denoted by f .

A closed subspace M of H^2 is said to be invariant if

$$zM \subset M \text{ and } wM \subset M.$$

Put $M_1 = [\bigcup_{n=0}^{\infty} \bar{z}^n M]$ and $M_2 = [\bigcup_{n=0}^{\infty} \bar{w}^n M]$ where $[\bigcup_{n=0}^{\infty} \bar{z}^n M]$ is the closed linear span of $\bigcup_{n=0}^{\infty} \bar{z}^n M$ in L^2 . Set

$$M^\times = M_1 \cap M_2$$

then M is also an invariant subspace of H^2 . Put

$$H_1 = [\bigcup_{n=0}^{\infty} \bar{z}^n H^2] \text{ and } H_2 = [\bigcup_{n=0}^{\infty} \bar{w}^n H^2]$$

then $H^2 = H_1 \cap H_2$ and hence $(H^2)^\times = H^2$. Therefore it is desirable to

know invariant subspaces M which have the following property : $M^\times = H^2$ or

$M^\times = M$. An invariant subspace M of H^2 has full range if $M_1 = H_1$ and

$M_2 = H_2$. Such an invariant subspace has been studied by Agrawal, Clark and Douglas [2]. It is clear that M has full range if and only if $M^\times = H^2$. An invariant subspace M with $M^\times = M$ has not been studied. In this paper we study invariant subspaces with $M^\times = M$. For an invariant subspace M of H^2 set

$$\mathcal{M}(M) = \{\phi \in L^\infty : \phi M \subseteq H^2\}.$$

It is essentially known [2] that if $M^\times = M$ then $\mathcal{M}(M) = H^\infty$. In this paper we give a necessary and sufficient condition for $\mathcal{M}(M) = H^\infty$.

Let K_0^2 denote the orthogonal complement of $\bar{H}^2 = \{\bar{f} : f \in H^2\}$ in L^2 .

The invariant subspace qH^2 for an inner function q is called a Beurling subspace.

The author would like to thank Professor K. Takahashi for his helpful discussions.

§ 2. Intersection of Beurling subspaces

A Beurling subspace M satisfies $M^\times = M$ trivially. In this section we show that the intersections of Beurling subspaces have this property.

Proposition 1. If M is a nonzero invariant subspace of H^2 then there

exist two unimodular functions Q_1 in H_1 and Q_2 in H_2 such that $M_1 = Q_1 H_1$ and $M_2 = Q_2 H_2$, and hence

$$M^\times = Q_1 H_1 \cap Q_2 H_2.$$

Proof. By [7, p164 ~ p165], $[\bigcup_{n=0}^{\infty} \bar{z}^n M] = \chi_E Q_1 H_1 +$

$(1 - \chi_E) L^2$ where χ_E is a characteristic function of some measurable set E in T^2 , $\chi_E \in H_1$ and $|Q_1| = 1$ a.e.. Since $M \subset H^2$ and H_1 has no reducing invariant subspaces under the multiplication of w , this implies that $M_1 = Q_1 H_1$ and $Q_1 \in H_1$ because $M_1 \subset H_1$. The same argument shows that $M_2 = Q_2 H_2$ and $Q_2 \in H_2$.

Theorem 2. Let ℓ be a finite positive integer. If $M = \bigcap_{j=1}^{\ell} q_j H^2$ and $\{q_j\}_{j=1}^{\ell}$ are inner functions then $M^\times = M$. Moreover

$$M^\times = Q_1 H_1 \cap Q_2 H_2$$

where $\{Q_i\}_{i=1}^2$ are unimodular functions in $\{H_i\}_{i=1}^2$ and

$$M_i = Q_i H_i = \bigcap_{j=1}^{\ell} q_j H_i \quad (i = 1, 2).$$

Proof. Let $N = \bar{M}^\perp$, that is, N is the orthogonal complement of $\bar{M} = \{\bar{f} ; f \in M\}$. Then $N = [\sum_{j=1}^{\ell} q_j K_0^2]$ because $(\bar{H}^2)^\perp = K_0^2$. Put $N_1 = [\sum_{j=1}^{\ell} q_j w H_1]$ and $N_2 = [\sum_{j=1}^{\ell} q_j z H_2]$ then $\{N_i\}_{i=1}^2$ are invariant subspaces and $N = [N_1 + N_2]$ because $K_0^2 = w H_1 + z H_2$. It is clear that $z N_1 = N_1$ and $w N_2 = N_2$. If $w N_1 = N_1$ then $N_1 \supset \bar{q}_j \bar{w}^n H_1$ for any positive n and so $N_1 = L^2$. This contradicts that $M \supset (\prod_{j=1}^{\ell} q_j) H^2$. Thus $w N_1 \neq N_1$ and

similarly $zN_2 \neq N_2$. By [7] there exist unimodular functions Q_1 and Q_2 such that $N_1 = Q_1 w H_1$ and $N_2 = Q_2 z H_2$. Since $N = [N_1 + N_2]$,

$$M = Q_1 H_1 \cap Q_2 H_2.$$

Since $Q_i H_i$ is the orthogonal complement of N_i and $N_i = (\bar{M}_i)^\perp$ for $i = 1, 2$,

$$M_i = Q_i H_i = \bigcap_{j=1}^n q_j H_i \quad (i = 1, 2).$$

In the case of one variable an intersection of two Beurling subspaces is also a Beurling subspace. This is not true in the case of two variables by [11, Theorem 2 and its proof]. In fact X_1 and X_2 in [11, Theorem 2] are intersections of two Beurling subspaces and $X_j \subseteq H^2$ for $j = 1, 2$.

If $X_1 = q_1 H^2$ and $X_2 = q_2 H^2$ for some inner functions q_1, q_2 then $\bar{q}_1 X_1 = \bar{q}_2 X_2 = H^2$. This contradicts [11, Theorem 2]. Hence our Theorem 2 is not

trivial. If an invariant subspace M is determined by vanishing conditions at finitely many points of U then $M^\times = M$ because M is a finite co-

dimensional subspace of H^2 . We are interested in an invariant subspace determined by vanishing conditions at infinitely many points of U . Let s

be an analytic function on U such that $s(U) \subset U$. Put

$$M_s = \{f \in H^2 : f(z, s(z)) = 0 \text{ for all } z \in U\}$$

then M_s is an invariant subspace of H^2 .

Proposition 3. Let s be an analytic function on U and $s(U) \subset U$. Then

$$M_s^* = QH_1 \cap H_2$$

where $Q = \frac{w-s}{1-\bar{s}w}$. (1) If $|s| = 1$ a.e. on T then $M_s^* = H^2$. (2) If

$|s| = r < 1$ a.e. on T for some constant r then

$$M_s^* = M_s = Q' H_1 \cap H_2 = Q' H^2 \cap H^2$$

where $Q' = \frac{w-s}{sr^{-1}-rw}$.

Proof. If $f \in M_s$ then for a.e. $z \in T$ $\frac{1-s\bar{w}}{w-s} f(z, w)$ is

w -analytic and hence f belongs to QH_1 . Therefore $M_{s,1} \subset QH_1$. Since

$(1-rsw)^{-1} \in H_1$ for any constant r with $0 < r < 1$ and $w-s \in M_s$,

$(w-s)(1-rsw)^{-1}$ belongs to $M_{s,1}$. Since

$$\left| \frac{w-s}{1-r\bar{s}w} - \frac{w-s}{1-\bar{s}w} \right| \leq \left| \frac{1-r}{1-r\bar{s}w} \right| \rightarrow 0 \quad (\text{as } r \rightarrow 1)$$

because $m\{(z, w) \in T^2 : \bar{s}(z)w = 1\} = 0$, Q belongs to $M_{s,1}$ and hence $M_{s,1}$

$= QH_1$. Since $(w-rs)^{-1} \in H_2$ for any r with $0 < r < 1$ and $w-s \in$

M_s , $(w-s)(w-rs)^{-1} \in M_{s,2}$. As $r \rightarrow 1$ the constant 1 belongs to $M_{s,2}$

and hence $M_{s,2} = H_2$. This implies $M_s^* = QH_1 \cap H_2$. (1) is clear because

$sH_1 = H_1$. (2) Since $\frac{w-s}{1-\bar{s}w} = \frac{sr^{-1}(w-s)}{sr^{-1}-rw}$, $M_s = Q' H_1 \cap H_2$. Put

$N = Q' H^2 \cap H^2$. If $Q' f = g$ for some f and g in H^2 then

$$(w-s)f(z, w) = (sr^{-1}-rw)g(z, w) \quad (z, w) \in U^2.$$

Thus g belongs to M_s and hence $M_s \supset N$. It is clear that $N_1 \subset Q' H_1$ and $N_2 \subset H_2$. The Q' belongs to N_1 because $(sr^{-1} - rw)^{-1} \in H_1$ and $w - s \in N$. The constant 1 belongs to N_2 because $(w - s)^{-1} \in H_2$ and $w - s \in N$. Thus $N = Q' H_1 \cap H_2$. Since $N^\times = N$ by the proof of Theorem 2, $M_s^\times = M_s = Q' H_1 \cap H_2 = Q' H^2 \cap H^2$.

§ 3. Multipliers of an invariant subspace

An invariant subspace M of H^2 is said to be podal if every invariant subspace N of H^2 is unitarily equivalent to M which is a subspace of M (cf. [3]). Agrawal, Clark and Douglas [2] showed that if an invariant subspace of H^2 has full range then it is podal. The following is a generalization of that. For if N is full range then $N^\times = H^2$.

Proposition 4. Let M and N be invariant subspaces of H^2 with $M \subset N^\times$.

If M is unitarily equivalent to N , then $M \subset N$.

Proof. By [2, Lemma 1] $M = qN$ for some unimodular function q . Then $M^\times = qN^\times$ and $M^\times \subset N^\times$. By [2, Proposition 3] the q is an inner function.

Proposition 5. Let M and N be invariant subspaces of H^2 with $M^\times = N^\times$.

Then M is unitarily equivalent to N if and only if $M = N$.

Proof. If M is unitarily equivalent to N then $M = qN$ for some unimodular function q . Then $M^\times = qN^\times$ and hence $M^\times = qM^\times$. Thus q is an inner function. By the same argument $N^\times = \bar{q}N^\times$ and hence q is also an inner function. Thus q is constant and $M = N$.

If M and N are invariant subspaces of finite codimension in H^2 , then we can show $M^\times = N^\times$ and hence by Proposition 5 $M = N$. This is Corollary 3 in [2] (see [5, Corollary 6]). Douglas and Yan [3] gave the following question : Can one characterize podal invariant subspaces ?

Proposition 6. Let M be a nonzero invariant subspace of H^2 . M is podal if and only if any unimodular functions in $\mathcal{M}(M)$ belong to H^∞ .

Proof. If ϕ is a unimodular function in $\mathcal{M}(M)$ then $\phi M \subset H^2$ and ϕM is unitarily equivalent to M . If M is podal then $\phi M \subset M$ and hence $\phi \in H^\infty$ by [2, Proposition 3]. Suppose any unimodular functions in $\mathcal{M}(M)$ belong to H^∞ . If M is unitarily equivalent to N which is an invariant subspace in H^2 then $N = \phi M$ for some unimodular function ϕ . Since $\phi \in \mathcal{M}(M)$, by the hypothesis ϕ belongs to H^∞ and so M is podal.

By the proposition above, if $\mathcal{M}(M) = H^\infty$ then M is podal. We will characterize an invariant subspace M with $\mathcal{M}(M) = H^\infty$. In general $\mathcal{M}(M)$ is an invariant subspace of L^∞ which contains H^∞ . The structure of $\mathcal{M}(M)$ is

simpler than that of M .

Theorem 7. If M is a nonzero invariant subspace of H^2 then

$$\mathcal{M}(M) = \bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty$$

where $M^\times = Q_1 H_1 \cap Q_2 H_2$ and Q_i is a unimodular function in H_1 for $i = 1, 2$. Hence $\mathcal{M}(M) = H^\infty$ if and only if

$$\bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty = H^\infty.$$

Proof. If $\phi \in \mathcal{M}(M)$ then $\phi M \subset H^2$. Hence for $i = 1, 2$

$$\phi M_i \subset H_i \quad \text{and} \quad M_i = Q_i H_i$$

where Q_i is a unimodular function in H_1 . Thus $\phi \in \bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty$ and hence $\mathcal{M}(M) \subset \bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty$. Conversely let ϕ be in $\bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty$ then

$$\phi = \bar{Q}_1 \phi_1 = \bar{Q}_2 \phi_2$$

where ϕ_i is in H_i for $i = 1, 2$. If $f \in M$ then $f \in M^\times$ and hence

$$f = Q_1 f_1 = Q_2 f_2$$

where f_i is in H_i for $i = 1, 2$. Therefore $\phi f = \phi_1 f_1 = \phi_2 f_2$. This implies that $\phi f \in H_1 \cap H_2 = H^2$ and hence $\phi \in \mathcal{M}(M)$. Thus $\bar{Q}_1 H_1 \cap \bar{Q}_2 H_2 \cap L^\infty \subset \mathcal{M}(M)$. This implies the theorem.

For $\phi \in L^\infty$, the Hankel operator determined by ϕ is

$$H_\phi = (I - P)M_\phi|_{H^2}$$

where P is the orthogonal projection from L^2 to H^2 .

Proposition 8. The kernel of a Hankel operator is an invariant subspace M with $M^\times = M$.

Proof. Suppose H_ϕ is the Hankel operator defined by $\phi \in L^\infty$ and let M be its kernel. It is clear that M is an invariant subspace. Since M is the kernel of H_ϕ , $\phi M \subset H^2$. By Theorem 7, $\mathcal{M}(M) = \mathcal{M}(M^\times)$ and $\phi M^\times \subset H^2$. This implies $M = M^\times$ because $M = \{\phi f \in H^2 : f \in H^2\}$.

References

1. P.R.Ahern and D.N.Clark, Invariant subspaces and analytic continuation in several variables, J. Math. Mech. 19(1970), 963-969.
2. O.P.Agrawal, D.N.Clark, and R.G.Douglas, Invariant subspaces in the polydisk, Pacific J. Math. 121(1986) 1-11.
3. R.G.Douglas and K.Yan, On the rigidity of Hardy submodules, Integral Eq. Oper. Th. 13(1990).
4. K.Hoffman, Banach Spaces of Analytic Functions, Prentice-Hall, 1962.
5. K.Izuchi, Unitary equivalence of invariant subspaces in the polydisk, Pacific J. Math. 130(1987), 351-358.

6. T.Nakazi, Nonmaximal weak-*Dirichlet algebra, Hokkaido Math. J.
5(1976), 88-96.
7. T.Nakazi, Invariant subspaces of weak-*Dirichlet algebras, Pacific J.
Math. 69(1977), 151-167.
8. T.Nakazi, Certain invariant subspaces of H^2 and L^2 on a bidisc, Can.
J. Math. XL (1988), 1272-1280.
9. T.Nakazi, Invariant subspaces in the bidisc and commutators, to appear
in J. Aust. Math. Soc.
10. T.Nakazi, Homogeneous polynomials and invariant subspaces in the
polydiscs. Archiv der Math. 58(1992), 56-63.
11. W.Rudin, Invariant subspaces of H^2 on a torus, J. Funct. Anal.
61(1985), 378-384.