HOKKAIDO UNIVERSITY

Title	Hochster's formula on Betti numbers and Buchsbaum complexes
Author(s)	Hibi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 209, 1-7
Issue Date	1993-07
DOI	https://doi.org/10.14943/83353
Doc URL	https://hdl.handle.net/2115/68955
Type	departmental bulletin paper
File Information	re209.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Hochster's formula on Betti numbers and Buchsbaum complexes

Takayuki Hibi

Series #209. July 1993

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

182: M. Yamada, Distance formulas of asymptotic Toeplitz and Hankel operators, 13 pages. 1993

183: G. Ishikawa, T. Ozawa, The genus of a connected compact real algebraic surface in the affine three space, 11 pages. 1993.

184: T. Hibi, Canonical modules and Cohen-Macaulay types of partially ordered sets, 6 pages. 1993.

185: Y. Giga, K. Yama-uchi, On a lower bound for the extinction time of surfaces moved by mean curvature, 16 pages. 1993.

186: Y. Kurokawa, On functional moduli for first order ordinary differential equations, 9 pages. 1993.

187: O. Ogurisu, Unitary equivalence between a spin 1/2 charged particle in a two-dimensional magnetic field and a spin 1/2 neutral particle with an anomalous magnetic moment in a two-dimensional electric field, 4 pages. 1993.

188: A. Jensen, T. Ozawa, Existence and non-existence results for wave operators for perturbations of the laplacian, 30 pages. 1993.

189: T. Nakazi, Multipliers of invariant subspaces in the bidisc, 12 pages. 1993.

190: S. Izumiya, G.T. Kossioris, Semi-local classification of geometric singularities for Hamilton-Jacobi equations, 24 pages. 1993.

191: N. Hayashi, T. Ozawa, Finite energy solutions of nonlinear Schrödinger equations of derivative type, 21 pages. 1993.

192: J. Seade, T. Suwa, A residue formula for the index of a holomorphic flow, 22 pages. 1993.

193: H. Kubo, Blow-up of solutions to semilinear wave equations with initial data of slow decay in low space dimensions, 8 pages. 1993.

194: F. Hiroshima, Scaling limit of a model of quantum electrodynamics, 52 pages. 1993.

195: T. Ozawa, Y. Tsutsumi, Global existence and asymptotic behavior of solutions for the Zakharov equations in three space dimensions, 34 pages. 1993.

196: H. Kubo, Asymptotic behaviors of solutions to semilinear wave equations with initial data of slow decay, 25 pages. 1993.

197: Y. Giga, Motion of a graph by convexified energy, 32 pages. 1993.

198: T. Ozawa, Local decay estimates for Schrödinger operators with long range potentials, 17 pages. 1993.

199: A. Arai, N. Tominaga, Quantization of angle-variables, 31 pages. 1993.

200: S. Izumiya, Y. Kurokawa, Holonomic systems of Clairaut type, 17 pages. 1993.

201: K.-S. Saito, Y. Watatani, Subdiagonal algebras for subfactors, 7 pages. 1993.

202: K. Iwata, On Markov properties of Gaussian generalized random fields, 7 pages. 1993.

203: A. Arai, Characterization of anticommutativity of self-adjoint operators in connection with Clifford algebra and applications, 13 pages. 1993.

204: J. Wierzbicki, An estimation of the depth from an intermediate subfactor, 7 pages. 1993.

205: N. Honda, Vanishing theorem for the tempered distributions, 11 pages. 1993.

206: T. Hibi, Betti number sequences of simplicial complexes, Cohen-Macaulay types and Möbius functions of partially ordered sets, and related topics, 25 pages. 1993.

207: A. Inoue, Regularly varying correlations, 23 pages. 1993.

208: S. Izumiya, B. Li, Overdetermined systems of first order partial differential equations with singular solution, 9 pages. 1993.

Hochster's formula on Betti numbers and Buchsbaum complexes

Takayuki Hibi

Department of Mathematics
Faculty of Science
Hokkaido University
Kita-ku, Sapporo 060, Japan

Abstract. The Betti numbers $\dim_k \underline{\mathrm{Tor}}_i^A(k[\Delta],k)$ with $i > v - d$ of the Stanley-Reisner ring $k[\Delta] = A/I_\Delta$ of a Buchsbaum complex Δ of dimension $d-1$ over a field k with v vertices are studied.

§1. Betti numbers of Stanley-Reisner rings

First, we recall some fundamental material for algebra, topology and combinatorics on simplicial complexes.

(1.1) Fix a finite set $V = \{x_1, x_2, \ldots, x_v\}$, called the vertex set, and let Δ be a simplicial complex on V. Thus Δ is a family of subsets of V such that (i) $\{x_i\} \in \Delta$ for each $1 \leq i \leq v$ and (ii) $\sigma \in \Delta$, $\tau \in \sigma$ imply $\tau \in \Delta$. Each element σ of Δ is called a *face* of Δ. Let $d := \max\{\#(\sigma) ; \sigma \in \Delta\}$. Here $\#(\sigma)$ is the cardinality of σ as a finite set. Then the dimension of Δ is defined by $\dim \Delta = d - 1$. A simplicial complex Δ is called *pure* if every maximal face has the same cardinality.

When W is a subset of V, we write Δ_W for the simplicial complex $\{\sigma \in \Delta ; \sigma \subset W\}$ on the vertex set W. On the other hand, given a face σ of Δ, we define the subcomplex $\mathrm{link}_\Delta(\sigma)$

and $\mathrm{star}_\Delta(\sigma)$ of Δ by

$$\mathrm{link}_\Delta(\sigma) := \{\tau \in \Delta ; \sigma \cap \tau = \emptyset \text{ and } \sigma \cup \tau \in \Delta\}$$

$$\mathrm{star}_\Delta(\sigma) := \{\tau \in \Delta ; \sigma \cup \tau \in \Delta\}.$$

Thus, in particular, $\mathrm{link}_\Delta(\emptyset) = \Delta$.

(1.2) Let $A = k[x_1, x_2, \ldots, x_v]$ be the polynomial ring over a field k whose indeterminates are the elements of V with the standard grading, i.e., each $\deg x_i = 1$. Define I_Δ to be the ideal of A which is generated by those square-free monomials $x_{i1}x_{i2}\ldots x_{ir}$, $1 \le i_1 < i_2 < \ldots < i_r \le v$, such that $\{x_{i1}, x_{i2}, \ldots, x_{ir}\} \notin \Delta$, and set $k[\Delta] := A/I_\Delta$. The algebra $k[\Delta]$ over k is called the ***Stanley-Reisner ring*** of Δ over k ([5], [6]). From now on, we regard $k[\Delta]$ as a graded module over A with the "quotient grading." Then $\dim_A(k[\Delta]) = d$.

Let $\underline{H}_m^i(k[\Delta])$ be the i-th local cohomology module of $k[\Delta]$ over A with respect to the irrelevant maximal ideal $m = (x_1, x_2, \ldots, x_v)$ of A, i.e.,

$$\underline{H}_m^i(k[\Delta]) := \varinjlim_n \underline{\mathrm{Ext}}_A^i(A/m^n, k[\Delta]),$$

and $t := \mathrm{depth}_A(k[\Delta])$. Then (i) $\underline{H}_m^i(k[\Delta]) = 0$ unless $t \le i \le d$ and (ii) $\underline{H}_m^t(k[\Delta]) \neq 0$, $\underline{H}_m^d(k[\Delta]) \neq 0$. Consult, e.g., [8] for basic facts on local cohomology modules $\underline{H}_m^i(k[\Delta])$.

(1.3) We say that a simplicial complex Δ is ***Cohen-Macaulay*** (resp. ***Buchsbaum***) over k if the module $k[\Delta]$ over A is Cohen-Macaulay (resp. Buchsbaum), i.e., $\underline{H}_m^i(k[\Delta]) = 0$ (resp. $\dim_k(\underline{H}_m^i(k[\Delta])) < \infty$) for every $0 \le i < d$. Let $\tilde{H}_i(\Delta;k)$ be the i-th reduced homology group of Δ with coefficients k. Then Δ is Cohen-Macaulay if and only if, for every face σ of Δ (possibly, $\sigma = \emptyset$) and for each $i \neq \dim(\mathrm{link}_\Delta(\sigma))$, we have $\tilde{H}_i(\mathrm{link}_\Delta(\sigma);k) = 0$. Every Cohen-Macaulay complex is pure. Moreover, a simplicial complex Δ is Buchsbaum if and only

if Δ is pure and $\mathrm{link}_\Delta(\sigma)$ is Cohen-Macaulay for every non-empty face σ of Δ. We refer the reader to, e.g., [3], [7] and [8] for further information on Cohen-Macaulay and Buchsbaum complexes. See also [1].

On the other hand, in [2], we study the integers $\alpha^*(\Delta) = \alpha^*(\Delta;k)$ and $\gamma^*(\Delta) = \gamma^*(\Delta;k)$ defined as follows:

$$\alpha^*(\Delta) := \max\{\, j \,;\, \underline{H}_m^i(k[\Delta]) = 0 \text{ for each } 0 \le i < j \ (\le d)\,\}$$

$$\gamma^*(\Delta) := \max\{\, j \,;\, \dim_k(\underline{H}_m^i(k[\Delta])) < \infty \text{ for each } 0 \le i < j \ (\le d)\,\}.$$

Thus $1 \le \alpha^*(\Delta) \le \gamma^*(\Delta) \le d$ and $\alpha^*(\Delta) = \mathrm{depth}_A(k[\Delta])$. Moreover, the simplicial complex Δ is Cohen-Macaulay (resp. Buchsbaum) if and only if $\alpha^*(\Delta) = d$ (resp. $\gamma^*(\Delta) = d$). Note that the integer $\alpha^*(\Delta)$ (resp. $\gamma^*(\Delta)$) is equal to the topological invariant $\alpha(\Delta) + 1$ (resp. $\gamma(\Delta) + 1$) in Munkres [4].

(1.4) The i-th *Betti number* $\beta_i^A(k[\Delta])$ of the module $k[\Delta]$ over A is defined to be

$$\beta_i^A(k[\Delta]) := \dim_k \underline{\mathrm{Tor}}_i^A(k[\Delta],k).$$

Let $\rho := v - \alpha^*(\Delta)$. Then $\beta_i^A(k[\Delta]) = 0$ unless $0 \le i \le \rho$. The following formula on Betti numbers $\beta_i^A(k[\Delta])$ is given by Hochster [3, Theorem (5.1)]:

$$\beta_i^A(k[\Delta]) = \sum_{W \subset V} \dim_k(\tilde{H}_{v-\#(W)-i-1}(\Delta_{V-W};k)). \qquad (1)$$

We are now in the position to state our main result in this paper.

(1.5) THEOREM. Let Δ be a simplicial complex on the vertex set $V = \{x_1, x_2, \ldots, x_v\}$ of dimension $d-1$, $A = k[x_1, \ldots, x_v]$ the polynomial ring over a field k, and $k[\Delta] = A/I_\Delta$. Suppose

that $(1 \leq)\ \alpha^*(\Delta) < \gamma^*(\Delta)\ (\leq d)$. Then, for each integer i with $v - \gamma^*(\Delta) < i \leq v - \alpha^*(\Delta)$, we have

$$\beta_i^A(k[\Delta]) = \sum_{j=0}^{v-\alpha^*(\Delta)-i} \binom{v}{j} \dim_k(\tilde{H}_{v-i-1-j}(\Delta;k)).$$

(1.6) COROLLARY. Let Δ be a simplicial complex on the vertex set $V = \{x_1, \ldots, x_v\}$ of dimension $d-1$, $A = k[x_1, \ldots, x_v]$ the polynomial ring over a field k, and $k[\Delta] = A/I_\Delta$. Suppose that Δ is Buchsbaum, but not Cohen-Macaulay. Then, for each integer i with $v - d < i \leq v - \mathrm{depth}_A(k[\Delta])$, we have

$$\beta_i^A(k[\Delta]) = \sum_{j=0}^{v-\mathrm{depth}_A(k[\Delta])-i} \binom{v}{j} \dim_k(\tilde{H}_{v-i-1-j}(\Delta;k)).$$

§2. Proof of Theorem (1.5)

We inherit the notation in the preceding section.

(2.1) LEMMA. $\alpha^*(\mathrm{star}_\Delta(\sigma)) \geq \gamma^*(\Delta)$ for every non-empty face σ of Δ.

Proof. See [2, Lemma (2.7)] for an algebraic proof based on [7, Theorem 4.1, p.70]. Also, consult [4, Lemma (6.1)] for a topological proof. Q.E.D.

(2.2) LEMMA. $\underline{H}_m^j(k[\Delta]) \cong \underline{H}_m^j(k[\Delta_{V-\{x\}}])$ for every $x \in V$ and for each $j < \gamma^*(\Delta) - 1$.

Proof. We have an exact sequence

$$0 \to k[\mathrm{star}_\Delta(\{x\})] \to k[\Delta] \to k[\Delta_{V-\{x\}}] \to 0$$

as graded modules over A. See, e.g., [2, Theorem (1.7)]. Hence, there exists a long exact sequence

$$\begin{aligned} 0 \to\ & \underline{H}_m^0(k[\mathrm{star}_\Delta(\{x\})]) \to \underline{H}_m^0(k[\Delta]) \to \underline{H}_m^0(k[\Delta_{V-\{x\}}]) \\ \to\ & \underline{H}_m^1(k[\mathrm{star}_\Delta(\{x\})]) \to \underline{H}_m^1(k[\Delta]) \to \underline{H}_m^1(k[\Delta_{V-\{x\}}]) \\ \to\ & \cdots \\ \to\ & \underline{H}_m^j(k[\mathrm{star}_\Delta(\{x\})]) \to \underline{H}_m^j(k[\Delta]) \to \underline{H}_m^j(k[\Delta_{V-\{x\}}]) \\ \to\ & \cdots \end{aligned}$$

of local cohomology modules. Since $\underline{H}_m^j(k[\mathrm{star}_\Delta(\{x\})]) = (0)$ for every $i < \alpha^*(\mathrm{star}_\Delta(\{x\}))$, Lemma (2.1) guarantees that $\underline{H}_m^j(k[\mathrm{star}_\Delta(\{x\})]) = (0)$ for every $i < \gamma^*(\Delta)$. Thus $\underline{H}_m^j(k[\Delta]) \cong \underline{H}_m^j(k[\Delta_{V-\{x\}}])$ for each $j < \gamma^*(\Delta) - 1$ as reqired. **Q.E.D.**

(2.3) LEMMA. $\gamma^*(\Delta_{V-W}) \geq \gamma^*(\Delta) - \#(W)$ for every $W \subset V$.

Proof. By Lemma (2.2), $\dim_k(\underline{H}_m^i(k[\Delta_{V-\{x\}}])) < \infty$ for each $0 \leq i < \gamma^*(\Delta) - 1$. Hence $\gamma^*(\Delta_{V-\{x\}}) \geq \gamma^*(\Delta) - 1$. Thus $\gamma^*(\Delta_{V-W}) \geq \gamma^*(\Delta_{V-(W-\{x\})}) - 1$ for every $x \in W$. Hence $\gamma^*(\Delta_{V-W}) \geq \gamma^*(\Delta) - \#(W-\{x\}) - 1 = \gamma^*(\Delta) - \#(W)$ as desired. **Q.E.D.**

(2.4) PROPOSITION. $\underline{H}_m^j(k[\Delta]) \cong \underline{H}_m^j(k[\Delta_{V-W}])$ for every $W \subset V$ and for each $j < \gamma^*(\Delta) - \#(W)$.

Proof. Let $W = \{x_{i1}, x_{i2}, \ldots, x_{is}\}$ and, for each $0 \leq \ell \leq s$, $W(\ell) = \{x_{i1}, x_{i2}, \ldots, x_{i\ell}\}$. Lemma (2.2) enables us to see $\underline{H}_m^j(k[\Delta_{V-W(\ell)}]) \cong \underline{H}_m^j(k[\Delta_{V-W(\ell+1)}])$ for each $0 \leq \ell < s$ and for every $j < \gamma^*(\Delta_{V-W(\ell)}) - 1$. On the other hand, by Lemma (2.3), $\gamma^*(\Delta_{V-W(\ell)}) \geq \gamma^*(\Delta) - \#(W(\ell))$ $(> \gamma^*(\Delta) - \#(W))$. Thus $\underline{H}_m^j(k[\Delta]) \cong \underline{H}_m^j(k[\Delta_{V-W}])$ for each $j < \gamma^*(\Delta) - \#(W)$. **Q.E.D.**

(2.5) COROLLARY. For every subset W of the vertex set V and for each $j < \gamma^*(\Delta) - \#(W)$, $\dim_k(\tilde{H}_{j-1}(\Delta;k))$ is equal to $\dim_k(\tilde{H}_{j-1}(\Delta_{V-W};k))$.

Proof. It follows from, e.g., [7, Theorem 4.1, p.70] that $\dim_k(\underline{H}_m^i(k[\Delta])) = \dim_k \tilde{H}_{i-1}(\Delta;k)$ if $\dim_k(\underline{H}_m^i(k[\Delta])) < \infty$.

Q.E.D.

We are now in the position to give our proof of Theorem (1.5). Suppose that $\alpha^*(\Delta) < \gamma^*(\Delta)$. Let i be an integer with $v - \gamma^*(\Delta) < i \leq v - \alpha^*(\Delta)$ and W a subset of V. By Corollary (2.5), we have the equality

$$\dim_k(\tilde{H}_{v-\#(W)-i-1}(\Delta;k)) = \dim_k(\tilde{H}_{v-\#(W)-i-1}(\Delta_{V-W};k))$$

since $v-\#(W)-i < \gamma^*(\Delta) - \#(W)$. Hence, by virtue of Eq. (1),

$$\begin{aligned}
\beta_i{}^A(k[\Delta]) &= \sum_{W \subset V} \dim_k(\tilde{H}_{v-\#(W)-i-1}(\Delta_{V-W};k)) \\
&= \sum_{W \subset V} \dim_k(\tilde{H}_{v-\#(W)-i-1}(\Delta;k)) \\
&= \sum_{j=0}^{v-\alpha^*(\Delta)-i} \binom{v}{j} \dim_k(\tilde{H}_{v-i-1-j}(\Delta;k))
\end{aligned}$$

as required. **Q.E.D.**

REFERENCES

[1] A. Björner and T. Hibi, Betti numbers of Buchsbaum complexes, Math. Scand. 67 (1990), 193-196.

[2] T. Hibi, Quotient algebras of Stanley-Reisner rings and local cohomology, J. Algebra 140 (1991), 336-343.

[3] M. Hochster, Cohen-Macaulay rings, combinatorics, and simplicial complexes, in "Ring Theory II" (B. R. McDonald and R. Morris, eds.), Lect. Notes in Pure and Appl. Math., No. 26, Dekker, New York, 1977, pp. 171-223.

[4] J. Munkres, Topological results in combinatorics, Michigan Math. J. 31 (1984), 113-128.

[5] G. Reisner, Cohen-Macaulay quotients of polynomial rings, Advances in Math. 21 (1976), 30-49.

[6] R. Stanley, The upper bound conjecture and Cohen-Macaulay rings, Stud. Appl. Math. 54 (1975), 135-142.

[7] _____, "Combinatorics and Commutative Algebra," Birkhäuser, Boston/Basel/Stuttgart, 1983.

[8] J. Stückrad and W. Vogel, "Buhsbaum Rings and Applications," Springer-Verlag, Berlin/Heidelberg/New York, 1986.