HOKKAIDO UNIVERSITY

| | |
|---|---|
| Title | A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature |
| Author(s) | Chen, Y. -G.; Giga, Y.; Hitaka, T. et al. |
| Citation | Hokkaido University Preprint Series in Mathematics, 258, 2-18 |
| Issue Date | 1994-08-01 |
| DOI | https://doi.org/10.14943/83405 |
| Doc URL | https://hdl.handle.net/2115/69009 |
| Type | departmental bulletin paper |
| File Information | re258.pdf |

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

# A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature

Y-G. Chen, Y. Giga

T. Hitaka and M. Honma

Series #258. August 1994

## HOKKAIDO UNIVERSITY

## PREPRINT SERIES IN MATHEMATICS

# 233: C. Dohmen, Y. Giga, N. Mizoguchi, Existence of selfsimilar shrinking curves for anisotropic curvature flow equations, 13 pages. 1994.

# 234: T. Nakazi, M. Yamada, Invertible Toeplitz operators and uniform algebras, 14 pages. 1994.

# 235: C. Dohmen, A Pohozaev-type inequality for a quasilinear Haraux-Weissler equation, 6 pages. 1994.

# 236: T. Mikami, Large deviations for the first exit time on small random perturbations of dynamical systems, 32 pages. 1994.

# 237 T. Ozawa, Characterization of Trudinger's inequality, 7 pages. 1994.

# 238 T. Hibi, Buchsbaum complexes with linear resolutions, 12 pages. 1994.

# 239 Y. Giga, N. Mizoguchi, On time periodic solutions of the Dirichlet problem for degenerate parabolic equations of nondivergence type, 23 pages. 1994.

# 240 C. Dohmen, Existence of Fast Decaying Solutions to a Haraux-Weissler Equation With a Prescribed Number of Zeroes, 12 pages. 1994.

# 241 K. Sugano, Note on H-separable Frobenius extensions, 8 pages. 1994.

# 242 J. Zhai, Some Estimates For The Blowing up Solutions of Semilinear Heat Equations, 11 pages. 1994.

# 243 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and Smoothing Effect of the Schrödinger Evolution Group, 10 pages. 1994.

# 244 D. Lehmann, T. Suwa, Residues of holomorphic vector fields relative to singular invariant subvarieties, 26 pages. 1994.

# 245 H. Kubo, Slowly decaying solutions for semilinear wave equations in odd space dimensions, 30 pages. 1994.

# 246 T. Nakazi, M. Yamada, $(A_2)$-Conditions and Carleson Inequalities, 27 pages. 1994.

# 247 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and smoothing Effect of Solutions to the Benjamin-ono Equation, 17 pages. 1994.

# 248 H. Kikuchi, Sheaf cohomology theory for measurable spaces, 12 pages. 1994.

# 249 A. Inoue, Tauberian theorems for Fourier cosine transforms, 9 pages. 1994.

# 250 S. Izumiya, G. T. Kossioris, Singularities for viscosity solutions of Hamilton-Jacobi equations, 23 pages. 1994.

# 251 H. Kubo, K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values $p$ in odd space dimensions, 51 pages. 1994.

# 252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.

# 253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.

# 254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.

# 255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.

# 256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.

# 257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.

# A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature

YUN-GANG CHEN [1], YOSHIKAZU GIGA[2]
TOSHIO HITAKA[2] AND MITSURU HONMA[2]

Abstract. A difference scheme is introduced for computing the motion of level surfaces moved by the mean curvature. This scheme is proved to be stable in the maximum norm so that the computation can be completed without overflow.

## §1. Introduction.

In the research fields of applied sciences like physics, engineering and biology, it is important to track the evolution (motion) of a surface, such as the interface between two kind of materials or two different phases of a certain kind of material. The problem how to track and compute the motion of a surface with a curvature-dependent speed is usually a key point in the studies. Here, we introduce a class of difference schemes for computing evolving surfaces. Our scheme is based on a level set method. Such a method is developed numerically by Osher and Sethian [OS], [S] and analytically by Chen, Giga and Goto [CGG1] and Evans and Spruck [ES]. See also [CGG2,3]. There are many works now available on this method but we do not try to mention all of them for lack of spaces. If the evolution depends on curvature, the scheme need not be monotone so the convergence to the analytic solution (viscosity solution) is not at all clear (cf. [CL]). In fact as explained later, some of the numerical solutions of the scheme in [OS] may not converge to analytic solutions in uniform topology.

In this paper, we introduce a little bit different scheme reflecting divergence structure and prove its stability in maximum norm for mean curvature flow problems.

We consider the Cauchy problem of the mean curvature flow equation

$$\text{(E)} \qquad u_t = |\nabla u| \operatorname{div}\left(\frac{\nabla u}{|\nabla u|}\right), \quad (t,x) \in Q = (0,\infty) \times \mathbb{R}^N$$

$$\text{(IV)} \qquad u(0,x) = u_0, \quad x \in \mathbb{R}^N.$$

Let $u = u(t,x)$ be a continuous viscosity solution of (E)–(IV) which takes a negative constant for large $|x|$, which holds true if the initial value $u_0$ is assumed to have such

[1] Hokkaido Tokai University, Sapporo 005, Japan.
[2] Faculty of Science, Hokkaido University, Sapporo 060, Japan.

property. If for each $t \in (0,\infty)$ there is a bounded open set $D(t)$ in $\mathbb{R}^N$ such that $u(t,x) > 0$ for $x \in D(t)$ and $u(t,x) < 0$ for $x \notin \overline{D(t)}$, then the 0-level set $\Gamma_t = \{x;\ u(t,x) = 0\}$ of $u(t,x)$ determines a closed surface which moves with a speed $\mathbf{V} = (n-1)\mathbf{H}$ at each point $x \in \Gamma_t$ where $\mathbf{H}(t,x)$ is the mean curvature vector at $x \in \Gamma_t$, provided that $\nabla u \neq 0$ on $\Gamma_t$. The global existence and uniqueness of the viscosity solution to (E)–(IV) have been proved by Chen, Giga & Goto [CGG1] and Evans & Spruck [ES]. And more important thing is that the level set $\Gamma_t$ is uniquely determined by its initial data $\Gamma_0$ which is independent of the choice of its *defining function* $u_0$, provided that $u_0$ is bounded, continuous and $u_0 > 0$ for $x \in D(0)$, $u_0 < 0$ for $x \notin \overline{D(0)}$ and $\Gamma_0 = \{x;\ u_0(x) = 0\}$. Moreover, in [CGG1] these results are proved for a general geometric evolution.

Here, we discuss the difference methods for computing $u(t,x)$, the viscosity solution of (E)–(IV). To overcome the difficulty of taking 0 value in the denominator which will cause errors in computers and stop the computation, we introduce a parameter $\delta > 0$ and consider the difference approximation of a modified equation

$$(\mathrm{E}_\delta) \qquad u_t = |\nabla u| \operatorname{div}\left(\frac{\nabla u}{(|\nabla u|^\sigma + \delta)^{1/\sigma}}\right), \quad (t,x) \in Q = (0,\infty) \times \mathbb{R}^N$$

with the same initial value (IV). Here, $\sigma \geqq 1$ is fixed. It can be shown that the viscosity solution of $(\mathrm{E}_\delta)$ with (IV) tends to that of (E) when $\delta \to 0$. Thus, it is reasonable to deal with the computation of the solution of $(\mathrm{E}_\delta)$ as an approximation of the solution of (E) with the same initial value (IV), for a sufficiently small $\delta > 0$ (say, $\delta = 10^{-50}$).

There are several methods now available to compute evolving surfaces moved by mean curvature. Using parametrization of surface a finite element method was studied by Deckelnick and Dziuk [DD]. See also [D]. Although the convergence of the scheme is proved, this scheme does not track the evolution after it develops singularities. A finite element method for level set equations was studied by Walkington [W] based on a co-volume method. Another way is to compute Allen-Cahn type reaction diffusion equation or its modification. Such a calculation is done by Nochetto and his collaborators [NPV] and it is very good to track the evolution after it experiences singularities. Another method related to the level set method is introduced by Bence, Merriman and Osher [BMO] where heat equation is used to study the motion by mean curvature. Its convergence is proved by Evans [E] and others.

We thank GARC for giving the opportunity for publication of this short note. The main part of this paper was completed in 1991 but because of personal problems of first two authors the completion of the paper had been delayed.

## §2. Difference schemes for the mean curvature flow equation.

Now we introduce our difference scheme for $(\mathrm{E}_\delta)$, and for simplicity we interpret the scheme here for the two dimensional case $N = 2$. Our difference equation for $(\mathrm{E}_\delta)$ is given

by

$$
\begin{aligned}
\frac{u_{jk}^{n+1}-u_{jk}^{n}}{\tau} &= g_{jk}^{n}\sum_{i=1}^{2}\mathrm{D}_i\left(\frac{\mathrm{D}_i\, u_{jk}^{n+\theta}}{((g_{jk}^{n})^{\sigma}+\delta)^{1/\sigma}}\right), \\
& \qquad j,k=0,\pm1,\pm2,\cdots;\ n=0,1,2,\cdots; \\
u_{jk}^{0} &= u_0(x_j,y_k), \quad j,k=0,\pm1,\pm2,\cdots.
\end{aligned}
\tag{1}
$$

Here, several notations have been introduced as below. Denoting by $x$ and $y$ the spatial variables in $\mathbb{R}^2$, we use $x_j$ and $y_k$ for the spatial coordinates of the net points.

NOTATION:
$\tau \equiv \Delta t > 0$: increment of the time variable $t$;
$t_n = n\tau$: $n$th time step;
$h_1,\ h_2$: mesh sizes of $x$ and $y$ directions, respectively;
$(x_j, y_k) = (jh_1, kh_2)$: net point in $\mathbb{R}^2$, $j,k = 0,\pm1,\pm2,\cdots$;
$u_{jk}^n$: value of the difference solution approximate to $u(t_n, x_j, y_k)$;
$\mathrm{D}_1 u_{jk}^n = (u_{j+\frac{1}{2},k}^n - u_{j-\frac{1}{2},k}^n)/h_1$, $\mathrm{D}_2 u_{jk}^n = (u_{j,k+\frac{1}{2}}^n - u_{j,k-\frac{1}{2}}^n)/h_2$: the approximations to $u_x(t_n, x_j, y_k)$ and $u_y(t_n, x_j, y_k)$ by central difference approach, respectively;
$g_{jk}^n \equiv g(\mathrm{D}\, u_{jk}^n)$: discretization of $|\nabla u|$ at $(t_n, x_j, y_k)$, which is chosen positive definite for $\{\mathrm{D}_i^{\pm} u_{jk}^n;\ i = 1,\ldots,N\}$, where $\mathrm{D}^+$ and $\mathrm{D}^-$ denote the standard forward and backward differences, respectively. For instance, we may take

$$
g_{jk}^n = \Big(\frac{1}{4}\sum_{i=1}^{N}\big(|\,\mathrm{D}_i^+ u_{jk}^n| + |\,\mathrm{D}_i^- u_{jk}^n|\big)^2\Big)^{\frac{1}{2}}, \qquad \text{or}
$$

$$
g_{jk}^n = \max_{1\leqq i\leqq N}\big\{|\,\mathrm{D}_i^+ u_{jk}^n|,\ |\,\mathrm{D}_i^- u_{jk}^n|\big\}.
$$

The notation $u^\theta$ here denotes $\theta u^{n+1} + (1-\theta)u^n$ for a fixed parameter $\theta \in [0,1]$, and the difference equation (1) is explicit for $u^{n+1}$ if $\theta = 0$, while implicit if $0 < \theta \leqq 1$.

We can prove a sufficient condition for the $L^\infty$ stability of maximum principle type for the difference scheme (1), as the following

THEOREM 1. *The difference scheme (1) is stable in the sense of* $\|u^n\|_\infty \leqq \|u^0\|_\infty$ *if either* $\theta = 1$ *or* $4\tau(1/h_1^2 + 1/h_2^2) \leqq 1/(1-\theta)$ *when* $0 \leqq \theta < 1$, *where* $\|u^n\|_\infty = \sup_{j,k}|u_{jk}^n|$.

It is sometimes convenient and economic to deal with a surface of rotation in a lower dimension space. Here, to compute the motion of a surface with axisymmetry in $\mathbb{R}^3$, i.e., surfaces of rotation, we rewrite (E) into

$$
u_t = |\widetilde{\nabla} u|\widetilde{\mathrm{div}}\Big(\frac{\widetilde{\nabla} u}{|\widetilde{\nabla} u|}\Big) + \frac{1}{r}u_r, \quad (t,r,z) \in Q = (0,\infty)\times(0,\infty)\times\mathbb{R} \tag{$\mathrm{E}_r$}
$$

where $r = \sqrt{x^2+y^2}$ for $(x,y,z) \in \mathbb{R}^3$, and the differential operators $\widetilde{\nabla}$ and $\widetilde{\mathrm{div}}$ are those with respect to $(r,z) \in [0,\infty) \times \mathbb{R}$.

For this equation, our difference scheme is constructed as

$$\frac{u_{0k}^{n+1}-u_{0k}^{n}}{\tau} = g_{0k}^{n}\sum_{i=1}^{2}\mathrm{D}_i\left(\frac{\mathrm{D}_i\, u_{0k}^{n+\theta}}{[(g_{0k}^{n})^{\sigma}+\delta]^{1/\sigma}}\right) + \frac{2(u_{1k}^{n+\theta}-u_{0k}^{n+\theta})}{h_1^2},$$

$$k = 0, \pm 1, \pm 2, \cdots;\ n = 0, 1, 2, \cdots;$$

(2)
$$\frac{u_{jk}^{n+1}-u_{jk}^{n}}{\tau} = g_{jk}^{n}\sum_{i=1}^{2}\mathrm{D}_i\left(\frac{\mathrm{D}_i\, u_{jk}^{n+\theta}}{[(g_{jk}^{n})^{\sigma}+\delta]^{1/\sigma}}\right) + \frac{1}{r_j}\cdot\frac{u_{j+1,k}^{n+\theta}-u_{jk}^{n+\theta}}{h_1},$$

$$k = 0, \pm 1, \pm 2, \cdots;\ j = 1, 2, \cdots;\ n = 0, 1, 2, \cdots;$$

$$u_{jk}^{0} = u_0(x_j, y_k), \quad k = 0, \pm 1, \pm 2, \cdots;\ j = 0, 1, 2, \cdots.$$

where $u_{jk}^{n}$ is the approximation of $u(t_n, r_j, z_k)$, $(r_j, z_k) = (jh_1, kh_2)$, and $\mathrm{D}_1$ and $\mathrm{D}_2$ denote the difference operators for $\partial_r$ and $\partial_z$. Here, the last term of the first equation is used for the approximation of $\dfrac{u_r}{r}$ on the points of $j = 0$ so that $r_j = 0$. The limit relation $\lim\limits_{r\to 0}\dfrac{u_r}{r} = u_{rr}$ and the symmetricity $u_{-1,k}^{n} = u_{1k}^{n}$ which follows from $u(t,r,z) = u(t,-r,z)$ and thus $u_r(t,0,z) = 0$ are also applied so that

$$\frac{u_{1k}^{n+\theta}-2u_{0k}^{n+\theta}+u_{-1,k}^{n+\theta}}{h_1^2} = \frac{2(u_{1k}^{n+\theta}-u_{0k}^{n+\theta})}{h_1^2}.$$

On the other hand, the last term of the second equation is derived by the upwind difference approach (with forward difference approximation here) to $\dfrac{u_r}{r}$ for $j > 0$.

This type of difference approximation is also used in [C] in the discretization of the Laplacian for the axisymmetric solutions.

In both cases, the schemes (1) and (2), the computation is carried out in a restricted domain, say, a rectangular domain. When the initial value function has a negative constant value $-C$ for large $|x|$, so dose the solution $u(t,x)$ with a same constant value. Then we can use this convenient property in our computation to make the work simpler. We have only to compute the value in a (sufficiently large) finite domain, on the boundary of which the solution has a negative constant value. So, we can deal with the boundary condition either as the Dirichlet condition or as the Neumann condition.

The stability condition for the difference scheme (2) is given by

THEOREM 2. *The difference equation (2) is stable if either*

$$\theta = 1 \quad \text{or} \quad 6\frac{\tau}{h_1^2} + 4\frac{\tau}{h_2^2} \leqq \frac{1}{1-\theta} \quad \text{when} \quad 0 \leqq \theta < 1.$$

It can be seen from Theorems 1 and 2 that the "linearly" full implicit scheme (the case when $\theta = 1$) is absolutely stable, with no restriction to the mesh size and the time increment.

## §3. The proof for the stability of the difference scheme.

Here we give the proof of Theorem 2, and Theorem 1 can be proved in the same way.

The difference equations (2) can be rewritten as

$$
\begin{aligned}
\frac{u_{0k}^{n+1}-u_{0k}^{n}}{\tau} =& g_{0k}^{n}\Bigg\{\frac{2}{h_1^2}\frac{u_{1k}^{n+\theta}-u_{0k}^{n+\theta}}{\left[(g_{\frac12,k}^{n})^{\sigma}+\delta\right]^{1/\sigma}}+\frac{1}{h_2^2}\Big(\frac{u_{0,k+1}^{n+\theta}-u_{0k}^{n+\theta}}{\left[(g_{0,k+\frac12}^{n})^{\sigma}+\delta\right]^{1/\sigma}}\\
&-\frac{u_{0k}^{n+\theta}-u_{0,k-1}^{n+\theta}}{\left[(g_{0,k-\frac12}^{n})^{\sigma}+\delta\right]^{1/\sigma}}\Big)\Bigg\}+\frac{2(u_{1k}^{n+\theta}-u_{0k}^{n+\theta})}{h_1^2}\\
&(n=0,1,2,\cdots;\ k=0,\pm1,\pm2,\cdots),\\
\frac{u_{jk}^{n+1}-u_{jk}^{n}}{\tau} =& g_{jk}^{n}\Bigg\{\frac{1}{h_1^2}\Big(\frac{u_{j+1,k}^{n+\theta}-u_{jk}^{n+\theta}}{\left[(g_{j+\frac12,k}^{n})^{\sigma}+\delta\right]^{1/\sigma}}-\frac{u_{jk}^{n+\theta}-u_{j-1,k}^{n+\theta}}{\left[(g_{j-\frac12,k}^{n})^{\sigma}+\delta\right]^{1/\sigma}}\\
&+\frac{1}{h_2^2}\Big(\frac{u_{j,k+1}^{n+\theta}-u_{jk}^{n+\theta}}{\left[(g_{j,k+\frac12}^{n})^{\sigma}+\delta\right]^{1/\sigma}}-\frac{u_{jk}^{n+\theta}-u_{j,k-1}^{n+\theta}}{\left[(g_{j,k-\frac12}^{n})^{\sigma}+\delta\right]^{1/\sigma}}\Big)\Bigg\}\\
&+\frac{1}{r_j}\cdot\frac{u_{j+1,k}^{n+\theta}-u_{jk}^{n+\theta}}{h_1}\\
&(n=0,1,2,\cdots;\ j=1,2,\cdots;\ k=0,\pm1,\pm2,\cdots),
\end{aligned}
\tag{3}
$$

where the values of $g$ at the fraction points are defined by

$$
g_{j\pm\frac12,k}^{n}=\frac12\left(g_{jk}^{n}+g_{j\pm1,k}^{n}\right),\qquad g_{j,k\pm\frac12}^{n}=\frac12\left(g_{jk}^{n}+g_{j,k\pm1}^{n}\right).
\tag{4}
$$

By introducing

$$
\begin{aligned}
&\alpha=\frac{\tau}{h_1^2},\qquad \beta=\frac{\tau}{h_2^2},\\
&a^{\pm}=a_{njk}^{\pm}=\frac{\alpha g_{jk}^{n}}{\left[(g_{j\pm\frac12,k}^{n})^{\sigma}+\delta\right]^{1/\sigma}},\\
&b^{\pm}=b_{njk}^{\pm}=\frac{\beta g_{jk}^{n}}{\left[(g_{j,k\pm\frac12}^{n})^{\sigma}+\delta\right]^{1/\sigma}}
\end{aligned}
\tag{5}
$$

at each $(t_n, r_j, z_k)$, the equations (3) can be changed to

$$
\begin{aligned}
u_{0k}^{n+1} =& \Big\{1-(1-\theta)\big(2a^+ + b^+ + b^- + 2\alpha\big)\Big\}u_{0k}^n \\
&+(1-\theta)\Big\{\big(2a^+ + 2\alpha\big)u_{1k}^n + b^+ u_{0,k+1}^n + b^- u_{0,k-1}^n\Big\} \\
&+\theta\Big\{\big(2a^+ + 2\alpha\big)u_{1k}^{n+1} + b^+ u_{0,k+1}^{n+1} + b^- u_{0,k-1}^{n+1} \\
&-\big(2a^+ + b^+ + b^- + 2\alpha\big)u_{0k}^{n+1}\Big\}, \\
&\qquad (n = 0,1,2,\cdots;\ k = 0,\pm1,\pm2,\cdots), \\
u_{jk}^{n+1} =& \Big\{1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}u_{jk}^n \\
&+(1-\theta)\Big\{\big(a^+ + \frac{\alpha}{j}\big)u_{j+1,k}^n + a^- u_{j-1,k}^n + b^+ u_{j,k+1}^n + b^- u_{j,k-1}^n\Big\} \\
&+\theta\Big\{\big(a^+ + \frac{\alpha}{j}\big)u_{j+1,k}^{n+1} + a^- u_{j-1,k}^{n+1} + b^+ u_{j,k+1}^{n+1} + b^- u_{j,k-1}^{n+1} \\
&-\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)u_{jk}^{n+1}\Big\}, \\
&\qquad (n = 0,1,2,\cdots;\ j = 1,2,\cdots;\ k = 0,\pm1,\pm2,\cdots), \\
u_{jk}^0 =& u_0(r_j, z_k) \quad (j = 0,1,2,\cdots;\ k = 0,\pm1,\pm2,\cdots).
\end{aligned} \tag{6}
$$

Noting that $g_{jk}^n \geqq 0$ and $\delta > 0$ we can get the following

LEMMA 1.

$$0 \leqq a^\pm = a_{njk}^\pm < 2\alpha, \qquad 0 \leqq b^\pm = b_{njk}^\pm < 2\beta.$$

Thus, we can show the following maximum results as

LEMMA 2. *If the values of* $\theta$, $a^\pm = a_{njk}^\pm$ *and* $b^\pm = b_{njk}^\pm$ *satisfies*

$$1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + 2\alpha\big) \geqq 0 \tag{7}$$

*then the difference solution of (2) satisfies*

$$
\begin{aligned}
\inf_{j,k} u_{jk}^n &\geqq \inf_{j,k} u_{jk}^0, \\
\max_{j,k} u_{jk}^n &\leqq \max_{j,k} u_{jk}^0, \qquad n = 1,2,\cdots.
\end{aligned}
$$

PROOF OF LEMMA 2: Rewriting the difference equations (6) and noting that $a^+ = a^-$

for $j=0$, we get

$$\begin{aligned}
&\Big\{1+\theta\big(a^+ + a^- + b^+ + b^- + 2\alpha\big)\Big\}u_{0k}^{n+1} \\
&\quad=\Big\{1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + 2\alpha\big)\Big\}u_{0k}^{n} \\
&\qquad+(1-\theta)\Big\{\big(a^+ + a^- + 2\alpha\big)u_{1k}^{n} + b^+ u_{0,k+1}^{n} + b^- u_{0,k-1}^{n}\Big\} \\
&\qquad+\theta\Big\{\big(a^+ + a^- + 2\alpha\big)u_{1k}^{n+1} + b^+ u_{0,k+1}^{n+1} + b^- u_{0,k-1}^{n+1}\Big\}, \\
&\qquad\qquad (n=0,1,2,\cdots;\ k=0,\pm1,\pm2,\cdots),
\end{aligned}$$

(8)
$$\begin{aligned}
&\Big\{1+\theta\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}u_{jk}^{n+1} \\
&\quad=\Big\{1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}u_{jk}^{n} \\
&\qquad+(1-\theta)\Big\{\big(a^+ + \frac{\alpha}{j}\big)u_{j+1,k}^{n} + a^- u_{j-1,k}^{n} + b^+ u_{j,k+1}^{n} + b^- u_{j,k-1}^{n}\Big\} \\
&\qquad+\theta\Big\{\big(a^+ + \frac{\alpha}{j}\big)u_{j+1,k}^{n+1} + a^- u_{j-1,k}^{n+1} + b^+ u_{j,k+1}^{n+1} + b^- u_{j,k-1}^{n+1}\Big\}, \\
&\qquad\qquad (n=0,1,2,\cdots;\ j=1,2,\cdots;\ k=0,\pm1,\pm2,\cdots), \\
&u_{jk}^{0}=u_0(r_j,z_k) \qquad (j=0,1,2,\cdots;\ k=0,\pm1,\pm2,\cdots).
\end{aligned}$$

Note that we have done our computation in a certain finite domain and on the boundary of the domain the value of the solution is a constant $-C$. Here, we claim that the maximum and minimum value of the difference solution $u_{jk}^n$ are reached on the boundary, or the initial plane (i.e. $n=0$). This is shown in the following way. If $j>0$, since $\alpha/j<2\alpha$, we have $\Big\{1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}\geqq 0$ and in a inner (mesh) point of the domain, say $(t_n, r_j, z_k)$, the difference solution satisfies

$$\begin{aligned}
&\Big\{1+\theta\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}u_{jk}^{n+1} \\
&\quad\leqq\Big\{1-(1-\theta)\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}\|u^n\| \\
&\qquad+(1-\theta)\Big\{\big(a^+ + \frac{\alpha}{j}\big)\|u^n\| + a^-\|u^n\| + b^+\|u^n\| + b^-\|u^n\|\Big\} \\
&\qquad+\theta\Big\{\big(a^+ + \frac{\alpha}{j}\big)\|u^{n+1}\| + a^-\|u^{n+1}\| + b^+\|u^{n+1}\| + b^-\|u^{n+1}\|\Big\}, \\
&\quad=\|u^n\| + \theta\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\|u^{n+1}\| \\
&\qquad\qquad (n=0,1,2,\cdots;\ j=1,2,\cdots;\ k=0,\pm1,\pm2,\cdots)
\end{aligned}$$

where $\|\cdot\| = \|\cdot\|_\infty$ here and hereafter. So, we get

$$\begin{aligned}
&\Big\{1+\theta\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\Big\}\|u^{n+1}\| \\
&\qquad\qquad\leqq\|u^n\| + \theta\big(a^+ + a^- + b^+ + b^- + \frac{\alpha}{j}\big)\|u^{n+1}\|,
\end{aligned}$$

which leads to

$$\|u^{n+1}\| \leqq \|u^n\|.$$

While if $j = 0$, then the estimates become

$$\begin{aligned}
&\Big\{1+\theta\big(2a^+ + b^+ + b^- + 2\alpha\big)\Big\}u_{0k}^{n+1} \\
&\qquad \leqq \Big\{1-(1-\theta)\big(2a^+ + b^+ + b^- + 2\alpha\big)\Big\}\|u^n\| \\
&\qquad\quad + (1-\theta)\Big\{\big(2a^+ + 2\alpha\big)\|u^n\| + b^+\|u^n\| + b^-\|u^n\|\Big\} \\
&\qquad\quad + \theta\Big\{\big(2a^+ + 2\alpha\big)\|u^{n+1}\| + b^+\|u^{n+1}\| + b^-\|u^{n+1}\|\Big\}, \\
&\qquad = \|u^n\| + \theta\big(2a^+ + b^+ + b^- + 2\alpha\big)\|u^{n+1}\| \\
&\qquad\qquad (n = 0,1,2,\cdots;\ k = 0,\pm1,\pm2,\cdots)
\end{aligned}$$

which also leads to

$$\|u^{n+1}\| \leqq \|u^n\|.$$

Here, if we take max instead of $\|\cdot\|$ in the above argument, then we can obtain

$$\max_{j,k} u_{jk}^n \leqq \max_{j,k} u_{jk}^0.$$

And similarly we can show another inequality

$$\inf_{j,k} u_{jk}^n \geqq \inf_{j,k} u_{jk}^0.$$

Thus we have completed the proof of Lemma 2. ∎

From Lemma 2, we can obtain our stability condition. Either of the following conditions leads to (7), the condition $\theta = 1$ or

$$a^+ + a^- + b^+ + b^- + 2\alpha \leqq \frac{1}{1-\theta} \qquad \text{if} \quad 0 \leqq \theta < 1. \tag{9}$$

While in virtue of Lemma 1, (9) holds true if

$$6\alpha + 4\beta \leqq \frac{1}{1-\theta}$$

is satisfied. This completes the proof for Theorem 2.

To prove Theorem 1, we have only to make a little change in the estimates: simply omit the terms $2\alpha$ and $\frac{\alpha}{j}$ in the statements.

## §4. For the case of generalized mean curvature flow equation.

With the above-mentioned methods, we can construct a stable difference scheme for the so-called generalized mean curvature flow equation

$$(E') \qquad u_t = |\nabla u| \operatorname{div}\left(\frac{\nabla u}{|\nabla u|}\right) + \nu|\nabla u|, \quad (t,x) \in Q = (0,\infty) \times \mathbb{R}^N$$

where $\nu$ is a constant (see [CGG1]).

The difference scheme for (E') is constructed by the following two parts:
(1) the first part of the scheme is constructed as that for (E) in the previous section;
(2) the second part of the scheme is constructed in the way of any kind of stable difference scheme with monotonicity for the Hamilton-Jacobi equation $u_t = \nu|\nabla u|$, such as Lax-Friedrichs scheme, Godunov scheme, etc. (see, for example [CL]).

Then, we can show that the obtained difference scheme is stable if the value of $\tau/h_i^2$ $(i = 1, 2)$ are taken sufficiently small ([CGH]).

## §5. Some remarks.

1. It is important to note that the stability conditions do not depend on $\delta > 0$ and $\sigma \geqq 1$.

2. If $g_{jk}^n$ is not positive definite for $\{D^{\pm} u_{jk}^n\}$, then the difference solution may not converge to the solution of $(E_\delta)$, nor to that of (E) when $\delta \to 0$.

**Example.** It is easy to see that

$$u(t,x) = \max\{1 - (2t + |x|^2), -1\}$$

is the unique viscosity solution of the level set equation (E) with initial data

$$u_0(x) = \max\{1 - |x|^2, -1\}.$$

Here, if $g_{jk}^n$ in the scheme is replaced by a general central difference approximation, then the value of numerical solution $u_{jk}^n$ at the origin is independent of $n \geqq 0$, because the symmetry forces the central difference at the origin equal to zero.

However, the value of the analytic solution $u(t,x)$ at the origin equals $1-2t$ (for $t \leqq \frac{1}{2}$) which is smaller than 1 for $t > 0$. Thus the numerical solution $u_{jk}^n$ does not pointwisely converge to the viscosity solution.

The same remark applies to the scheme in [OS]. Since the central difference approximation is used in [OS], the numerical solution there may not converge to the analytic solution. In that paper, in order to avoid the problem they shifted the grids so that non

of the net points agrees with the center of symmetry, for the case when symmetric data are started with.

3. Osher and Sethian discussed some difference schemes constructed in a different way with level surface approach ([OS], [S]). They computed several interesting examples including the torus and dumbbells without discussing the fundamental theory such as stability, etc. In [S], an example of unstable computation of a torus was presented with a quite large $\Delta t$ but no condition for the stability was given there.

4. In [OS] and [S], the axisymmetric surfaces are computed under the rectilinear coordinates instead of the cylindrical coordinates.

With our stable difference schemes and level surface approach, we investigated motions of several typical surfaces, including the shrink of a torus (surface of a doughnut) and the break of a dumbbell. With this method we can track motions of a surface even after the time when a singularity occurs.

## §6. Numerical results.

We present several computation results on the evolution of axisymmetric surfaces here: the "fat torus" and "Hamilton's dumbbell". The computation is carried out by our difference scheme (2) in its explicit form (with $\theta = 0$) while the "safety parameter" $\delta$ is taken $\delta = 10^{-50}$ and $\sigma = 2$. We restricted the domain to $(r, z) \in [0, 1.5] \times [-1.5, 1.5]$ which is discreted into net points with the mesh sizes $h_1 = h_2 = 0.015$ and the time increment $\tau = 0.00001125$. On the axis of symmetry $r = 0$, the boundary condition is treated with the symmetry property of the solution.

### 1. Fat Torus

As the analytic theory predicts ([SS]), no fattening occurs for generalized solution after it collapses. According to numerical simulation, it becomes convex and shrinks to a point. If the torus is thin, it converges to a "marriage ring" ([AI]).

### 2. Hamilton's Dumbbell

We consider the initial surface of the form

$$\Gamma_0^\lambda = \{(x, y, z);\ x^2 + y^2 = (1 - z^2)(1 - \lambda + \lambda z^2)^2\}$$

where $0 \leqq \lambda \leqq 1$ is a parameter. We take the initial data

$$u_0 = (1 - z^2)(1 - \lambda + \lambda z^2)^2 - r^2$$

and select the zero level surface to observe. If $\lambda$ is close to 1, $\Gamma_t^\lambda$ pinches into two pieces in a finite time. Then both pieces become convex. Actually, according to Altshuler, Angenent and Giga [AAG], no fattening occurs and $\Gamma_t^\lambda$ becomes smooth instantaneously after it pinches. At last each of the pieces shrinks to one point, respectively. See the cases $\lambda = 0.8$ and $\lambda = 0.65$.

From the numerical results we can see that $\Gamma_t^\lambda$ becomes convex and then shrinks to a single point for $\lambda = 0.63$.

In between $\lambda = 0.63$ and $\lambda = 0.65$, there may exists a critical value of $\lambda$ for which the behavior of the shrinking dumbbell is exceptional. The surface does not become convex nor does it pinch. It stays smooth and shrinks to a point. The existence of such a dumbbell is proved analytically in [AAG] by topological argument. See the picture of the case $\lambda = 0.64$, which seems to be the critical value.

Recently, Nochetto et al. [NPV] calculated the evolution of $\Gamma_t^\lambda$ by a different method. Their calculation suggests that the critical value be $\lambda = 0.654$, which is very close to ours.

**Acknowledgement.**

This work is partially supported by the Japan Ministry of Education, Science and Culture, and partially supported by Environmental Research Institute of Hokkaido Tokai University. Here, we express our gratitude to Professors Tsutomu Ikeda and Hitoshi Ishii for helpful advices.

## References

[AI]. K.Ahara and N.Ishimura, *On the mean curvature flow of "thin" doughnuts*, Lecture Notes in Num.Appl.Anal. (Ed. by K.Masuda, M.Mimura and T.Nishida) Kinokuniya, Tokyo **12** (1993), 1–33.

[AAG]. S.Altshuler, S.Angenent and Y.Giga, *Motion by mean curvature through singularities for surfaces of rotation*, J. Geom. Anal., to appear.

[BMO]. J.Bence, B.Merriman and S.Osher, *Diffusion generated motion by mean curvature*, preprint (1992).

[C]. Yun-Gang Chen, *Blow-up solutions to a finite difference analogue of* $u_t = \Delta u + u^{1+\alpha}$ *in N-dimensional balls*, Hokkaido Mathematical Journal **21** (1992), 447–474.

[CGG1]. Y.-G.Chen, Y.Giga and S.Goto, *Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations*, J.Differential Geom. **33** (1991), 749–786.

[CGG2]. Y.-G.Chen, Y.Giga and S.Goto, *Analysis toward snow crystal growth*, Proceedings of International Symposium on Functional Analysis and Related Topics, Sapporo, Ed. S.Koshi (1991), 43–57.

[CGG3]. Y.-G.Chen, Y.Giga and S.Goto, *Remarks on viscosity solutions for evolution equations*, Proc. Japan Acad. Ser.A **67** (1991), 323–328.

[CGH]. Y.-G.Chen, Y.Giga and M.Honma, *On difference schemes for generalized mean curvature flow equations*, in preparation.

[CL]. M. G. Crandall and P. L. Lions, *Two approximations of solutions of Hamilton-Jacobi equations*, Math. Comp. **43** (1984), 1–19.

[D]. G. Dziuk, *An algorithm for evolutionary surfaces*, Numer. Math. **58** (1991), 603–611.

[DD]. K.Deckelnick and G.Dziuk, *Convergence of a finite element method for non-parametric mean curvature flow*, preprint.

[E]. L.C.Evans, *Convergence of an algorithm for mean curvature motion*, Indiana Univ. Math. J. **42** (1993), 533–557.

[ES]. L.C.Evans and J.Spruck, *Motion of level sets by mean curvature, I.*, J.Differential Geom. **33** (1991), 635–681.

[NPV]. R. H. Nochetto, M. Paolini and C. Verdi, *Double obstacle formulation with variable relaxation parameter for smooth geometric front evolutions: asymptotic interface error estimates*, preprint.

[OS]. S.Osher and J.A.Sethian, *Fronts propagating with curvature dependent speed: Algorithms based on Halmiton-Jacobi formulations*, J. Comput. Phys. **79** (1988), 12–49.

[S]. J. A. Sethian, *Numerical algorithms for propagating interfaces: Hamilton-Jacobi equation and conservation laws*, J. Differential Geom. **31** (1990), 131–161.

[SS]. H.M.Soner and P.E.Souganidis, *Singularities and uniqueness of cylindrically symmetric surfaces moving by mean curvature*, Commun. in Partial Differential Equations **18** (1993), 859–894.

[W]. N.J.Walkington, *Algorithm for computing motion by mean curvature*, preprint.

Hokkaido Tokai University

Sapporo 005, Japan

and

Department of Mathematics

Hokkaido University

Sapporo 060, Japan

The present address of Toshio HITAKA:
NTT Corporation, Tokyo, Japan

The present address of Mitsuru HONMA:
Kao Institute for Knowledge and Intelligence Science
Kao Corporation, Tokyo 131, Japan




Time= 0.00000000




Time= 0.00036032




Time= 0.04556250

**Figure 1.** Evolution of a fat torus.




**Figure 2.** Evolution of Hamilton's dumbbell of the case $\lambda = 0.80$.




**Figure 3.** Evolution of Hamilton's dumbbell of the case $\lambda = 0.65$.




**Figure 4.** Evolution of Hamilton's dumbbell of the case $\lambda = 0.63$.




**Figure 5.** Evolution of Hamilton's dumbbell of the case $\lambda = 0.64$. The scale of the coordinate is changed in the last two pictures to show the details of the shrinking.