HOKKAIDO UNIVERSITY

Title	An Abel-Tauber theorem for Fourier sine transforms
Author(s)	Inoue, A.
Citation	Hokkaido University Preprint Series in Mathematics, 261, 1-6
Issue Date	1994-08-01
DOI	https://doi.org/10.14943/83408
Doc URL	https://hdl.handle.net/2115/69012
Type	departmental bulletin paper
File Information	re261.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

An Abel-Tauber theorem for Fourier sine transforms

Akihiko Inoue

Series #261. August 1994

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

236: T. Mikami, Large deviations for the first exit time on small random perturbations of dynamical systems, 32 pages. 1994.

237 T. Ozawa, Characterization of Trudinger's inequality, 7 pages. 1994.

238 T. Hibi, Buchsbaum complexes with linear resolutions, 12 pages. 1994.

239 Y. Giga, N. Mizoguchi, On time periodic solutions of the Dirichlet problem for degenerate parabolic equations of nondivergence type, 23 pages. 1994.

240 C. Dohmen, Existence of Fast Decaying Solutions to a Haraux-Weissler Equation With a Prescribed Number of Zeroes, 12 pages. 1994.

241 K. Sugano, Note on H-separable Frobenius extensions, 8 pages. 1994.

242 J. Zhai, Some Estimates For The Blowing up Solutions of Semilinear Heat Equations, 11 pages. 1994.

243 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and Smoothing Effect of the Schrödinger Evolution Group, 10 pages. 1994.

244 D. Lehmann, T. Suwa, Residues of holomorphic vector fields relative to singular invariant subvarieties, 26 pages. 1994.

245 H. Kubo, Slowly decaying solutions for semilinear wave equations in odd space dimensions, 30 pages. 1994.

246 T. Nakazi, M. Yamada, (A_2)-Conditions and Carleson Inequalities, 27 pages. 1994.

247 N. Hayashi, K. Kato and T. Ozawa, Dilation Method and smoothing Effect of Solutions to the Benjamin-ono Equation, 17 pages. 1994.

248 H. Kikuchi, Sheaf cohomology theory for measurable spaces, 12 pages. 1994.

249 A. Inoue, Tauberian theorems for Fourier cosine transforms, 9 pages. 1994.

250 S. Izumiya, G. T. Kossioris, Singularities for viscosity solutions of Hamilton-Jacobi equations, 23 pages. 1994.

251 H. Kubo, K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values p in odd space dimensions, 51 pages. 1994.

252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.

253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.

254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.

255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.

256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.

257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.

258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.

259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.

260 T. Mikami, Copula fields and its applications, 14 pages. 1994.

An Abel-Tauber theorem for Fourier sine transforms

By Akihiko INOUE

Abstract. We prove an Abel-Tauberian theorem for Fourier sine transforms. It can be considered as the analogue of the Abel-Tauber theorem of Pitman in the boundary case. We apply it to Fourier sine series as well as to the tail behavior of a probability distribution.

1. Introduction and results

The aim of this paper is to prove an Abel-Tauber theorem for Fourier sine transforms. It characterizes, for example, the asymptotic behavior $f(t) \sim 1/t^2$ as $t \to \infty$ in terms of the Fourier sine transform of f, where f is a locally integrable, eventually non-increasing function on $[0, \infty)$ such that $\lim_{t\to\infty} f(t) = 0$. A similar result for Fourier sine series will be obtained as a corollary.

To state our results, we recall and introduce some notation. We denote by R_0 the whole class of slowly varying functions at infinity, that is, R_0 is the class of positive measurable l, defined on some neighborhood of infinity, satisfying

$$\forall \lambda > 0, \qquad \lim_{x\to\infty} l(\lambda x)/l(x) = 1.$$

For $l \in R_0$, the class Π_l is the class of measurable g satisfying

$$\forall \lambda \geq 1, \qquad \lim_{x\to\infty} \{g(\lambda x) - g(x)\}/l(x) = c \log \lambda$$

for some constant c called the *l-index* of g. It is useful to name the class of functions of which we define the Fourier sine transforms. The function $f : [0, \infty) \to \mathbb{R}$ belongs to $DL^1_{loc}[0, \infty)$ if it is locally integrable and eventually non-increasing on $[0, \infty)$, $\lim_{t\to\infty} f(t) = 0$. For $f \in DL^1_{loc}[0, \infty)$, we define its *Fourier sine transform* F_s by

$$F_s(\xi) = \int_0^{\infty-} f(t) \sin t\xi \, dt \qquad (0 < \xi < \infty), \tag{1.1}$$

where we write $\int_0^{\infty-}$ to denote an improper integral obtained from $\int_0^M$ by letting $M \uparrow \infty$. Since the improper integral on the right converges uniformly on each (a, ∞) with $a > 0$, F_s is a continuous function on $(0, \infty)$. See the proof of Theorem 6 of Titchmarsh [6].

Here is the main theorem of this paper:

1991 *Mathematics Subject Classification.* Primary 40E05; Secondary 42A32, 60E10.

THEOREM 1.1. *Let $l \in R_0$ and $f \in DL^1_{loc}[0,\infty)$. Let F_s be the Fourier sine transform of f. Then*

$$f(t) \sim t^{-2} l(t) \qquad (t \to \infty) \tag{1.2}$$

if and only if

$$xF_s(1/x) \in \Pi_l \text{ in } x \text{ with } l\text{-index } 1. \tag{1.3}$$

The analogue for Fourier sine series is:

THEOREM 1.2. *Let $l \in R_0$. Suppose that the real sequence $\{b_n\}$ is eventually non-increasing, and tends to 0 as $n \to \infty$. We set*

$$g_s(\xi) = \sum_{n=1}^{\infty} b_n \sin n\xi \qquad (0 < \xi < 2\pi). \tag{1.4}$$

Then

$$b_n \sim n^{-2} l(n) \qquad (n \to \infty) \tag{1.5}$$

if and only if

$$xg_s(1/x) \in \Pi_l \text{ in } x \text{ with } l\text{-index } 1. \tag{1.6}$$

Now we recall the Abel-Tauber theorem of Pitman [4] which is closely related to Theorem 1.1. Let $l \in R_0$ and $0 < \alpha < 2$. Let $f \in DL^1_{loc}[0,\infty)$, and let F_s be the Fourier sine transform of f. Then, by Pitman [4],

$$f(t) \sim t^{-\alpha} l(t) \qquad (t \to \infty) \tag{1.7}$$

if and only if

$$F_s(\xi) \sim \xi^{\alpha-1} l(1/\xi)\Gamma(1-\alpha)\cos(\pi\alpha/2) \qquad (\xi \to 0+). \tag{1.8}$$

From this and Theorem 1.1, we see that the behavior (1.2) is a critical case; we need Π-variation to characterize (1.2) in terms of the Fourier sine transform of f. For the related Abel-Tauber theorems for integral transforms, we refer to Chapter 4 of Bingham, Goldie and Teugels [1].

In the proof of Theorem 1.1, we will find that it is enough to prove the theorem for $f \in DL^1_{loc}[0,\infty)$ finite and non-increasing on $[0,\infty)$. For such f, we have the inversion formula which represents f by the Fourier sine transform of f (see Theorem 7 of Titchmarsh [6]). However it is difficult to use to prove the Tauberian implication (1.3) $\Rightarrow$ (1.2). The difficulty is in that the problem we have to deal with involves both Π-variation and improper integrals. The key to the proof is to reduce the problem to the analogous result of Inoue [2], [3] for Fourier cosine transforms.

The proofs of Theorems 1.1 and 1.2 will be given in section 2. In section 3, we apply Theorem 1.1 to the tail behavior of a probability distribution.

2. Proofs of Theorems 1.1 and 1.2

For $f \in DL^1_{loc}[0,\infty)$, we define its *Fourier cosine transform* F_c by

$$F_c(\xi) = \int_0^{\infty-} f(t)\cos t\xi dt \qquad (0 < \xi < \infty). \tag{2.1}$$

We have the following Abel-Tauber theorem for Fourier cosine transforms:

THEOREM 2.1 (Pitman [4] and Inoue [2], [3]). *Let $l \in R_0$ and $f \in DL^1_{loc}[0,\infty)$. Let F_c be the Fourier cosine transform of f. Then*

$$f(t) \sim t^{-1} l(t) \qquad (t \to \infty) \tag{2.2}$$

if and only if

$$F_c(1/\cdot) \in \Pi_l \text{ with } l\text{-index } 1. \tag{2.3}$$

The Abelian implication (2.2) $\Rightarrow$ (2.3) is essentially Theorem 7 (iii) of Pitman [4], while the Tauberian assertion (2.3) $\Rightarrow$ (2.2) is due to Inoue [2], [3].

Using Theorem 2.1, we shall prove Theorem 1.1.

PROOF OF THEOREM 1.1. Choose M so large that f is positive, finite and non-increasing on $[M,\infty)$. We set

$$g(t) = \begin{cases} f(M) & (0 \le t < M), \\ f(t) & (M \le t < \infty). \end{cases}$$

Let G_s be the Fourier sine transform of g. We set

$$a(x) = x^{-1} \sin x \qquad (0 < x < \infty). \tag{2.4}$$

Then by the mean-value theorem,

$$|a(x) - a(y)| \le \text{const.}|x - y| \qquad (0 < x, y < 1), \tag{2.5}$$

so for any $\lambda > 1$,

$$\begin{aligned} &|\lambda x F_s(1/(\lambda x)) - x F_s(1/x) - \lambda x G_s(1/(\lambda x)) + x G_s(1/x)|/l(x) \\ &= \frac{1}{l(x)} \left| \int_0^M t \{f(t) - f(M)\} \{a(t/(\lambda x)) - a(t/x)\} \, dt \right| \\ &\le \text{const.} \frac{(1 - \lambda^{-1})}{x l(x)} \int_0^M t^2 |f(t) - f(M)| dt \to 0 \qquad (x \to \infty), \end{aligned}$$

whence (1.3) holds if and only if $xG_s(1/x) \in \Pi_l$ in x with l-index 1. Therefore we may assume that f is positive, finite and non-increasing on $[0,\infty)$.

First we assume (1.2). Then f is integrable over $[0,\infty)$, and so, by integration by parts,

$$F_s(\xi)/\xi = \int_0^{\infty -} h(t) \cos t\xi dt \qquad (0 < \xi < \infty) \tag{2.6}$$

with

$$h(t) = \int_t^\infty f(s) ds \qquad (0 \le t < \infty). \tag{2.7}$$

Since (1.2) is equivalent to $h(t) \sim t^{-1}l(t)$ as $t \to \infty$, we immediately obtain (1.3) by the Abelian implication of Theorem 2.1.

Next, we prove that (1.3) implies (1.2). By the second mean-value theorem for integrals,

$$F_s(\xi) = \lim_{U\to\infty} \int_0^U f(t) \sin t\xi dt = \lim_{U\to\infty} f(0+) \int_0^\zeta \sin t\xi dt$$

for some $\zeta \in (0, U)$, and so $\xi F_s(\xi)$ is bounded on $(0, \infty)$. In particular, for any $x > 0$, $F_s(\xi)(1 - \cos x\xi)/\xi$ is integrable over $(0, \infty)$. Then by applying Theorem 38 of Titchmarsh [6] to the pair of $\mathrm{I}_{[0,x]}$ and f, we obtain

$$\int_0^x f(t)dt = \frac{2}{\pi} \int_0^\infty F_s(\xi) \frac{1 - \cos x\xi}{\xi} d\xi \qquad (0 < x < \infty).$$

By Theorem 3.7.4 of Bingham, Goldie and Teugels [1], (1.3) implies $|xF_s(1/x)| \in R_0$ in x, so $F_s(\xi)/\xi$ is integrable on $(0, 1)$ whence on $(0, \infty)$. Thus for any $x > 0$,

$$\int_0^x f(t)dt \le \frac{4}{\pi} \int_0^\infty \frac{|F_s(\xi)|}{\xi} d\xi < \infty,$$

whence f is integrable over $(0, \infty)$. Again we arrive at (2.6) with (2.7), and similarly we obtain (1.2) from (1.3) by the Tauberian implication of Theorem 2.1. □

Following the method of Soni and Soni [5], we shall prove Theorem 1.2 as a corollary of Theorem 1.1.

PROOF OF THEOREM 1.2. We set

$$f(t) = \begin{cases} 0 & (0 \le t < 1/2), \\ b_n & (n - 1/2 \le t < n + 1/2, \quad n = 1, 2, \cdots). \end{cases}$$

Then f is in $DL^1_{loc}[0, \infty)$, and (1.5) is equivalent to

$$f(t) \sim t^{-2}l(t) \qquad (t \to \infty). \tag{2.8}$$

Let F_s be the Fourier sine transform of f. Then by a simple calculation,

$$F_s(\xi) = a(\xi/2)g_s(\xi) \qquad (0 < \xi < 2\pi),$$

where a is defined by (2.4). So we have for any $\lambda > 1$ and $x > 0$,

$$\begin{aligned} &\{\lambda x F_s(1/(\lambda x)) - xF_s(1/x)\}/l(x) \\ &\quad = a(1/(2\lambda x))\{\lambda x g_s(1/(\lambda x)) - x g_s(1/x)\}/l(x) \\ &\qquad + x g_s(1/x)\{a(1/(2\lambda x)) - a(1/(2x))\}/l(x). \end{aligned} \tag{2.9}$$

By Theorem 3.7.4 of Bingham, Goldie and Teugels [1], (1.6) implies $|xg_s(1/x)| \in R_0$ in x, so by (2.5) the second term on the right of (2.9) tends to 0 as $x \to \infty$. Therefore

(1.6) implies $xF_s(1/x) \in \Pi_l$ in x with l-index 1, which, by the Tauberian implication of Theorem 1.1, implies (2.8) whence (1.5).

Conversely, we set $c(x) = 1/a(x)$ for $x > 0$. Then we also have

$$|c(x) - c(y)| \leq \text{const.}|x - y| \qquad (0 < x, y < 1).$$

Arguing similarly, we obtain (1.6) from (1.5), which completes the proof. □

3. Application to the tail behavior

In this section, we apply Theorem 1.1 to the tail behavior of a probability distribution. For the related results, we refer to Pitman [4] as well as pp. 336-337 of Bingham, Goldie and Teugels [1], and Inoue [3].

Let X be a real random variable defined on a probability space $(\Omega, \mathcal{F}, P)$. The *tail-sum* of X is the function T defined by

$$T(x) = P(X \leq -x) + P(X > x) \qquad (0 \leq x < \infty).$$

Let U be the real part of the characteristic function of X:

$$U(\xi) = E[\cos \xi X] \qquad (\xi \in \mathbb{R}).$$

Then we have

$$\{1 - U(\xi)\}/\xi = \int_0^{\infty -} T(x) \sin \xi x dx \qquad (0 < \xi < \infty).$$

Since T is finite and non-increasing on $[0, \infty)$, $\lim_{x \to \infty} T(x) = 0$, by Theorem 1.1 we immediately obtain

Theorem 3.1. *Let $l \in R_0$. Then $T(x) \sim x^{-2} l(x)$ as $x \to \infty$ if and only if $x^2\{1 - U(1/x)\} \in \Pi_l$ in x with l-index 1.*

References

[1] Bingham, N. H., Goldie, C. M. and J. L. Teugels, Regular variation, Cambridge University Press, Cambridge, 1987.

[2] Inoue, A., Regularly varying correlation functions of solutions of delay Langevin equations, Preprint (1994).

[3] Inoue, A., Tauberian theorems for Fourier cosine transforms, Preprint (1994).

[4] Pitman, E. J. G., On the behaviour of the characteristic function of a probability distribution in the neighbourhood of the origin, J. Austral. Math. Soc. 8 (1968), 422–443.

[5] Soni, K. and R. P. Soni, Slowly varying functions and asymptotic behaviour of a class of integral transforms III, J. Math. Anal. Appl. 49 (1975), 612–628.

[6] Titchmarsh, E. C., Introduction to the Theory of Fourier Integrals, 2nd ed., Oxford University Press, London, 1948.

Department of Mathematics
Hokkaido University
Sapporo
060 Japan