



HOKKAIDO UNIVERSITY

Title	Computation of Betti numbers of monomial ideals associated with stacked polytopes
Author(s)	Terai, N.; Hibi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 278, 1-8
Issue Date	1995-01-01
DOI	https://doi.org/10.14943/83425
Doc URL	https://hdl.handle.net/2115/69029
Type	departmental bulletin paper
File Information	re278.pdf



**Computation of Betti
numbers of monomial ideals
associated with stacked polytopes**

N. Terai and T. Hibi

Series #278. January 1995

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 252 T. Mikami, Large Deviations and Central Limit Theorems for Eyraud-Farlie-Gumbel-Morgenstern Processes, 9 pages. 1994.
- # 253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.
- # 254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.
- # 255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.
- # 256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.
- # 257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.
- # 258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.
- # 259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.
- # 260 T. Mikami, Copula fields and its applications, 14 pages. 1994.
- # 261 A. Inoue, An Abel-Tauber theorem for Fourier sine transforms, 6 pages. 1994.
- # 262 N. Kawazumi, Homology of hyperelliptic mapping class groups for surfaces, 13 pages. 1994.
- # 263 Y. Giga, M. E. Gurtin, A comparison theorem for crystalline evolution in the plane, 14 pages. 1994.
- # 264 J. Wierzbicki, On Commutativity of Diagrams of Type II_1 Factors, 26 pages. 1994.
- # 265 N. Hayashi, T. Ozawa, Schrödinger Equations with nonlinearity of integral type, 12 pages. 1994.
- # 266 T. Ozawa, On the resonance equations of long and short waves, 8 pages. 1994.
- # 267 T. Mikami, A sufficient condition for the uniqueness of solutions to a class of integro-differential equations, 9 pages. 1994.
- # 268 Y. Giga, Evolving curves with boundary conditions, 10 pages. 1994.
- # 269 A. Arai, Operator-theoretical analysis of representation of a supersymmetry algebra in Hilbert space, 12 pages. 1994.
- # 270 A. Arai, Gauge theory on a non-simply-connected domain and representations of canonical commutation relations, 18 pages. 1994.
- # 271 S. Jimbo, Y. Morita and J. Zhai, Ginzburg Landau equation and stable steady state solutions in a non-trivial domain, 17 pages. 1994.
- # 272 S. Izumiya, A. Takiyama, A time-like surface in Minkowski 3-space which contains light-like lines, 7 pages. 1994.
- # 273 K. Tsutaya, Global existence of small amplitude solutions for the Klein-Gordon-Zakharov equations, 11 pages. 1994.
- # 274 H. Kubo, On the critical decay and power for semilinear wave equations in odd space dimensions, 22 pages. 1994.
- # 275 N. Terai, T. Hibi, Alexander duality theorem and second Betti numbers of Stanley-Reisner rings, 2 pages. 1995.
- # 276 N. Terai, T. Hibi, Stanley-Reisner rings whose Betti numbers are independent of the base field, 12 pages. 1995.
- # 277 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with cyclic polytopes, 11 pages. 1995.

Computation of Betti numbers of monomial ideals associated with stacked polytopes

Naoki Terai Takayuki Hibi

Abstract

Let $P(v, d)$ be a stacked d -polytope with v vertices, $\Delta(P(v, d))$ the boundary complex of $P(v, d)$, and $k[\Delta(P(v, d))] = A/I_{\Delta(P(v, d))}$ the Stanley–Reisner ring of $\Delta(P(v, d))$ over a field k . We compute the Betti numbers which appear in a minimal free resolution of $k[\Delta(P(v, d))]$ over A , and show that every Betti number depends only on v and d and is independent of the base field k .

Introduction

Let Δ be a simplicial complex on the vertex set $V = \{x_1, x_2, \dots, x_v\}$. Thus Δ is a collection of subsets of V such that (i) $\{x_i\} \in \Delta$ for every $1 \leq i \leq v$ and (ii) $\sigma \in \Delta, \tau \subset \sigma \Rightarrow \tau \in \Delta$. Each element σ of Δ is called a *face* of Δ . Set $d = \max\{\#\sigma; \sigma \in \Delta\}$ and define the dimension of Δ to be $\dim \Delta = d - 1$. Here $\#\sigma$ is the cardinality of a finite set σ . A maximal face of Δ is also called a *facet* of Δ .

Let $A = k[x_1, x_2, \dots, x_v]$ denote the polynomial ring in v -variables over a field k with the standard grading, i.e., each $\deg x_i = 1$. We identify each $x_i \in V$ with the indeterminate x_i of A . Define I_Δ to be the ideal of A which is generated by square-free monomials $x_{i_1}x_{i_2}\cdots x_{i_r}$, $1 \leq i_1 < i_2 < \cdots < i_r \leq v$, with $\{x_{i_1}, x_{i_2}, \dots, x_{i_r}\} \notin \Delta$. The quotient algebra $k[\Delta] := A/I_\Delta$ is called the *Stanley–Reisner ring* of Δ over k . We may regard $k[\Delta] = \bigoplus_{n \geq 0} (k[\Delta])_n$ as a graded module over A with the quotient grading. We refer the reader to [Bru–Her], [H], [Hoc], [Sta] for the detailed information about Stanley–Reisner rings.

We are interested in a minimal free resolution of $k[\Delta]$ over A . Let $A(j)$, $j \in \mathbb{Z}$, denote the graded module $A(j) = \bigoplus_{n \in \mathbb{Z}} [A(j)]_n$ over A with $[A(j)]_n := A_{n+j}$. Here $\mathbb{Z}$ is the set of integers.

A graded *finite free resolution* of $k[\Delta]$ over A is an exact sequence

$$0 \longrightarrow \bigoplus_{j \in \mathbb{Z}} A(-j)^{\beta_{h,j}} \xrightarrow{\varphi_h} \dots \xrightarrow{\varphi_2} \bigoplus_{j \in \mathbb{Z}} A(-j)^{\beta_{1,j}} \xrightarrow{\varphi_1} A \xrightarrow{\varphi_0} k[\Delta] \longrightarrow 0 \quad (1)$$

of graded modules over A , where each $\bigoplus_{j \in \mathbb{Z}} A(-j)^{\beta_{i,j}}$ is a graded free module of rank $\sum_{j \in \mathbb{Z}} \beta_{i,j}$ ($< \infty$), and where every φ_i is degree-preserving. The *homological dimension* $\text{hd}_A(k[\Delta])$ of $k[\Delta]$ over A is the minimal h possible in (1). It is known (see, e.g., [Bru-Her, Theorem (1.3.3), Corollary (2.2.14)]) that $v - d \leq \text{hd}_A(k[\Delta]) \leq v$. A finite free resolution (1) is called *minimal* if each $\beta_{i,j}$ is smallest possible. A minimal free resolution of $k[\Delta]$ over A exists and is essentially unique. See, e.g., [Bru-Her, p. 35]. When a finite free resolution (1) is minimal with $h = \text{hd}_A(k[\Delta])$, we say that $\beta_{i,j} = \beta_{i,j}^A(k[\Delta])$ is the i_j -th *Betti number* of $k[\Delta]$ over A . Let $\beta_i = \beta_i^A(k[\Delta])$ denote the rank of the i -th free module which appears in a minimal free resolution of $k[\Delta]$ over A ; viz, $\beta_i = \sum_{j \in \mathbb{Z}} \beta_{i,j}$.

In the paper [T-H], we give a combinatorial formula to compute the Betti numbers of the Stanley-Reisner ring of the boundary complex of the cyclic d -polytope $C(v, d)$ with v vertices and show that these Betti numbers do not depend on the base field. The purpose of the present paper is to study the same problem for a stacked d -polytope $P(v, d)$ with v vertices. In the combinatorial theory of convex polytopes, the cyclic polytope appears in the upper bound theorem and the stacked polytope appears in the lower bound theorem. See, e.g., [Bay-Lee] for background information about cyclic polytopes and stacked polytopes.

§1. Stacked polytopes and Betti numbers

We refer the reader to, e.g., [Brø] for foundations on convex polytopes. Starting with a d -simplex, one can add new vertices by building shallow pyramids over facets to obtain a simplicial convex d -polytope with v vertices, called a *stacked polytope*. Recall that the boundary complex $\Delta(P)$ of a simplicial d -polytope $P \subset \mathbb{R}^N$ with the vertex set V is the simplicial complex on V of dimension $d - 1$ whose faces are those subsets σ of V such that the convex hull of σ in $\mathbb{R}^N$ is a face of P .

Our main result in this paper is to present a combinatorial formula for the computation of the Betti numbers of the Stanley-Reisner ring associated with the boundary complex of a stacked d -polytope $P(v, d)$ with v vertices.

(1.1) THEOREM. *Fix $v > d \geq 3$. Let $P(v, d)$ be a stacked d -polytope with v vertices and $\Delta(P(v, d))$ its boundary complex. Then, a minimal free*

resolution of the Stanley-Reisner ring $k[\Delta(P(v, d))] = A/I_{\Delta(P(v, d))}$ over A is of the form

$$\begin{aligned} 0 &\longrightarrow A(-v) \longrightarrow A(-v + d)^{b_{v-d-1}} \oplus A(-v + 2)^{b_1} \\ &\longrightarrow A(-v + d + 1)^{b_{v-d-2}} \oplus A(-v + 3)^{b_2} \longrightarrow \dots \\ &\longrightarrow A(-3)^{b_2} \oplus A(-d - 1)^{b_{v-d-2}} \\ &\longrightarrow A(-2)^{b_1} \oplus A(-d)^{b_{v-d-1}} \longrightarrow A \longrightarrow k[\Delta(P(v, d))] \longrightarrow 0, \end{aligned}$$

where

$$b_i = (v - d - 1) \binom{v - d - 1}{i} - \sum_{j=i}^{v-d-2} \binom{j}{i}$$

for each $1 \leq i \leq v - d - 1$.

When $d = 2$, $\Delta(P(v, 2))$ is the cycle C_v with v vertices. A minimal free resolution of $k[C_v] = A/I_{C_v}$ over A is a *pure resolution*, which is discussed in, e.g., [B-H₁] and [B-H₂].

(1.2) COROLLARY. *Every Betti number of $k[\Delta(P(v, d))] = A/I_{\Delta(P(v, d))}$ over A is independent of the base field k and of the combinatorial type of $P(v, d)$.*

§2. Proof of Theorem (1.1)

(2.1) Given a subset W of the vertex set V of a simplicial complex Δ , the subcomplex Δ_W is defined to be $\Delta_W = \{\sigma \in \Delta \mid \sigma \subset W\}$. In particular, $\Delta_V = \Delta$ and $\Delta_\emptyset = \{\emptyset\}$. Let $\tilde{H}_i(\Delta; k)$ denote the i -th reduced simplicial homology group of Δ with the coefficient field k . Note that $\tilde{H}_{-1}(\Delta; k) = 0$ if $\Delta \neq \{\emptyset\}$ and

$$\tilde{H}_i(\{\emptyset\}; k) = \begin{cases} 0 & (i \geq 0) \\ k & (i = -1). \end{cases}$$

Suppose that a graded finite free resolution (1) of $k[\Delta]$ over A is minimal with $h = \text{hd}_A(k[\Delta])$. Then, Hochster's formula [Hoc, Theorem (5.1)] guarantees that

$$\beta_{i,j} = \sum_{W \subset V, \#(W)=j} \dim_k \tilde{H}_{j-i-1}(\Delta_W; k). \quad (2)$$

Thus, in particular,

$$\beta_i^A(k[\Delta]) = \sum_{W \subset V} \dim_k \tilde{H}_{\#(W)-i-1}(\Delta_W; k).$$

(2.2) Let $P = P(v, d)$ be a stacked d -polytope with the vertex set V , $\sharp(V) = v$, $\Delta = \Delta(P)$ the boundary complex of P , and $\mathcal{F}$ a facet of P with the vertex set X . Let P' denote a stacked d -polytope with $(v + 1)$ -vertices which is obtained by building a shallow pyramid over $\mathcal{F}$ with a new vertex α , and Δ' the boundary complex of P' . Let $V' = V \cup \{\alpha\}$ be the vertex set of Δ' and W a subset of V' . We fix a base field k .

(2.2.1) LEMMA (a) *If $\alpha \notin W$ and $X \not\subset W$, then*

$$\Delta'_W = \Delta_W.$$

(b) *If $\alpha \notin W$, $W \neq V$ and $X \subset W$, then*

$$\dim_k \tilde{H}_i(\Delta'_W; k) = \begin{cases} \dim_k \tilde{H}_i(\Delta_W; k) & (i \neq d - 2) \\ \dim_k \tilde{H}_i(\Delta_W; k) + 1 & (i = d - 2). \end{cases}$$

(c) *If $\alpha \in W$ and $X \cap W \neq \emptyset$, then, for each i , we have*

$$\tilde{H}_i(\Delta'_W; k) \cong \tilde{H}_i(\Delta_{W - \{\alpha\}}; k).$$

(d) *If $\alpha \in W$, $W \neq \{\alpha\}$ and $X \cap W = \emptyset$, then*

$$\dim_k \tilde{H}_i(\Delta'_W; k) = \begin{cases} \dim_k \tilde{H}_i(\Delta_{W - \{\alpha\}}; k) & (i \neq 0) \\ \dim_k \tilde{H}_i(\Delta_{W - \{\alpha\}}; k) + 1 & (i = 0). \end{cases}$$

Proof. (a) In general, $\Delta' = (\Delta - \{X\}) \cup \{\sigma \subset V' \mid \sigma \subset X \cup \{\alpha\}, \sigma \neq X\}$. Hence, we have $\Delta'_W = \Delta_W$ if $\alpha \notin W$ and $X \not\subset W$.

(b) Let Γ denote the set of all subsets of X and set $\partial\Gamma = \Gamma - \{X\}$. Then $\Delta'_W \cup \Gamma = \Delta_W$ and $\Delta'_W \cap \Gamma = \partial\Gamma$. Since Γ is a simplicial $(d - 1)$ -ball, $\partial\Gamma$ is a simplicial $(d - 2)$ -sphere and $\tilde{H}_{d-1}(\Delta_W; k) = 0$, the required equalities follow from the reduced Mayer-Vietoris exact sequence

$$\begin{aligned} \dots &\longrightarrow \tilde{H}_i(\partial\Gamma; k) \longrightarrow \tilde{H}_i(\Gamma; k) \oplus \tilde{H}_i(\Delta'_W; k) \longrightarrow \tilde{H}_i(\Delta_W; k) \\ &\longrightarrow \tilde{H}_{i-1}(\partial\Gamma; k) \longrightarrow \tilde{H}_{i-1}(\Gamma; k) \oplus \tilde{H}_{i-1}(\Delta'_W; k) \longrightarrow \tilde{H}_{i-1}(\Delta_W; k) \\ &\longrightarrow \dots \end{aligned}$$

(c) If $X \subset W$, then the geometric realization of Δ'_W is homeomorphic to that of $\Delta_{W - \{\alpha\}}$. Thus $\tilde{H}_i(\Delta'_W; k) \cong \tilde{H}_i(\Delta_{W - \{\alpha\}}; k)$ for each i . On the other hand, if $X \cap W \neq X$, then $\Delta_{W - \{\alpha\}} \cup \Delta'_{W \cap (\{\alpha\} \cup X)} = \Delta'_W$ and $\Delta_{W - \{\alpha\}} \cap \Delta'_{W \cap (\{\alpha\} \cup X)} = \Delta_{W \cap X}$. Since both $\Delta'_{W \cap (\{\alpha\} \cup X)}$ and $\Delta_{W \cap X}$ are contractible, again the reduced Mayer-Vietoris exact sequence guarantees the desired equalities.

(d) Since Δ'_W is the disjoint union of $\Delta_{W-\{\alpha\}}$ and one point $\{\alpha\}$, we immediately have the required equalities. Q. E. D.

(2.2.2) COROLLARY. Let $\Delta = \Delta(P)$ denote the boundary complex of a stacked d -polytope $P = P(v, d)$ with the vertex set V , $\sharp(V) = v$. Then, for every non-empty subset W of V with $W \neq V$ and for each $i \neq 0, d-2$, we have

$$\tilde{H}_i(\Delta_W; k) = 0.$$

Proof. If $v = d + 1$, i.e., P is a d -simplex, then Δ_W is contractible. Hence, $\tilde{H}_i(\Delta_W; k) = 0$ for each i . We now work with the same situation as in the above Lemma (2.2.1) and suppose that $\tilde{H}_i(\Delta_W; k) = 0$ for every non-empty subset W of V with $W \neq V$ and for each $i \neq 0, d-2$. Let W be a non-empty subset of V' with $W \neq V'$. If $W = V' - \{\alpha\}$, then Δ'_W is a simplicial $(d-1)$ -ball. Hence, $\tilde{H}_i(\Delta_W; k) = 0$ for each i . Moreover, if $W = \{\alpha\}$, then $\tilde{H}_i(\Delta_W; k) = 0$ for each i . On the other hand, if W is a non-empty subset of V' with $W \neq V'$ such that $W \neq V$ and $W \neq \{\alpha\}$, and if $i \neq 0, d-2$, then $\dim_k \tilde{H}_i(\Delta'_W; k) = \dim_k \tilde{H}_i(\Delta_{W-\{\alpha\}}; k)$ by Lemma (2.2.1). Hence, $\tilde{H}_i(\Delta_W; k) = 0$ as desired. Q. E. D.

(2.3) Fix $d \geq 3$, and keep the notation P, P', Δ and Δ' in (2.2). Let β_i be the i_j -th Betti number of $k[\Delta]$ and β'_i the i_j -th Betti number of $k[\Delta']$.

(2.3.1) LEMMA. For each $i \geq 1$ we have

$$\beta'_{i+1} = \beta_{i+1} + \beta_{i-1} + \binom{v-d}{i}.$$

Proof. By virtue of Eq. (2) as well as Lemma (2.2.1), we have

$$\begin{aligned} \beta'_{i+1} &= \sum_{W \subset V', \sharp(W)=i+1} \dim_k \tilde{H}_0(\Delta'_W; k) \\ &= \sum_{\alpha \notin W \subset V', \sharp(W)=i+1} \dim_k \tilde{H}_0(\Delta'_W; k) + \sum_{\alpha \in W \subset V', \sharp(W)=i+1} \dim_k \tilde{H}_0(\Delta'_W; k) \\ &= \sum_{W \subset V, \sharp(W)=i+1} \dim_k \tilde{H}_0(\Delta_W; k) + \sum_{W \subset V, \sharp(W)=i} \dim_k \tilde{H}_0(\Delta_W; k) + \binom{v-d}{i} \\ &= \beta_{i+1} + \beta_{i-1} + \binom{v-d}{i} \end{aligned}$$

as desired. Q. E. D.

(2.3.2) COROLLARY. Let $\Delta = \Delta(P)$ denote the boundary complex of a stacked d -polytope $P = P(v, d)$, $d \geq 3$, with v vertices. Then, for each $1 \leq i \leq v - d - 1$, the i_{i+1} -th Betti number of $k[\Delta] = A/I_\Delta$ over A is

$$\beta_{i_{i+1}}^A(k[\Delta]) = (v - d - 1) \binom{v - d - 1}{i} - \sum_{j=i}^{v-d-2} \binom{j}{i}.$$

Proof. In this proof, we set $\binom{a}{0} = 0$ for every integer $a \geq 0$. Thanks to Lemma (2.3.1), we have

$$\begin{aligned} \beta_{i_{i+1}}^A(k[\Delta]) &= (v - d - 2) \binom{v - d - 2}{i} - \sum_{j=i}^{v-d-3} \binom{j}{i} \\ &\quad + (v - d - 2) \binom{v - d - 2}{i-1} - \sum_{j=i-1}^{v-d-3} \binom{j}{i-1} + \binom{v - d - 1}{i} \\ &= (v - d - 2) \left(\binom{v - d - 2}{i} + \binom{v - d - 2}{i-1} \right) + \binom{v - d - 1}{i} \\ &\quad - \left(\binom{i-1}{i-1} + \sum_{j=i}^{v-d-3} \left(\binom{j}{i} + \binom{j}{i-1} \right) \right) \\ &= (v - d - 2) \binom{v - d - 1}{i} + \binom{v - d - 1}{i} \\ &\quad - \left(\binom{i}{i} + \sum_{j=i}^{v-d-3} \binom{j+1}{i} \right) \\ &= (v - d - 1) \binom{v - d - 1}{i} - \sum_{j=i}^{v-d-2} \binom{j}{i} \end{aligned}$$

as required.

Q. E. D.

(2.4) We are now in the position to give a proof of Theorem (1.1). Since $\Delta = \Delta(P(v, d))$ is a simplicial $(d-1)$ -sphere with v vertices, we know that the homological dimension of $k[\Delta] = A/I_\Delta$ over A is $\text{hd}_A(k[\Delta]) = v - d$ and that $\beta_{i_j}^A(k[\Delta]) = \beta_{v-d-i_{v-j}}^A(k[\Delta])$ for every i and j . By Corollary (2.2.2), we have $\beta_{i_j}^A(k[\Delta]) = 0$ for each $1 \leq i \leq v - d - 1$ and for each $j \neq i + 1, i + d - 1$. On the other hand, Corollary (2.3.2) enables us to compute $b_i = \beta_{i_{i+1}}^A(k[\Delta]) = \beta_{v-d-i_{v-i-1}}^A(k[\Delta])$ for each $1 \leq i \leq v - d - 1$. Hence, we obtain a desired minimal free resolution of $k[\Delta]$ over A .

§3. Unimodality of Betti number sequences

Let β_i , $0 \leq i \leq v - d$, denote the rank of the i -th free module which appears in a minimal free resolution of $k[\Delta(P(v, d))] = A/I_{\Delta(P(v, d))}$ over A . Then, $\beta_0 = \beta_{v-d} = 1$ and $\beta_i = b_i + b_{v-d-i}$ for each $1 \leq i \leq v - d - 1$ with the notation of Theorem (1.1). The sequence $(\beta_0, \beta_1, \dots, \beta_{v-d})$ is called the *Betti number sequence* of $k[\Delta(P(v, d))]$ over A . This sequence is *symmetric*, i.e., $\beta_i = \beta_{v-d-i}$ for every $0 \leq i \leq v - d$. We now show that the symmetric sequence $(\beta_0, \beta_1, \dots, \beta_{v-d})$ is *unimodal*, i.e., $\beta_0 \leq \beta_1 \leq \dots \leq \beta_{\lfloor (v-d)/2 \rfloor}$.

The following Lemma (3.1) follows from a simple combinatorial argument based on Lemma (2.3.1).

(3.1) LEMMA. (a) If $v - d$ is even, then

$$b_{\frac{v-d}{2}} \geq b_{\frac{v-d}{2}-1} \geq b_{\frac{v-d}{2}+1} \geq \dots \geq b_1 \geq b_{v-d-1}.$$

(b) If $v - d$ is odd, then

$$b_{\frac{v-d-1}{2}} \geq b_{\frac{v-d-1}{2}+1} \geq b_{\frac{v-d-1}{2}-1} \geq \dots \geq b_1 \geq b_{v-d-1}.$$

(3.2) COROLLARY. Fix $v > d \geq 3$. Let $P = P(v, d)$ a stacked d -polytope with v vertices and $\Delta = \Delta(P)$ its boundary complex. Then, the Betti number sequence $(\beta_0, \beta_1, \dots, \beta_{v-d})$ of $k[\Delta] = A/I_{\Delta}$ over A is unimodal.

Proof. By Lemma (3.1), we have

$$1 \leq b_1 \leq b_2 \leq \dots \leq b_{\lfloor \frac{v-d}{2} \rfloor} \geq b_{\lfloor \frac{v-d}{2} \rfloor + 1} \geq \dots \geq b_{v-d-1} \geq 1.$$

Hence, the Betti number sequence $(\beta_0, \beta_1, \dots, \beta_{v-d})$ of $k[\Delta] = A/I_{\Delta}$ over A is unimodal. Q. E. D.

References

- [Bay-Lee] M. Bayer and C. Lee, *Combinatorial aspects of convex polytopes*, in "Handbook of Convex Geometry" (P. Gruber and J. Wills, eds.), North-Holland, Amsterdam / New York / Tokyo, 1993, pp. 485–534.
- [Brø] A. Brøndsted, "An Introduction to Convex Polytopes," Graduate Texts in Math., Vol. 90, Springer-Verlag, Berlin / New York / Tokyo, 1983.

- [Bru-Her] W. Bruns and J. Herzog, "Cohen-Macaulay Rings," Cambridge University Press, Cambridge / New York / Sydney, 1993.
- [B-H₁] W. Bruns and T. Hibi, *Cohen-Macaulay partially ordered sets with pure resolutions*, preprint.
- [B-H₂] W. Bruns and T. Hibi, *Stanley-Reisner rings with pure resolutions*, Comm. in Algebra, to appear.
- [H] T. Hibi, "Algebraic Combinatorics on Convex Polytopes," Carslaw Publications, Glebe, N.S.W., Australia, 1992.
- [Hoc] M. Hochster, *Cohen-Macaulay rings, combinatorics, and simplicial complexes*, in "Ring Theory II" (B. R. McDonald and R. Morris, eds.), Lect. Notes in Pure and Appl. Math., No. 26, Dekker, New York, 1977, pp.171 - 223.
- [Sta] R. P. Stanley, "Combinatorics and Commutative Algebra," Birkhäuser, Boston / Basel / Stuttgart, 1983.
- [T-H] N. Terai and T. Hibi, *Computation of Betti numbers of monomial ideals associated with cyclic polytopes*, Discrete Comput. Geom., to appear.

DEPARTMENT OF MATHEMATICS
 NAGANO NATIONAL COLLEGE OF TECHNOLOGY
 NAGANO 381, JAPAN
 E-mail address: terai@cc.nagano-nct.ac.jp

DEPARTMENT OF MATHEMATICS
 HOKKAIDO UNIVERSITY
 KITA-KU, SAPPORO 060, JAPAN
 E-mail address: hibi@math.hokudai.ac.jp