HOKKAIDO UNIVERSITY

Title	Finite free resolutions and 1-skeletons of simplicial(d - l)-spheres
Author(s)	Terai, N.; Hibi, T.
Citation	Hokkaido University Preprint Series in Mathematics, 279, 1-3
Issue Date	1995-01-01
DOI	https://doi.org/10.14943/83426
Doc URL	https://hdl.handle.net/2115/69030
Type	departmental bulletin paper
File Information	re279.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Finite free resolutions and 1-skeletons of simplicial$(d-1)$-spheres

N. Terai and T. Hibi

Series ♯279. January 1995

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

253 T. Nishimori, Some remarks in a qualitative theory of similarity pseudogroups, 19 pages. 1994.

254 T. Suwa, Residues of complex analytic foliations relative to singular invariant subvarieties, 15 pages. 1994.

255 T. Tsujishita, On Triple Mutual Information, 7 pages. 1994.

256 T. Tsujishita, Construction of Universal Modal World based on Hyperset Theory, 15 pages. 1994.

257 A. Arai, Trace Formulas, a Golden-Thompson Inequality and Classical Limit in Boson Fock Space, 35 pages. 1994.

258 Y-G. Chen, Y. Giga, T. Hitaka and M. Honma, A Stable Difference Scheme for Computing Motion of Level Surfaces by the Mean Curvature, 18 pages. 1994.

259 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 7 pages. 1994.

260 T. Mikami, Copula fields and its applications, 14 pages. 1994.

261 A. Inoue, An Abel-Tauber theorem for Fourier sine transforms, 6 pages. 1994.

262 N. Kawazumi, Homology of hyperelliptic mapping class groups for surfaces, 13 pages. 1994.

263 Y. Giga, M. E. Gurtin, A comparison theorem for crystalline evolution in the plane, 14 pages. 1994.

264 J. Wierzbicki, On Commutativity of Diagrams of Type II_1 Factors, 26 pages. 1994.

265 N. Hayashi, T. Ozawa, Schrödinger Equations with nonlinearity of integral type, 12 pages. 1994.

266 T. Ozawa, On the resonance equations of long and short waves, 8 pages. 1994.

267 T. Mikami, A sufficient condition for the uniqueness of solutions to a class of integro-differential equations, 9 pages. 1994.

268 Y. Giga, Evolving curves with boundary conditions, 10 pages. 1994.

269 A. Arai, Operator-theoretical analysis of representation of a supersymmetry algebra in Hilbert space, 12 pages. 1994.

270 A. Arai, Gauge theory on a non-simply-connected domain and representations of canonical commutation relations, 18 pages. 1994.

271 S. Jimbo, Y. Morita and J. Zhai, Ginzburg landau equation and stable steady state solutions in a non-trivial domain, 17 pages. 1994.

272 S. Izumiya, A. Takiyama, A time-like surface in Minkowski 3-space which contains light-like lines, 7 pages. 1994.

273 K. Tsutaya, Global existence of small amplitude solutions for the Klein-Gordon-Zakharov equations, 11 pages. 1994.

274 H. Kubo, On the critical decay and power for semilinear wave equations in odd space dimensions, 22 pages. 1994.

275 N. Terai, T. Hibi, Alexander duality theorem and second Betti numbers of Stanley-Reisner rings, 2 pages. 1995.

276 N. Terai, T. Hibi, Stanley-Reisner rings whose Betti numbers are independent of the base field, 12 pages. 1995.

277 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with cyclic polytopes, 11 pages. 1995.

278 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with stacked polytopes, 8 pages. 1995.

Finite free resolutions and 1-skeletons of simplicial $(d-1)$-spheres

Naoki Terai Takayuki Hibi

A simplicial complex Δ on the vertex set V is a collection of subsets of V such that (i) $\{x\} \in \Delta$ for every $x \in V$ and (ii) $\sigma \in \Delta$, $\tau \subset \sigma \Rightarrow \tau \in \Delta$. The 1-*skeleton* $\Delta^{(1)}$ of Δ is a graph on V whose edges are all 2-element subsets $\{x, y\}$ of V with $\{x, y\} \in \Delta$. We say that a simplicial complex Δ is a *simplicial* $(d-1)$-*sphere* if the geometric realization of Δ is homeomorphic to the $(d-1)$-sphere.

The purpose of the present paper is to give a ring-theoretical short proof of the following result, which was first proved by Barnette [1].

THEOREM. *The* 1-*skeleton of a simplicial* $(d-1)$-*sphere is* d-*connected.*

Given a subset W of the vertex set $V = \{x_1, x_2, \ldots, x_v\}$ of a simplicial complex Δ, the restriction of Δ to W is the subcomplex $\Delta_W = \{\sigma \in \Delta \mid \sigma \subset W\}$ of Δ. Let $\tilde{H}_i(\Delta; k)$ denote the i-th reduced simplicial homology group of Δ with the coefficient field k. Note that $\tilde{H}_{-1}(\Delta; k) = 0$ if $\Delta \neq \{\emptyset\}$, $\tilde{H}_{-1}(\{\emptyset\}; k) \cong k$, and $\tilde{H}_i(\{\emptyset\}; k) = 0$ for each $i \geq 0$.

Let $A = k[x_1, x_2, \ldots, x_v]$ be the polynomial ring in v-variables over a field k. Here, we identify each $x_i \in V$ with the indeterminate x_i of A. Define I_Δ to be the ideal of A which is generated by square-free monomials $x_{i_1} x_{i_2} \cdots x_{i_r}$, $1 \leq i_1 < i_2 < \cdots < i_r \leq v$, with $\{x_{i_1}, x_{i_2}, \cdots, x_{i_r}\} \notin \Delta$. We say that the quotient algebra $k[\Delta] := A/I_\Delta$ is the *Stanley–Reisner ring* of Δ over k. In what follows, we consider A to be the graded algebra $A = \oplus_{n \geq 0} A_n$ with the standard grading, i.e., each $\deg x_i = 1$, and may regard $k[\Delta] = \oplus_{n \geq 0}(k[\Delta])_n$ as a graded module over A with the quotient grading. Let $\mathbf{Z}$ denote the set of integers. We write $A(j)$, $j \in \mathbf{Z}$, for the graded module $A(j) = \oplus_{n \in \mathbf{Z}}[A(j)]_n$ over A with $[A(j)]_n := A_{n+j}$.

We study a graded minimal free resolution

$$0 \longrightarrow \bigoplus_{j \in \mathbf{Z}} A(-j)^{\beta_{h_j}} \xrightarrow{\varphi_h} \cdots \xrightarrow{\varphi_2} \bigoplus_{j \in \mathbf{Z}} A(-j)^{\beta_{1_j}} \xrightarrow{\varphi_1} A \xrightarrow{\varphi_0} k[\Delta] \longrightarrow 0$$

of $k[\Delta]$ over A. Here $h = \mathrm{hd}_A(k[\Delta])$ is the homological dimension of $k[\Delta]$ over A. It is known [3, Theorem (5.1)] that

$$\beta_{i_j} = \sum_{W \subset V,\ \sharp(W)=j} \dim_k \tilde{H}_{j-i-1}(\Delta_W; k), \tag{1}$$

where $\sharp(W)$ is the cardinality of a finite set W.

Now, suppose that Δ is a simplicial $(d-1)$-sphere on the vertex set V with $\sharp(V) = v$. Then $k[\Delta]$ is Cohen–Macaulay, i.e., $\mathrm{hd}_A(k[\Delta]) = v - d$. Moreover, since $k[\Delta]$ is Gorenstein, $\beta_{h_j} = 0$ if $j \neq v$ and $\beta_{h_v} = 1$. In particular, $\beta_{(v-d)_{v-(d-1)}} = 0$. Hence, it follows from Eq. (1) that $\tilde{H}_0(\Delta_W; k) = 0$ for every subset W of V with $\sharp(W) = v - (d-1)$. Thus $\mid \Delta_{V-W} \mid$ is connected for every subset W of V with $\sharp(W) = d - 1$. Hence, the 1-skeleton $\Delta^{(1)}$ of Δ is d-connected.

The above ring-theoretical technique enables us to show that the 1-skeleton of a level complex Δ ([3], [6]) of dimension $d-1$ with v vertices is d-connected if $\sharp(\{\sigma \in \Delta \mid \sharp(\sigma) = d\}) \neq v - d + 1$.

We refer the reader to [2], [4], [5] and [7] for the detailed information about algebra and combinatorics on Stanley–Reisner rings.

References

[1] D. Barnette, *Graph theorems for manifolds*, Israel J. of Math. **16** (1973), 62 – 72.

[2] W. Bruns and J. Herzog, "Cohen–Macaulay Rings," Cambridge University Press, Cambridge / New York / Sydney, 1993.

[3] T. Hibi, *Level rings and algebras with straightening laws*, J. Algebra **117** (1988), 343 –362.

[4] T. Hibi, "Algebraic Combinatorics on Convex Polytopes," Carslaw Publications, Glebe, N.S.W., Australia, 1992.

[5] M. Hochster, *Cohen–Macaulay rings, combinatorics, and simplicial complexes*, in "Ring Theory II" (B. R. McDonald and R. Morris, eds.), Lect. Notes in Pure and Appl. Math., No. 26, Dekker, New York, 1977, pp.171 – 223.

[6] R. P. Stanley, *Cohen–Macaulay complexes*, in "Higher Combinatorics" (M. Aigner, Ed.), Reidel, Dordrecht / Boston, 1977, pp. 51 – 62.

[7] R. P. Stanley, "Combinatorics and Commutative Algebra," Birkhäuser, Boston / Basel / Stuttgart, 1983.

Department of Mathematics
Nagano National College of Technology
Nagano 381, Japan
E-mail address: terai@cc.nagano-nct.ac.jp

Department of Mathematics
Hokkaido University
Kita-ku, Sapporo 060, Japan
E-mail address: hibi@math.hokudai.ac.jp