



HOKKAIDO UNIVERSITY

Title	Strong coupling limit of the zero-energy-state density of the Dirac-Weyl operator with a singular vector potential
Author(s)	Arai, A.
Citation	Hokkaido University Preprint Series in Mathematics, 286, 2-8
Issue Date	1995-03-01
DOI	https://doi.org/10.14943/83433
Doc URL	https://hdl.handle.net/2115/69037
Type	departmental bulletin paper
File Information	re286.pdf



**Strong Coupling Limit of the Zero-
Energy-State Density of the Dirac
-Weyl Operator with a Singular
Vector Potential**

Asao Arai

Series #286. March 1995

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- # 261 A. Inoue, An Abel-Tauber theorem for Fourier sine transforms, 6 pages. 1994.
- # 262 N. Kawazumi, Homology of hyperelliptic mapping class groups for surfaces, 13 pages. 1994.
- # 263 Y. Giga, M. E. Gurtin, A comparison theorem for crystalline evolution in the plane, 14 pages. 1994.
- # 264 J. Wierzbicki, On Commutativity of Diagrams of Type II_1 Factors, 26 pages. 1994.
- # 265 N. Hayashi, T. Ozawa, Schrödinger Equations with nonlinearity of integral type, 12 pages. 1994.
- # 266 T. Ozawa, On the resonance equations of long and short waves, 8 pages. 1994.
- # 267 T. Mikami, A sufficient condition for the uniqueness of solutions to a class of integro-differential equations, 9 pages. 1994.
- # 268 Y. Giga, Evolving curves with boundary conditions, 10 pages. 1994.
- # 269 A. Arai, Operator-theoretical analysis of representation of a supersymmetry algebra in Hilbert space, 12 pages. 1994.
- # 270 A. Arai, Gauge theory on a non-simply-connected domain and representations of canonical commutation relations, 18 pages. 1994.
- # 271 S. Jimbo, Y. Morita and J. Zhai, Ginzburg landau equation and stable steady state solutions in a non-trivial domain, 17 pages. 1994.
- # 272 S. Izumiya, A. Takiyama, A time-like surface in Minkowski 3-space which contains light-like lines, 7 pages. 1994.
- # 273 K. Tsutaya, Global existence of small amplitude solutions for the Klein-Gordon-Zakharov equations, 11 pages. 1994.
- # 274 H. Kubo, On the critical decay and power for semilinear wave equations in odd space dimensions, 22 pages. 1994.
- # 275 N. Terai, T. Hibi, Alexander duality theorem and second Betti numbers of Stanley-Reisner rings, 2 pages. 1995.
- # 276 N. Terai, T. Hibi, Stanley-Reisner rings whose Betti numbers are independent of the base field, 12 pages. 1995.
- # 277 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with cyclic polytopes, 11 pages. 1995.
- # 278 N. Terai, T. Hibi, Computation of Betti numbers of monomial ideals associated with stacked polytopes, 8 pages. 1995.
- # 279 N. Terai, T. Hibi, Finite free resolutions and 1-skeletons of simplicial $(d - 1)$ -spheres, 3 pages. 1995.
- # 280 N. Terai, T. Hibi, Monomial ideals and minimal non-faces of Cohen-Macaulay complexes, 6 pages. 1995.
- # 281 A. Arai, N. Tominaga, Analysis of a family of strongly commuting self-adjoint operators with applications to perturbed d'Alembertians and the external field problem in quantum field theory, 44 pages. 1995.
- # 282 T. Mikami, Asymptotic behavior of the first exit time of randomly perturbed dynamical systems with a repulsive equilibrium point, 29 pages. 1995.
- # 283 K. Iwata, J. Schäfer, Markov property and cokernels of local operators, 17 pages. 1995.
- # 284 T. Nakazi, M. Yamada, Riesz's Functions In Weighted Hardy And Bergman Spaces, 20 pages. 1995.
- # 285 K. Hidano, K. Tsutaya, Scattering theory for nonlinear wave equations in the invariant Sobolev space, 32 pages. 1995.

Strong Coupling Limit of the Zero-Energy-State Density of the Dirac-Weyl Operator with a Singular Vector Potential

ASAO ARAI*

Department of Mathematics, Hokkaido University, Sapporo 060, JAPAN

Abstract. For a self-adjoint extension of the Dirac-Weyl operator with a singular vector potential, the strong coupling limit of the zero-energy-state density (ZESD) is computed. The result shows that a limit theorem of L.Erdős (Lett.Math.Phys.29:219–240,1993) concerning the ZESD of the Dirac-Weyl operator with a “regular” magnetic field does not hold in the present singular case.

1. Introduction

In a paper [3], Erdős considered the two-dimensional Pauli operator H with a “regular” magnetic field B and showed that the naturally rescaled ground-state density of H with respect to the coupling constant L converges to B as $L \rightarrow \infty$.

The ground states of the Pauli operator considered in [3] are in fact zero-energy states. Hence they are also the zero-energy states of the corresponding Dirac-Weyl operator. Thus the Erdős’ limit theorem mentioned above can also be regarded as one concerning the zero-energy-state density (ZESD) of the Dirac-Weyl operator.

It is interesting to examine if the Erdős’ limit theorem holds for wider classes of magnetic fields. With this motivation, we consider in this paper the strong coupling limit of the ZESD of the Dirac-Weyl operator with a singular vector potential such that the magnetic field is a distribution concentrated on two points in $\mathbf{R}^2$ and show that the Erdős’ limit theorem does not hold in this singular case.

Mathematical aspects of a quantum system with a magnetic field concentrated on isolated points in $\mathbf{R}^2$ were discussed in [1,2]. We begin with a brief review of some results obtained in these papers.

2. The Dirac-Weyl Operator with a Singular Vector Potential and the ZESD

We consider a quantum system of a charged particle with charge $q \in \mathbf{R} \setminus \{0\}$ moving in the plane $\mathbf{R}^2$ under the influence of a perpendicular magnetic field $B(\mathbf{r})$ ($\mathbf{r} = (x, y) \in \mathbf{R}^2$) concentrated on two points $\mathbf{a}_\nu = (a_{\nu 1}, a_{\nu 2}) \in \mathbf{R}^2, \nu = 1, 2$. A general form of such a magnetic field is given by

$$B(\mathbf{r}) = \sum_{\nu=1}^2 \sum_{0 \leq \alpha + \beta \leq \ell} C_{\alpha, \beta}^{(\nu)} D_x^\alpha D_y^\beta \delta(\mathbf{r} - \mathbf{a}_\nu),$$

*Supported by Grant-In-Aid 06640188 for science research from the Ministry of Education, Japan.

where ℓ and $C_{\alpha,\beta}^{(\nu)}$ are a nonnegative integer and a real constant, respectively, D_x and D_y denote the distributional partial differential operators in x and y , respectively, and $\delta(\mathbf{r})$ is the Dirac delta distribution on $\mathbf{R}^2$. A vector (gauge) potential of the magnetic field is a vector-valued distribution $\mathbf{A} = (A_1, A_2)$ satisfying $B = D_x A_2 - D_y A_1$. One of such distributions $\mathbf{A}$ is given by

$$A_1(\mathbf{r}) = - \sum_{\nu=1}^2 \sum_{0 \leq \alpha+\beta \leq \ell} \frac{C_{\alpha,\beta}^{(\nu)}}{2\pi} D_x^\alpha D_y^\beta \left(\frac{y - a_{\nu 2}}{|\mathbf{r} - \mathbf{a}_\nu|^2} \right),$$

$$A_2(\mathbf{r}) = \sum_{\nu=1}^2 \sum_{0 \leq \alpha+\beta \leq \ell} \frac{C_{\alpha,\beta}^{(\nu)}}{2\pi} D_x^\alpha D_y^\beta \left(\frac{x - a_{\nu 1}}{|\mathbf{r} - \mathbf{a}_\nu|^2} \right).$$

We use a system of units where the light speed c and $\hbar$ (the Planck constant divided by 2π) are equal to 1. The kinetic momentum operator $\mathbf{P} = (P_1, P_2)$ with the vector potential $\mathbf{A}$ is defined by

$$P_1 = -iD_x - qA_1, \quad P_2 = -iD_y - qA_2,$$

in $L^2(\mathbf{R}^2)$ with $D(P_1) = D(D_x) \cap D(A_1)$ and $D(P_2) = D(D_y) \cap D(A_2)$ (we denote by $D(\cdot)$ operator domain). Let

$$M = \mathbf{R}^2 \setminus \{\mathbf{a}_1, \mathbf{a}_2\}.$$

It was shown in [1] that each P_j is essentially self-adjoint on $C_0^\infty(M)$.

The Dirac-Weyl operator with the vector potential $\mathbf{A}$ is given by

$$\mathcal{D} := \sigma_1 P_1 + \sigma_2 P_2 = \begin{pmatrix} 0 & D_- \\ D_+ & 0 \end{pmatrix}$$

acting in $L^2(\mathbf{R}^2; \mathbf{C}^2)$, where $\sigma_j, j = 1, 2$, are the first two of the Pauli matrices and

$$D_\pm := P_1 \pm iP_2.$$

It is an interesting and important problem to discuss essential self-adjointness of $\mathcal{D}$, which is non-trivial because of the singularity of the vector potential. But here we only define a self-adjoint extension of the “minimal” Dirac-Weyl operator $\mathcal{D}_{\min} := \mathcal{D} \upharpoonright C_0^\infty(M)$ [2].

Let $D_{-, \min}$ be the closure of $D_- \upharpoonright C_0^\infty(M)$. Then the operator

$$\tilde{\mathcal{D}} = \begin{pmatrix} 0 & D_{-, \min} \\ D_{-, \min}^* & 0 \end{pmatrix}$$

with $D(\tilde{\mathcal{D}}) = D(D_{-, \min}^*) \oplus D(D_{-, \min})$ is a self-adjoint extension of $\mathcal{D}_{\min}$ (see Section IV-B in [2], where $\tilde{\mathcal{D}}$ is denoted $Q_{\min}^{(2)}$). The set of zero-energy states of $\tilde{\mathcal{D}}$ is given by $\ker \tilde{\mathcal{D}} := \{\psi \in D(\tilde{\mathcal{D}}) | \tilde{\mathcal{D}}\psi = 0\}$.

The constants

$$\gamma_\nu := C_{0,0}^{(\nu)}, \quad \nu = 1, 2,$$

physically mean the magnetic fluxes passing through the points $\mathbf{a}_1, \mathbf{a}_2$, respectively. For simplicity, we assume that

$$q > 0, \quad \gamma_\nu > 0, \quad \nu = 1, 2,$$

and set

$$\varepsilon_\nu(q) = \frac{q\gamma_\nu}{2\pi} - \left[\frac{q\gamma_\nu}{2\pi} \right], \quad \nu = 1, 2,$$

where $[x]$ denotes the largest integer less than or equal to x . It follows that $0 \leq \varepsilon_\nu(q) < 1$.

For each point $\mathbf{r} = (x, y) \in \mathbf{R}^2$, we denote by $z = x + iy$ the complex number corresponding to $\mathbf{r}$. Let

$$a_\nu := a_{\nu 1} + ia_{\nu 2}$$

and

$$F(z) = -\frac{1}{2\pi} \sum_{\nu=1}^2 \sum_{k=1}^{\ell} \frac{C_k^{(\nu)}}{k(z - a_\nu)^k},$$

with $C_k^{(\nu)} = (-1)^k k! \sum_{\alpha=0}^k i^{k-\alpha} C_{\alpha, k-\alpha}^{(\nu)}$. The function F is related to $\mathbf{A}$ [2]:

$$A_2(\mathbf{r}) + iA_1(\mathbf{r}) = \frac{\partial F(z)}{\partial z} + \frac{1}{2\pi} \sum_{\nu=1}^2 \frac{\gamma_\nu}{z - a_\nu}.$$

Let

$$\Omega_q(\mathbf{r}) = \left(\prod_{\nu=1}^2 |z - a_\nu|^{-q\gamma_\nu/2\pi} (z - a_\nu)^{[q\gamma_\nu/2\pi]} \right) e^{iq \operatorname{Im} F(z)}.$$

LEMMA 2.1. $\dim \ker \tilde{\mathcal{D}} = 1$ if and only if

$$\varepsilon_1(q) + \varepsilon_2(q) > 1. \quad (2.1)$$

In that case, the zero-energy state of $\tilde{\mathcal{D}}$ is given by $\begin{pmatrix} \Omega_q(\mathbf{r}) \\ 0 \end{pmatrix}$ (up to constant multiples).

Proof. This is a special case of Theorem 4.7 in [2] (the case $n = 2$). ■

Remark. Under condition (2.1), $q\gamma_1/2\pi$ and $q\gamma_2/2\pi$ are not integers, i.e., the magnetic flux is not locally quantized (see [1,2]).

In what follows, we consider only the case where (2.1) is satisfied. Under this condition, the ZESD of $\tilde{\mathcal{D}}$ is given by

$$\varrho_q(\mathbf{r}) := \frac{|\Omega_q(\mathbf{r})|^2}{\|\Omega_q\|_{L^2(\mathbf{R}^2)}^2} = \frac{|\mathbf{r} - \mathbf{a}_1|^{-2\varepsilon_1(q)} |\mathbf{r} - \mathbf{a}_2|^{-2\varepsilon_2(q)}}{\int_{\mathbf{R}^2} |\mathbf{r} - \mathbf{a}_1|^{-2\varepsilon_1(q)} |\mathbf{r} - \mathbf{a}_2|^{-2\varepsilon_2(q)} d\mathbf{r}}. \quad (2.2)$$

3. Strong Coupling Limit of the ZESD

We now consider the strong coupling limit $q \rightarrow \infty$ of the ZESD $\varrho_q(\mathbf{r})$ given by (2.2). Since the magnetic field B is not a function, but a distribution, it is natural to take the strong coupling limit in the distribution sense. For this purpose, for each $\mu, \lambda \in (0, 1)$ satisfying $\mu + \lambda > 1$, we define a functional $\Phi_{\mu, \lambda}$ in $L^\infty(\mathbf{R}^2)^*$, the topological dual space of $L^\infty(\mathbf{R}^2)$, by

$$\Phi_{\mu, \lambda}(f) = \int_{\mathbf{R}^2} \frac{f(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2\mu} |\mathbf{r} - \mathbf{a}_2|^{2\lambda}} d\mathbf{r}, \quad f \in L^\infty(\mathbf{R}^2).$$

In terms of this functional, the zero-energy-state functional

$$\varrho_q(f) := \int_{\mathbf{R}^2} \varrho_q(\mathbf{r}) f(\mathbf{r}) d\mathbf{r}$$

is written

$$\varrho_q(f) = \frac{\Phi_{\varepsilon_1(q), \varepsilon_2(q)}(f)}{\Phi_{\varepsilon_1(q), \varepsilon_2(q)}(1)}. \quad (3.1)$$

As (2.2) (or (3.1)) shows, the q -dependence of ϱ_q comes only from the factors $\varepsilon_\nu(q)$, $\nu = 1, 2$. It is obvious that $\lim_{q \rightarrow \infty} \varepsilon_\nu(q)$ does not exist. But, for suitable monotone increasing sequences $\{q_n\}_{n=1}^\infty$ of positive numbers satisfying

$$q_n \rightarrow \infty \ (n \rightarrow \infty), \quad \varepsilon_1(q_n) + \varepsilon_2(q_n) > 1, \quad n \geq 1, \quad (3.2)$$

the limits

$$\lambda_\nu := \lim_{n \rightarrow \infty} \varepsilon_\nu(q_n), \quad \nu = 1, 2, \quad (3.3)$$

may exist, depending on the choice of $\{q_n\}_{n=1}^\infty$. For this reason, we discuss the strong coupling limit of the ZESD according to the magnitude of λ_ν , $\nu = 1, 2$.

We denote by $\mathfrak{B}(\mathbf{R}^2)$ the set of bounded continuous functions on $\mathbf{R}^2$. Limiting behaviors of the functional $\Phi_{\mu, \lambda}$ in μ and λ are given in the following lemma.

LEMMA 3.1.

(i) Let $\mu_0, \lambda_0 \in (0, 1)$ such that $\mu_0 + \lambda_0 > 1$. Then, for all $f \in L^\infty(\mathbf{R}^2)$,

$$\lim_{\mu \rightarrow \mu_0, \lambda \rightarrow \lambda_0} \Phi_{\mu, \lambda}(f) = \Phi_{\mu_0, \lambda_0}(f). \quad (3.4)$$

(ii) Let $\lambda_0 \in (0, 1)$. Then, for all $f \in \mathfrak{B}(\mathbf{R}^2)$,

$$\lim_{\mu \rightarrow 1, \lambda \rightarrow \lambda_0} (1 - \mu) \Phi_{\mu, \lambda}(f) = \frac{\pi f(\mathbf{a}_1)}{|\mathbf{a}_1 - \mathbf{a}_2|^{2\lambda_0}}. \quad (3.5)$$

(iii) Let $\tau > 0$. Then, for all $f \in \mathfrak{B}(\mathbf{R}^2)$,

$$\lim_{\mu \rightarrow 1} (1 - \mu) \Phi_{\mu, (\tau - 1 + \mu)/\tau}(f) = \frac{\pi}{|\mathbf{a}_1 - \mathbf{a}_2|^2} (f(\mathbf{a}_1) + \tau f(\mathbf{a}_2)). \quad (3.6)$$

Proof. (i) For $\delta > 0$, we define

$$B_\delta(\mathbf{a}_\nu) = \{\mathbf{r} \in \mathbf{R}^2 \mid 0 \leq |\mathbf{r} - \mathbf{a}_\nu| \leq \delta\}.$$

Without loss of generality, we can assume that $\mu_0 - \varepsilon < \mu < \mu_0 + \varepsilon < 1$ and

$$\lambda_0 - \varepsilon < \lambda < \lambda_0 + \varepsilon < 1, \quad (3.7)$$

where $\varepsilon > 0$ is a constant such that $\mu_0 + \lambda_0 - 2\varepsilon > 1$. Then we have for $\mathbf{r} \in M$

$$\begin{aligned} \frac{1}{|\mathbf{r} - \mathbf{a}_1|^{2\mu}} &\leq \frac{\chi_{B_1(\mathbf{a}_1)}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\mu_0+\varepsilon)}} + \frac{\chi_{B_1(\mathbf{a}_1)^c}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\mu_0-\varepsilon)}}, \\ \frac{1}{|\mathbf{r} - \mathbf{a}_2|^{2\lambda}} &\leq \frac{\chi_{B_1(\mathbf{a}_2)}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\lambda_0+\varepsilon)}} + \frac{\chi_{B_1(\mathbf{a}_2)^c}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\lambda_0-\varepsilon)}}, \end{aligned}$$

where χ_S denotes the characteristic function of the set S . Hence, for all $\mathbf{r} \in M$,

$$\begin{aligned} &\frac{|f(\mathbf{r})|}{|\mathbf{r} - \mathbf{a}_1|^{2\mu}|\mathbf{r} - \mathbf{a}_2|^{2\lambda}} \\ &\leq \|f\|_\infty \left(\frac{\chi_{B_1(\mathbf{a}_1)}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\mu_0+\varepsilon)}} + \frac{\chi_{B_1(\mathbf{a}_1)^c}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2(\mu_0-\varepsilon)}} \right) \left(\frac{\chi_{B_1(\mathbf{a}_2)}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_2|^{2(\lambda_0+\varepsilon)}} + \frac{\chi_{B_1(\mathbf{a}_2)^c}(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_2|^{2(\lambda_0-\varepsilon)}} \right). \end{aligned}$$

The function on the right hand side (RHS) is independent of μ, λ and integrable on $\mathbf{R}^2$. Hence, by the dominated convergence theorem, we obtain (3.4).

(ii) Let $\mathbf{b} = \mathbf{a}_1 - \mathbf{a}_2 = (b_1, b_2)$ and $\delta > 0$ be a constant such that $0 < \delta < \min\{1, |\mathbf{b}|/2\}$. Then we can write

$$\Phi_{\mu,\lambda}(f) = I_{\mu,\lambda}(f) + J_{\mu,\lambda}(f) + K_{\mu,\lambda}(f) \quad (3.8)$$

with

$$\begin{aligned} I_{\mu,\lambda}(f) &= \int_{B_\delta(\mathbf{a}_1)} \frac{f(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2\mu}|\mathbf{r} - \mathbf{a}_2|^{2\lambda}} d\mathbf{r}, \\ J_{\mu,\lambda}(f) &= \int_{B_\delta(\mathbf{a}_2)} \frac{f(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2\mu}|\mathbf{r} - \mathbf{a}_2|^{2\lambda}} d\mathbf{r}, \\ K_{\mu,\lambda}(f) &= \int_{B_\delta(\mathbf{a}_1)^c \cap B_\delta(\mathbf{a}_2)^c} \frac{f(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^{2\mu}|\mathbf{r} - \mathbf{a}_2|^{2\lambda}} d\mathbf{r}. \end{aligned}$$

By the change of variables $x \rightarrow a_{11} + r \cos \theta, y \rightarrow a_{12} + r \sin \theta, 0 < r \leq \delta, 0 \leq \theta < 2\pi$, we have

$$I_{\mu,\lambda}(f) = \int_0^\delta dr \int_0^{2\pi} d\theta \frac{f(a_{11} + r \cos \theta, a_{12} + r \sin \theta)}{r^{2\mu-1}(r^2 + |\mathbf{b}|^2 + 2rb(\theta))^\lambda},$$

where $b(\theta) = b_1 \cos \theta + b_2 \sin \theta$. Moreover, the change of variable $s = r^{2(1-\mu)}$ gives

$$(1 - \mu)I_{\mu,\lambda}(f) = \frac{1}{2} \int_0^1 ds \int_0^{2\pi} d\theta F_{\mu,\lambda}(s, \theta)$$

with

$$F_{\mu,\lambda}(s, \theta) = \frac{\chi_{[0, \delta(\mu)]}(s) f(a_{11} + s^{\frac{1}{2(1-\mu)}} \cos \theta, a_{12} + s^{\frac{1}{2(1-\mu)}} \sin \theta)}{(s^{\frac{1}{1-\mu}} + |b|^2 + 2s^{\frac{1}{2(1-\mu)}} b(\theta))^\lambda},$$

where $\delta(\mu) = \delta^{2(1-\mu)} < 1$. We have

$$\lim_{\mu \rightarrow 1, \lambda \rightarrow \lambda_0} F_{\mu,\lambda}(s, \theta) = \begin{cases} \frac{f(a_1)}{|b|^{2\lambda_0}} & ; \text{for } 0 < s < 1 \\ 0 & ; \text{for } s > 1 \end{cases}.$$

Moreover, it is easy to see that

$$|F_{\mu,\lambda}(s, \theta)| \leq \frac{\|f\|_\infty}{(|b| - \delta)^\lambda} \leq C_0 \|f\|_\infty$$

with $C_0 = \sup_{0 < x < 1} (|b| - \delta)^{-x} < \infty$. Hence, by the dominated convergence theorem, we obtain

$$\lim_{\mu \rightarrow 1, \lambda \rightarrow \lambda_0} (1 - \mu)I_{\mu,\lambda}(f) = \frac{\pi f(a_1)}{|b|^{2\lambda_0}}. \quad (3.9)$$

To estimate $J_{\mu,\lambda}(f)$, we note that there exist positive constants c_1, c_2 such that

$$c_1 \leq |r - a_1| \leq c_2, \quad r \in B_\delta(a_2),$$

so that, for all $r \in B_\delta(a_2)$,

$$\frac{1}{|r - a_1|^{2\mu}} \leq \frac{1}{c_1^{2\mu}} \leq C_1$$

with $C_1 = \sup_{0 < x < 1} c_1^{-2x} < \infty$. Let $\varepsilon > 0$ be such that (3.7) holds. Then we have

$$|r - a_2|^{2\lambda} \geq |r - a_2|^{2(\lambda_0 + \varepsilon)}, \quad r \in B_\delta(a_2),$$

which implies that

$$\frac{|f(r)|}{|r - a_1|^{2\mu} |r - a_2|^{2\lambda}} \leq \frac{C_1 \|f\|_\infty}{|r - a_2|^{2(\lambda_0 + \varepsilon)}}, \quad r \in B_\delta(a_2).$$

The function on the RHS is integrable on $B_\delta(a_2)$, since $\lambda_0 + \varepsilon < 1$. Thus we can apply the dominated convergence theorem to obtain

$$\lim_{\mu \rightarrow 1, \lambda \rightarrow \lambda_0} J_{\mu,\lambda}(f) = \int_{B_\delta(a_2)} \frac{f(r)}{|r - a_1|^2 |r - a_2|^{2\lambda_0}} dr. \quad (3.10)$$

By a method similar to that of the proof of part (i), one can easily show that

$$\lim_{\mu \rightarrow 1, \lambda \rightarrow \lambda_0} K_{\mu, \lambda}(f) = \int_{B_\delta(\mathbf{a}_1)^c \cap B_\delta(\mathbf{a}_2)^c} \frac{f(\mathbf{r})}{|\mathbf{r} - \mathbf{a}_1|^2 |\mathbf{r} - \mathbf{a}_2|^{2\lambda_0}} d\mathbf{r}. \quad (3.11)$$

Formula (3.5) now follows from (3.8)–(3.11).

(iii) Let $\lambda = (\tau - 1 + \mu)/\tau$ and $\max\{1 - \tau, (1 + \tau)^{-1}\} < \mu < 1$. Then, $\lambda \in (0, 1)$, $\lambda + \mu > 1$, and $\lambda \rightarrow 1$ as $\mu \rightarrow 1$. As is seen from the proof of (3.9), (3.9) holds also in the case $\lambda_0 = 1$. Hence

$$\lim_{\mu \rightarrow 1} (1 - \mu) I_{\mu, \lambda}(f) = \frac{\pi f(\mathbf{a}_1)}{|\mathbf{b}|^2}.$$

Similarly we have

$$\lim_{\mu \rightarrow 1} (1 - \lambda) J_{\mu, \lambda}(f) = \frac{\pi f(\mathbf{a}_2)}{|\mathbf{b}|^2}.$$

Formula (3.11) holds also in the case $\lambda_0 = 1$. Since $1 - \mu = \tau(1 - \lambda)$, (3.6) follows. ■

A simple application of Lemma 3.1 to $\varrho_q(f)$ gives the following result.

THEOREM 3.2. *Let $\{q_n\}_{n=1}^\infty$ be a sequence satisfying (3.2) and (3.3).*

(i) *Suppose that $\lambda_1, \lambda_2 \in (0, 1)$ such that $\lambda_1 + \lambda_2 > 1$. Then, for all $f \in L^\infty(\mathbb{R}^2)$,*

$$\lim_{n \rightarrow \infty} \varrho_{q_n}(f) = \frac{\Phi_{\lambda_1, \lambda_2}(f)}{\Phi_{\lambda_1, \lambda_2}(1)}.$$

(ii) *Suppose that $\lambda_1 = 1, \lambda_2 \in (0, 1)$. Then, for all $f \in \mathfrak{B}(\mathbb{R}^2)$,*

$$\lim_{n \rightarrow \infty} \varrho_{q_n}(f) = f(\mathbf{a}_1).$$

(iii) *Let $\tau > 0$ and suppose that, for all sufficiently large n , $1 - \varepsilon_1(q_n) = \tau(1 - \varepsilon_2(q_n))$ and $\lambda_1 = 1$ (hence $\lambda_2 = 1$). Then, for all $f \in \mathfrak{B}(\mathbb{R}^2)$,*

$$\lim_{n \rightarrow \infty} \varrho_{q_n}(f) = \frac{f(\mathbf{a}_1) + \tau f(\mathbf{a}_2)}{1 + \tau}. \quad (3.12)$$

Remark. One can easily find examples of $\{q_n\}_{n=1}^\infty$ for each case in Theorem 3.2.

As a corollary, we obtain the following:

COROLLARY 3.3. *Let $\{q_n\}_{n=1}^\infty$ be a sequence satisfying (3.2) and (3.3). Then, for all cases (i)–(iii) in Theorem 3.2,*

$$\lim_{n \rightarrow \infty} \frac{\varrho_{q_n}(f)}{q_n} = 0, \quad f \in \mathfrak{B}(\mathbb{R}^2).$$

This result shows that the Erdős' theorem established in [3] does not hold in the present case.

In a special case of part (iii) of Theorem 3.2, a strong coupling limit of the ZESD yields the magnetic field :

COROLLARY 3.4. *Let $\{q_n\}_{n=1}^{\infty}$ be a sequence satisfying (3.2) and (3.3) with $\lambda_{\nu} = 1, \nu = 1, 2$. Suppose that, for all sufficiently large n , $\gamma_1(1 - \varepsilon_1(q_n)) = \gamma_2(1 - \varepsilon_2(q_n))$. Then,*

$$\lim_{n \rightarrow \infty} (\gamma_1 + \gamma_2) \varrho_{q_n}(\mathbf{r}) = \gamma_1 \delta(\mathbf{r} - \mathbf{a}_1) + \gamma_2 \delta(\mathbf{r} - \mathbf{a}_2)$$

in the distribution sense. In particular, if $C_{\alpha, \beta}^{(\nu)} = 0$ for $\nu = 1, 2$ and all α, β with $\alpha + \beta \geq 1$, then

$$\lim_{n \rightarrow \infty} (\gamma_1 + \gamma_2) \varrho_{q_n}(\mathbf{r}) = B(\mathbf{r})$$

in the distribution sense.

Proof. We need only to apply part (iii) of Theorem 3.2 with $\tau = \gamma_2/\gamma_1$. ■

References

1. Arai, A., *Momentum operators with gauge potentials, local quantization of magnetic flux, and representation of canonical commutation relations*, J.Math.Phys. 34 (1992), 3374-3378.
2. Arai, A., *Properties of the Dirac-Weyl operator with a strongly singular gauge potential*, J.Math.Phys. 34 (1993), 915-935.
3. Erdős, L., *Ground-state density of the Pauli operator in the large field limit*, Lett. Math.Phys. 29 (1993), 219-240.