HOKKAIDO UNIVERSITY

Title	A remark on estimates of bilinear forms of gradients in Hardy space
Author(s)	Shimizu, Y.
Citation	Hokkaido University Preprint Series in Mathematics, 342, 1-8
Issue Date	1996-07-01
DOI	https://doi.org/10.14943/83488
Doc URL	https://hdl.handle.net/2115/69092
Type	departmental bulletin paper
File Information	re342.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

A REMARK ON ESTIMATES OF BILINEAR FORMS OF GRADIENTS IN HARDY SPACE

Yasuyuki Shimizu

Series #342. July 1996

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

#318 Wei-Zhi Sun, Shadows of moving surfaces, 19 pages. 1995.

#319 S. Izumiya and G.T. Kossioris, Bifurcations of shock waves for viscosity solutions of Hamilton-Jacobi equations of one space variable, 39 pages. 1995.

#320 T. Teruya, Normal intermediate subfactors, 44 pages. 1995.

#321 M. Ohnuma, Axisymmetric solutions and singular parabolic equations in the theory of viscosity solutions, 26 pages. 1995.

#322 T. Nakazi, An outer function and several important functions in two variables, 12 pages. 1995.

#323 N. Kawazumi, An infinitesimal approach to the stable cohomology of the moduli of Riemann surfaces, 22 pages. 1995.

#324 A. Arai, Factorization of self-adjoint operators by abstract Dirac operators and its application to second quantizations on Boson Fermion Fock spaces, 15 pages. 1995.

#325 K. Sugano, On strongly separable Frobenius extensions, 11 pages. 1995.

#326 D. Lehmann and T. Suwa, Residues of holomorphic vector fields on singular varieties, 21 pages. 1995.

#327 K. Tsutaya, Local regularity of non-resonant nonlinear wave equations, 23 pages. 1996.

#328 T. Ozawa and Y. Tsutsumi, Space-time estimates for null gauge forms and nonlinear Schrödinger equations, 25 pages. 1996.

#329 O. Ogurisu, Anticommutativity and spin 1/2 Schrödinger operators with magnetic fields, 12 pages. 1996.

#330 Y. Kurokawa, Singularities for projections of contour lines of surfaces onto planes, 24 pages. 1996.

#331 M.-H. Giga and Y. Giga, Evolving graphs by singular weighted curvature, 94 pages. 1996.

#332 M. Ohnuma and K. Sato, Singular degenerate parabolic equations with applications to the p-laplace diffusion equation, 20 pages. 1996.

#333 T. Nakazi, The spectra of Toeplitz operators with unimodular symbols, 9 pages. 1996.

#334 B. Khanedani and T. Suwa, First variation of horomorphic forms and some applications, 11 pages. 1996.

#335 J. Seade and T. Suwa, Residues and topological invariants of singular holomorphic foliations[1], 28 pages. 1996.

#336 Y. Giga, M.E. Gurtin and J. Matias, On the dynamics of crystalline motions, 67 pages. 1996.

#337 I. Tsuda, A new type of self-organization associated with chaotic dynamics in neural networks, 22 pages. 1996.

#338 F. Hiroshima, A scaling limit of a Hamiltonian of many nonrelativistic particles interacting with a quantized radiation field, 34 pages. 1996.

#339 N. Tominaga, Analysis of a family of strongly commuting self-adjoint operators with applications to perturbed Dirac operators, 29 pages. 1996.

#340 A. Inoue, Abel-Tauber theorems for Fourier-Stieltjes coefficients, 17 pages. 1996.

#341 G. Ishikawa, Topological classification of the tangent developables of space curves, 19 pages. 1996.

A REMARK ON ESTIMATES OF BILINEAR FORMS OF GRADIENTS IN HARDY SPACE

YASUYUKI SHIMIZU

Department of Mathematics
Hokkaido University
Sapporo 060,Japan

§0. Introduction.

In a recent interesting paper[1] L.C.Evans and S.Müller established the estimate of local Hardy space norm of gradients ψ_{x_1}, ψ_{x_2}:

$$(0.1) \quad \|\phi\psi_{x_1}\psi_{x_2}\|_{h^1} + \|\phi(\psi_{x_1}^2 - \psi_{x_2}^2)\|_{h^1} \leqq C(\|\psi_{x_1}\|^2_{L^2(B(0,R))} + \|\psi_{x_2}\|^2_{L^2(B(0,R))})$$

provided that

$$(0.2) \qquad -\Delta\psi = \omega \geqq 0 \text{ in } \mathbb{R}^2.$$

Here ϕ is in $C_0^\infty(\mathbb{R}^2)$ and the constants C and R depends only on ϕ ; h^1 is a local Hardy space defined in §1 and $B(x,R)$ denotes the closed ball of radius R centered at $x \in \mathbb{R}^2$. (Another proof based on harmonic analysis is given by Semmes[2].)

This estimate is useful in proving the existence of weak solutions for the initial value problem of the two-dimensional Euler equation when the vorticity of the initial value is nonnegative measure ([1] and Delort[3]). The assumption $\omega \geqq 0$ in (0.2) is essential for the estimate (0.1); in fact, Evans and Müller[1] gave a counterexample for (0.1) when the condition $\omega \geqq 0$ is violated. However, in their example the set where ω is nonnegative may be complicated.

In this paper we give another counterexample for (0.1) even when ω is odd in the second variable x_2 i.e. $\omega(x_1,x_2) = -\omega(x_1,-x_2)$ and $\omega(x_1,x_2) \geqq 0$ for $x_2 \geqq 0$. This suggests that it is difficult to extend weak solutions for the initial-boundary value problem of the Euler equation when the domain is a half space $\mathbb{R}^2_+$ even if initial value is nonnegative in $\mathbb{R}^2_+$.

To get our counterexample we construct a sequence ψ^ϵ of form $\psi^\epsilon(x) = \psi(x/\epsilon)$. A key observation is the existence of function ψ that satisfies

$$\int_{\mathbb{R}^2} \psi_{x_1}^2\,dx \neq \int_{\mathbb{R}^2} \psi_{x_2}^2\,dx$$

with $-\Delta\psi = \omega$, where $\omega \in C_0^\infty(\mathbb{R}^2)$ is odd in the second variable x_2 and $\omega \geqq 0$ in $\mathbb{R}^2_+$, and $\psi \in H^1(\mathbb{R}^2)$; $H^1(\mathbb{R}^2)$ denotes the Sobolev space, i.e. the space of $f \in L^2(\mathbb{R}^2)$ with $f_{x_1}, f_{x_2} \in L^2(\mathbb{R}^2)$.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

§1. Definition and Main Theorem.

We begin with definition of local Hardy space as in [1].

Definition 1.1. Let η be in $C_0^\infty(\mathbb{R}^n)$ with $\mathrm{supp}\eta \subset B(0,1)$ and $\int_{\mathbb{R}^n} \eta dx = 1$. For a function$f$ in $L^1_{loc}(\mathbb{R}^n)$, f^{**} is defined by

$$f^{**}(x) = \sup_{0<r<1} \left| \int_{\mathbb{R}^n} \eta\left(\frac{x-y}{r}\right) f(y)dy \right|. \tag{1.1}$$

The local Hardy space $\mathcal{H}^1_{loc}$ is defined by

$$\mathcal{H}^1_{loc}(\mathbb{R}^n) = \{f \in L^1_{loc}(\mathbb{R}^n) |\ f^{**} \in L^1_{loc}(\mathbb{R}^n)\}. \tag{1.2}$$

We recall the normed local Hardy space h^1 defined by

$$h^1(\mathbb{R}^n) = \{f \in L^1(\mathbb{R}^n) |\ f^{**} \in L^1(\mathbb{R}^n)\} \tag{1.3}$$

with the norm

$$\|f\|_{h^1(\mathbb{R}^n)} = \|f^{**}\|_{L^1(\mathbb{R}^n)}.$$

Definition 1.2. For a function f in $C_0^\infty(\mathbb{R}^2)$, we define the operator $(-\Delta)^{-1}$ by

$$(-\Delta)^{-1} f(x) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(y) \log|x-y| dy. \tag{1.4}$$

Theorem 1.3. *Let T and S be the spaces of form*

$$T = \left\{\omega \in C_0^\infty(\mathbb{R}^2) |\ \omega(x_1,x_2) \geqq 0 \text{ for } x_2 \geqq 0,\ \omega(x_1,x_2) = -\omega(x_1,-x_2)\right\},$$
$$S = \left\{(-\Delta)^{-1}\omega |\ \omega \in T\right\}.$$

Then there exists a sequence $\{\psi^\epsilon\}_{0<\epsilon<1}$ such that

$$\sup_{0<\epsilon<1} \|\psi^\epsilon\|_{H^1(\mathbb{R}^2)} < \infty$$

and

$$\lim_{\epsilon \downarrow 0} \left\|\phi\left\{(\psi^\epsilon_{x_1})^2 - (\psi^\epsilon_{x_2})^2\right\}\right\|_{h^1(\mathbb{R}^2)} = \infty \tag{1.5}$$

where $\phi \in C_0^\infty(\mathbb{R}^2), 0 \leqq \phi \leqq 1, \phi|_{B(0,1/8)} \equiv 1$ and $\mathrm{supp}\phi \subset B(0,1/2)$.

§2. Proof of Theorem

At first, we show a fundamental estimate in normed local Hardy space; this is an extension of a result of Evans and Müller[1].

Lemma 2.1. *Assume that f is in $L^1(\mathbb{R}^n)$, and $\int_{\mathbb{R}^n} f(x)dx \neq 0$. Let $f^\epsilon = \frac{1}{\epsilon^n} f\left(\frac{x}{\epsilon}\right)$. Then $\|f^\epsilon\|_{L^1} = \|f\|_{L^1} < \infty$, and*

$$\lim_{\epsilon \downarrow 0} \|\phi f^\epsilon\|_{h^1(\mathbb{R}^n)} = \infty \tag{2.1}$$

for a function ϕ in $C_0^\infty(\mathbb{R}^n)$ with $0 \leqq \phi \leqq 1, \phi|_{B(0,1/8)} \equiv 1$, and $\mathrm{supp}\phi \subset B(0,1/4)$.

Proof. L^1-estimate is easily obtained by scaling variables. To show the estimate (2.1), assume that the function η in (1.1) satisfies $0 \leqq \eta \leqq 1$ and $\eta|_{B(0,1/2)} \equiv 1$. Now we estimate the function $(\phi f^\epsilon)^{**}$:

$$\begin{aligned}(\phi f^\epsilon)^{**}(x) &= \sup_{0<r<1} \left| \frac{1}{r^n} \int_{\mathbb{R}^n} \eta\left(\frac{x-y}{r}\right) \phi(y) f^\epsilon(y) dy \right| \\ &= \sup_{0<r<1} \left| \frac{1}{r^n} \int_{\mathbb{R}^n} \eta\left(\frac{x-\epsilon z}{r}\right) \phi(\epsilon z) f(z) dz \right|.\end{aligned} \tag{2.2}$$

Now take a parameter $R > 0$, and let $r = 4|x|$. Then

$$\begin{aligned}(\phi f^\epsilon)^{**}(x) &\geqq \frac{1}{4^n|x|^n} \left| \int_{B(0,R)} \eta\left(\frac{x-\epsilon z}{4|x|}\right) \phi(\epsilon z) f(z) dz \right| \\ &\quad - \frac{1}{4^n|x|^n} \left| \int_{\mathbb{R}^n \setminus B(0,R)} \eta\left(\frac{x-\epsilon z}{4|x|}\right) \phi(\epsilon z) f(z) dz \right| \\ &= I_1^\epsilon - I_2^\epsilon \text{ for } \epsilon R < |x| < 1/4.\end{aligned} \tag{2.3}$$

We show that $\lim_{\epsilon \downarrow 0} \|I_2^\epsilon\|_{L^1(B(0,1/4))} = 0$ and $\lim_{\epsilon \downarrow 0} \|I_1^\epsilon\|_{L^1(B(0,1/4))} = \infty$ to complete the proof. Firstly, we estimate the term I_2^ϵ :

$$\begin{aligned}I_2^\epsilon(x) &\leqq \frac{1}{(4\epsilon R)^n} \int_{\mathbb{R}^n \setminus B(0,R)} |f(z)|\, dz = \frac{F(R)}{(4\epsilon R)^n} \\ &\text{with } F(R) = \int_{\mathbb{R}^n \setminus B(0,R)} |f(z)|\, dz.\end{aligned}$$

Since $F(R)$ is continuous, nonincreasing, and $\lim_{R \to \infty} F(R) = 0$, for sufficiently small ϵ, there exists $R = R(\epsilon)$ such that

$$\frac{\{F(R)\}^{1/2}}{R^n} = \epsilon^n. \tag{2.4}$$

By (2.4), we get

$$I_2^\epsilon(x) \leqq \frac{\{F(R)\}^{1/2}}{4^n} \longrightarrow 0 \text{ as } \epsilon \downarrow 0 \tag{2.5}$$

and get $\lim_{\epsilon \downarrow 0} \|I_2^\epsilon\|_{L^1(B(0,1/4))} = 0$.

Secondly, we estimate the term I_1^ϵ. As $\left|\frac{x-\epsilon z}{4|x|}\right| \leqq \frac{|x|}{4|x|} + \frac{\epsilon R}{4|x|} \leqq \frac{1}{2}$,

$$I_1^\epsilon = \frac{1}{4^n|x|^n}\left|\int_{B(0,R)} \phi(\epsilon z) f(z) dz\right| \tag{2.6}$$
$$= \frac{1}{4^n|x|^n}\left|\int_{B(0,R)} f(z) dz\right| \text{ for } \epsilon R \leqq |x| \leqq 1/4.$$

for $\epsilon R < 1/8$.

Now let $\epsilon \downarrow 0$. For $\epsilon R = \{F(R)\}^{1/2n} \longrightarrow 0$ by (2.4),

$$\lim_{\epsilon \downarrow 0} I_1^\epsilon = \frac{1}{4^n|x|^n}\left|\int_{\mathbb{R}^n} f(z) dz\right| = \frac{\|f\|_{L^1(\mathbb{R}^n)}}{4^n|x|^n} \text{ for } 0 \leqq |x| \leqq 1/4. \tag{2.7}$$

Since $\frac{1}{|x|^n}$ is not in $L^1(\mathbb{R}^n)$, we get a conclusion. □

Next lemma is important to show Lemma 2.3 which is the key to show Theorem 1.3:

Lemma 2.2. *For any function ψ in S,there exists a constant C depending only on ψ such that*

$$|\psi(x)| \leqq \frac{C}{1+|x|} \tag{2.8}$$

$$|\psi_{x_j}(x)| \leqq \frac{C}{1+|x|^2},\ j = 1, 2 \tag{2.9}$$

for $x \in \mathbb{R}^2$.

Proof. If ψ is in S, then there exists a function ω in T such that

$$\psi(x) = \frac{1}{2\pi}\int_{\mathbb{R}^2} \omega(y) \log|x-y| dy. \tag{2.10}$$

Notice that ω is in $C_0^\infty(\mathbb{R}^2)$, so there is a constant $R = R(\omega)$ such that $\mathrm{supp}\omega \subset B(0,R)$. Since ω is odd in x_2, we get

$$\psi(x) = \frac{1}{2\pi}\int_{B(0,R)} \omega(y) \log|x-y| dy \tag{2.11}$$
$$= \frac{1}{2\pi}\int_{B_+(0,R)} \omega(y) \log\frac{|x-y|}{|x-\bar{y}|} dy,$$

$$\psi_{x_1}(x) = \frac{1}{2\pi}\int_{B(0,R)} \omega(y)\frac{x_1-y_1}{|x-y|^2} dy \tag{2.12}$$
$$= \frac{2x_2}{\pi}\int_{B_+(0,R)} \omega(y)\frac{y_2(x_1-y_1)}{|x-y|^2|x-\bar{y}|^2} dy,$$

$$\psi_{x_2}(x) = \frac{1}{2\pi}\int_{B(0,R)} \omega(y)\frac{x_2-y_2}{|x-y|^2} dy \tag{2.13}$$
$$= \frac{1}{\pi}\int_{B_+(0,R)} \omega(y)\frac{y_2\left\{(x_2-y_2)(x_2+y_2)-(x_1-y_1)^2\right\}}{|x-y|^2|x-\bar{y}|^2} dy,$$

where $B_+(0,R)$ denotes $B(0,R)\cap\mathbb{R}^2_+$ and $\bar{y}=(y_1,-y_2)$. Now we show that ψ and ψ_{x_j} is bounded in $B(0,R)$ and that there exists a constant C such that

$$
\begin{aligned}
|\psi(x)| &\leqq C|x|^{-1}, \\
|\psi_{x_j}(x)| &\leqq C|x|^{-2} \text{ for } |x|\geqq 2R.
\end{aligned} \tag{2.14}
$$

The boundedness of ψ and ψ_j on $B(0,2R)$ is obtained by the fact that $\log|x|$ and $|x|^{-1}$ are in $L^1_{loc}(\mathbb{R}^2)$:

$$
\begin{aligned}
|\psi(x)| &\leqq \frac{\sup|\omega|}{2\pi}\int_{B(0,R)}|\log|x-y||\,dy \\
&= C_\omega\int_{B(x,R)}|\log|y||\,dy \\
&\leqq C_\omega\int_{B(0,3R)}|\log|y||\,dy \\
&= C_{\omega,R}<\infty,
\end{aligned} \tag{2.15}
$$

$$
\begin{aligned}
|\psi_{x_j}(x)| &\leqq \frac{\sup|\omega|}{2\pi}\int_{B(0,R)}\frac{1}{|x-y|}dy \\
&\leqq C_\omega\int_{B(0,3R)}|y|^{-1}dy \\
&= C_{\omega,R}<\infty
\end{aligned} \tag{2.16}
$$

for $|x|\leqq 2R$. (Notice that $|y|\leqq|x-y|+|x|\leqq 3R$.)

Now we show (2.14) to complete the proof. We may assume $x_2\geqq 0$ to estimate ψ, because ψ is odd in x_2. By this assumption, we get the following inequality:

$$
1\leqq\frac{|x-\bar{y}|}{|x-y|}\leqq\frac{|x-y|+|y-\bar{y}|}{|x-y|}\leqq 1+\frac{2R}{|x|-R}\leqq 1+\frac{4R}{|x|} \tag{2.17}
$$

for $|x|\geqq 2R$ and $|y|\leqq R$. The inequality (2.17) leads the estimate of ψ :

$$
\begin{aligned}
|\psi(x)| &\leqq \frac{1}{2\pi}\log\left(1+\frac{4R}{|x|}\right)\int_{B_+(0,R)}\omega(y)dy \\
&\leqq \frac{1}{2\pi}\frac{4R}{|x|}\log\left(1+\frac{4R}{|x|}\right)^{\frac{|x|}{4R}}\|\omega\|_{L^1(B_+(0,R))} \\
&\leqq \frac{2R}{\pi}\|\omega\|_{L^1(\mathbb{R}^2)}|x|^{-1}.
\end{aligned} \tag{2.18}
$$

Notice that the inequality

$$
\frac{|x|}{2}\leqq|x-y|,\ \frac{|x|}{2}\leqq|x-\bar{y}|
$$

holds for $|x| \geqq 2R, |y| \leqq R$. This inequality leads the estimate of ψ_{x_j} :

$$|\psi_{x_1}(x)| \leqq \frac{2|x|}{\pi} \int_{B_+(0,R)} |\omega(y)| \frac{|y|}{|x-y|^2|x-\bar{y}|} dy \leqq \frac{16R\|\omega\|_{L^1}}{\pi|x|^2}, \tag{2.19}$$

$$|\psi_{x_2}(x)| \leqq \frac{1}{\pi} \int_{B_+(0,R)} |\omega(y)| \frac{2|y|}{|x-y||x-\bar{y}|} dy \leqq \frac{8R\|\omega\|_{L^1}}{\pi|x|^2}. \tag{2.20}$$

Combining the estimate (2.15), (2.16), (2.18), (2.19), and (2.20) leads the conclusion (2.8) and (2.9). □

Now we are ready to show the key lemma.

Lemma 2.3. *There exists a function ψ in S such that*

$$\int_{\mathbb{R}^2} \{\psi_{x_1}(x)\}^2 dx \neq \int_{\mathbb{R}^2} \{\psi_{x_2}(x)\}^2 dx. \tag{2.21}$$

Proof. Assume that the conclusion is not true, i.e.

$$\int_{\mathbb{R}^2} \{\psi_{x_1}(x)\}^2 dx = \int_{\mathbb{R}^2} \{\psi_{x_2}(x)\}^2 dx \tag{2.22}$$

for any ψ in S. Let ψ is in S, and let $\psi^h(x) = \psi(x_1 - h, x_2)$. Then the function ψ^h and $\psi + \psi^h$ are in S. In fact,there exists a function ω in T such that $\psi = (-\Delta)^{-1}\omega$, and we can write $\psi^h = (-\Delta)^{-1}\omega^h$ and $\psi + \psi^h = (-\Delta)^{-1}(\omega + \omega^h)$, where $\omega^h(x) = \omega(x_1 - h, x_2)$. It is obvious that ω^h and $\omega + \omega^h$ are in T.

By the assumption (2.22), we get

$$\begin{aligned} \int_{\mathbb{R}^2} \{\psi_{x_1}(x)\}^2 dx &= \int_{\mathbb{R}^2} \{\psi_{x_2}(x)\}^2 dx, \\ \int_{\mathbb{R}^2} \{(\psi^h)_{x_1}(x)\}^2 dx &= \int_{\mathbb{R}^2} \{(\psi^h)_{x_2}(x)\}^2 dx, \\ \int_{\mathbb{R}^2} \{(\psi + \psi^h)_{x_1}(x)\}^2 dx &= \int_{\mathbb{R}^2} \{(\psi + \psi^h)_{x_2}(x)\}^2 dx. \end{aligned} \tag{2.23}$$

Combining the equalities (2.23), we get

$$\int_{\mathbb{R}^2} \psi_{x_1}(x)(\psi^h)_{x_1}(x) dx = \int_{\mathbb{R}^2} \psi_{x_2}(x)(\psi^h)_{x_2}(x) dx. \tag{2.24}$$

Now we integrate the equality (2.24) by h:

$$\int_{-\infty}^{\infty} \int_{\mathbb{R}^2} \psi_{x_1}(x)(\psi^h)_{x_1}(x) dx = \int_{-\infty}^{\infty} \int_{\mathbb{R}^2} \psi_{x_2}(x)(\psi^h)_{x_2}(x) dx. \tag{2.25}$$

Notice that we can change the order of integration by the estimate (2.9). Firstly, we compute the left term of (2.25):

(2.26)
$$\begin{aligned}\int_{-\infty}^{\infty}\int_{\mathbb{R}^2}\psi_{x_1}(x)(\psi^h)_{x_1}(x)dx\,dh &= \int_{\mathbb{R}^2}\psi_{x_1}(x)\int_{-\infty}^{\infty}\psi_{x_1}(x_1-h,x_2)dh\,dx\\ &= \int_{\mathbb{R}^2}\psi_{x_1}(x)\Big[\psi(h,x_2)\Big]_{-\infty}^{\infty}dx\\ &= 0.\end{aligned}$$

The equality is obvious by the estimate (2.8). Secondly, we compute the right term of (2.25):

(2.27)
$$\begin{aligned}\int_{-\infty}^{\infty}\int_{\mathbb{R}^2}\psi_{x_2}(x)(\psi^h)_{x_2}(x)dx\,dh &= \int_{\mathbb{R}^2}\psi_{x_2}(x)\int_{-\infty}^{\infty}\psi_{x_2}(x_1-h,x_2)dh\,dx\\ &= \int_{-\infty}^{\infty}\left\{\int_{\infty}^{\infty}\psi_{x_2}(x_1,x_2)dx_1\right\}^2 dx_2\\ &> 0.\end{aligned}$$

The positivity of integrals is shown by computing the form of ψ_{x_2}. Notice that ψ_{x_2} is continuous, so it is sufficient to check that $\psi_{x_2}(x_1,0)$ is not zero. By (2.13), we can write ψ_{x_2} as

$$\psi_{x_2}(x)=\frac{1}{\pi}\int_{B_+(0,R)}\omega(y)\frac{y_2\left\{(x_2-y_2)(x_2+y_2)-(x_1-y_1)^2\right\}}{|x-y|^2|x-\bar{y}|^2}dy \tag{2.28}$$

where ω is a function in T. Putting $x_2=0$ in (2.28) leads

$$\psi_{x_2}(x_1,0)=-\frac{1}{\pi}\int_{B_+(0,R)}\omega(y)\frac{y_2}{(x_1-y_1)^2+y_2^2}dy. \tag{2.29}$$

Since ω and the integral kernel are positive in $B_+(0,R)$, $\psi_{x_2}(x_1,0)$ is always nonzero. The results of compution (2.26) and (2.27) leads a contradiction, and we get a conclusion of the lemma. □

Proof of Theorem 1.3. Let ψ the function in S that satisfies (2.20). Let $\psi^\epsilon(x)=\psi(x/\epsilon)$. Then

$$\left\{(\psi^\epsilon)_{x_1}(x)\right\}^2-\left\{(\psi^\epsilon)_{x_2}(x)\right\}^2=\frac{1}{\epsilon^2}\left\{(\psi_{x_1})^2-(\psi_{x_2})^2\right\}\left(\frac{x}{\epsilon}\right)$$

and Lemma 2.1 is appliable. (Notice that $\psi_{x_1}^2-\psi_{x_2}^2$ is in $L^1(\mathbb{R}^2)$: the estimate (2.9) leads this fact.) □

REFERENCES

1. L.C.Evans and S.Müller, *Hardy Spaces and the Two-Dimensional Euler Equations with Nonnegative Volticity*, J. Amer. Math. Soc. **7** (1994), p.199-p.219.
2. S.Semmes, *A Primer on Hardy Spaces and Some Remarks on a Theorem of Evans and Müller*, Commun. in Partial Differential Equations **19** (1994), p.277-p.319.
3. J.M.Delort, *Existence de napples de tourbillon en dimension deux*, J. Amer. Math. Soc. **4** (1991), p.553-p.586.
4. C.Fefferman and E.Stein, $\mathcal{H}^p$ *Spaces of Several Variables*, Acta. Math. **129** (1972), p.137-p.197.
5. A.Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Academic Press, 1986.