



HOKKAIDO UNIVERSITY

Title	An Abel-Tauber theorem for Hankel transforms
Author(s)	Bingham, N. H; Inoue, A.
Citation	Hokkaido University Preprint Series in Mathematics, 370, 1-8
Issue Date	1997-02-01
DOI	https://doi.org/10.14943/83516
Doc URL	https://hdl.handle.net/2115/69120
Type	departmental bulletin paper
File Information	re370.pdf



**AN ABEL-TAUBER THEOREM
FOR HANKEL TRANSFORMS**

N.H. Bingham and A. Inoue

Series #370. February 1997

HOKKAIDO UNIVERSITY

PREPRINT SERIES IN MATHEMATICS

- #346 H. Kubo and K. Kubota, Asymptotic behaviors of radially symmetric solutions of $\square u = |u|^p$ for super critical values p in even space dimensions, 66 pages. 1996.
- #347 T. Nakazi and T. Yamamoto, Weighted Norm Inequalities For Some Singular Integral Operators, 17 pages. 1996.
- #348 Y. Ito and I. Nakamura, Hilbert schemes and simple singularities A_n and D_n , 22 pages. 1996
- #349 R. Agemi and K. Yokoyama, The null condition and global existence of solutions to systems of wave equations with different speeds, 42 pages. 1996
- #350 F. Hiroshima, Weak coupling limit with a removal of an ultraviolet cut-off for a Hamiltonian of particles interacting with a massive scalar field, 21 pages. 1996.
- #351 T. Nakazi and Y. Watatani, Invariant subspace theorems for subdiagonal algebras, 22 pages. 1996.
- #352 Y. Nishiura and H. Suzuki, Nonexistence of stable turing patterns with smooth limiting interfacial configurations in higher dimensional spaces, 21 pages. 1996.
- #353 Y.-G. Chen, Y. Giga and K. Sato, On instant extinction for very fast diffusion equations, 9 pages. 1996.
- #354 A. Gyoja and H. Yamashita, Associated variety, Kostant-Sekiguchi correspondence, and locally free $U(n)$ -action on Harish-Chandra modules, 25 pages. 1996.
- #355 G. Ishikawa, Topology of plane trigonometric curves and the strangeness of plane curves derived from real pseudo-line arrangements, 18 pages. 1996.
- #356 N.H. Bingham and A. Inoue, The Drasin-Shea-Jordan theorem for Hankel transforms of arbitrarily large order, 13 pages. 1996.
- #357 S. Izumiya, Singularities of solutions for first order partial differential equations, 20 pages. 1996.
- #358 N. Hayashi, P.I. Naumkin and T. Ozawa, Scattering theory for the Hartree equation, 14 pages. 1996.
- #359 I. Tsuda and K. Tadaki, A logic-based dynamical theory for a genesis of biological threshold, 49 pages. 1996.
- #360 I. Tsuda and A. Yamaguchi, Singular-continuous nowhere-differentiable attractors in neural systems, 40 pages. 1996.
- #361 M. Nakamura and T. Ozawa, Low energy scattering for nonlinear Schrödinger equations in fractional order Sobolev spaces, 17 pages. 1996.
- #362 I. Nakamura, Hilbert schemes and simple singularities E_6 , E_7 and E_8 , 21 pages. 1996.
- #363 T. Mikami, Equivalent conditions on the central limit theorem for a sequence of probability measures on R , 7 pages. 1996.
- #364 S. Izumiya and T. Sano, Generic affine differential geometry of space curves, 23 pages. 1996.
- #365 T. Tsukada, Stability of reticular optical caustics, 12 pages. 1996.
- #366 A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of a generalized spin-boson model, 40 pages. 1996.
- #367 A. Arai, A class of representations of the $*$ -algebra of the canonical commutation relations over a Hilbert space and instability of embedded eigenvalues in quantum field models, 12 pages. 1996.
- #368 K. Ito, BV-solutions of a hyperbolic-elliptic system for a radiating gas, 33 pages. 1997.
- #369 M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, 20 pages. 1997.

AN ABEL-TAUBER THEOREM FOR HANKEL TRANSFORMS

N. H. BINGHAM and A. INOUE

§1. Abel-Tauber theorems

We write R_0 for the class of slowly varying functions at infinity, that is, the class of positive measurable ℓ , defined on some neighbourhood of infinity, satisfying

$$\ell(\lambda x)/\ell(x) \rightarrow 1 \quad (x \rightarrow \infty) \quad \forall \lambda > 0.$$

For $\ell \in R_0$, the class Π_ℓ is the class of measurable f satisfying

$$\{f(\lambda x) - f(x)\}/\ell(x) \rightarrow c \log \lambda \quad (x \rightarrow \infty) \quad \forall \lambda > 0$$

for some constant c , called the ℓ -index of f . For details, we refer to [BGT].

Let $\nu \geq -1/2$, $t^{\nu+\frac{1}{2}}f(t) \in L^1_{loc}[0, \infty)$, and let f be ultimately decreasing to zero at infinity (this plays the role of our Tauberian condition; see [BI1], §2 for discussion of the possibility of weakening it). We consider the *Hankel transform*

$$F_\nu(x) := \int_0^{\infty-} f(t)(xt)^{\frac{1}{2}} J_\nu(xt) dt \quad (x > 0), \quad (1.1)$$

where $\int_0^{\infty-}$ denotes an improper integral $\lim_{M \rightarrow \infty} \int_0^M$ and J_ν is the Bessel function

$$J_\nu(x) = \left(\frac{x}{2}\right)^\nu \sum_{n=0}^{\infty} \frac{(-1)^n}{n! \Gamma(\nu + n + 1)} \left(\frac{x}{2}\right)^{2n} \quad (0 \leq x < \infty).$$

Since the improper integral converges uniformly on each (a, ∞) with $a > 0$, F_ν is finite and continuous on $(0, \infty)$.

Here is our main theorem:

THEOREM 1.1. *Let $\ell \in R_0$, and let ν, f and F_ν be as above. We set $\bar{F}_\nu(x) := x^{\nu+\frac{1}{2}}F_\nu(1/x)$ for $x > 0$. Then*

$$f(t) \sim t^{-(\nu+\frac{3}{2})}\ell(t) \quad (t \rightarrow \infty) \quad (1.2)$$

if and only if

$$\bar{F}_\nu \in \Pi_\ell \quad \text{with} \quad \ell\text{-index} \quad \frac{1}{2\nu\Gamma(\nu+1)}. \quad (1.3)$$

Both imply

$$\begin{aligned} & \{\bar{F}_\nu(x) - \frac{1}{2^\nu \Gamma(\nu+1)} \int_0^x t^{\nu+\frac{1}{2}} f(t) dt\} / (x^{\nu+\frac{3}{2}} f(x)) \\ & \rightarrow \frac{1}{2^\nu \Gamma(\nu+1)} \left\{ \log 2 - \frac{\gamma}{2} + \frac{1}{2} \psi(\nu+1) \right\} \quad (x \rightarrow \infty), \end{aligned} \quad (1.4)$$

where γ is Euler's constant and ψ is the digamma function: $\psi(x) := \Gamma'(x)/\Gamma(x)$ for $x > 0$.

The theorem above treats the boundary case of the following known Abel-Tauber theorem for Hankel transforms:

THEOREM 1.2 ([B1], [RS]). *Let ν, f, F_ν and ℓ be as in Theorem 1.1. Then for $\rho \in (-\nu - \frac{3}{2}, 0)$,*

$$f(t) \sim t^\rho \ell(t) \quad (t \rightarrow \infty) \quad (1.5)$$

if and only if

$$F_\nu(x) \sim x^{-\rho-1} \ell(1/x) \cdot 2^{\rho+\frac{1}{2}} \frac{\Gamma(\frac{3}{4} + \frac{\nu}{2} + \frac{\rho}{2})}{\Gamma(\frac{1}{4} + \frac{\nu}{2} - \frac{\rho}{2})} \quad (x \rightarrow 0+).$$

See also [SS, pp. 617-619].

The point of Theorem 1.1 is that one obtains Π -variation rather than regular variation - or in the terminology of [BGT], one uses de Haan theory rather than Karamata theory - in the boundary case $\rho = -\nu - \frac{3}{2}$. The cosine case $\nu = -1/2$ and sine case $\nu = 1/2$ of Theorem 1.1 were recently proved in [I1] and [I2] respectively. However, the proofs of [I1] and [I2] do not apply to general Hankel transforms directly. Our proof of Theorem 1.1 in §2 is closer to those of [B2, Th. 3b] and [BGT, Th. 4.10.1] than those of [I1], [I2].

§2. Proof of Theorem 1.1.

Step 1. Choose X so large that f is positive and non-increasing on $[X, \infty)$. We first show that we lose no generality by supposing that f vanishes on $[0, X)$.

Set $\tilde{f}(t) := I_{[X, \infty)}(t)f(t)$, and let $\tilde{F}_\nu$ be its Hankel transform:

$$\tilde{F}_\nu(x) := \int_0^{\infty-} \tilde{f}(t)(xt)^{1/2} J_\nu(xt) dt \quad (x > 0).$$

By the mean-value theorem, there exists $c_1 \in (0, \infty)$ such that

$$|x^{-\nu} J_\nu(x) - y^{-\nu} J_\nu(y)| \leq c_1 |x - y| \quad (0 \leq x, y \leq 1).$$

So for $\lambda > 1$,

$$\frac{1}{\ell(x)} |\{\bar{F}_\nu(\lambda x) - \bar{F}_\nu(x)\} - \{(\lambda x)^{\nu+\frac{1}{2}} \tilde{F}_\nu(1/\lambda x) - x^{\nu+\frac{1}{2}} \tilde{F}_\nu(1/x)\}|$$

$$\begin{aligned}
&= \frac{1}{\ell(x)} \left| \int_0^X t^{\nu+\frac{1}{2}} f(t) \{ (t/\lambda x)^{-\nu} J_\nu(t/\lambda x) - (t/x)^{-\nu} J_\nu(t/x) \} dt \right| \\
&\leq c_1 \frac{(1-\lambda^{-1})}{x\ell(x)} \cdot X \cdot \int_0^X t^{\nu+\frac{1}{2}} |f(t)| dt \rightarrow 0 \quad (x \rightarrow \infty).
\end{aligned}$$

So (1.3) holds if and only if $x^{\nu+\frac{1}{2}} \tilde{F}_\nu(1/x) \in \Pi_\ell$ with ℓ -index $1/\{2^\nu \Gamma(\nu+1)\}$.

On the other hand, there exists $c_2 \in (0, \infty)$ such that

$$|x^{-\nu} J_\nu(x) - \frac{1}{2^\nu \Gamma(\nu+1)}| \leq c_2 x^2 \quad (0 \leq x \leq 1). \quad (2.1)$$

So for $x > X$,

$$\begin{aligned}
&\frac{1}{x^{\nu+\frac{3}{2}} f(x)} \left| x^{\nu+\frac{3}{2}} \int_0^X f(t) (t/x)^{1/2} J_\nu(t/x) dt - \frac{1}{2^\nu \Gamma(\nu+1)} \int_0^X f(t) t^{\nu+\frac{1}{2}} dt \right| \\
&\leq \frac{1}{x^{\nu+\frac{3}{2}} f(x)} \int_0^X t^{\nu+\frac{1}{2}} |f(t)| \cdot \left| (t/x)^{-\nu} J_\nu(t/x) - \frac{1}{2^\nu \Gamma(\nu+1)} \right| dt \\
&\leq \frac{c_2 X^2}{x^{\nu+\frac{3}{2}+2} f(x)} \int_0^X t^{\nu+\frac{1}{2}} |f(t)| dt \rightarrow 0 \quad (x \rightarrow \infty),
\end{aligned}$$

under (1.2). Thus when proving that (1.2) is equivalent to (1.3) and implies (1.4), we may replace f by $\tilde{f}$ - that is, we may assume that f vanishes on $[0, X]$.

Step 2: Abelian part. First we assume (1.2) and show (1.4) and (1.3). For $x > 0$,

$$\frac{1}{x^{\nu+\frac{3}{2}} f(x)} \left\{ \tilde{F}_\nu(x) - \frac{1}{2^\nu \Gamma(\nu+1)} \int_0^x f(t) t^{\nu+\frac{1}{2}} dt \right\} = I(x) + II(x), \quad (2.2)$$

where

$$\begin{aligned}
I(x) &:= \int_0^1 \frac{(xu)^{\nu+2} f(xu)}{x^{\nu+2} f(x)} \left\{ u^{1/2} J_\nu(u) - \frac{u^{\nu+\frac{1}{2}}}{2^\nu \Gamma(\nu+1)} \right\} \frac{du}{u^{\nu+2}}, \\
II(x) &:= \int_1^{\infty} \frac{f(xu)}{f(x)} u^{1/2} J_\nu(u) du.
\end{aligned}$$

By the uniform convergence theorem for regularly varying functions ([BGT, Th. 1.5.2]), $(xu)^{\nu+2} f(xu)/x^{\nu+2} f(x)$ converges to $u^{1/2}$ as $x \rightarrow \infty$ uniformly in $u \in (0, 1]$, whence using (2.1) we find that

$$I(x) \rightarrow \int_0^1 \frac{1}{u^{\nu+\frac{3}{2}}} \left\{ u^{1/2} J_\nu(u) - \frac{u^{\nu+\frac{1}{2}}}{2^\nu \Gamma(\nu+1)} \right\} du \quad (x \rightarrow \infty).$$

In the same way, for any $Y > X$,

$$\int_1^Y \frac{f(xu)}{f(x)} u^{1/2} J_\nu(u) du \rightarrow \int_1^Y \frac{1}{u^{\nu+\frac{3}{2}}} \cdot u^{\frac{1}{2}} J_\nu(u) du \quad (x \rightarrow \infty).$$

By the second integral mean-value theorem ([WW, §4.14]), for $x > 1$,

$$\int_Y^{\infty-} \frac{f(xu)}{f(x)} u^{1/2} J_\nu(u) du = \frac{f(xY+)}{f(x)} \int_Y^{Y'} u^{1/2} J_\nu(u) du$$

for some $Y' \in (Y, \infty)$. If we set

$$c_3 := \sup\{|\int_x^y u^{1/2} J_\nu(u) du| : 0 < x < y < \infty\},$$

then

$$\limsup_{x \rightarrow \infty} |\int_Y^{\infty-} \frac{f(xu)}{f(x)} u^{1/2} J_\nu(u) du| \leq \limsup_{x \rightarrow \infty} \frac{f(xY)}{f(x)} \cdot c_3 = c_3 / Y^{\nu+\frac{3}{2}},$$

which can be made arbitrarily small by choosing Y large enough. So

$$II(x) \rightarrow \int_1^\infty \frac{1}{u^{\nu+\frac{3}{2}}} \cdot u^{1/2} J_\nu(u) du \quad (x \rightarrow \infty).$$

The sum of the limits of $I(x)$ and $II(x)$ is identified in the Appendix by classical results on special functions, and we obtain (1.4).

We can prove the implication from (1.4) to (1.3) by a standard argument (see e.g. [I1, p. 767]).

Step 3: Tauberian part. We now prove the implication from (1.3) to (1.2). By a formula of Gegenbauer,

$$\int_0^\infty t^{\nu+\frac{1}{2}} e^{-xt} (yt)^{1/2} J_\nu(yt) dt = d_\nu \cdot \frac{xy^{\nu+\frac{1}{2}}}{(x^2 + y^2)^{\nu+\frac{3}{2}}} \quad (x > 0),$$

where

$$d_\nu := \pi^{-1/2} 2^{\nu+1} \Gamma(\nu + \frac{3}{2})$$

([WW, §13.2 (5)]; cf. [BI1, §2]). So by Parseval's formula for Hankel transforms ([RS]; cf. [BI1, §1]), for $x > 0$

$$\int_0^\infty f(t) t^{\nu+\frac{1}{2}} e^{-xt} dt = d_\nu \int_{0+}^\infty F_\nu(y) \frac{xy^{\nu+\frac{1}{2}}}{(x^2 + y^2)^{\nu+\frac{3}{2}}} dy \quad (2.3)$$

(for $\int_{0+}$ on the right, recall the assumption at the end of Step 1). By the second integral mean-value theorem,

$$|F_\nu(y)| \leq c_3 f(X)/y \quad (0 < y < \infty), \quad (2.4)$$

whence

$$\int_1^\infty |F_\nu(y)| \cdot \frac{y^{\nu+\frac{1}{2}}}{(x^2 + y^2)^{\nu+\frac{3}{2}}} dy < \infty.$$

On the other hand, by [BGT, Th. 3.7.4], (1.3) implies $|\bar{F}_\nu| \in R_0$, whence

$$\int_0^1 |F_\nu(y)| \cdot \frac{y^{\nu+\frac{1}{2}}}{(x^2+y^2)^{\nu+\frac{3}{2}}} dy = \int_1^\infty \frac{|\bar{F}_\nu(u)|}{(x^2u^2+1)^{\nu+\frac{3}{2}}} du < \infty.$$

Thus the integral on the right of (2.3) converges absolutely - and so the results of [BGT Ch. 4] apply.

We use Laplace transforms. Write

$$U(t) := \int_0^t f(u)u^{\nu+\frac{1}{2}} du \quad (t \geq 0),$$

$$\hat{U}(x) := \int_{[0,\infty)} e^{-xt} dU(t) = \int_0^\infty f(t)t^{\nu+\frac{1}{2}} e^{-xt} dt \quad (x > 0).$$

Then by (2.3),

$$\hat{U}(1/x) = (k * \bar{F}_\nu)(x) \quad (x > 0),$$

where

$$k(x) := d_\nu \frac{x^{2\nu+2}}{(1+x^2)^{\nu+\frac{3}{2}}} \quad (x > 0)$$

and $k * \bar{F}_\nu$ denotes the Mellin convolution

$$(k * \bar{F}_\nu)(x) = \int_0^\infty k(x/t) \bar{F}_\nu(t) dt/t.$$

The absolute convergence strip of the Mellin transform

$$\check{k}(z) := \int_0^\infty t^{-z} k(t) dt/t$$

is $-1 < \Re z < 2\nu + 2$, and for z in the strip

$$\check{k}(z) = \frac{2^\nu}{\pi^{1/2}} \Gamma(\nu + 1 - \frac{z}{2}) \Gamma(\frac{1}{2} + \frac{z}{2}),$$

in particular,

$$\check{k}(0) = 2^\nu \Gamma(\nu + 1).$$

By (2.4), $\bar{F}_\nu$ is locally bounded on $[0, \infty)$. So by the argument of [BGT, p. 242] (Abelian theorem for differences), we find that (1.3) implies $k * \bar{F}_\nu \in \Pi_\ell$ with ℓ -index 1 - i.e., so is $\hat{U}(1/\cdot)$. So by a Tauberian theorem of de Haan (cf. [BGT, Th. 3.9.1]), we see that $U \in \Pi_\ell$ with ℓ -index 1. Finally, by [BGT, Th. 3.6.10] (with slow decrease replaced by slow increase), we obtain (1.2), completing the proof. •

§3. Mercerian theorems.

For positive f , the *upper order* $\rho(f)$ is defined by

$$\rho(f) := \limsup_{x \rightarrow \infty} \frac{\log f(x)}{\log x}.$$

In [BI1, BI2] we recently proved the following Mercerian counterpart to Theorem 1.2:

THEOREM 3.1. *Let ν, f, F_ν be as in Theorem 1.1. We write $\rho := \rho(f)$. If*

$$-\nu - \frac{3}{2} < \rho < 0, \quad \rho \neq -\frac{1}{2}$$

and

$$x^{-1} F_\nu(1/x)/f(x) \rightarrow c > 0 \quad (x \rightarrow \infty),$$

then $c = 2^{\rho+\frac{1}{2}} \Gamma(\frac{3}{4} + \frac{\nu}{2} + \frac{\rho}{2}) / \Gamma(\frac{1}{4} + \frac{\nu}{2} - \frac{\rho}{2})$ and $f \in R_\rho$: f varies regularly with index ρ .

The question arises of proving the Mercerian counterpart of Theorem 1.1, that is, the implication from (1.4) to (1.2), under mild assumptions. We leave this question open here, but we think that the techniques of [BI1, BI2] will be helpful in this problem too. For corresponding results for Laplace transforms, see Embrechts [E] and [BGT, Th. 5.4.1].

Appendix

We complete the proof above by showing

$$\begin{aligned} & \int_0^1 \frac{1}{u^{\nu+\frac{3}{2}}} \{u^{1/2} J_\nu(u) - \frac{u^{\nu+\frac{1}{2}}}{2^\nu \Gamma(\nu+1)}\} du + \int_1^\infty \frac{1}{u^{\nu+\frac{3}{2}}} u^{1/2} J_\nu(u) du \\ &= \frac{1}{2^\nu \Gamma(\nu+1)} \left\{ \log 2 - \frac{\gamma}{2} + \frac{\psi(\nu+1)}{2} \right\}. \end{aligned} \quad (A1)$$

By Weber's integral ([W, §13.24 (1); cf. [BI1, §1]) for the Mellin transform of the Bessel function, for $x \in (-\nu - \frac{3}{2}, \nu + \frac{1}{2})$,

$$\begin{aligned} & \int_0^1 u^x \{u^{1/2} J_\nu(u) - \frac{u^{\nu+\frac{1}{2}}}{2^\nu \Gamma(\nu+1)}\} du + \int_1^\infty u^x u^{1/2} J_\nu(u) du \\ &= \int_0^\infty u^{-x} \cdot u^{-3/2} J_\nu(1/u) du / u - \frac{1}{2^\nu \Gamma(\nu+1)} \int_0^1 u^{x+\nu+\frac{1}{2}} du \\ &= \frac{1}{2^\nu \Gamma(\nu+1)(x+\nu+\frac{3}{2})} \{g_1(x+\nu+\frac{3}{2}) - 1\}, \end{aligned} \quad (A2)$$

where

$$g_1(x) := \Gamma(\nu+1) \frac{2^x \Gamma(\frac{x}{2} + 1)}{\Gamma(-\frac{x}{2} + \nu + 1)}.$$

We have $g_1(0) = 1$. For $x \in (-2, 2\nu+2)$, we write

$$g_2(x) := \log g_1(x) = x \log 2 + \log \Gamma(\nu+1) + \log \Gamma(1 + \frac{x}{2}) - \log \Gamma(-\frac{x}{2} + \nu + 1).$$

Then

$$g'_1(x)/g_1(x) = g'_2(x) = \log 2 + \frac{1}{2} \psi(1 + \frac{x}{2}) + \frac{1}{2} \psi(-\frac{x}{2} + \nu + 1),$$

and so

$$g_1'(0) = g_2'(0) = \log 2 - \frac{\gamma}{2} + \frac{1}{2}\psi(\nu + 1).$$

Letting $x \downarrow -\nu - \frac{3}{2}$ in (A2), we obtain (A1), as required.

References

- [B1] BINGHAM, N. H.: A Tauberian theorem for integral transforms of Hankel type, *J. London Math. Soc.* (2) **5** (1972), 493-503.
- [B2] BINGHAM, N. H.: Tauberian theorems for Jacobi series, *Proc. London Math. Soc.* (3) **36** (1978), 285-309.
- [BGT] BINGHAM, N. H., GOLDIE, C. M. and TEUGELS, J. L.: *Regular variation* (2nd ed.), *Encycl. Math. Appl.* **27**, Cambridge University Press, 1989 (1st ed. 1987).
- [BI1] BINGHAM, N. H. and INOUE, A.: The Drasin-Shea-Jordan theorem for Fourier and Hankel transforms, *Quart. J. Math.* (2) **48** (1997), to appear (Statistics Reports 95/1, Birkbeck College, University of London).
- [BI2] BINGHAM, N. H. and INOUE, A.: The Drasin-Shea-Jordan theorem for Hankel transforms of arbitrarily large order, *Statistics Reports 96/9*, Birkbeck College, University of London; submitted.
- [E] EMBRECHTS, P.: A second-order theorem for Laplace transforms, *J. London Math. Soc.* (2) **17** (1978), 102-106.
- [I1] INOUE, A.: On Abel-Tauber theorems for Fourier cosine transforms, *J. Math. Anal. Appl.* **196** (1995), 764-776.
- [I2] INOUE, A.: An Abel-Tauber theorem for Fourier sine transforms, *J. Math. Sci. Univ. Tokyo* **2** (1995), 303-309.
- [RS] RIDENHOUR, J. R. and SONI, R. P.: Parseval relation and Tauberian theorems for the Hankel transform, *SIAM J. Math. Anal.* **5** (1974), 809-821.
- [SS] SONI, K. and SONI, R. P.: Slowly varying functions and asymptotic behaviour of a class of integral transforms, III, *J. Math. Anal. Appl.* **49** (1975), 612-628.
- [W] WATSON, G. N.: *A treatise on the theory of Bessel functions*, 2nd ed., Cambridge University Press, 1944.
- [WW] WHITTAKER, E. T. and WATSON, G. N.: *Modern analysis*, 4th ed., Cambridge University Press, 1927.

Statistics Department, Birkbeck College (University of London), Malet Street, London
WC1E 7HX, England
n.bingham@stat.bbk.ac.uk

Department of Mathematics, Hokkaido University, Sapporo 060, Japan
inoue@math.hokudai.ac.jp