HOKKAIDO UNIVERSITY

Title	On estimates in Hardy spaces for the Stokes flow in a half space
Author(s)	Giga, Y.; Matsui, S.; Shimizu, Y.
Citation	Hokkaido University Preprint Series in Mathematics, 381, 1-12
Issue Date	1997-05-01
DOI	https://doi.org/10.14943/83527
Doc URL	https://hdl.handle.net/2115/69131
Type	departmental bulletin paper
File Information	re381.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

On Estimates in Hardy Spaces for the Stokes Flow in a Half Space

Y. Giga, S. Matsui and Y. Shimizu

Series #381. May 1997

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

#357 S. Izumiya, Singularities of solutions for first order partial differential equations, 20 pages. 1996.

#358 N. Hayashi, P.I. Naumkin and T. Ozawa, Scattering theory for the Hartree equation, 14 pages. 1996.

#359 I. Tsuda and K. Tadaki, A logic-based dynamical theory for a genesis of biological threshold, 49 pages. 1996.

#360 I. Tsuda and A. Yamaguchi, Singular-continuous nowhere-differentiable attractors in neural systems, 40 pages. 1996.

#361 M. Nakamura and T. Ozawa, Low energy scattering for nonlinear Schrödinger equations in fractional order Sobolev spaces, 17 pages. 1996.

#362 I. Nakamura, Hilbert schemes and simple singularities E_6, E_7 and E_8, 21 pages. 1996.

#363 T. Mikami, Equivalent conditions on the central limit theorem for a sequence of probability measures on R, 7 pages. 1996.

#364 S. Izumiya and T. Sano, Generic affine differential geometry of space curves, 23 pages. 1996.

#365 T. Tsukada, Stability of reticular optical caustics, 12 pages. 1996.

#366 A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of a generalized spin-boson model, 40 pages. 1996.

#367 A. Arai, A class of representations of the *-algebra of the canonical commutation relations over a Hilbert space and instability of embedded eigenvalues in quantum field models, 12 pages. 1996.

#368 K. Ito, BV-solutions of a hyperbolic-elliptic system for a radiating gas, 33 pages. 1997.

#369 M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, 20 pages. 1997.

#370 N.H. Bingham and A. Inoue, An Abel-Tauber theorem for Hankel transforms, 8 pages. 1997.

#371 T. Nakazi and H. Sawada, The commutator ideal in Toeplitz algebras for uniform algebras and the analytic structure, 9 pages. 1997.

#372 M.-H. Giga and Y. Giga, Stability for evolving graphs by nonlocal weighted curvature, 70 pages. 1997.

#373 T. Nakazi, Brown-Halmos type theorems of weighted Toeplitz operators, 14 pages. 1997.

#374 J. Inoue and S.-E. Takahashi, On characterizations of the image of Gelfand transform of commutative Banach algebras, 30 pages. 1997.

#375 L. Solomon and H. Terao, The double coxeter arrangement, 21 pages. 1997.

#376 G. Ishikawa and T. Morimoto, Solution surfaces of Monge-Ampère equations, 15 pages. 1997.

#377 G. Ishikawa, A relative transversality theorem and its applications, 16 pages. 1997.

#378 J. Inoue and T. Nakazi, On the zeroes of solutions of an extremal problem in H^1, 14 pages. 1997.

#379 Y. Giga and K. Ito, On pintching of curves moved by surface diffusion, 12 pages. 1997.

#380 F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field, 39 pages. 1997.

On Estimates in Hardy Spaces for the Stokes Flow in a Half Space

YOSHIKAZU GIGA†, SHIN'YA MATSUI‡, YASUYUKI SHIMIZU

Department of Mathematics, Hokkaido University, Sapporo 060, Japan
‡Faculty of Buisiness Adminstration and Information Science,
Hokkaido Information University, Nishi-Nopporo, Ebetshu 069, Japan

§0 Introduction

We consider the Stokes equation

$$
\begin{aligned}
(0.1) \qquad & u_t - \Delta u + \nabla p = 0, \operatorname{div} u = 0 \text{ in } \Omega \times (0,\infty), \\
& u = u_0 \text{ at } t = 0, \\
& u = 0 \text{ on } \partial\Omega \times (0,\infty)
\end{aligned}
$$

in a domain Ω in $\mathbb{R}^n (n \geq 2)$ with smooth boundary. Here $u = (u^1, \ldots, u^n)$ is unknown velocity field and p is unknown pressure field. Initial data u_0 is assumed to satisfy a *compatibility condition* : $\operatorname{div} u_0 = 0$ in Ω and the normal component of u_0 equals zero on $\partial\Omega$. This system is a typical parabolic equation and it has several properties resembling to the heat equation.

If $\Omega = \mathbb{R}^n$, u is reduced to a solution of the heat equation with initial data u_0 because there is no boundary condition. For example regularity-decay estimate

$$
(0.2) \qquad \|\nabla u(t)\|_p \leq C t^{-1/2} \|u_0\|_p \text{ for } t > 0
$$

holds for all $1 \leq p \leq \infty$ with C independent of t and u_0, where $\|f(t)\|_p := \left(\int_\Omega |f(t,x)|^p dx\right)^{1/p}$ and ∇ denotes the gradient in space variables. If $p = 2$, the estimate (0.2) is still valid for any domain. Indeed, since the Stokes operator A is self–adjoint and nonnegative, the operator A generates an analytic semigroup e^{-tA}. This yields

$$
\|A^{1/2} e^{-tA} u_0\|_2 \leq C t^{-1/2} \|u_0\|_2.
$$

Since $u = e^{-tA} u_0$ and $\|A^{1/2} u\|_2 = \|\nabla u\|_2$, (0.2) follows for $p = 2$.(See Borchers and Miyakawa [3] for applications.) For $1 < p < \infty$, (0.2) is valid for bounded domains (Giga [7]) and for a half space (Ukai [13]). The estimate (0.2) is also valid for exterior domain with $n \geq 3$, with extra restriction $1 < p < n$.(See Borchers and Miyakawa [2], Giga and Sohr [8], Iwashita [10].)

†Partly supported by NISSAN SCIENCE FOUNDATION and the Japan Ministry of Education, Science, Sports, and Culture through Grant No.08874005 ‡Partly supported by the Japan Ministry of Education, Science, Sports, and Culture through Grant No.08640135

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

However, there was no result for $p = 1$ or $p = \infty$ where the boundary of Ω is not empty. The main difficulty lies in the fact that the projection associated with the Helmholtz decomposition is not bounded in L^1 type spaces, because it involves the singular integral operator such as Riesz operators. Nevertheless in this paper, we prove (0.2) for $p = 1$ where Ω is a half space $\mathbb{R}^n_+ = \{x = (x_1, \cdots, x_n); x_n > 0\}$.

Theorem 0.1. *Let u be the solution of the Stokes equation (0.1) in $\Omega = \mathbb{R}^n_+$ with initial data $u_0 \in L^1(\mathbb{R}^n)$, which satisfies the compatibility condition. Then there is a constant C independent of u_0 such that*

$$\|\nabla u(t)\|_1 \le Ct^{-1/2}\|u_0\|_1 \tag{0.3}$$

for all $t > 0$.

This is rather surprising since we do not expect $\|u(t)\|_1 \le C\|u_0\|_1$ for $\Omega = \mathbb{R}^n_+$. Actually, the estimate (0.3) follows from a stronger estimate:

Theorem 0.2. *Under the same hypothesis of the Theorem 0.1, there is a constant C' independent of u_0 such that*

$$\|\nabla u(t)\|_{\mathcal{H}^1(\mathbb{R}^n_+)} \le C't^{-1/2}\|u_0\|_1 \tag{0.4}$$

for all $t > 0$.

Here

$$\|f\|_{\mathcal{H}^1(\mathbb{R}^n_+)} = \inf\{\|\tilde{f}\|_{\mathcal{H}^1(\mathbb{R}^n)}; \tilde{f} \in \mathcal{H}^1(\mathbb{R}^n),\ \tilde{f}|_{\mathbb{R}^n_+} \equiv f\},$$

where $\mathcal{H}^1(\mathbb{R}^n)$ is the Hardy space in $\mathbb{R}^n$ defined later.

Combining the Sobolev inequality with (0.3), we have

$$\|u(t)\|_{n/(n-1)} \le C_0 t^{-1/2}\|u_0\|_1 \tag{0.5}$$

with C_0 independent of $t > 0$ and u_0. This has been already proved by Borchers and Miyakawa [1] where a general $L^p - L^q$ estimate

$$\|u(t)\|_p \le C_0 t^{-\alpha}\|u_0\|_q$$

with $\alpha = (n/2)(1/q - 1/p)$ has been proved for all $1 \le q < p \le \infty$ where $\Omega = \mathbb{R}^n_+$. Their method does not depend on (0.3). For $1 < q < p < \infty$, such estimate has been proved by Ukai [12]. There is an extensive literature on $L^p - L^q$ estimate for exterior domain Ω ($n \ge 3$) (e.g. Giga and Sohr [9], Borchers and Miyakawa [2], Iwashita [10], Chen [4]) but the case $q = 1$ and $p = \infty$ is included only in Chen [4] for $n = 3$.

To show (0.4), we recall the solution formula obtained by Ukai [13]. The solution is represented by the Gauss kernel and various Riesz operators. It is known by Carpio [4] that the solution $u = G_t * u_0$ of the heat equation with initial data $u_0 \in L^1(\mathbb{R}^n)$ enjoys

$$\|\nabla u(t)\|_{\mathcal{H}^1(\mathbb{R}^n)} \le C_1 t^{-1/2}\|u_0\|_1 \tag{0.6}$$

where G_t is the Gauss kernel. If the solution of (0.1) were represented only by G_t and a Riesz operator in $\mathbb{R}^n$, (0.6) could yield (0.4) since the Riesz operator is bounded in $\mathcal{H}^1$. Unfortunately, the formula contains the Riesz operator in tangential variables $x' = (x_1, \dots, x_{n-1})$ to $\partial\mathbb{R}^n_+$, it is not clear that such operators are bounded in $\mathcal{H}^1(\mathbb{R}^n)$. To overcome this difficulty, we rewrite Ukai's formula so that ∇u does not have tangential Riesz operators with use of the operator Λ whose symbol equals $|\xi'|$, where $(\xi', \xi_n) = \xi \in \mathbb{R}^n$. Because of this, we need to prove

$$\|\Lambda u(t)\|_{\mathcal{H}^1(\mathbb{R}^n)} \le C_2 t^{-1/2} \|u_0\|_1 \tag{0.7}$$

in addition to (0.5). Although there are several extra technical difficulty, because of the formula, this is a rough idea for the proof of (0.4).

§1. The solution formula

In this section we rearrange the solution formula for (0.1) obtained by Ukai [13] for later use.

First, we establish conventions of notations. For an n-dimensional vector a, we denote a tangential component $(a_1, \dots, a_{n-1})$ by $a' \in \mathbb{R}^{n-1}$, so that $a = (a', a_n)$. We set $\partial_j = \partial/\partial x_j$ and let $\nabla' = (\partial_1, \cdots, \partial_{n-1})$. Hereafter, C denotes a positive constant which may differ from one occasion to another.

Let $\mathcal{F}$ be the Fourier transform in $\mathbb{R}^n$:

$$\mathcal{F}f(\xi) = \int_{\mathbb{R}^n} e^{-ix\cdot\xi} f(x)dx,$$

and let $\hat{f}$ be the Fourier tranform of f in the tangential space:

$$\hat{f}(\xi', x_n) = \int_{\mathbb{R}^{n-1}} e^{-ix'\cdot\xi'} f(x', x_n)dx'.$$

The Riesz operators R_j $(j = 1, \dots, n)$, S_j $(j = 1, \dots, n-1)$, and the operator Λ are defined by

$$\mathcal{F}(R_j f)(\xi) = \frac{i\xi_j}{|\xi|}\mathcal{F}f(\xi),$$
$$\mathcal{F}(S_j f)(\xi) = \frac{i\xi_j}{|\xi'|}\mathcal{F}f(\xi),$$
$$\mathcal{F}(\Lambda f)(\xi) = |\xi'|\mathcal{F}f(\xi).$$

We set $R' = (R_1, \dots, R_{n-1})$, $S = (S_1, \dots, S_{n-1})$ and define U by

$$Uf = rR' \cdot S(R' \cdot S + R_n)e,$$

where r is the restriction operator from $\mathbb{R}^n$ to $\mathbb{R}^n_+$, and e is the extension operator from $\mathbb{R}^n_+$ over $\mathbb{R}^n$ with value 0, that is,

$$ef = \begin{cases} f & \text{for } x_n \ge 0, \\ 0 & \text{for } x_n < 0. \end{cases}$$

We also define the operator $E(t)$ and $F(t)$ by

$$[E(t)f](x) = \int_{\mathbb{R}^n_+} \{G_t(x-y) - G_t(x'-y', x_n+y_n)\} f(y)dy,$$

$$[F(t)f](x) = \int_{\mathbb{R}^n_+} \{G_t(x-y) + G_t(x'-y', x_n+y_n)\} f(y)dy,$$

where G_t is the Gauss kernel $G_t(x) = (4\pi t)^{-n/2} e^{-|x|^2/4t}$. Note that $E(t)f$ (resp. $F(t)f$) is the solution to the heat equation in $\mathbb{R}^n_+$ with Dirichlet (resp. Neumann) data;

$$\begin{aligned} z_t - \Delta z &= 0 \text{ in } \mathbb{R}^n_+ \times (0,T), \\ z|_{t=0} &= f, \\ z|_{x_n=0} &\equiv 0. \text{ (resp. } \partial_n z|_{x_n=0} = 0.) \end{aligned}$$

We recall the formula obtained by Ukai.

Theorem 1.1(Ukai). *The solution to* (1.1) *can be expressed as*

$$u^n = UE(t)V_1u_0, \tag{1.1a}$$

$$u' = E(t)V_2u_0 - SUE(t)V_1u_0, \tag{1.1b}$$

where $V_1u_0 = -S \cdot u_0' + u_0^n$ *and* $V_2u_0 = u_0' + Su_0^n$.

We give a formal proof of Theorem 1.1 for the reader's convenience. By (0.1a) and (0.1b), we get $\Delta p = 0$ in $\mathbb{R}^n_+$. Applying the tangential Fourier transform, the equation $\Delta p = 0$ is reduced to an ordinary differential equation $(\partial_n^2 - |\xi'|^2)\hat{p} = 0$. Assuming that p is bounded, we get $(\partial_n + |\xi'|)\hat{p} = 0$. We set $v^n = (\partial_n + \Lambda)u^n$ and $v' = V_2u = u' + Su^n$. Then v satisfies $v_t - \Delta v = 0$, $v^n|_{t=0} = \Lambda V_1u_0$, $v'|_{t=0} = V_2u_0$, and $v|_{x_n=0} = 0$. Thus v solves the heat equation in $\mathbb{R}^n_+$ with zero Dirichlet data. Solving v with some manipulations leads (1.1).

To solve our problem, we rewrite the formula (1.1). Note that the vector field u in (1.2) is given as a restriction $r\bar{u}$ of a vector field $\bar{u} = (\bar{u}', \bar{u}_n)$ of form

$$\bar{u}^n = R' \cdot S(R' \cdot S + R_n)eE(t)V_1u_0, \tag{1.2a}$$

$$\bar{u}' = E(t)V_2u_0 - SR' \cdot S(R' \cdot S + R_n)eE(t)V_1u_0. \tag{1.2b}$$

Lemma 1.2. *Let* j *be an integer with* $1 \le j \le n$. *Assume that* $\mathrm{div}u_0 = 0$ *in* $\mathbb{R}^n_+$ *when* $j = n$. *Then the first space derivative of* $\bar{u}$ *are expressed as*

$$\begin{aligned} \partial_j\bar{u}^n = &- R_j\{R' \cdot \Lambda eE(t)u_0' - R_n\nabla' \cdot eE(t)u_0' \\ &+ R' \cdot \nabla' eE(t)u_0^n + R_n\Lambda eE(t)u_0^n\}, \end{aligned} \tag{1.3a}$$

$$\begin{aligned} \partial_j\bar{u}' = &\partial_j E(t)u_0' + w_j \\ &+ R_j\{R'(\nabla' \cdot eE(t)u_0') - R_n\nabla'(\nabla'\Lambda^{-1} \cdot eE(t)u_0') \\ &- R'\Lambda eE(t)u_0^n + R_n\nabla' eE(t)u_0^n\}, \end{aligned} \tag{1.3b}$$

where

$$w_j = \begin{cases} \partial_j \nabla' \Lambda^{-1} E(t) u_0^n & \text{for } 1 \le j \le n-1, \\ -\nabla'(\nabla' \cdot \Lambda^{-1} F(t) u_0') & \text{for } j = n. \end{cases} \tag{1.4}$$

Proof. To show (1.3), it is convenient to use the Fourier transformation by $\partial_j \bar{u}$ in (1.2). Note that the operator S_j and $eE(t)$ are commutable. Then we get

$$\begin{aligned}
\mathcal{F}(\partial_j \bar{u}^n) &= i\xi_j \frac{i\xi'}{|\xi|} \cdot \frac{i\xi'}{|\xi'|} \left(\frac{i\xi'}{|\xi|} \cdot \frac{i\xi'}{|\xi'|} + \frac{i\xi_n}{|\xi|} \right) \left(-\frac{i\xi'}{|\xi'|} \cdot \mathcal{F}(eE(t)u_0') + \mathcal{F}(eE(t)u_0^n) \right) \\
&= -\frac{i\xi_j}{|\xi|} \left\{ \left(\frac{i\xi'}{|\xi|} |\xi'| - \frac{i\xi_n}{|\xi|} i\xi' \right) \cdot \mathcal{F}(eE(t)u_0') + \left(\frac{i\xi'}{|\xi|} \cdot i\xi' + \frac{i\xi_n}{|\xi|} |\xi'| \right) \mathcal{F}(eE(t)u_0^n) \right\}, \\
\mathcal{F}(\partial_j \bar{u}') &= i\xi_j \left(\mathcal{F}(E(t)u_0') + \frac{i\xi'}{|\xi'|} \mathcal{F}(E(t)u_0^n) \right) \\
&\quad - i\xi_j \frac{i\xi'}{|\xi'|} \left(\frac{i\xi'}{|\xi|} \cdot \frac{i\xi'}{|\xi'|} \right) \left(\frac{i\xi'}{|\xi|} \cdot \frac{i\xi'}{|\xi'|} + \frac{i\xi_n}{|\xi|} \right) \left(-\frac{i\xi'}{|\xi'|} \cdot \mathcal{F}(eE(t)u_0') + \mathcal{F}(eE(t)u_0^n) \right) \\
&= i\xi_j \mathcal{F}(eE(t)u_0') - \frac{\xi_j \xi'}{|\xi'|} \mathcal{F}(eE(t)u_0^n) \\
&\quad + \frac{i\xi_j}{|\xi|} \left\{ \frac{i\xi'}{|\xi|} \xi' \cdot \mathcal{F}(eE(t)u_0) - \frac{i\xi_n}{|\xi|} i\xi' \left(i\xi' \cdot \frac{1}{|\xi'|} \mathcal{F}(eE(t)u_0') \right) \right. \\
&\quad \left. - \left(\frac{i\xi'}{|\xi|} |\xi'| - \frac{i\xi_n}{|\xi|} i\xi' \right) \mathcal{F}(eE(t)u_0^n) \right\}.
\end{aligned}$$

By the inverse Fourier transform,the first identity implies (1.3a). To show (1.3b), we must handle the term $i\xi_j(i\xi'/|\xi'|)\mathcal{F}[E(t)u_0^n]$. Taking the inverse Fourier transform, this term is transformed to $\partial_j \nabla' \Lambda^{-1} E(t) u_0^n$. For $1 \le j \le n-1$, this equals to w_j. For $j = n$, we invoke the assumption $\mathrm{div} u_0 = 0$, so that $\partial_n u_0^n = -\nabla' \cdot u_0'$:

$$\begin{aligned}
\partial_n \nabla' \Lambda^{-1} E(t) u_0^n &= \partial_n \nabla' \Lambda^{-1} \int_{\mathbb{R}^n_+} \{ G_t(x-y) - G_t(x'-y', x_n + y_n) \} u_0^n(y) dy \\
&= \nabla' \Lambda^{-1} \int_{\mathbb{R}^n_+} \left\{ \frac{\partial}{\partial x_n} G_t(x-y) - \frac{\partial}{\partial x_n} G_t(x'-y', x_n+y_n) \right\} u_0^n(y) dy \\
&= \nabla' \Lambda^{-1} \int_{\mathbb{R}^n_+} \left\{ -\frac{x_n - y_n}{2t} G_t(x-y) + \frac{x_n + y_n}{2t} G_t(x'-y', x_n+y_n) \right\} u_0^n(y) dy \\
&= \nabla' \Lambda^{-1} \left\{ \int_{\mathbb{R}^{n-1}} \left[\{ -G_t(x-y) - G_t(x'-y', x_n+y_n) \} u_0^n(y) \right]_{y_n=0}^{y_n=+\infty} dy' \right. \\
&\quad \left. + \int_{\mathbb{R}^n_+} \{ G_t(x-y) + G_t(x'-y', x_n+y_n) \} \partial_n u_0^n(y) dy \right\} \\
&= -\nabla' \Lambda^{-1} \int_{\mathbb{R}^n_+} \{ G_t(x-y) + G_t(x'-y', x_n+y_n) \} \nabla' \cdot u_0'(y) dy \\
&= -\nabla'(\nabla' \cdot \Lambda^{-1} F(t) u_0') = w_n. \quad \square
\end{aligned}$$

§2. Proof of theorem

To prove Theorem 0.1, we need to estimate the right hand side of (1.3) in $L^1(\mathbb{R}^n)$. In this section we estimate these terms in the Hardy space $\mathcal{H}^1$, which is the subspace of L^1, instead of L^1. We recall the definition of the Hardy space $\mathcal{H}^1$. Note that the following definition is one of many equivalent definitions of the Hardy space. (See Fefferman and Stein [6].)

Definition 2.1. A function $f \in L^1(\mathbb{R}^n)$ belongs to the Hardy space $\mathcal{H}^1 = \mathcal{H}^1(\mathbb{R}^n)$ if

$$f^*(x) = \sup_{s>0} |G_s * f(x)| \in L^1(\mathbb{R}^n),$$

where the symbol $*$ denotes the convolution with respect to the space variable x. The norm of $f \in \mathcal{H}^1(\mathbb{R}^n)$ is defined by

$$\|f\|_{\mathcal{H}^1} := \|f^*\|_{L^1(\mathbb{R}^n)}$$

Here, we remark that a L^1 function f belongs to $\mathcal{H}^1$ if and only if its Riesz transform $R_j f$ belongs to $L^1(\mathbb{R}^n)$ for all j, and that

$$\|f\|_{\mathcal{H}^1} \cong \|f\|_{L^1(\mathbb{R}^n)} + \sum_{j=1}^{n} \|R_j f\|_{L^1(\mathbb{R}^n)} \texttt{(equivalent norm)}.$$

For the convenience, we denote the operator norm of R_j in $\mathcal{H}^1$ by $|||\cdot|||_{\mathcal{H}^1}$.

To estimate (1.3) in $\mathcal{H}^1$, we require the following lemma.

Lemma 2.2. *Let K be an integral operator of form*

$$Kf(x) = \int_{\mathbb{R}^n} k(x,y) f(y) dy \text{ for } x \in \mathbb{R}^n. \tag{2.1}$$

If the kernel $k(x,y)$ satisfies that

$$\sup_{y \in \mathbb{R}^n} \|k(\cdot, y)\|_{\mathcal{H}^1} = k_0 < \infty,$$

then K is a bounded operator from $L^1(\mathbb{R}^n)$ to $\mathcal{H}^1(\mathbb{R}^n)$ i.e.

$$\|Kf\|_{\mathcal{H}^1} \le k_0 \|f\|_{L^1(\mathbb{R}^n)}. \tag{2.2}$$

Proof. By definition of $\mathcal{H}^1$,

$$\begin{aligned} (Kf)^*(x) &= \sup_{s>0} \left| \int_{\mathbb{R}^n} G_s(x-z) \int_{\mathbb{R}^n} k(z,y) f(y) dy dz \right| \\ &\le \sup_{s>0} \left| \int_{\mathbb{R}^n} f(y) \int_{\mathbb{R}^n} G_s(x-z) k(z,y) dz dy \right| \\ &\le \int_{\mathbb{R}^n} |f(y)| \left\{ \sup_{s>0} \left| \int_{\mathbb{R}^n} G_s(x-z) k(z,y) dz \right| \right\} dy. \end{aligned} \tag{2.3}$$

Integrating (2.3) by x,

$$\begin{aligned} \|Kf\|_{\mathcal{H}^1} &\le \int_{\mathbb{R}^n} |f(y)| |||k(\cdot, y)|||_{\mathcal{H}^1} dy \\ &\le k_0 \|f\|_{L^1(\mathbb{R}^n)}. \quad \square \end{aligned}$$

We next show several pointwise estimates on the heat kernel.

Lemma 2.3. *Assume that real parameters l and m satisfy $0 \le l \le n$ and $m \ge 0$. Then there exists a constant $C = C_{l,m}$ which does not depend on $x \in \mathbb{R}^n$ and $t \ge 0$ such that*

(2.4a) $|\partial_j G_t(x)| \le Ct^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}$ *for* $1 \le j \le n$ *with* $n \ge 2$,

(2.4b)
$|\partial_j \partial_k \Lambda^{-1} G_t(x)| \le Ct^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}$ *for* $1 \le j,k \le n-1$ *with* $n \ge 3$,

(2.4c) $|\Lambda G_t(x)| \le Ct^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}$ *with* $n \ge 2$.

In (2.4a), *the restriction* $l \le n$ *is unnecessary.*

Proof. We first prove (2.1a). Since $\partial_j G_t(x) = -(x_j/2t)G_t(x)$ and $e^{-|x|^2/4t} \le C|t^{-1/2}x|^{-\alpha}$ for $\alpha \ge 0$, we have

$$\begin{aligned}(2.5)\qquad \partial_j G_t(x) &= -\frac{x_j}{2t}G_t(x)\\ &= -\frac{x_j}{2t^{n/2+1}}e^{-|x'|^2/4t}e^{-|x_n|^2/4t}\\ &\le Ct^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}.\end{aligned}$$

We next show (2.4b). Note that Λ^{-1} is equal to $(-\Delta')^{-1/2} = \left(\sum_{k=1}^{n-1}\partial_k^2\right)^{-1/2}$, so the integral kernel of Λ^{-1} is $c_n|x'|^{-n+2}$ for $n \ge 3$, where c_n is some positive constant. Therefore we have

$$(2.6)\qquad \partial_j\partial_k\Lambda^{-1}G_t(x) = c_n\partial_j\partial_k\int_{\mathbb{R}^{n-1}}|x'-y'|^{-n+2}G_t(y',x_n)dy'.$$

Set $x = t^{1/2}z$ to get

$$\partial_{x_j}\partial_{x_k}\Lambda^{-1}G_t(x) = t^{-(n+1)/2}\partial_{z_j}\partial_{z_k}\Lambda^{-1}G_1(z).$$

So it is sufficient to show (2.4b) for $t = 1$, i.e.

$$(2.7)\qquad |\partial_j\partial_k\Lambda^{-1}G_1(z)| \le C|z'|^{-l}|z_n|^{-m}.$$

In fact, if (2.7) is valid, then we have

$$\begin{aligned}|\partial_{x_j}\partial_{x_k}\Lambda^{-1}G_t(x)| &= t^{-(n+1)/2}|\partial_{z_j}\partial_{z_k}\Lambda^{-1}G_1(z)|\\ &\le Ct^{-(n+1)/2}|z'|^{-l}|z_n|^{-m}\\ &= Ct^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}\end{aligned}$$

for any $t > 0$.

Let ψ_1 be a smooth function in $\mathbb{R}^{n-1}$ such that $0 \le \psi_1 \le 1$, supp $\psi \subset \{|z'| \le 1\}$, and $\psi_1|_{|z'|<1/2} \equiv 1$. Set $\psi_2 = 1 - \psi_1$. Then

(2.8)
$$\begin{aligned}\partial_j\partial_k\Lambda^{-1}G_1(z) = \frac{C}{(4\pi)^{n/2}}e^{-z_n^2/4}\Big\{&\partial_j\partial_k\int_{\mathbb{R}^{n-1}}\frac{\psi_1(z'-y')}{|z'-y'|^{n-2}}e^{-|y'|^2/4}dy'\\ &+\partial_j\partial_k\int_{\mathbb{R}^{n-1}}\frac{\psi_2(z'-y')}{|z'-y'|^{n-2}}e^{-|y'|^2/4}dy'\Big\}\\ = Ce^{-z_n^2/4}\{&I_1(z')+I_2(z')\}.\end{aligned}$$

The estimate of the term I_1: We have

$$I_1(z') = \partial_j\partial_k \int_{|y'|\le 1} \frac{\psi_1(y')}{|y'|^{n-2}} e^{-|z'-y'|^2/4} dy' \tag{2.9}$$
$$= \int_{|y'|\le 1} \frac{\psi_1(y')}{|y'|^{n-2}} K_{j,k}(z'-y')dy,$$

where

$$K_{j,k}(z') = \left(\frac{z_j z_k}{4} - \frac{\delta_{j,k}}{2}\right) e^{-|z'|^2/4}$$

and $\delta_{j,k}$ is Kronecker's delta. Recalling $|z'-y'| \le |z'|+1$ and $|z'-y'|^2 \ge |z'|^2/2-1$ holds for $|y'| \le 1$, we get

$$\begin{aligned} |K_{j,k}(z'-y')| &\le \left\{\frac{(|z'|+1)^2}{4} + \frac{1}{2}\right\} e^{-(|z'|^2-2)/8} \\ &= \frac{e^{1/4}}{4}\left\{(|z'|+1)^2+2\right\} e^{-|z'|^2/8} \\ &\le C|z'|^{-l}. \end{aligned}$$

Hence we have

$$\begin{aligned} |I_1(z')| &\le C\int_{|y'|\le 1} \frac{1}{|y'|^{n-2}} |z'|^{-l} dy' \\ &\le C|z'|^{-l} \end{aligned}$$

The estimate of the term I_2: We have

$$I_2(z') = \int_{\mathbb{R}^{n-1}} \frac{(\partial_j\partial_k\psi_2)(z'-y')}{|z'-y'|^{n-2}} e^{-|y'|^2/4} dy' \tag{2.10}$$
$$- (n-2)\Big\{\int_{\mathbb{R}^{n-1}} (\partial_j\psi_2)(z'-y') \frac{z_k-y_k}{|z'-y'|^n} e^{-|y'|^2/4} dy'$$
$$+ \int_{\mathbb{R}^{n-1}} (\partial_k\psi_2)(x'-y') \frac{z_j-y_j}{|z'-y'|^n} e^{-|y'|^2/4} dy'\Big\}$$
$$+ \int_{\mathbb{R}^{n-1}} \psi_2(z'-y') L_{j,k}(z'-y') e^{-|y'|^2/4} dy'$$
$$= J_1(z') - (n-1)J_2(z') + J_3(z'),$$

where

$$L_{j,k}(z') = (n-2)\left\{n\frac{x_j x_k}{|z'|^{n+2}} - \frac{\delta_{j,k}}{|z'|^n}\right\}.$$

Since the support of $\partial_j\psi_2$ and $\partial_j\partial_k\psi_2$ are included in $1/2 \le |z| \le 1$, the estimates

of J_1 and J_2 can be obtained like as the estimate of I_1:

$$|J_1(z')| = \left| \int_{1/2\le|y'|\le 1} \frac{(\partial_j\partial_k\psi_2)(y')}{|y'|^{n-2}} e^{-|z'-y'|^2/4} dy' \right| \tag{2.11}$$
$$\le \|\nabla^2\psi_2\|_{L^\infty} \int_{1/2\le|y'|\le 1} \frac{1}{|y'|^{n-2}} e^{-(|z'|^2-2)/8} dy'$$
$$\le C|z'|^{-l},$$

$$|J_2(z')| \le \|\nabla\psi_2\|_{L^\infty} \int_{1/2\le|y'|\le 1} \frac{1}{|y'|^{n-1}} e^{-(|z'|^2-2)/8} dy' \tag{2.12}$$
$$\le C|z'|^{-l}.$$

To estimate the term J_3, we use the inequality $|z'|^l \le C_l(|z'-y'|^l + |y'|^l)$. Since $|L_{j,k}(z')| \le \frac{C}{|z'-y'|^{n+1}}$, we get

(2.13)
$$|J_3(z')| \le C|z'|^{-l} \int_{|z'-y'|\ge 1/2} \left(\frac{|z'-y'|^l}{|z'-y'|^n} + \frac{|y'|^l}{|z'-y'|^n} \right) e^{-|y'|^2/4t} dy'$$
$$\le C|z'|^{-l} \int_{|z'-y'|\ge 1/2} (2^{l-n} + 2^n|y'|^l) e^{-|y'|^2/4} dy'$$
$$= C|z'|^{-l}.$$

Combining the estimate (2.11), (2.12), and (2.13), we get $|I_2(z')| \le C|z'|^{-l}$ and

$$|\partial_j\partial_k\Lambda^{-1}G_1(z)| \le Ce^{-x_n^2/4}|z'|^{-l} \tag{2.14}$$
$$\le C_{l,m}|z'|^{-l}|z_n|^{-m}.$$

This proves (2.7) for $n \ge 3$.

The estimate (2.4c) for $n \ge 3$ is easily obtained by the fact that Λ is equal to $(-\Delta')\Lambda^{-1} = -(\partial_1^2 + \cdots \partial_{n-1}^2)\Lambda^{-1}$ and by applying (2.4b).

Finally, we show (2.4c) for $n = 2$. Note that Λ is equal to $|\partial_1| = \partial_1 S_1$. So we have

$$\Lambda G_t(x) = \partial_1 S_1 G_t(x) \tag{2.15}$$
$$= \partial_1 \frac{1}{\pi} \lim_{\epsilon\downarrow 0} \int_{|y_1|>\epsilon} \frac{1}{y_1} G_t(x_1 - y_1, x_2) dy_1.$$

(See Torchinsky [12], p.266.) Integrating by parts, we get

$$\int_{|y_1|>\epsilon} \frac{1}{y_1} G_t(x_1 - y_1, x_2) dy_1 = \Big[\log|y_1| G_t(x_1 - y_1, x_2)\Big]_\epsilon^\infty$$
$$+ \Big[\log|y_1| G_t(x_1 - y_1, x_2)\Big]_{-\infty}^{-\epsilon}$$
$$- \int_{|y_1|>\epsilon} \log|y_1| \partial_{y_1} G_t(x_1 - y_1, x_2) dy_1$$
$$= \log\epsilon\big(G_t(x_1+\epsilon, x_2) - G_t(x_1-\epsilon, x_2)\big)$$
$$+ \int_{|y_1|>\epsilon} \log|y_1| \frac{x_1 - y_1}{2t} G_t(x_1 - y_1, x_2) dy_1.$$

Sending $\epsilon \downarrow 0$, we get

$$\Lambda G_t(x) = \frac{1}{\pi}\partial_1 \int_{-\infty}^{\infty} \log|y_1| \frac{x_1 - y_1}{2t} G_t(x_1 - y_1, x_2) dy_1. \tag{2.16}$$

Set $x = t^{1/2}z$ and $y = t^{1/2}w$. Then we have

$$\begin{aligned}(\Lambda G_t)(x) &= \frac{1}{\pi} t^{-1/2}\partial_{z_1} \int_{-\infty}^{\infty} (\log|w_1| + \log t^{1/2}) \frac{z_1 - w_1}{2t^{1/2}} t^{-1} G_1(z_1 - w_1, w_2) t^{1/2} dw_1 \\ &= t^{-3/2}(\Lambda G_1)(z).\end{aligned}$$

So it is sufficient to show (2.4c) for $t = 1$.

$$\begin{aligned}\Lambda G_1(z) &= \frac{1}{\pi}\frac{1}{4\pi} e^{-z_2^2/4}\partial_1 \Big\{ \int_{|y_1|<1} \log|y_1| \frac{z_1 - y_1}{2} e^{-(z_1-y_1)^2/4} dy_1 \\ &\quad + \int_{|y_1|>1} \log|y_1| \frac{z_1 - y_1}{2} e^{-(z_1-y_1)^2/4} dy_1 \Big\} \\ &= \frac{1}{4\pi^2} e^{-z_2^2/4}(I_1(z_1) + I_2(z_1)).\end{aligned} \tag{2.17}$$

The estimate of I_1: We have

$$I_1(z_1) = \int_{-1}^{1} \log|y_1| \frac{1}{2}\Big(1 - \frac{|z_1 - y_1|^2}{2}\Big) e^{-(z_1-y_1)^2/4} dy_1.$$

As the same suggestion to (2.11), we obtain

$$\begin{aligned}|I_1(z_1)| &\le \frac{1}{2}\int_{-1}^{1} |\log|y_1|| \Big(1 + \frac{(|z_1|+1)^2}{4}\Big) e^{-\frac{|z_1|^2}{8} + \frac{1}{4}} dy_1 \\ &\le C(1 + |z_1|^2) e^{-|z_1|^2/8}.\end{aligned} \tag{2.18}$$

The estimate of I_2: The method is similar to the case $n \ge 3$. Integrating by parts,

$$\begin{aligned}I_2(z_1) &= \partial_1 \Big\{ \Big[\log|y_1| e^{-(z_1-y_1)^2/4}\Big]_1^{+\infty} \\ &\quad + \Big[\log|y_1| e^{-(z_1-y_1)^2/4}\Big]_{-\infty}^{-1} \\ &\quad - \int_{|y_1|>1} \frac{1}{y_1} e^{-(z_1-y_1)^2/4} dy_1 \Big\} \\ &= \int_{|y_1|>1} \frac{1}{y_1}\frac{z_1 - y_1}{2} e^{-(z_1-y_1)^2/4} dy_1 \\ &= e^{-(z_1+1)^2/4} - e^{-(z_1-1)^2/4} + \int_{|y_1|>1} \frac{1}{y_1^2} e^{-(z_1-y_1)^2/4} dy_1.\end{aligned}$$

We set $w_1 = z_1 - y_1$ and obtain

$$I_2(z_1) = e^{-(z_1+1)^2/4} - e^{-(z_1-1)^2/4} + \int_{|z_1-w_1|>1} \frac{1}{(z_1-w_1)^2} e^{-w_1^2/4} dw_1.$$

Using $|z_1|^l \le C(|z_1-w_1|^l + |w_1|^l)$, we obtain

(2.19)
$$\begin{aligned} |I_2(z_1)| &\le |e^{-(z_1+1)^2/4}| + |e^{-(z_1-1)^2/4}| \\ &\quad + C\int_{|z_1-w_1|>1} \frac{1}{|z_1|^l} \left(|z_1-w_1|^{l-2} + \frac{|w_1|^l}{|z_1-w_1|^2} \right) e^{-|w_1|^2/4} dw_1 \\ &\le C|z|^{-l} \end{aligned}$$

since $l \le 2$ so that $|z_1 - w_1|^{l-2} \le 1$. Combining the estimate (2.18) and (2.19),we obtain (2.4c) for $n = 2$. □

We are now ready to show the key lemma for the main theorem.

Lemma 2.4. *Assume a function $a = a(x)$ is in $L^1(\mathbb{R}^n_+)$. Then*

(2.20a) $$\|\partial_j E(t)a\|_{\mathcal{H}^1} \le Ct^{-1/2}\|a\|_{L^1(\mathbb{R}^n_+)} \text{ for } 1 \le j \le n,$$

(2.20b) $$\|\partial_j\partial_k \Lambda^{-1} eE(t)a\|_{\mathcal{H}^1} \le Ct^{-1/2}\|a\|_{L^1(\mathbb{R}^n_+)} \text{ for } 1 \le j,k \le n-1,$$

(2.20c) $$\|\Lambda eE(t)a\|_{\mathcal{H}^1} \le Ct^{-1/2}\|a\|_{L^1(\mathbb{R}^n_+)},$$

(2.20d) $$\|\partial_j\partial_k \Lambda^{-1} F(t)a\|_{\mathcal{H}^1} \le Ct^{-1/2}\|a\|_{L^1(\mathbb{R}^n_+)} \text{ for } 1 \le j,k \le n-1.$$

Proof. To show (2.20a,b,c), we extend the function $a(x)$ from $\mathbb{R}^n_+$ over $\mathbb{R}^n$ with $a(x', x_n) = -a(x', -x_n)$ for $x_n < 0$. Then

$$\begin{aligned} [E(t)a](x) &= G_t * a(x) \\ &= \int_{\mathbb{R}^n} G_t(x-y)a(y)dy, \\ [eE(t)a](x) &= \theta(x_n)[E(t)a](x), \end{aligned}$$

where θ is the Heaviside function i.e.

$$\theta(x_n) = \begin{cases} 1 & \text{for } x_n \ge 0, \\ 0 & \text{for } x_n < 0. \end{cases}$$

Since $G_s * (\partial_j G_t)(x) = \partial_j G_{s+t}(x)$, the estimate (2.4a) implies

$$|G_s * (\partial_j G_t)(x)| \le C(s+t)^{(l+m-n-1)/2}|x'|^{-l}|x_n|^{-m}$$

for any nonnegative l and m. Thus, for $0 \le l+m \le n+1$ we have

$$(\partial_j G_t)^*(x) \le Ct^{(l+m-n-1)/2}|x'|^{-l}|x'|^{-m}.$$

Therefore we obtain

$$\|\partial_j G_t\|_{\mathcal{H}^1} \le \sum_{k=1}^{4} C_{l,m} t^{(l+m-n-1)/2} \int_{\Omega_k} |x'|^{-l}|x_n|^{-m}dx, \tag{2.21}$$

where $\Omega_1 = \{|x'| \le t^{1/2}, |x_n| \le t^{1/2}\}$, $\Omega_2 = \{|x'| > t^{1/2}, |x_n| \le t^{1/2}\}$, $\Omega_3 = \{|x'| \le t^{1/2}, |x_n| > t^{1/2}\}$ and $\Omega_4 = \{|x'| > t^{1/2}, |x_n| > t^{1/2}\}$. For each integration of (2.21), we take suitable l and m such that $l = m = 0$ in Ω_1, $l = n$, $m = 0$ in Ω_2, $l = 0$, $m = 2$ in Ω_3 and $l = n - 1/2$, $m = 3/2$ in Ω_4. We thus observe that the right hand side of (2.21) is estimated from above by constant times $t^{-1/2}$. Thus (2.20a) is obtained. The estimate is obtained by Carpio [3, Lemma 2.1] but the proof is misprinted in [3, p.457 line 4], so we gave the proof.

To prove (2.20b), we put $k(x,y) = \partial_j\partial_k\Lambda^{-1}\theta(x_n)G_t(x-y)$. Then

(2.22)

$$(G_s * k(\cdot,y)(x)) = \int_{\mathbb{R}^n} G_s(z-x)k(z,y)dz$$

$$= \frac{1}{(4\pi s)^{n/2}}\frac{1}{(4\pi t)^{n/2}} \int_{\mathbb{R}^{n-1}} e^{-|z'-x'|^2/4s}\partial_j\partial_k\Lambda^{-1}e^{-|z'-y'|^2/4t}dz' \times$$

$$\int_0^{+\infty} e^{-|z_n-x_n|^2/4s}e^{-|z_n-y_n|^2/4t}dz_n.$$

Since the integrand in the last integral in (2.22) is nonnegative, we get

$$|(G_s * k(\cdot,y)(x)| \le |\partial_j\partial_k\Lambda^{-1}G_{s+t}(x)|.$$

By (2.4b) calculation similar to derive (2.21) yields

$$\sup_y \|k(\cdot,y)\|_{\mathcal{H}^1} \le Ct^{-1/2}$$

for $n \ge 3$ and for $n = 2$ with $j = k = 1$. Applying Lemma 2.2 we get (2.20b,c). Note that (2.20b) agrees with (2.20c) if $n = 2$.

The estimate (2.20d) is obtained in the same way as above but this time we have to extend $a(x)$ as an even function in x_n, i.e. $a(x', x_n) = a(x', -x_n)$ for $x_n < 0$. □

We are now ready to prove Theorem 0.2. By Lemma 1.2 and Lemma 2.4,

$$\begin{aligned}
\|\partial_j\bar{u}_n\|_{\mathcal{H}^1} &\le |||R_j|||_{\mathcal{H}^1}\Big\{\sum_{k=1}^{n-1}|||R_k|||_{\mathcal{H}^1}(\|\Lambda eE(t)u_0^k\|_{\mathcal{H}^1} + \|\partial_k eE(t)u_0^n\|_{\mathcal{H}^1}) \\
&\quad + |||R_n|||_{\mathcal{H}^1}(\|\nabla\cdot eE(t)u_0'\|_{\mathcal{H}^1} + \|\Lambda eE(t)u_0^n\|_{\mathcal{H}^1})\Big\} \\
&\le Ct^{-1/2}\|u_0\|_{L^1(\mathbb{R}^n_+)}, \\
\|\partial_j\bar{u}'\|_{\mathcal{H}^1} &\le Ct^{-1/2}\|u_0\|_{L^1(\mathbb{R}^n_+)}.
\end{aligned}$$

Since $u = \bar{u}|_{\mathbb{R}^n_+}$, we now get

$$\|\nabla u\|_{L^1(\mathbb{R}^n_+)} \le \|\nabla u\|_{\mathcal{H}^1(\mathbb{R}^n_+)} \le \|\nabla\bar{u}\|_{\mathcal{H}^1} \le Ct^{-1/2}\|u_0\|_{L^1(\mathbb{R}^n_+)}.$$

The proof is complete.

REFERENCES

1. W. Borchers and T. Miyakaya, *L^2 decay for the Navier-Stokes flow in halfspaces*, Math. Ann. **282** (1988), 139–155.
2. ———, *Algebraic L^2 decay for Navier-Stokes flows in exterior domains*, Acta Math. **165** (1990), 189–227.
3. ———, *Algebraic L^2 decay for Navier-Stokes flows in exterior domains, II*, Hiroshima Math. J. **21** (1991), 621–640.
4. A.Carpio, *Large time behavior in incompressible Navier-Stokes equations*, SIAM J. Math. Anal. **27** (1996), 449-475.
5. Z.-M.Chen, *Solution of the stationary and nonstationary Navier-Stokes equations in exterior domains*, Pacific J. Math. **159** (1993), 227–240.
6. C.Fefferman and E.Stein, *$\mathcal{H}^p$ spaces of several variavles*, Acta Math. **129** (1972), 137–197.
7. Y.Giga, *Analyticity of the semigroup generated by the Stokes operator in L^r spaces*, Math. Z. **178** (1981), 297–329.
8. Y.Giga and H.Sohr, *On the Stokes operator in exterior domains*, J. Fac. Sci. Univ. Tokyo, Sect. IA Math. **36** (1989), 103–130.
9. Y.Giga and H.Sohr, *Abstract L^p estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains*, J. of Functional Analysis **102** (1991), 72–94.
10. H.Iwashita, *$L_q - L_r$ estimate for solutions of the nonstationary Stokes equations in an exterior domain and the Navier-Stokes initial value problems in L_q spaces*, Math. Ann. **285** (1989), 265–288.
11. T.Miyakawa, *Hardy spaces of solenoidal vector fields, with application to the Navier-Stokes equations*, Kyushu J. Math. **50** (1997), 1–64.
12. A.Torchinsky, *Real-variable methods in harmonic anarysis*, Academic Press, 1986.
13. S.Ukai, *A solution formula for the Stokes equation in $\mathbb{R}^n_+$*, Comm. Pure Appl. Math. **XL** (1987), 611–621.