



# HOKKAIDO UNIVERSITY

|                  |                                                                                   |
|------------------|-----------------------------------------------------------------------------------|
| Title            | Remarks on convergence of evolving graphs by nonlocal curvature                   |
| Author(s)        | Giga, M.-H; Giga, Y.                                                              |
| Citation         | Hokkaido University Preprint Series in Mathematics, 388, 1-18                     |
| Issue Date       | 1997-09-01                                                                        |
| DOI              | <a href="https://doi.org/10.14943/83534">https://doi.org/10.14943/83534</a>       |
| Doc URL          | <a href="https://hdl.handle.net/2115/69138">https://hdl.handle.net/2115/69138</a> |
| Type             | departmental bulletin paper                                                       |
| File Information | re388.pdf                                                                         |



**Remarks on convergence of  
evolving graphs by nonlocal curvature**

**M.-H. Giga and Y. Giga**

**Series #388. September 1997**

**HOKKAIDO UNIVERSITY**  
**PREPRINT SERIES IN MATHEMATICS**

- #363 T. Mikami, Equivalent conditions on the central limit theorem for a sequence of probability measures on  $R$ . 7 pages. 1996.
- #364 S. Izumiya and T. Sano, Generic affine differential geometry of space curves, 23 pages. 1996.
- #365 T. Tsukada, Stability of reticular optical caustics, 12 pages. 1996.
- #366 A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of a generalized spin-boson model, 40 pages. 1996.
- #367 A. Arai, A class of representations of the  $*$ -algebra of the canonical commutation relations over a Hilbert space and instability of embedded eigenvalues in quantum field models, 12 pages. 1996.
- #368 K. Ito, BV-solutions of a hyperbolic-elliptic system for a radiating gas, 33 pages. 1997.
- #369 M. Nakamura and T. Ozawa, Nonlinear Schrödinger equations in the Sobolev space of critical order, 20 pages. 1997.
- #370 N.H. Bingham and A. Inoue, An Abel-Tauber theorem for Hankel transforms, 8 pages. 1997.
- #371 T. Nakazi and H. Sawada, The commutator ideal in Toeplitz algebras for uniform algebras and the analytic structure, 9 pages. 1997.
- #372 M.-H. Giga and Y. Giga, Stability for evolving graphs by nonlocal weighted curvature, 70 pages. 1997.
- #373 T. Nakazi, Brown-Halmos type theorems of weighted Toeplitz operators, 14 pages. 1997.
- #374 J. Inoue and S.-E. Takahashi, On characterizations of the image of Gelfand transform of commutative Banach algebras, 30 pages. 1997.
- #375 L. Solomon and H. Terao, The double coxeter arrangement, 21 pages. 1997.
- #376 G. Ishikawa and T. Morimoto, Solution surfaces of Monge-Ampère equations, 15 pages. 1997.
- #377 G. Ishikawa, A relative transversality theorem and its applications, 16 pages. 1997.
- #378 J. Inoue and T. Nakazi, On the zeroes of solutions of an extremal problem in  $H^1$ , 14 pages. 1997.
- #379 Y. Giga and K. Ito, On pinching of curves moved by surface diffusion, 12 pages. 1997.
- #380 F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field, 39 pages. 1997.
- #381 Y. Giga, S. Matsui and Y. Shimizu, On estimates in Hardy spaces for the Stokes flow in a half space, 13 pages. 1997.
- #382 A. Arai, A new estimate for the ground state energy of Schrödinger operators, 12 pages. 1997.
- #383 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order, 24 pages. 1997.
- #384 K. Ito, Asymptotic stability of planar rarefaction wave for scalar viscous conservation law, 8 pages. 1997.
- #385 A. Arai, Representation-theoretic aspects of two-dimensional quantum systems in singular vector potentials: canonical commutation relations, quantum algebras, and reduction to lattice quantum systems, 32 pages. 1997.
- #386 P. Aviles and Y. Giga, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields, 21 pages. 1997.
- #387 T. Nakazi and T. Yamamoto, Norms of some singular integral operators and their inverse operators, 28 pages. 1997.

MI-HO GIGA and YOSHIKAZU GIGA

## Remarks on convergence of evolving graphs by nonlocal curvature

### 1. Introduction

This is a continuation of the work of the authors [GG1-5] on graph-like solutions for motion driven by nonlocal weighted curvature in the plane.

We consider a fully nonlinear evolution equation in one space dimension:

$$u_t + F(u_x, \Lambda_W(u)) = 0 \quad (1.1)$$

with  $\Lambda_W(u) = (W'(u_x))_x$ . Here  $W$  is a given convex function on  $\mathbf{R}$  and its derivative  $W'$  may have jumps ;  $F$  is a given continuous function satisfying monotonicity condition:

$$F(p, X) \leq F(p, Y) \quad \text{for } X \geq Y \quad (1.2)$$

so that the equation (1.1) is at least degenerate parabolic. The subscripts  $t$  and  $x$  denote partial differentiation in time and space variables, respectively. If  $W(p) = p^2/2$  and  $F(p, X) = -X$ , (1.1) is the heat equation. If  $W(p) = (1 + p^2)^{1/2}$ ,  $\Lambda_W(u)$  is the curvature of the graph of  $u(t, \cdot)$  so  $\Lambda_W$  is often called the weighted curvature. The value  $\Lambda_W$  is actually unchanged by adding an affine function to  $W$  but we still denote it by  $\Lambda_W$  rather than  $\Lambda_{W''}$ . If  $W(p) = (1 + p^2)^{1/2}$  and  $F(p, X) = -(1 + p^2)^{1/2}X$ , then (1.1) is the curve shortening equation for the graph of  $u(t, \cdot)$ . We are interested in studying the case when  $W'$  has jump discontinuities. A typical example is  $W(p) = |p|/2$ . In this case the meaning of solutions of (1.1) is unclear even if  $u$  is smooth, since  $\Lambda_W(u) = \delta(u_x)u_{xx}$  is not a well-defined distribution because of the presence of the Dirac  $\delta$  measure.

To circumvent this inconvenience we introduced in [GG1, 3] a notion of generalized solution for (1.1) by extending the theory of viscosity solutions [CIL]. In [GG3] we established comparison principles and the unique global existence of generalized solutions for a given periodic continuous initial data, although  $\Lambda_W(u)$  turns to be a nonlocal quantity. As anticipated, our solution is obtained by the limit of solutions of smoother strictly parabolic problem approximating the original problem [GG5].

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

We shall first state it in a rigorous way. Let  $\mathcal{E}$  be the subset of the space of convex functions of form

$$\mathcal{E} = \{W; W \text{ is convex in } \mathbf{R} \text{ and } W \text{ is } C^2 \text{ except some discrete set } P.$$

$$\text{Moreover, } \sup_{K \setminus P} W'' = c_K < \infty \text{ for every compact set } K \text{ in } \mathbf{R}\}. \quad (1.3)$$

Of course,  $\mathcal{E}$  includes all piecewise linear convex functions. The set  $P$  is called the *singularity* of  $W$ . Let  $\mathcal{F} = \mathcal{F}_T$  be the set of all continuous function  $F : [0, T) \times \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$  satisfying the monotonicity condition (1.2) for  $F(t, p, X)$  with respect to the last variable  $X$ . Let  $\mathcal{F}_\# = \mathcal{F}_{\#T}$  be the set of  $F \in \mathcal{F}$  which is either time independent or uniformly continuous on  $[0, T'] \times [-K_0, K_0] \times \mathbf{R}$  for each  $T' < T$ ,  $K_0 > 0$ . The last assumption together with boundedness of  $c_K$  in (1.3) seems to be technical in the next convergence theorem but we do not attempt to remove it. We consider the initial value problem with periodic boundary condition to avoid extra technicality; see [GG3] for other boundary conditions. Let  $C(\mathbf{T})$  denote the space of all continuous functions on  $\mathbf{T} = \mathbf{R}/\omega\mathbf{Z}$ , i.e. the space of  $\omega$ -periodic continuous functions.

**Convergence Theorem 1.1.** ([GG5]) *For  $F_\varepsilon \in \mathcal{F}_{\#T}, W_\varepsilon \in \mathcal{E}$  with  $\varepsilon \geq 0$  assume that  $F_\varepsilon \rightarrow F_0, W_\varepsilon \rightarrow W_0$  locally uniformly as  $\varepsilon \rightarrow 0$  in  $[0, T) \times \mathbf{R} \times \mathbf{R}$  and  $\mathbf{R}$ , respectively. For  $\varepsilon > 0$  let  $u^\varepsilon \in C([0, T) \times \mathbf{T})$  be the generalized solution of*

$$\begin{cases} u_t + F_\varepsilon(t, u_x, \Lambda_{W_\varepsilon}(u)) = 0 & \text{in } (0, T) \times \mathbf{R}, \\ u(0, x) = u_0^\varepsilon(x), & x \in \mathbf{R} \end{cases} \quad (1.4)$$

*with  $u_0^\varepsilon \in C(\mathbf{T})$ . If  $u_0^\varepsilon \rightarrow u_0$  in  $C(\mathbf{T})$ , then  $u^\varepsilon$  converges to a function  $u \in C([0, T) \times \mathbf{T})$  locally uniformly in  $[0, T) \times \mathbf{T}$  and  $u$  is a generalized solution of*

$$\begin{cases} u_t + F_0(t, u_x, \Lambda_{W_0}(u)) = 0 & \text{in } (0, T) \times \mathbf{R}, \\ u(0, x) = u_0(x), & x \in \mathbf{R}. \end{cases} \quad (1.5)$$

*The constant  $T$  may be taken as  $+\infty$ .*

Theorem 1.1 is quite general because it allows any degeneracy of  $W_0'', W_\varepsilon''$ . This justifies a way to construct a solution  $u$  of (1.5) with general  $W_0 \in \mathcal{E}, F_0 \in \mathcal{F}_\#$  by approximating  $W_0$  by  $W_\varepsilon \in C^\infty \cap \mathcal{E}$  with  $W_\varepsilon'' > 0$  and  $F_0$  by  $F_\varepsilon \in C^\infty \cap \mathcal{F}_\#$  for  $\varepsilon > 0$ , so that the problem (1.4) is strictly parabolic and it is solvable by the classical theory (cf. Proposition 2.3).

In this note as an application of Theorem 1.1 we study the convergence of derivatives  $u_x^\epsilon$ . The key ingredient is the  $L^1(\mathbf{T})$  estimate of second derivatives, or more generally

$$\|(\xi(u_x^\epsilon))_x\|_{L^1(\mathbf{T})}(t) = \int_0^\omega |(\xi(u_x^\epsilon))_x(t, x)| dx \leq \|\xi(u_{0x}^\epsilon)_x\|_{L^1(T)} \quad (1.6)$$

for any  $C^1$  nondecreasing function  $\xi$ . It is not difficult to prove this estimate for smooth solution with smooth  $F_\epsilon$  and  $W_\epsilon$  but Theorem 1.1 passes such an estimate to a general problem. This shows that Theorem 1.1 is useful. As noted in [GG5] Theorem 1.1 is also essential to prove the convergence of crystalline algorithm (initiated by S. Angenent and M. Gurtin [AG] and J. Taylor [T] and analyzed by P. Girão and R. Kohn [GK1-2]) for a general equation (1.5). The convergence results by [GK1-2] as well as [FG] apply, for example, the heat equation and curve shortening equation. In [GK1]  $L^2$  convergence of derivatives of solutions by crystalline algorithm is proved with rate of convergence for these examples with both the Neumann and Dirichlet boundary conditions. Our convergence of derivatives is  $L^p$  convergence for any  $p \geq 1$  for a general problem with no rate of convergence.

It turns out that Theorem 1.1 is powerful to get a priori estimates (1.6) with  $\xi(\sigma) = \sigma$  of solutions by analyzing crystalline algorithm. In this paper we extend crystalline algorithm in [GK1] for a general problem (1.5). We moreover derive a priori estimate (1.6) with  $\xi(\sigma) = \sigma$ , which yields the convergence of derivative  $u_x^\epsilon$ , without using any analytic theory of parabolic equations other than Theorem 1.1. This is one of main goals of this paper.

In crystalline algorithm one restricts a class of solutions to a special class of piecewise linear function to construct solutions [AG], [T]. The problem roughly corresponds to the case when  $W$  is piecewise linear. The main ansatz is that each piecewise linear part (called a facet) of the graph of solution  $u(t, \cdot)$  stays as a facet and does not change its slope and that only its end points move. This ansatz is natural and it is regarded as a generalized solution in our sense [GG2]. However, if  $F$  depends on space variable, such an assumption that a facet stays as a facet of the same slope (unless it disappears) is turns to be unnatural. In this paper we give an example that the comparison principle may not hold if 'facet-stay-facet assumption' is imposed.

For physical background of problems the reader is referred to [GG3] and [GG5] and references cited there especially a book of M. Gurtin [Gu], reviews of J. Taylor,

J. Cahn and C. Handwerker [TCH] and of P. Girão and R. Kohn [GK2]. The bibliography of [GG3] and [GG5] includes many references related to recent work on the motion by crystalline energy and nonlocal curvature. We won't repeat to cite these articles unless necessary in this note. This note is written so that no explicit definition of generalized solutions is necessary. To read this note we only need to admit that classical solutions for strictly parabolic problems are generalized solutions.

We are grateful to Professor D. Hilhorst for informing us of the reference [O]. The second author was partly supported by the NISSAN SCIENCE FOUNDATION and the Japanese Ministry of Education, Science, Sports and Culture through grant no. 08874005, 09242201 for scientific research.

## 2. Convergence of derivatives

We state a priori estimate (1.6) in its general form. For a finite (signed) Radon measure  $\mu$  we denote its total variation by  $\|\mu\|_1$  since  $\|\mu\|_1$  equals  $L^1(\mathbf{T})$  norm if  $\mu$  is absolutely continuous with respect to the Lebesgue measure. We shall simply write  $L^p(\mathbf{T})$  norm of  $f$  by  $\|f\|_p$ . The partial derivative in  $\mathbf{x}$  is taken in the distribution sense.

**Theorem 2.1 (A priori estimate).** *Let  $u \in C([0, T) \times \mathbf{T})$  be a generalized solution (1.5) with  $u_0 \in C(\mathbf{T})$ , where  $W_0 \in \mathcal{E}$ ,  $F_0 \in \mathcal{F}_{\#T}$ . Let  $\xi$  be a nondecreasing  $C^1$  function on  $\mathbf{R}$ . Assume that  $u_{0\mathbf{x}\mathbf{x}}$  is a finite Radon measure (so that  $u_{0\mathbf{x}} \in L^\infty(\mathbf{T})$  i.e.,  $u_0$  is Lipschitz). Then*

$$\|u_{\mathbf{x}}\|_\infty(t) \leq \|u_{0\mathbf{x}}\|_\infty, \quad 0 \leq t < T, \quad (2.1)$$

$$\|(\xi(u_{\mathbf{x}}))_{\mathbf{x}}\|_1(t) \leq \|(\xi(u_{0\mathbf{x}}))_{\mathbf{x}}\|_\infty, \quad 0 \leq t < T. \quad (2.2)$$

*In particular,*

$$\|\Lambda_{W_0}(u)\|_1(t) \leq \|\Lambda_{W_0}(u_0)\|_1, \quad 0 \leq t < T$$

*if  $W_0 \in C^2$ , where  $\Lambda_{W_0}(u) = W_0'(u_{\mathbf{x}})_{\mathbf{x}}$ .*

To show (2.1)-(2.2) we first prove them for smooth solutions with smooth  $W_0$  and  $F_0$ .

**Proposition 2.2.** *In addition to the hypotheses of Theorem 2.1 assume that  $W_0$  is smooth and that  $F_0(t, \cdot, \cdot)$  is smooth in  $\mathbf{R} \times \mathbf{R}$  for each  $t$ . Assume that  $u$  is smooth in  $[0, T) \times \mathbf{T}$ . Assume that  $\xi' > 0$ . Then the estimates (2.1)-(2.2) are valid.*

We approximate the original problem (1.5) by smooth strictly parabolic problem so that there exists a unique smooth solution.

**Proposition 2.3.** *Assume that  $W_0 \in \mathcal{E}$  is smooth and  $W_0'' > 0$  everywhere. Assume that  $F_0 = F_0(t, p, X) \in \mathcal{F}_{\#T}$  is smooth in each variables and that for each  $T' < T$ ,  $K_0 > 0$  there is  $\lambda > 0$  such that  $\partial F_0 / \partial X \leq -\lambda$  on  $[0, T'] \times [-K_0, K_0] \times \mathbf{R}$ . Assume that  $u_0$  is smooth. Then there is a unique classical solution  $u$  of (1.5) which is smooth in  $[0, T) \times \mathbf{T}$ .*

*Proof of Proposition 2.3.* We differentiate (1.5) in  $x$  and set  $v = u_x$  to get

$$\begin{cases} v_t + \partial_x(F_0(t, v, W_0'(v)_x)) = 0, \\ v(0, x) = v_0, \quad v_0 = u_{0x}. \end{cases} \quad (2.3)$$

By the maximum principle we have a priori bound

$$\|v\|_\infty(t) \leq \|v_0\|_\infty, \quad 0 \leq t < T. \quad (2.4)$$

By (2.4) and assumptions of  $F_0$  and  $W_0$  we now apply the theory of uniformly parabolic equations [LSU] to get a unique global smooth solution  $v$  of (2.3) for smooth  $v_0$ . Since the condition  $\int_0^\omega v_0 = 0$  implies  $\int_0^\omega v(t, \cdot) = 0$ , we get smooth periodic solution  $u$  of (1.5) by integrating  $v$  in the space variable  $u(t, x) = \int_0^x v(t, y) dy + u(t, 0)$ .  $\square$

**Lemma 2.4 (Approximation).** *Assume that  $u_0 \in C(\mathbf{T})$  and that  $u_{0xx}$  is a finite Radon measure. Let  $\xi$  be a nondecreasing  $C^\infty$  function on  $\mathbf{R}$  with  $\xi' \geq \delta$  uniformly for some  $\delta > 0$ . Then there is  $u_0^\epsilon \in C^\infty(\mathbf{T})$  ( $\epsilon > 0$ ) that satisfies*

$$\lim_{\epsilon \rightarrow 0} \|(\xi(u_0^\epsilon))_x\|_1 = \|(\xi(u_{0x}))_x\|_1, \quad (2.5)$$

$$\lim_{\epsilon \rightarrow 0} \|u_0^\epsilon - u_{0x}\|_p = 0, \quad 1 \leq p < \infty, \quad (2.6)$$

$$\lim_{\epsilon \rightarrow 0} \|u_0^\epsilon\|_\infty = \|u_{0x}\|_\infty. \quad (2.7)$$

*Sketch of Proof of Lemma 2.4.* We first note that  $\|u_{0xx}\|_1 < \infty$  with  $u_0 \in C(\mathbf{T})$  implies  $u_{0x} \in L^\infty(\mathbf{T})$ . Indeed, by a fundamental formula of calculus we have  $u_{0x}(x) = \int_a^x u_{0xx}(y) dy$  for  $u_0 \in C^2(T)$  if  $u_{0x}(a) = 0$ ; such  $a \in \mathbf{T}$  exists since

$\int_0^\omega u_{0x} = 0$ . This yields  $\|u_{0x}\|_\infty \leq \|u_{0xx}\|_1$ . Extension to general functions is standard by approximation argument as in [Giu].

Since  $\xi(u_{0x}) \in L^\infty(\mathbf{T})$ , we approximate  $\xi(u_{0x})$  by  $\theta_\varepsilon \in C^\infty(\mathbf{T})$  (see e.g. [Giu]) so that  $\theta_\varepsilon \rightarrow \xi(u_{0x})$  in  $L^p(\mathbf{T})$  ( $1 \leq p < \infty$ ),  $\|\theta_{\varepsilon x}\|_1 \rightarrow \|(\xi(u_{0x}))_x\|_1$ ,  $\|\theta_\varepsilon\|_\infty \rightarrow \|\xi(u_{0x})\|_\infty$  as  $\varepsilon \rightarrow 0$ . We take  $w_\varepsilon = \xi^{-1}(\theta_\varepsilon)$  to get  $w_\varepsilon \rightarrow u_{0x}$  in  $L^p(\mathbf{T})$  ( $p \geq 1$ ) since  $\xi' \geq \delta$  implies  $|\xi(v) - \xi(w)| \geq \delta|v - w|$ . Unfortunately  $w_\varepsilon$  may not satisfy  $\int_0^\omega w_\varepsilon = 0$ . However, since  $\int_0^\omega u_{0x} = 0$ , the convergence  $w_\varepsilon \rightarrow u_{0x}$  implies  $\lim_{\varepsilon \rightarrow 0} \int_0^\omega w_\varepsilon = 0$ . We get  $v_\varepsilon = w_\varepsilon - c_\varepsilon$  with  $c_\varepsilon = \omega^{-1} \int_0^\omega w_\varepsilon$  to get  $\int_0^\omega v_\varepsilon = 0$  and  $v_\varepsilon \rightarrow u_{0x}$  in  $L^p(\mathbf{T})$  with  $\|v_\varepsilon\|_\infty \rightarrow \|u_{0x}\|_\infty$ . Moreover,  $\|(\xi(v_\varepsilon))_x\|_1 \rightarrow \|(\xi(u_{0x}))_x\|_1$  as  $\varepsilon \rightarrow 0$ , since  $\|(\xi(w_\varepsilon))_x\|_1 \rightarrow \|(\xi(u_{0x}))_x\|_1$  and

$$\|(\xi(w_\varepsilon))_x - (\xi(v_\varepsilon))_x\|_1 \leq \|\xi'(v_\varepsilon + c_\varepsilon) - \xi'(v_\varepsilon)\|_\infty \|v_{\varepsilon x}\|_1 \rightarrow 0.$$

The last convergence follows since  $\|v_\varepsilon\|_\infty$  and  $\|v_{\varepsilon x}\|_1$  is bounded and  $c_\varepsilon \rightarrow 0$ . If we set  $u_0^\varepsilon(x) = \int_0^x v_\varepsilon(y)dy + u_0(0)$ , then  $\int_0^\omega v_\varepsilon = 0$  implies  $u_0^\varepsilon \in C^\infty(\mathbf{T})$ . By constructions it is clear that  $u_0^\varepsilon$  satisfies (2.5)-(2.7).  $\square$

*Proof of Theorem 2.1.* We first note that  $\|u_{0x}\|_\infty \leq \|u_{0xx}\|_1$  so that  $u_{0x} \in L^\infty$ . Since  $\|u_{0xx}\|_1 < \infty$  we may assume that  $\xi' \geq \delta > 0$  uniformly for some  $\delta > 0$ . Indeed, assume that  $\|(\xi_\delta(u_x))_x\|_1(t) \leq \|(\xi_\delta(u_{0x}))_x\|_1$  for  $\xi_\delta(\sigma) = \xi(\sigma) + \delta\sigma$ ,  $\delta > 0$ . The right hand side converges to  $\|(\xi(u_{0x}))_x\|_1$  since  $\delta\|u_{0xx}\|_1 \rightarrow 0$  as  $\delta \rightarrow 0$ . By lowersemicontinuity  $\|(\xi(u_x))_x\|_1(t) \leq \liminf_{\delta \rightarrow 0} \|(\xi_\delta(u_x))_x\|_1$ . Thus (2.2) follows for general  $\xi \in C^1$  with  $\xi' \geq 0$ .

For given  $W_0 \in \mathcal{E}$ ,  $F_0 \in \mathcal{F}_{\#T}$  it is not difficult to construct a sequence  $W_\varepsilon, F_\varepsilon$  converging to  $W_0, F_0$  locally uniformly in  $\mathbf{R}$  and  $[0, T) \times \mathbf{R} \times \mathbf{R}$ , respectively as  $\varepsilon \rightarrow 0$ , such that  $W_\varepsilon$  and  $F_\varepsilon$  satisfy all assumptions on  $W_0$  and  $F_0$  in Proposition 2.3. We approximate  $u_0$  by  $u_0^\varepsilon$  by Lemma 2.4. By Proposition 2.2 we get (2.1) and (2.2) for  $u_0^\varepsilon$  and  $u^\varepsilon$ . By Theorem 1.1 and lowersemicontinuity of norms we get (2.1) and (2.2) by sending  $\varepsilon \rightarrow 0$ .  $\square$

*Proof of Proposition 2.2.* We consider (2.3) instead of (1.5). Estimate  $\|u_{xx}\|_1(t) \leq \|u_{0xx}\|_1$  is proved by proving  $L^1$ -contraction property of solutions which is standard; see references of [O] for example. We rather estimate directly, although the idea of approximating sign function by  $\text{sgn}_\delta(p)$  (which equals  $p/\delta$  for  $|p| \leq \delta$  and  $\text{sgn}(p)$  for  $|p| \geq \delta$ ) with  $\delta > 0$  is standard. We set  $\psi_\delta(p) = \int_0^p \text{sgn}_\delta(q)dq$ . For

$w = \xi(v)_x$  we calculate, by integration by parts, to get

$$\frac{d}{dt} \int_0^\omega \psi_\delta(w) dx = \int_0^\omega \operatorname{sgn}_\delta(w) \xi(v)_{xt} dx = - \int_0^\omega \operatorname{sgn}'_\delta(w) w_x \xi(v)_t dx. \quad (2.8)$$

Using the equation (2.3) we get

$$\xi'(v)_t = -\xi'(v) \partial_x (F(t, v, z_x)) = -w F_p(t, v, z_x) - \xi'(v) F_X(t, v, z_x) z_{xx}$$

with  $z = W'(v)$ . Since  $\xi' > 0$ , we rewrite  $z_{xx}$  by using  $w_x$  and  $w$  to get

$$\begin{aligned} \xi(v)_t &= -w A - F_X(t, v, z_x) W''(v) w_x, \\ A &= F_p(t, v, z_x) + \xi'(v) F_X(t, v, z_x) (W''(v) / \xi'(v))_x. \end{aligned}$$

We use this formula in (2.8) to get

$$\begin{aligned} \frac{d}{dt} \int_0^\omega \psi_\delta(w) dx &= \int_0^\omega \operatorname{sgn}'_\delta(w) w_x w A dx \\ &\quad + \int_0^\omega \operatorname{sgn}'_\delta(w) w_x^2 F_X(t, v, z_x) W''(v) dx \\ &\leq \int_0^\omega \operatorname{sgn}'_\delta(w) w_x w A dx =: \sigma_\delta(t) \end{aligned} \quad (2.9)$$

since  $W'' \geq 0$ ,  $F_X \leq 0$  and  $\operatorname{sgn}'_\delta \geq 0$ . By definition of  $\operatorname{sgn}_\delta$  we see  $\|\operatorname{sgn}'_\delta(w) w\|_1(t) \rightarrow 0$  as  $\delta \rightarrow 0$  uniformly in  $[0, T]$ . Since  $A$  and  $w_x$  is smooth in  $[0, T] \times \mathbb{T}$ , this now yields  $\sigma_\delta(t) \rightarrow 0$  uniformly for  $t \in [0, T']$  for each  $T' < T$ . Thus integrating (2.9) yields

$$\int_0^\omega |w|(t, x) dx - \int_0^\omega |w|(0, x) dx \leq \lim_{\delta \downarrow 0} \int_0^t \sigma_\delta(s) ds = 0$$

for all  $0 \leq t < T$ . This yields (2.2). The estimate (2.1) follows from the maximum principle (2.4).  $\square$

**Theorem 2.5 (Convergence of derivatives).** *Assume the same hypotheses of Theorem 1.1. Assume moreover that  $u_{0xx}^\varepsilon$  ( $\varepsilon > 0$ ) is a finite Radon measure with  $\lim_{\varepsilon \rightarrow 0} \|u_{0xx}^\varepsilon\|_1 = \gamma < \infty$ . Then  $\|u_{0xx}\|_1 \leq \gamma$ ,  $\|u_x^\varepsilon\|_\infty(t) \leq \gamma$ ,  $\|u_x\|_\infty(t) \leq \gamma$ . Moreover, for every  $r$ ,  $1 \leq r < \infty$  and  $0 < T' < T$*

$$\lim_{\varepsilon \rightarrow 0} \sup_{0 \leq t \leq T'} \|u_x^\varepsilon - u_x\|_r(t) = 0. \quad (2.10)$$

*Proof.* Since  $\|u_{0x}^\varepsilon\|_\infty \leq \|u_{0xx}^\varepsilon\|_1$  the estimate  $\|u_x^\varepsilon\|_\infty(t) \leq \gamma$  follows from (2.1). Since  $u_0^\varepsilon \rightarrow u_0$  in  $C(\mathbf{T})$  implies  $\|u_{0xx}^\varepsilon\|_1 \leq \lim_{\varepsilon \rightarrow 0} \|u_{0xx}^\varepsilon\|_1$  by duality representation of norms, we have  $\|u_{0xx}\|_1 \leq \gamma$ , so that  $\|u_{0x}\|_\infty \leq \gamma$ , which yields  $\|u_x\|_\infty(t) \leq \gamma$  by (2.1).

If (2.10) were false, then for some  $r, 1 \leq r < \infty$  there would exist sequences  $t_j \in [0, T']$  ( $j = 1, 2, \dots$ ),  $\varepsilon_j \rightarrow 0$  ( $\varepsilon_j > 0$ ) and a constant  $\delta > 0$  and  $t_* \in [0, T']$  such that

$$\|u_x^{\varepsilon_j}(t_j, \cdot) - u_x(t_*, \cdot)\|_r \geq \delta \quad \text{and} \quad t_j \rightarrow t_* \quad (\text{as } j \rightarrow \infty). \quad (2.11)$$

By (2.2) and BV version of Rellich's compactness (see [Giu])  $f_j := u_x^{\varepsilon_j}(t_j, \cdot)$  has a convergent subsequence (still denoted  $f_j$ ) in  $L^r(\mathbf{T})$ . Since  $u_x^{\varepsilon_j}(t_j, \cdot) \rightarrow u_x(t_*, \cdot)$  in  $C(\mathbf{T})$  by Theorem 1.1, the limit of  $f_j$  in  $L^r(\mathbf{T})$  should equal  $u_x(t_*, \cdot)$ . This contradicts (2.11).  $\square$

*Remark 2.6 (Uniqueness of solution of (1.5)).* There are a number of papers on (2.3) even if  $W'$  has jumps. The uniqueness of solutions as well as  $L^1$  contraction property of solutions is well-studied. Although the boundary condition is different, the reader is referred to the article [O] of F. Otto and references cited there. However, if  $W'$  allows jumps together with degeneracy of  $W''$ , the uniqueness of our generalized solution of (1.5) does not follow from existing uniqueness theory (cf. [O] and references cited there) for (2.3) when  $F_0(t, p, X)$  depends on  $X$  nonlinearly even if  $u_0$  is smooth.

### 3. Method by crystalline algorithm

A typical example of an energy  $W$  in  $\mathcal{E}$  is a piecewise linear convex function. Such an energy is called a *crystalline* energy (density). For simplicity we assume that the singularity  $P$  is a finite set, i.e.,  $P = \{p_1 < p_2 < \dots < p_m\}$ . To analyse (1.5) with crystalline  $W = W_0$ , one often considers solutions in a special class of piecewise linear functions. To explain the method we recall a special class of piecewise linear solutions as in [GK1] and [GG2].

For a piecewise linear continuous function  $f$ , a (bounded) closed (nontrivial) interval is called a *faceted region* of  $f$  if it is a maximal interval on which  $f$  is affine. We say that a piecewise linear continuous function is an *admissible crystal* if

- (a) slope  $f_x$  belongs to  $P$ ;

- (b) slope  $f_x$  in adjacent faceted regions should be adjacent in  $P$ , i.e. if  $f_x = p_i$  on a faceted region, then  $f_x$  on each adjacent faceted region equals either  $p_{i-1}$  or  $p_{i+1}$  with  $i+1 \leq m, i-1 \geq 1$ .

We say that  $u \in C(J \times \mathbf{T})$  is an *admissible evolving crystal* (with respect to  $P$ ) in a time interval  $J$  if  $u(t, \cdot)$  is an admissible crystal and (the abscissa of) a jump of  $u_x(t, \cdot)$  moves smoothly in time  $t \in J$ . We have implicitly assumed that each jump does not collide each other. We now recall a system of ordinary differential equations (ODEs) so that an admissible evolving crystal solves (1.5). By periodicity of  $u(t, x)$  in  $x$  there are only finitely many points (for each  $t \in J$ )  $\{x_1(t) < x_2(t) < \dots < x_d(t)\}$  in  $[0, \omega)$  for which any faceted region of  $u(t, \cdot)$  is represented as  $I_j(t) = [x_{j-1}(t), x_j(t)]$ , ( $j = 1, \dots, d$ ) with convention that  $x_j + \omega = x_{j+d}$ . On each  $I_j(t)$ ,  $u_t$  is independent of  $x$  so its value is denoted  $(u_t)_j(t)$  and  $u_x$  is independent of both  $x$  and  $t$  so its value is denoted  $(u_x)_j$ . We say that an admissible evolving crystal  $u \in C(J \times \mathbf{T})$  is an *admissible solution* of (1.5)<sub>1</sub> (with  $W = W_0, F = F_0$ ) if

$$(u_t)_j(t) + F(t, (u_x)_j, \chi_j \Delta((u_x)_j)/L_j(t)) = 0, \quad j = 1, \dots, d, \quad t \in J \quad (3.1)$$

with

$$\Delta(p_i) = W'(p_i + 0) - W'(p_i - 0), \quad p_i \in P,$$

$$L_j(t) = \text{the length of faceted region } I_j(t),$$

where  $\chi_j = 1$  (resp.  $-1$ ) if  $u(t, \cdot)$  is convex (resp. concave) around  $I_j$ ; otherwise  $\chi_j = 0$ . The quantity  $\chi_j \Delta((u_x)_j)/L_j(t)$  corresponding to  $\Lambda_W(u)$  is *nonlocal weighted curvature*. An elementary geometric consideration shows that

$$dL_j(t)/dt = \rho_j^0(u_t)_j + \rho_j^{-1}(u_t)_{j-1} + \rho_j^1(u_t)_{j+1}, \quad j = 1, \dots, d \quad (3.2)$$

with

$$\begin{aligned} \rho_j^0 &= ((u_x)_j - (u_x)_{j-1})^{-1} + ((u_x)_{j+1} - (u_x)_j)^{-1}, \\ \rho_j^{-1} &= -((u_x)_j - (u_x)_{j-1})^{-1}, \quad \rho_j^1 = -((u_x)_{j+1} - (u_x)_j)^{-1}. \end{aligned} \quad (3.3)$$

Since  $\chi_j$  and  $(u_x)_j$  are time independent, the system (3.1), (3.2) yields a system of ODEs for  $L_j$ 's, so that its initial value problem is solvable locally in time at least when  $F \in \mathcal{F}_T$  is  $C^1$ .

We shall prove fundamental properties for admissible solutions.

**Theorem 3.1 (Maximum principle for  $u_t$ ).** Assume that  $F = F(t, p, X) \in C([0, T] \times P \times \mathbf{R})$  satisfies the following: (a)  $F_X \in C([0, T] \times P \times \mathbf{R})$  (so that (3.1)-(3.2) is locally solvable); (b)  $F$  satisfies the monotonicity condition (1.2) with respect to  $X$ ; (c)  $F_t \in C([0, T] \times P \times \mathbf{R})$ . Let  $C_0$  be  $\sup\{|F_t|; [0, T_0] \times P \times \mathbf{R}\}$  with  $0 < T_0 < T$ . Let  $u \in C([0, T_0] \times \mathbf{T})$  be an admissible solution of (1.5) with an admissible crystal  $u_0 \in C(\mathbf{T})$ . Then

$$(i) \min_{1 \leq i \leq d} (u_t)_i(0) - C_0 t \leq (u_t)_j(t) \leq \max_{1 \leq i \leq d} (u_t)_i(0) + C_0 t$$

$$\text{for all } 1 \leq j \leq d, t \in [0, T_0].$$

(ii) If  $\chi_j \neq 0$ , then  $j$ -th faceted region does not disappear at  $t = T_0$ , i.e.,  $x_{j+1}(t) - x_j(t)$  does not converges to zero as  $t \uparrow T_0$  provided that  $\lim_{X \rightarrow \pm\infty} F(t, p, X) = \mp\infty$  for all  $t \in [0, T], p \in P$ .

**Remark 3.2.** (i) Results in Theorem 3.1 generalize the corresponding results in [GK1] when  $F(t, p, X) = -X$ . At some time, say  $t = T_0 > 0$  some facet may disappear but by (ii) only a faceted region with  $\chi_j = 0$  may disappear. Thus as in [GK1] there is no chance that three consecutive faceted regions disappear simultaneously and  $u(T_0 - 0, \cdot)$  is still admissible; see [GG2] for detail. One can solve (3.1)-(3.2) with initial data  $u(T_0 - 0, \cdot)$  which has less numbers of faceted regions. We repeat this procedure renumbering indices of faceted region  $I_j$  and construct  $u$  globally in time. Such  $u$  is called a *weakly admissible* solution of (1.5). (ii) Assumption (a) on  $F$  can be weakened. For regularity in  $X$  we only need to assume that  $F$  is locally Lipschitz in  $X$  with a uniform bound for  $|F_X|$  on  $[0, T'] \times P \times K$  for any  $T' < T$  and any compact set  $K$  in  $\mathbf{R}$  (so that (3.1)-(3.2) is uniquely locally solvable). For assumption (c) of regularity in  $t$  we only need to assume Lipschitz continuity in  $t$  and a bound for  $F_t$  in  $[0, T'] \times P \times \mathbf{R}$  for any  $T' < T$ . In this case  $(u_t)_j(t)$  may not be  $C^1$  but is still locally Lipschitz. However, the proof below can be easily adjusted under this weaker assumption.

**Proof of Theorem 3.1.** The idea of the proof is similar to that of [GK1]; we give it for completeness. Part (ii) follows from (i). Indeed, if a faceted region  $I_j$  with  $\chi_j \neq 0$  disappeared, then  $F(t, (u_x)_j, \chi_j \Delta((u_x)_j)/L_j(t)) \rightarrow +\infty$  or  $-\infty$  as  $t \rightarrow T_0$ . By (3.1) this contradicts (i).

Since  $(u_t)_j$  solves (3.1)-(3.2),  $(u_t)_j(t) \in C^1[0, T_0]$ . For  $U(t) := \max_{1 \leq j \leq d} (u_t)_j(t)$  and  $\hat{t} \in [0, T_0]$ , let  $\hat{j} = j(\hat{t})$  be a number that satisfies  $(u_t)_{\hat{j}}(\hat{t}) = U(\hat{t})$ . Then

$$d(u_t)_{\hat{j}}/dt \leq C_0 \quad \text{at } t = \hat{t}. \quad (3.4)$$

Indeed, if  $\chi_j = 1$  (resp.  $\chi_j = -1$ ) then  $(u_x)_{j-1} < (u_x)_j < (u_x)_{j+1}$  (resp.  $(u_x)_{j-1} > (u_x)_j > (u_x)_{j+1}$ ), so that  $\rho_j^0 > 0, \rho_j^{-1} < 0, \rho_j^1 < 0$  (resp.  $\rho_j^0 < 0, \rho_j^{-1} > 0, \rho_j^1 > 0$ ) for  $\rho_j$ 's in (3.3). By (3.2) and maximality of  $(u_t)_j(\hat{t})$  we now observe that

$$\begin{aligned} (dL_j/dt)(\hat{t}) &= \rho_j^0(u_t)_j(\hat{t}) + \rho_j^{-1}(u_t)_{j-1}(\hat{t}) + \rho_j^1(u_t)_{j+1}(\hat{t}) \\ &\geq (\text{resp. } \leq) \rho_j^0(u_t)_j(\hat{t}) + \rho_j^{-1}(u_t)_j(\hat{t}) + \rho_j^1(u_t)_j(\hat{t}) = 0, \end{aligned}$$

since  $\rho_j^{-1} + \rho_j^1 + \rho_j^0 = 0$ . Thus

$$(d\Lambda_j/dt)(\hat{t}) \leq 0 \quad \text{for} \quad \Lambda_j(t) = \chi_j \Delta((u_x)_j)/L_j(t). \quad (3.5)$$

Differentiating (3.1) in  $t$  and using (3.5) we obtain

$$(d(u_t)_j/dt)(\hat{t}) = -F_t(\hat{t}, (u_x)_j, \Lambda_j(\hat{t})) - F_X(\hat{t}, (u_x)_j, \Lambda_j(\hat{t}))(d\Lambda_j/dt)(\hat{t}) \leq C_0 + 0$$

since  $F_X \leq 0$  by monotonicity of  $F$  in  $X$ . This yields (3.4).

The next lemma (which is not explicit in [GK1]) with  $f_j(t) := (u_t)_j(t) - C_0 t$  implies that  $U(t) - C_0 t$  is a nonincreasing function. Thus  $(u_t)_j(t) \leq \max_{1 \leq i \leq d} (u_t)_i(0) + C_0 t$  for all  $j, 1 \leq j \leq d, t \in [0, T_0]$ . A symmetric argument gives the estimate from below for  $(u_t)_j$  in (i).  $\square$

**Lemma 3.3.** For  $t_0 > 0$  and  $j = 1, 2, \dots, d$  let  $f_j$  be a real-valued locally Lipschitz function on  $[0, t_0]$ . Let  $f$  be  $f(t) = \max_{1 \leq j \leq d} f_j(t)$ . For each  $j$  let  $\Sigma_j$  be of form  $\Sigma_j = \{t \in [0, t_0]; f(t) = f_j(t)\}$ . Assume that  $f'_j(\hat{t}) \leq 0$  for all  $j \in \{1, \dots, d\}$  and a.e.  $\hat{t} \in \Sigma_j$ . Then  $f$  is nonincreasing in  $[0, t_0]$ .

*Proof.* If  $f$  is not nonincreasing, then  $f(a) < f(b)$  for some  $a < b, a, b \in [0, t_0]$ . Since  $f$  is locally Lipschitz continuous, it is almost differentiable and  $0 < f(b) - f(a) = \int_a^b f'(x) dx$ . Thus there is a point  $\hat{s} \in (a, b)$  such that  $f$  together with all  $f_j$ 's is differentiable at  $\hat{s}$  and  $\hat{p} := f'(\hat{s}) > 0$ .

There exist  $i \in \{1, \dots, d\}$  and  $x_\ell \downarrow \hat{s}, x_\ell \in (\hat{s}, b)$  that satisfies  $f(x_\ell) = f_i(x_\ell)$  for all  $\ell = 1, 2, \dots$ . If not, for each  $i$  there is  $\delta_i$  such that  $0 < x - \hat{s} \leq \delta_i$  implies  $f(x) \neq f_i(x)$ . However, this implies  $f(x) \neq f_i(x)$  for all  $i \in \{1, \dots, d\}$  and  $x$  with  $0 < x - \hat{s} \leq \min_{1 \leq i \leq d} \delta_i$ . This contradicts the definition of  $f$ .

Since  $f$  and  $f_i$  are continuous,  $f(\hat{s}) = f_i(\hat{s})$ . Since  $f'_i(\hat{s}) \leq 0$ , for any  $0 < q < \hat{p} = f'(\hat{s})$  it holds that  $f_i(x) - f_i(\hat{s}) \leq q(x - \hat{s})$  for  $x > \hat{s}$  sufficiently close to  $\hat{s}$ . We take  $x = x_\ell$  for sufficiently large  $\ell$  to get

$$q(x_\ell - \hat{s}) \geq f_i(x_\ell) - f_i(\hat{s}) = f(x_\ell) - f(\hat{s}),$$

which contradicts that  $f'(\hat{s}) = \hat{p} > q > 0$ .  $\square$

It is easy to derive a priori estimates like (2.1), (2.2) for weakly admissible solutions.

**Lemma 3.4.** *Assume the same hypotheses of Theorem 3.1 (including (ii)) concerning  $F$ . Let  $u \in C([0, T) \times \mathbb{T})$  be a weakly admissible solution of (1.5) such that  $u$  is admissible on  $[t_\ell, t_{\ell+1})$  for some  $t_0 = 0 < t_1 < \dots < t_h < t_{h+1} = T$  ( $\ell = 0, \dots, h-1$ ) and some faceted region disappears as  $t \uparrow t_\ell$  ( $\ell \geq 1$ ).*

- (i)  $\|u_x\|_\infty(t)$  is constant in  $[t_\ell, t_{\ell+1})$  ( $\ell \geq 0$ ) and may decrease at  $t = t_\ell$  ( $\ell \geq 1$ ).  
In particular,

$$\|u_x\|_\infty(t) \leq \|u_{0x}\|_\infty, \quad t \in [0, T). \quad (3.6)$$

- (ii)  $\|u_{xx}\|_1(t)$  is constant in  $[t_\ell, t_{\ell+1})$  ( $\ell \geq 0$ ) and  $\|u_{xx}\|_1(t_\ell) = \|u_{xx}\|_1(t_\ell + 0) < \|u_{xx}\|_1(t_\ell - 0)$  for  $\ell \geq 1$ . In particular,

$$\|u_{xx}\|_1(t) \leq \|u_{0xx}\|_1, \quad t \in [0, T). \quad (3.7)$$

- (iii)  $\|\Lambda(u)\|_1(t)$  is constant in  $[t_\ell, t_{\ell+1})$  ( $\ell \geq 0$ ) and  $\|\Lambda(u)\|_1(t_\ell) = \|\Lambda(u)\|_1(t_\ell + 0) \leq \|\Lambda(u)\|_1(t_\ell - 0)$  for  $\ell \geq 1$ , where  $\Lambda(u)(t, x) = \Lambda_j(t)$  in (3.5) for  $x \in I_j(t)$ .  
In particular,

$$\|\Lambda(u)\|_1(t) \leq \|\Lambda(u_0)\|_1, \quad t \in [0, T). \quad (3.8)$$

*Proof.* (i) This is clear since  $(u_x)_j$  does not depend upon  $t \in [t_\ell, t_{\ell+1})$  and no faceted regions are created as  $t \uparrow t_\ell$  ( $\ell \geq 1$ ).

(ii) Since  $\|u_{xx}\|_1(t) = \sum_{j=1}^d |(u_x)_{j+1} - (u_x)_j|$ ,  $t \in [0, t_1)$ , it is easy to see that  $\|u_{xx}\|_1(t)$  is independent of  $t \in [t_\ell, t_{\ell+1})$  ( $\ell \geq 0$ ). As  $t \uparrow t_\ell$  ( $\ell \geq 1$ ) jumps of  $u_x(t, \cdot)$  of  $u$  actually disappear and  $u(t_\ell, \cdot)$  is still an admissible crystal. Thus  $\|u_{xx}\|_1$  actually drops at  $t = t_\ell$  ( $\ell \geq 1$ ) as  $t \uparrow t_\ell$ .

(iii) Since  $\|\Lambda(u)\|_1(t) = \sum_{j=1}^d L_j(t) |\Lambda_j(t)| = \sum_{j=1}^d |\chi_j| \Delta((u_x)_j)$ ,  $t \in [0, t_1)$  it is easy to see that  $\|\Lambda(u)\|_1(t)$  is independent of  $t \in [t_\ell, t_{\ell+1})$  ( $\ell \geq 0$ ). Examining the pictures of possible disappearance of faceted region in [GG2], which is still applicable for our general function  $F$ , we conclude that  $\|\Lambda(u)\|_1$  may drop at  $t = t_\ell$  ( $\ell \geq 1$ ) as  $t \uparrow t_\ell$ ; note that in the case that our two consecutive faceted regions disappears  $\|\Lambda(u)\|_1(t)$  does not drop at  $t = t_\ell$  as  $t \uparrow t_\ell$  ( $\ell \geq 1$ ).  $\square$

**Remark 3.5.** The estimates (3.6) and (3.7) follow from Theorem 2.1 once we know that a weakly admissible solution is a generalized solution. This idea can be implemented if we modify the value of  $F(t, p, 0)$  for  $p \neq P$  so that the next proposition applies. We note that our proof for (3.6)-(3.8) does not depend on the theory of partial differential equations. It is not known whether (3.8) follows from Theorem 2.1 since  $\Lambda(u)$  is not a local quantity.

**Proposition 3.6 (Consistency).** *Assume the same hypothesis of Theorem 3.1 (including (ii)) concerning  $F$ . Then a weakly admissible solution  $u \in C([0, T] \times \mathbf{T})$  of (1.5) is a generalized solution of (1.5) provided that*

$$F(t, p, 0) = \theta F(t, p_k, 0) + (1 - \theta) F(t, p_{k+1}, 0) \quad (3.8)$$

for  $p = \theta p_k + (1 - \theta) p_{k+1}$  with  $0 \leq \theta \leq 1$ ,  $k = 1, \dots, m - 1$ .

This is a main topic of [GG2] for  $F(t, p, X) = -a(p)(X - C(t))$  with a nonnegative continuous function  $a$  and a continuous function  $C$ . However, the proof can be easily extended in our general setting. We remark that assumptions on  $F$  in Lemma 3.4 and Proposition 3.6 can be weakened as in Remark 3.2 (ii).

**A priori estimates by crystalline algorithm.** We shall prove (2.1) and (2.2) with  $\xi(\sigma) = \sigma$  by Theorem 1.1 without using the theory of uniformly parabolic equations (Proposition 2.3). For this purpose we approximate  $u_0 \in C(\mathbf{T})$  by admissible crystals; this approximation is not intended for numerical computation.

**Lemma 3.7 (Approximation by admissible crystals).** *Assume that  $u_0 \in C(\mathbf{T})$  and  $\|u_{0xx}\|_1 < \infty$ . Then there is a sequence of finite set  $P_n$  in  $\{p; |p| \leq \|u_{0x}\|_\infty + 1\}$  ( $n = 1, 2, \dots$ ) and a sequence of  $u_0^n$  which is an admissible crystal with respect to  $P_n$  such that  $u_0^n \rightarrow u_0$  in  $C(\mathbf{T})$  with  $\|u_{0x}^n\|_\infty \rightarrow \|u_{0x}\|_\infty$  and*

$$\lim_{n \rightarrow \infty} \|u_{0xx}^n\|_1 = \|u_{0xx}\|_1, \quad (3.9)$$

$$\lim_{n \rightarrow \infty} |P_n| = 0 \quad \text{with} \quad |P_n| = \max\{|r - s|, r, s \in P_n, r < s, (r, s) \cap P_n = \emptyset\}. \quad (3.10)$$

*Proof.* There is a sequence of smooth functions  $v_\ell$  such that  $v_\ell \rightarrow u_0$  in  $C(\mathbf{T})$  with  $\|v_{\ell x}\|_\infty \rightarrow \|u_{0x}\|_\infty$  and  $\|v_{\ell xx}\|_1 \rightarrow \|u_{0xx}\|_1$  as  $\ell \rightarrow \infty$  (cf. [Giu]). By a diagonal argument we may assume  $u_0 \in C^\infty(\mathbf{T})$ . Similarly, we may still assume that  $u_0$  is real analytic by truncating  $\omega$ -periodic Fourier expansion of  $u_0$ .

If  $u_{0x} \neq 0$  is real analytic, the set of points  $Z = \{x \in \mathbf{T}; u_{0xx}(x) = 0\}$  is finite (and nonempty by periodicity). We express the set  $Z$  by  $Z = \{0 \leq z_1 < z_2 < \dots < z_{i_0} < \omega\}$  and set  $q_i = u_{0x}(z_i)$  ( $i = 1, 2, \dots, i_0$ ), so that

$$\|u_{0xx}\|_1 = \sum_{j=1}^{i_0} |q_{j+1} - q_j| \quad \text{with} \quad q_{i_0+1} = q_1. \quad (3.11)$$

For  $r_+ := \|u_{0x}\|_\infty + 1$  and  $r_- := -r_+$ , we set

$$P_n = \{p_j \in \mathbf{R}; p_j = (r_+ - r_-) \cdot j/2^n + r_-, j = 0, 1, 2, \dots, 2^n\} \cup \{q_i; i = 1, 2, \dots, i_0\},$$

so that (3.10) holds. If  $u_{0x} \equiv 0$  we take  $P_n$  as above by interpreting that the set of  $q_i$ 's is empty and we set  $u_0^n = u_0$ . If  $u_{0xx} > 0$  in  $(z_i, z_{i+1})$  then we approximate  $u_{0x}$  by a piecewise constant nondecreasing functions  $f_n$  in  $[z_i, z_{i+1}]$  with values in  $P_n$  that satisfies

$$\lim_{n \rightarrow \infty} \sup_{z_i \leq x \leq z_{i+1}} |u_{0x} - f_n|(x) = 0, \quad (3.12)$$

$$\begin{cases} f_n \text{ takes all values of } P_n \cap [q_i, q_{i+1}] \text{ and its inverse } f_n^{-1}(r) \\ \text{for } r \in P_n \cap [q_i, q_{i+1}] \text{ has an interior in } [z_i, z_{i+1}]; \\ \text{moreover } f_n(z_i) = q_i, f_n(z_{i+1}) = q_{i+1}, \end{cases} \quad (3.13)$$

$$\int_{z_i}^{z_{i+1}} f_n(x) dx = \int_{z_i}^{z_{i+1}} u_{0x}(x) dx. \quad (3.14)$$

If  $u_{xx} < 0$  in  $(z_i, z_{i+1})$  we take  $f_n$  in  $[z_i, z_{i+1}]$  as a piecewise constant nonincreasing function satisfying (3.12)-(3.14). In each piece  $[z_i, z_{i+1}]$  we assign  $u_0^n(x) = \int_{z_i}^x f_n(y) dy + u_0(z_i)$ . Then, (3.13) and (3.14) guarantee that  $u_0^n$  is an admissible crystal with respect to  $P_n$ . The convergence  $u_0^n \rightarrow u_0$  in  $C(\mathbf{T})$  with  $\|u_{0x}^n\|_\infty \rightarrow \|u_{0x}\|_\infty$  follows from (3.12). Since  $\|u_{0xx}^n\|_1 = \sum_{i=1}^{i_0} |q_{i+1} - q_i|$  by construction we have  $\|u_{0xx}^n\|_1 = \|u_{0xx}\|_1$  for all  $n$ . Thus (3.9) follows from (3.11).  $\square$

We introduce our crystalline algorithm to approximate solutions of (1.5).

- (i) For given  $u_0 \in C(\mathbf{T})$  with  $\|u_{0xx}\|_1 < \infty$  we take  $u_0^n$  and  $P_n$  as in Lemma 3.7.
- (ii) For  $W_0 \in \mathcal{E}$  we take a piecewise linear convex function  $W_n$  such that  $W_n = W_0$  on  $P_n$  and  $W_n$  is affine outside  $P_n$  with  $W_n'(p_1 - 0) = W_0'(p_1 - 0)$ ,  $W_n'(p_m + 0) = W_0'(p_m + 0)$ , where  $p_1 = \min P_n$ ,  $p_m = \max P_n$ .
- (iii) For  $F_0 \in \mathcal{F}_{\#T}$ , we may assume that  $F_0$  is  $C^\infty$  by mollifying. We take  $F_{0n}(t, p, X) = F_0(t, p, X) + G_n(X)$  with  $G_n(X) = G(X/n)$  with a nonincreasing

$C^\infty$  function  $G$  (with bounded derivative) that satisfies  $G(X) = 0$  for  $|X| \leq 1$  and  $\lim_{X \rightarrow \pm\infty} G(X) = \mp\infty$ . We then take

$$F_n(t, p, X) = \begin{cases} F_{0n}(t, p, X), & p \in P_n \text{ or } p \notin [p_1, p_m], \\ \theta F_{0n}(t, p_k, X) + (1 - \theta) F_{0n}(t, p_{k+1}, X), & p = \theta p_k + (1 - \theta) p_{k+1}, 1 \leq k \leq m - 1, 0 \leq \theta \leq 1. \end{cases}$$

For such  $F_n$  and  $W_n$  we consider a weakly admissible solution  $u^n$  of (1.4) with  $F_\varepsilon = F_n$ ,  $W_\varepsilon = W_n$ ,  $u_0^\varepsilon = u_0^n$ ,  $\varepsilon = 1/n$ ,  $n \geq 1$ ; modification by  $G$  guarantees the global existence of weakly admissible solutions. By definition  $F_n$  satisfies (3.8). Thus, by Proposition 3.6  $u^n$  is a generalized solution of (1.4). Since (3.10) implies that  $W_n \rightarrow W_0$  uniformly in  $[r_-, r_+]$ ,  $F_n \rightarrow F_0$  locally uniformly and  $u_0^n \rightarrow u_0$  in  $C(\mathbf{T})$  as  $n \rightarrow \infty$ , Theorem 1.1 guarantees the local uniform convergence  $u^n$  to the generalized solution  $u$  of (1.5) in  $[0, T] \times \mathbf{T}$ . Note that the limit of  $W_n$  may not agree with  $W_0$  outside  $[p_1, p_m](= [r_-, r_+])$ . However, since  $\|u_x^n\|_\infty < r_+$ , our  $u$  actually solves (1.5) with original  $W_0$ . Since  $u^n$  approximates  $u$  and  $u^n$  is constructed by solving ODEs (3.1)-(3.2), the method (i)-(iii) is a kind of crystalline algorithm. Our way of approximation of  $W_0$  is different from [GK1, 2] since our  $W_n$  and  $P_n$  also depend on initial data  $u_0$ .

It is by now clear that (2.1) and (2.2) with  $\xi(\sigma) = \sigma$  are obtained by Lemma 3.4 and Theorem 1.1 since we have  $\|u_{0xx}^n\|_1 \rightarrow \|u_{0xx}\|_1$  and  $\|u_{0x}^n\|_\infty \rightarrow \|u_{0x}\|_\infty$  as  $n \rightarrow \infty$  by Lemma 3.7. The convergence (Theorem 2.5) also follows since it needs (2.2) only for  $\xi(\sigma) = \sigma$ .

#### 4. Counterexample when $F$ depends on the space variable

As we observed by Proposition 3.6 and Theorem 1.1 a weakly admissible solution is obtained by approximation by a smooth strict parabolic problem.

When  $F$  depends on the space variable, it is still possible to derive a system like (3.1)-(3.2) if we admit the ansatz that a facet stays as a facet with same slope. However, such an 'admissible solution' may not fulfill a comparison principle. We give an example below. This observation shows that such a solution cannot be obtained by approximation by smoother strict parabolic problem. In [R] and [GG4] a reasonable way to interpret solutions is proposed.

Let us consider an example given in [GG4] when  $F$  depends on the space variable. We consider an equation of form

$$u_t - \{W'(u_x)_x - C(x)\} = 0, \quad x \in \mathbf{R}, \quad t > 0, \quad (4.1)$$

where  $W(p) = \Delta|p|/2$ ,  $C(x) = \max\{0, 1 - |x|\}$  with positive number  $\Delta$ . We take

$$u_0(x; \alpha_0) = \begin{cases} 0 & |x| \leq \alpha_0, \\ q(x - \alpha_0) & x > \alpha_0, \\ -q(x + \alpha_0) & x < -\alpha_0 \end{cases} \quad (4.2)$$

as an initial data, where  $q$  is a positive number and  $\alpha_0 > 1$ . Let us recall that the exact solution  $u$  of this initial value problem in [GG4] obtained by subdifferential formulation.

(i) the case  $\Delta \geq 1$ .

$$u(t, x; \alpha_0) = \begin{cases} u_0(\alpha(t; \alpha_0)) & |x| < \alpha(t; \alpha_0), \\ u_0(x) & \text{otherwise} \end{cases} \quad (4.3)$$

with  $\alpha(t; \alpha_0) = (\alpha_0^2 + (\Delta - 1)t/q)^{1/2}$ . For later convenience we set the right hand side of (4.3) as  $f(t, x; \alpha_0)$ .

(ii) the case  $\Delta < 1$ .

$$u(t, x; \alpha_0) = \begin{cases} -C(\Delta^{1/2})t & |x| \leq \Delta^{1/2}, \\ -C(x)t & \Delta^{1/2} < |x| < 1, \\ u_0(x; \alpha_0) & \text{otherwise.} \end{cases}$$

We observe that in the latter case a facet of  $u_0$  on  $[-\alpha_0, \alpha_0]$  is split into five facets. We also observe that for any  $\bar{\alpha}_0 > \alpha_0 > 1$  we have  $u(t, x; \bar{\alpha}_0) \leq u(t, x; \alpha_0)$  for  $t > 0$  and  $x \in \mathbb{R}$  for the both cases.

We shall discuss what would happen if we assume 'facet-stay-facet assumption.' Let us assume that a function  $v(t, x)$  solves (4.1) under facet-stay-facet assumption with initial data  $u_0(x; \alpha_0)$ . Integrating  $v_t$  from  $\alpha_0 - \sigma$  to  $\alpha_0 + \sigma$  at  $t = +0$  with small positive  $\sigma$ , we have

$$\int_{-\alpha_0 - \sigma}^{\alpha_0 + \sigma} v_t \, dx = \int_{-\alpha_0 - \sigma}^{\alpha_0 + \sigma} \{W'(u_x)_x - C(x)\} dx = W'(u_x)|_{x=-\alpha_0 - \sigma}^{\alpha_0 + \sigma} - 1$$

at  $t = +0$ , since  $\alpha_0 > 1$ . Facet-stay-facet assumption implies that the left hand side equals  $2(\alpha_0 + \sigma)v_t(+0, x)$  which is independent of  $x \in (-\alpha_0, \alpha_0)$ . Sending  $\sigma$  to 0 implies that

$$v_t(+0, x) = \frac{\Delta - 1}{2\alpha_0} = \Lambda_W(v(+0, \cdot), 0) - \frac{1}{2\alpha_0} \quad \text{for } x \in (-\alpha_0, \alpha_0).$$

We observe that the right hand side equals weighted curvature minus the average of  $C(x)$  on the faceted region  $(-\alpha_0, \alpha_0)$ . Thus we obtain that  $v(t, x; \alpha_0) = f(t, x; \alpha_0)$  for  $x \in \mathbb{R}$  and  $t > 0$  with  $\alpha(t; \alpha_0) > 0$ .

When  $\Delta < 1$ , comparison principle is not valid for  $v$ . In fact, for  $\bar{\alpha}_0 > \alpha_0 > 0$  it holds that  $u_0(x; \bar{\alpha}_0) \leq u_0(x; \alpha_0)$  for  $x \in \mathbb{R}$ . However, we have  $v(t, x; \bar{\alpha}_0) > v(t, x; \alpha_0)$  at least for  $t \in (0, \alpha_0^2 q / (1 - \Delta))$  and  $x \in (-\alpha(t; \alpha_0), \alpha(t; \alpha_0))$ , since  $\alpha(t; \bar{\alpha}_0) - \bar{\alpha}_0 > \alpha(t; \alpha_0) - \alpha_0$  implies that  $u_0(\alpha(t; \bar{\alpha}_0)) > u_0(\alpha(t; \alpha_0))$ .

When  $C$  is independent of  $x$ , the subdifferential formulation is consistent with our generalized solutions [GG5]. It is desirable that the notion of generalized solutions is extended to the case when  $C$  depends on  $x$  so that it is compatible with the subdifferential formulation as suggested in [GG4].

#### REFERENCES

- [AG] S. B. Angenent and M. E. Gurtin, *Multiphase thermomechanics with interfacial structure 2. Evolution of an isothermal interface*, Arch. Rational Mech. Anal. **108** (1989), 323-391.
- [CIL] M. G. Crandall, H. Ishii and P. L. Lions, *User's guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc. **27** (1992), 1-67.
- [FG] T. Fukui and Y. Giga, *Motion of a graph by nonsmooth weighted curvature*, In: World Congress of Nonlinear Analysts '92 (ed. V. Lakshmikantham), Walter de Gruyter, Berlin, vol. I (1996), pp.47-56.
- [GG1] M.-H. Giga and Y. Giga, *Geometric evolution by nonsmooth interfacial energy*, In: Nonlinear Analysis and Applications (eds. N. Kenmochi, A. Damlamian, M. Niezgódko and A. Visintin) Proceedings of Banach Center Minisemester, November-December, 1994, Gakuto International Series Mathematical Sciences and Applications, **7** (1995) pp.125-140, Gakuto, Tokyo.
- [GG2] M.-H. Giga and Y. Giga, *Consistency in evolutions by crystalline curvature*, In: Free Boundary Problems '95, Theory and Applications (eds. M. Niezgódko and P. Strzelecki), Proc. of the Zakopane Congress, Poland (1995), pp.186-202.
- [GG3] M.-H. Giga and Y. Giga, *Evolving graphs by singular weighted curvature*, Arch. Rational. Mech. Anal., to appear.
- [GG4] M.-H. Giga and Y. Giga, *A subdifferential interpretation of crystalline motion under nonuniform driving force*, Hokkaido University Preprint Series in Mathematics #344 (1996).
- [GG5] M.-H. Giga and Y. Giga, *Stability for evolving graphs by nonlocal weighted curvature*, Hokkaido University Preprint Series in Mathematics #372

- (1997).
- [GK1] P. M. Girão and R. V. Kohn, *Convergence of a crystalline algorithm for the heat equation in one dimension and for the motion of a graph by weighted curvature*, Numer. Math. **67** (1994), 41-70.
  - [GK2] P. M. Girão and R. V. Kohn, *The crystalline algorithm for computing motion by curvature*, In: Variational Methods for Discontinuous Structure (eds. R. Serapioni and F. Tomarelli) Birkhauser, Progress in Nonlinear Differential Equations and Applications, **25** (1996), 7-18.
  - [Giu] E. Giusti, *Minimal Surfaces and Functions of Bounded Variations*, Birkhauser, Boston, 1984.
  - [Gu] M. E. Gurtin, *Thermomechanics of Evolving Phase Boundaries in the Plane*, Oxford, Clarendon Press, 1993.
  - [LSU] O. A. Ladyzhenskaya, V. Solonnikov and N. Ural'ceva, *Linear and Quasilinear Equations of Parabolic Type*, Translations of Mathematical Monographs, Amer. Math. Soc. **23** (1968).
  - [O] F. Otto,  *$L^1$ -Contraction and uniqueness for quasilinear elliptic-parabolic equations*, J. Differential Equations **131** (1996), 20-38.
  - [R] A. R. Roosen, *Crystalline curvature and flat flow in a diffusion field*, preprint (1994).
  - [T] J. Taylor, *Constructions and conjectures in crystalline nondifferential geometry*, In: Differential Geometry (eds. B. Lawson and K. Tanenblat), Proceedings of the Conference on Differential Geometry, Rio de Janeiro, Pitman Monographs in Pure and Applied Math. **52** (1991) pp.321-336, Pitman, London.
  - [TCH] J. Taylor, J. Cahn and C. Handwerker, *Geometric models of crystal growth*, Acta Metall. **40** (1992), 1443-1474.

Department of Mathematics  
Hokkaido University  
Sapporo 060, Japan