

HOKKAIDO UNIVERSITY

Title	Generalized Numerical Radius And Unitary p-Dilation
Author(s)	Nakazi, T.; Okubo, K.
Citation	Hokkaido University Preprint Series in Mathematics, 408, 1-12
Issue Date	1998-04-01
DOI	https://doi.org/10.14943/83554
Doc URL	https://hdl.handle.net/2115/69158
Type	departmental bulletin paper
File Information	re408.pdf



Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Generalized Numerical Radius And Unitary ρ -Dilation

T. Nakazi and K. Okubo

Series #408. April 1998

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

#383 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the Sobolev space of critical order, 24 pages. 1997.

#384 K. Ito, Asymptotic stability of planar rarefaction wave for scalar viscous conservation law, 8 pages. 1997.

#385 A. Arai, Representation-theoretic aspects of two-dimensional quantum systems in singular vector potentials: canonical commutation relations, quantum algebras, and reduction to lattice quantum systems, 32 pages. 1997.

#386 P. Aviles and Y. Giga, On lower semicontinuity of a defect energy obtained by a singular limit of the Ginzburg-Landau type energy for gradient fields, 21 pages. 1997.

#387 T. Nakazi and T. Yamamoto, Norms of some singular integral operators and their inverse operators, 28 pages. 1997.

#388 M.-H. Giga and Y. Giga, Remarks on convergence of evolving graphs by nonlocal curvature, 18 pages. 1997.

#389 T. Tsukada, Reticular Lagrangian singularities, 41 pages. 1997.

#390 M. Nakamura and T. Ozawa, The Cauchy problem for nonlinear wave equations in the homogeneous Sobolev space, 12 pages. 1997.

#391 Y. Giga, M. Ohnuma and M.-H. Sato, On strong maximum principle and large time behaviour of generalized mean curvature flow with the Neumann boundary condition, 24 pages. 1997.

#392 T. Tsujishita and H. Watanabe, Monoidal closedness of the category of simulations, 24 pages. 1997.

#393 T. Arase, A remark on the quantale structure of multisets, 10 pages. 1997.

#394 N. H. Bingham and A. Inoue, Extension of the Drasin-Shea-Jordan theorem, 16 pages. 1997.

#395 N. H. Bingham and A. Inoue, Ratio Mercerian theorems with applications to Hankel and Fourier transforms, 30 pages. 1997.

#396 Y. Nishiura and D. Ueyama, A skeleton structure of self-replicating dynamics, 27 pages. 1997.

#397 K. Hirata and K. Sugano, On semisimple extensions of serial rings, 6 pages. 1997.

#398 D. Pei, Singularities of $\mathbb{R}P^2$-valued Gauss maps of surfaces in Minkowski 3-space, 15 pages. 1997.

#399 T. Mikami, Markov marginal problems and their applications to Markov optimal control, 28 pages. 1997.

#400 M. Tsujii, A simple proof for monotonicity of entropy in the quadratic family, 8 pages. 1998.

#401 M. Nakamura and T. Ozawa, Global solutions in the critical Sobolev space for the wave equations with nonlinearity of exponential growth, 9 pages. 1998.

#402 D. Lehmann and T. Suwa, Generalization of variations and Baum-Bott residues for holomorphic foliations on singular varieties, 19 pages. 1998.

#403 T. Nakazi and K. Okubo, ρ-contraction and 2×2 matrix, 6 pages. 1998.

#404 Y. Kohsaka, Free boundary problem for quasilinear parabolic equation with fixed angle of contact to a boundary, 28 pages. 1998.

#405 K. Yokoyama, Global existence of classical solutions to systems of wave equations with critical nonlinearity in three space dimensions, 25 pages. 1998.

#406 F. Hiroshima, Ground states and spectrum of quantum electrodynamics of non-relativistic particles, 58pages. 1998

#407 N. Kawazumi and T. Uemura, Riemann-Hurwitz formula for Morita-Mumford classes and surface symmetries, 9pages. 1998.

Generalized Numerical Radius And Unitary ρ-Dilation

by

Takahiko Nakazi*

and

Kazuyoshi Okubo*

* This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education

1991 Mathematics Subject Classification. Primary 47 A 20

Key words and phrases : numerical radius, norm, ρ-dilation

Abstract. In this paper, we study an operator A on a Hilbert space H which satisfies one of the following inequalities :
For some λ with $0 \leq \lambda \leq 1$

$$|(Ay, y)| \leq \lambda \|y\|^2 + (1-\lambda)\|Ay\|^2 \quad (y \in H)$$

or

$$\lambda \|Ay\|^2 + (1-\lambda)|(Ay, y)| \leq \|y\|^2 \quad (y \in H).$$

These two inequalities can be regarded as special cases of generalized numerical ranges. If A has a ρ-dilation with $\rho > 0$, then it satisfies one of them. We show that the operator radii $w_\rho(A)$ of A are calculated using $|(Ay, y)|$ and $\|Ay\|$. Several applications are given.

§1. Introduction

According to Sz.-Nagy and Foias [7], a bounded linear operator A on a complex Hilbert space H is said to be of class $\mathcal{C}_\rho$ with $\rho > 0$ if there exists a unitary operator U on some Hilbert space K such that K contains H as a subspace and such that

$$A^n = \rho P_H U^n | H \quad \text{for } n = 1, 2, \cdots,$$

where P_H is the orthogonal projection of K onto H. For $\rho = 2$ it is known (see [7, Chapter I, Proposition 11.2]) that A is of class $\mathcal{C}_2$ if and only if its numerical radius of A :

$$w(T) = \sup\{|(Ay, y)| : \|y\| \leq 1\}$$

is not greater than one. Of course, A is of class $\mathcal{C}_1$ if and only if the usual norm $\|A\|$ of A is not greater than one. Holbrook [3] introduced the operator radii $w_\rho(A)$ of an operator A, relative to $\mathcal{C}_\rho$, by the formula :

$$w_\rho(A) = \inf\{\gamma \ ; \ \gamma > 0, \gamma^{-1} A \in \mathcal{C}_\rho\}.$$

Then $w_1(A) = \|A\|$, $w_2(A) = w(A)$, and $\lim_{\rho \to \infty} w_\rho(A) = r(A)$: the spectral radius of A.

Let S be a positive bounded operator on H. We define $V_S^+(A)$ and $v_S(A)$ as

$$V_S^+(A) = \{(Ay, y) \ ; \ y \in H, (Sy, y) = 1\}.$$

and

$$v_S(A) = \sup\{|(Ay, y)| \ ; \ y \in H, (Sy, y) = 1\}.$$

$V_S^+(A)$ is defined and studied in [4] for a self-adjoint S. If S is the identity operator I on H, then $V_S^+(A)$ is the numerical range of A and $v_S(A) = w(A)$. In Section 2, when $0 < \rho \leq 2$ and $\rho \neq 1$, we show that A is of class $\mathcal{C}_\rho$ if and only if

$$S = \frac{\rho}{2|\rho - 1|} I + \frac{\rho - 2}{2|\rho - 1|} |A|^2$$

is nonnegative and $v_S(A) \leq 1$. As results, several corollaries are given. In Section 3, when $0 < \rho \leq 2$ and $\rho \neq 1$, we give two formulae for $w_\rho(A)$ using $|(Ay, y)|$ and $\|Ay\|$. In Section 4, we try to generalize results in Sections 2 and 3 for $2 < \rho < \infty$.

A result in this paper is the following : For $0 < \rho \leq 2$ and $\rho = 2/(\lambda + 1)$,

$$\sup_{\|y\|=1} \{\lambda \|Ay\|^2 + |1 - \lambda| \cdot |(Ay, y)|\} \leq 1$$

if and only if $w_\rho(A) \leq 1$ (see Corollay 2). This shows the following as $A/w_\rho(A)$. If $w_\rho(A) \leq 1$, then

$$\sup_{\|y\|=1} \{\lambda \|Ay\|^2 + |1 - \lambda| \cdot |(Ay, y)|\} \leq w_\rho(A)$$

and if $w_\rho(A) \geq 1$, then

$$\sup_{\|y\|=1} \{\lambda\|Ay\|^2 + |1-\lambda| \cdot |(Ay,y)|\}^{1/2} \leq w_\rho(A).$$

$w_\rho(A)$ is calculated using $|(Ay,y)|$ and $\|Ay\|$ (see (1) of Theorem 2). This shows the following general inequality (see Corollary 3). For any $w_\rho(A)$,

$$w_\rho(A) \leq \sup_{\|y\|=1} \{\sqrt{\lambda}\|Ay\| + |1-\lambda| \cdot |(Ay,y)|\}.$$

§2. ρ-dilation for $0 < \rho \leq 2$ and generalized numerical radius

In this section, we are interested in operators with $v_S(A) \leq 1$ when S is a special positive operator. If $S = |A|$ and $v_S(A) \leq 1$, then A is normal (*cf.* [2],[8]). We consider A when $S = \lambda I + \mu|A|^2, \lambda + \mu = \pm 1$ and $v_S(A) \leq 1$ where $\lambda \geq 0$ and μ are constants.

For $0 \leq \mu < \infty$

$$\mathbf{w}_\mu(A) = \sup\{\mu\|Ay\|^2 + |1-\mu| \cdot (Ay,y)| \ ; \ \|y\| = 1\}.$$

Then $\mathbf{w}_0(A) = w(A)$ and $\mathbf{w}_1(A) = \|A\|^2$.

Theorem 1. *Suppose $0 < \rho \leq 2$ and $\rho \neq 1$. Then A is of class C_ρ if and only if $S \geq 0$ and $v_S(A) \leq 1$ where $S = (\rho I + (\rho-2)|A|^2)/2|\rho - 1|$.*

Proof. It is known that A is of class C_ρ if and only if

$$\|y\|^2 + \left(1 - \frac{2}{\rho}\right)|\zeta|^2\|Ay\|^2 - 2\left(1 - \frac{1}{\rho}\right) Re\zeta(Ay,y) \geq 0$$

for $\zeta \in D$ and $y \in H$ where $D = \{z \in \mathbf{C} : |z| < 1\}$. This inequality is equivalent to

$$Re\zeta(Ay,y) \leq \frac{\rho}{2|\rho-1|}\|y\|^2 + \frac{\rho-2}{2|\rho-1|}|\zeta|^2\|Ay\|^2$$

for $\zeta \in D$ and $y \in H$, and

$$|\zeta|(Ay,y)| \leq \frac{\rho}{2|\rho-1|}\|y\|^2 + \frac{\rho-2}{2|\rho-1|}|\zeta|^2\|Ay\|^2$$

for $\zeta \in D$ and $y \in H$. Since $0 < \rho \leq 2$, the last inequality is equivalent to

$$|(Ay,y)| \leq \frac{\rho}{2|\rho-1|}\|y\|^2 + \frac{\rho-2}{2|\rho-1|}\|Ay\|^2$$

for $y \in H$.

Corollary 1. *Suppose* $0 < \rho < 2$ *and* $\rho \neq 1$. *A is of class* C_ρ *if and only if A admits a factorization :* $A = S^{1/2}BS^{1/2}$ *where* $S = (\rho I + (\rho - 2)|A|^2)/2|\rho - 1|$ *and* $w(B) \leq 1$.

Proof. Suppose A is of class C_ρ. If there exists $y \in H$ such that $Sy = 0$, then $|A|^2 y = \rho y/(2-\rho)$. Since $\rho w_\rho(A) \geq \|A\|$ [1],

$$1 \geq w_\rho(A) \geq \frac{\|A\|}{\rho} \geq \frac{1}{\sqrt{\rho(2-\rho)}}.$$

and hence $\rho = 1$. This contradiction implies that $\ker S = \{0\}$. Since $\ker S = \{0\}, v_S(A) \leq 1$ if and only if

$$|(S^{-1/2}AS^{-1/2}x, x)| \leq (x, x) \quad (x \in H).$$

Let $B = S^{-1/2}AS^{-1/2}$, then this inequality is equivalent to that

$$A = S^{1/2}BS^{1/2} \text{ and } w(B) \leq 1.$$

Now Theorem 1 implies the corollary.

Corollary 2. *Let A be a bounded linear operator on H and* $0 < \rho = \dfrac{2}{\mu + 1} \leq 2$. *Then the following conditions are mutually equivalent :*

(1) $w_\rho(A) \leq 1$.

(2) $\mathbf{w}_\mu(A) \leq 1$

and

(3) $w(\mu|A|^2 + |1 - \mu|e^{i\theta}A) \leq 1$

for any $\theta \in R$.

Proof. By Theorem 1,$w_\rho(A) \leq 1$ if and only if

$$\frac{2-\rho}{2|\rho-1|}\|Ay\|^2 + |(Ay, y)| \leq \frac{\rho}{2|\rho-1|}\|y\|^2$$

for $y \in H$. The last inequality is equivalent to

$$\mu\|Ay\|^2 + |1 - \mu| \cdot |(Ay, y)| \leq \|y\|^2$$

where $\mu = (2-\rho)/\rho$. This implies the equivalence of (1) and (2). The equivalence of (2) and (3) is trivial.

(1),(2) of Corollary 2 implies that if $w_\rho(A) \leq 1$ then $\mathbf{w}_\mu(A) \leq w_\rho(A)$ and if $w_\rho(A) \geq 1$ then $\mathbf{w}_\mu(A)^{1/2} \leq w_\rho(A)$. By (1),(3) of Corollary 2, if $\mu\|A\|^2 + |1 - \mu|w(A) \leq 1$

then $w_\rho(A) \leq 1$. However the converse is not true. For if $A = \begin{bmatrix} 0 & \rho \\ 0 & 0 \end{bmatrix}$, then $\|A\| = \rho, w(A) = \rho/2$ and $w_\rho(A) = 1$. Hence $\mu = (2-\rho)/\rho$ and

$$\mu\|A\|^2 + |1-\mu|w(A) = \frac{2-\rho}{\rho} \times \rho^2 + \frac{2|\rho-1|}{\rho} \times \frac{\rho}{2} > 1.$$

§3. Operator radii for $0 < \rho \leq 2$

In this section, we give two exact formulae and useful estimates for $w_\rho(A)$ when $0 < \rho \leq 2$ and $\rho \neq 1$. Put

$$D = D(A, \rho, y) = |(Ay, y)|^2 - \frac{\rho(\rho-2)}{(\rho-1)^2}\|Ay\|^2$$

for a bounded linear operator $A, \rho > 0$ and y in H.

Theorem 2. *Let A be a bounded linear operator on H.*
(1) If $0 < \rho \leq 2$ and $\rho \neq 1$, then

$$w_\rho(A) = \frac{|\rho-1|}{\rho} \sup_{\|y\|=1} \{|(Ay, y)| + \sqrt{D}\}.$$

(2) If $0 < \rho \leq 2$, and $\rho \neq 1$ then

$$w_\rho(A) = \frac{2}{\rho} \sup_{\|y\|=1} \sup_{0 \leq t \leq 1} \{\sqrt{\rho(2-\rho)}\|Ay\|\sqrt{t(1-t)} + |\rho-1| \cdot |(Ay, y)|t\}.$$

Proof. (1) By Theorem 1, if $t \geq w_\rho(A)$ then $\lambda\|y\|^2t^2 - |(Ay, y)|t + (1-\lambda)\|Ay\|^2 \geq 0$ for $y \in H$ where $\lambda = \rho/2|\rho-1|$. Hence

$$w_\rho(A) \leq \frac{|(Ay, y)| - \sqrt{D}}{2\lambda\|y\|^2}$$

or

$$w_\rho(A) \geq \frac{|(Ay, y)| + \sqrt{D}}{2\lambda\|y\|^2}.$$

Put

$$t_0 = \sup_{y \neq 0} \frac{|(Ay, y)| + \sqrt{D}}{2\lambda\|y\|^2}.$$

If $t \geq t_0$, then

$$\lambda\|y\|^2 t^2 - |(Ay,y)|t + (1-\lambda)\|Ay\|^2 \geq 0.$$

for $y \in H$ and so by Theorem 1 $w_\rho(A) \leq t_0$. When $0 < \rho \leq 2, |(Ay,y)| - \sqrt{D} \leq 0$ and so $w_\rho(A) \geq t_0$. Thus $w_\rho(A) = t_0$.

(2) Put $g(t,y) = \sqrt{\rho(2-\rho)}\|Ay\|\sqrt{t(1-t)} + |\rho-1| \cdot |(Ay,y)|t$ for each y with $\|y\| = 1$. Then

$$\begin{aligned} & 2\left(\frac{d}{dt}g(t,y)\right)\sqrt{t(1-t)} \\ = \; & \sqrt{\rho(2-\rho)}\|Ay\|(1-2t) + 2|\rho-1| \cdot |(Ay,y)|\sqrt{t(1-t)}. \end{aligned}$$

Hence $\dfrac{d}{dt}g(t,y)|_{t=t_0} = 0$ and $0 \leq t_0 \leq 1$ if and only if

$$t_0 = \frac{1}{2} + \frac{|(Ay,y)|}{2\sqrt{D}}$$

and so

$$\sqrt{t_0(1-t_0)} = \frac{\sqrt{\rho(2-\rho)}\|Ay\|}{2|\rho-1|\sqrt{D}}.$$

Therefore

$$\begin{aligned} & \frac{2}{\rho}\sup_{\|y\|=1}\sup_{0\leq t\leq 1}\{\sqrt{\rho(2-\rho)}\|Ay\|\sqrt{t(1-t)} + |\rho-1| \cdot |(Ay,y)|t\} \\ & = \frac{2}{\rho}\sup_{\|y\|=1}\left\{\sqrt{\rho(2-\rho)}\|Ay\| \times \frac{\sqrt{\rho(2-\rho)}\|Ay\|}{2|\rho-1| \cdot \sqrt{D}} + |\rho-1| \cdot |(Ay,y)| \times \left(\frac{1}{2} + \frac{|(Ay,y)|}{2\sqrt{D}}\right)\right\} \\ & = \frac{|\rho-1|}{\rho}\sup_{\|y\|=1}\{|(Ay,y)| + \sqrt{D}\}. \quad \square \end{aligned}$$

Corollary 3. *If* $0 < \rho \leq 2$, *then*

$$w_\rho(A) \leq \sup_{\|y\|=1}\{\sqrt{\frac{2-\rho}{\rho}}\|Ay\| + 2\left|1 - \frac{1}{\rho}\right| \cdot |(Ay,y)|\}.$$

Proof. Since $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}, (a,b>0)$ for $\rho \neq 1$

$$\sqrt{D} \leq |(Ay,y)| + \frac{\sqrt{\rho(2-\rho)}}{|\rho-1|}\|Ay\|.$$

This inequality and (1) of Theorem 2 imply the corollary.

Corollary 4. *If* $0 < \rho \leq 2$, *then*

$$\max\left\{2\left|1-\frac{1}{\rho}\right| w(A), \sqrt{\frac{2-\rho}{\rho}}\|A\|\right\} \leq w_\rho(A) \leq 2\left|1-\frac{1}{\rho}\right| w(A) + \sqrt{\frac{2-\rho}{\rho}}\|A\|$$

Proof. Since $-\rho(\rho-2)\|Ay\|^2/(\rho-1)^2 \geq 0$. by (1) of Theorem 2

$$w_\rho(A) \geq 2\frac{|\rho-1|}{\rho}|(Ay,y)|,\ \frac{\sqrt{\rho(2-\rho)}}{\rho}\|Ay\|$$

for $y \in H$. Hence

$$w_\rho(A) \geq \max\left\{2\left|1-\frac{1}{\rho}\right| w(A), \sqrt{\frac{2-\rho}{\rho}}\|A\|\right\}.$$

We can get the upper estimate of $w_\rho(A)$ using Corollary 3.

Corollary 5. *If* $0 < \rho \leq 2$, *and* $\rho \neq 1$ *then*

$$\begin{aligned} w_\rho(A) &\leq \frac{2}{\rho}\sup_{0\leq t\leq 1}\{\sqrt{\rho(2-\rho)}\|A\|\sqrt{t(1-t)} + |\rho-1|\cdot w(A)t\} \\ &= \frac{|\rho-1|}{\rho}\left\{w(A) + \sqrt{w(A)^2 - \frac{\rho(\rho-2)}{(\rho-1)^2}\|A\|^2}\right\} \end{aligned}$$

and for any $\|y\| = 1$ and $0 \leq t \leq 1$

$$w_\rho(A) \geq \frac{2}{\rho}\{\sqrt{\rho(2-\rho)}\|Ay\|\sqrt{t(1-t)} + |\rho-1|\cdot|(Ay,y)|t\}.$$

Corollary 6. [5] *If* $0 < \rho \leq 2$, *then*

$$\rho w_\rho(A) = 2w\left(\begin{bmatrix} 0 & \sqrt{\rho(2-\rho)}\ A \\ 0 & (1-\rho)\ \ A \end{bmatrix}\right).$$

Proof. For $0 < \rho \leq 2$,

$$\begin{aligned} & 2w\left(\begin{bmatrix} 0 & \sqrt{\rho(2-\rho)}\ A \\ 0 & (1-\rho)\ \ A \end{bmatrix}\right) \\ &= 2\sup_{\|x\|^2+\|z\|^2=1}|\sqrt{\rho(2-\rho)}(Az,x) + (1-\rho)(Az,z)| \\ &= 2\sup_{\|x\|^2+\|z\|^2=1}\{\sqrt{\rho(2-\rho)}|(Az,x)| + |1-\rho|\cdot|(Az,z)|\} \\ &= 2\sup_{0\leq\lambda\leq 1}\sup_{\|z\|=\sqrt{\lambda}}\{\sqrt{\rho(2-\rho)}\|Az\|\sqrt{1-\lambda} + |1-\rho|\cdot|(Az,z)|\} \\ &= 2\sup_{0\leq\lambda\leq 1}\sup_{\|y\|=1}\{\sqrt{\rho(2-\rho)}\|Ay\|\sqrt{\lambda(1-\lambda)} + |1-\rho|\cdot|(Ay,y)|\sqrt{\lambda}\} \\ &= \rho\ w_\rho(A). \end{aligned}$$

We used (2) of Theorem 2 to show the last eqnality. □

§4. The case of $2 < \rho < \infty$

In this section, we consider Theorems 1 and 2 for $2 < \rho < \infty$. Unfortunately the results for $2 < \rho < \infty$ are more complicated than those for $0 < \rho \leq 2$.

For $0 \leq \lambda \leq 1$, put

$$\mathbf{w}'_\lambda(A) = \sup\{|(Ay, y)| \ ; \ \lambda\|y\|^2 + (1-\lambda)\|Ay\|^2 \leq 1\}.$$

Then $\mathbf{w}'_1(A) = w(A)$, and $\mathbf{w}'_0(A) = w(A^{-1})$ if A is invertible.

Proposition 3. *Suppose* $2 < \rho < \infty$ *and for* $0 < t \leq 1$

$$S_t = \frac{1}{t}\ \frac{\rho}{2|\rho-1|} I + t\ \frac{\rho-2}{|\rho-1|}\ |A|^2.$$

A is of class C_ρ if and only if $S_t \geq 0$ and $v_{S_t}(A) \leq 1$ for $0 < t \leq 1$.

Proof is almost same to that of Theorem 1.

Using Proposition 3 we can show a version of Corollary 1 for $2 < \rho < \infty$. For $0 < \lambda < 1$ and arbitrary bounded operator A on H,

$$\mathbf{w}'_\lambda(A) \leq 1/\sqrt{\lambda(1-\lambda)}.$$

In fact, for any constant $t > 0$

$$|(Ay, y)| \leq (\sqrt{t}\|y\|)(\frac{1}{\sqrt{t}}\|Ay\|) \leq 2t\|y\|^2 + \frac{1}{2t}\|Ay\|^2.$$

Assuming $\lambda \neq 0, 1$, if $k \geq 1/\sqrt{\lambda(1-\lambda)}$ then $k\lambda \geq 1/k(1-\lambda)$. Hence if $k\lambda = 2t$ then $k(1-\lambda) \geq 1/2t$. Therefore if $k \geq 1/\sqrt{\lambda(1-\lambda)}$, then

$$|(Ay, y)| \leq k(\lambda|y\|^2 + (1-\lambda)\|Ay\|^2)$$

and so $\mathbf{w}'_\lambda \leq 1/\sqrt{\lambda(1-\lambda)}$.

$\mathbf{w}'_0(A) < \infty$ if and only if there exists a bounded operator B such that $BA = P$ where P is an orthogonal projection to $(\ker A)^\perp$. For if $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$ is the matrix

of A with respect to the decomposition $H = (\ker A)^{\perp} \oplus \ker A$, then $A_{12} = A_{22} = 0$. If $k = \mathbf{w}_0'(A) < \infty$, then for $z = (x, y) \in (\ker A)^{\perp} \oplus \ker A$

$$|(A_{11}x, x) + (A_{21}x, y)| \leq k(\|A_{11}x\|^2 + \|A_{21}x\|^2).$$

This implies $A_{21} = 0, A_{11}$ is one to one on $(\ker A)^{\perp}$ and $|(A_{11}x, x)| \leq k\|A_{11}x\|^2$. If $B_{11}A_{11}x = x$, then by the inequality above B_{11} is bounded and so $B = B_{11} \oplus O$ satisfies $BA = P$. The proof is reversible.

Corollary 7. *Let A be a bounded linear operator and let $0 \leq \lambda \leq 1$.*

(1) Suppose $\frac{1}{2} < \lambda \leq 1$ and $\rho = 2\lambda/(2\lambda - 1) \geq 2$. Then $\mathbf{w}_{\lambda}'(tA) \leq 1$ for any $0 < t \leq 1$ and if and only if $w_{\rho}(A) \leq 1$

(2) Suppose $0 \leq \lambda < \frac{1}{2}, \rho = 2(\lambda - 1)/(2\lambda - 1) \geq 2$ and A is invertible. Then $\mathbf{w}_{\lambda}'(tA) \leq 1$ for any $t \geq 1$ if and only if $w_{\rho}(A^{-1}) \leq 1$.

Proof. It is clear by Proposition 3. □

Proposition 4. *Let A be a bounded linear operator on H*

(1) If $2 < \rho < \infty$, then

$$w_{\rho}(A) = \frac{\rho - 1}{\rho} \sup_{\|y\|=1} \{|(Ay, y)| + \sqrt{D} \ ; \ D \geq 0\}.$$

(2) If $2 < \rho < \infty$, then

$$w_{\rho}(A) = \frac{2}{\rho} \sup_{\|y\|=1} \inf_{t \geq 1} \{-\sqrt{\rho(\rho - 2)}\|Ay\|\sqrt{t(t-1)} + (\rho - 1)|(Ay, y)|t \ ; \ D \geq 0\}.$$

Proof. (1) The proof is almost same to that of (1) of Theorem 2.

(2) Put $f(t, y) = -\sqrt{\rho(\rho - 2)}\|Ay\|\sqrt{t(t-1)} + (\rho - 1)|(Ay, y)|t$ for each y with $\|y\| = 1$. Then

$$\begin{aligned} &2\left(\frac{d}{dt}f(t, y)\right)\sqrt{t(t-1)} \\ &\quad = -\sqrt{\rho(\rho - 2)}\|Ay\|(2t - 1) + 2(\rho - 1)|(Ay, y)|\sqrt{t(t-1)}. \end{aligned}$$

Hence $\frac{d}{dt}f(t, y)|_{t=t_0} = 0$ and $t_0 \geq 1$ if and only if

$$t_0 = \frac{1}{2} + \frac{|(Ay, y)|}{2\sqrt{D}}$$

and so

$$\sqrt{t_0(t_0-1)} = \frac{\sqrt{\rho(\rho-2)}\|Ay\|}{2(\rho-1)\sqrt{D}}$$

Therefore

$$\begin{aligned}
&\frac{2}{\rho}\sup_{\substack{\|y\|=1\\ D\geq 0}}\inf_{t\geq 1}\{-\sqrt{\rho(\rho-2)}\|Ay\|\sqrt{t(t-1)}+(\rho-1)|(Ay,y)|t\}\\
&\quad=\frac{2}{\rho}\sup_{\substack{\|y\|=1\\ D\geq 0}}\left\{-\sqrt{\rho(\rho-2)}\|Ay\|\times\frac{\sqrt{\rho(\rho-2)}\|Ay\|}{2(\rho-1)\sqrt{D}}+(\rho-1)|(Ay,y)|\times\left(\frac{1}{2}+\frac{|(Ay,y)|}{2\sqrt{D}}\right)\right\}\\
&\quad=\frac{\rho-1}{\rho}\sup_{\substack{\|y\|=1\\ D\geq 0}}\{|(Ay,y)|+\sqrt{D}\}
\end{aligned}$$

Corollary 8. *Suppose $\rho \geq 2$ and $w(A) \geq \frac{\sqrt{\rho(\rho-2)}}{\rho-1}\|A\|$. Then*

$$\begin{aligned}
&2(1-\frac{1}{\rho})w(A)-\sqrt{\frac{\rho-2}{\rho}}\|A\|\\
&\leq\ (1-\frac{1}{\rho})\left\{w(A)+\left(w(A)^2-\frac{\rho(\rho-2)}{(\rho-1)^2}\|A\|^2\right)^{1/2}\right\}\\
&\leq\ w_\rho(A)\\
&\leq\ (1-\frac{1}{\rho})\left\{w(A)+\left(w(A)^2+\frac{\rho(\rho-2)}{(\rho-1)^2}\|A\|^2\right)^{1/2}\right\}\\
&\leq\ 2(1-\frac{1}{\rho})w(A)+\sqrt{\frac{\rho-2}{\rho}}\|A\|.
\end{aligned}$$

References

1. T.Ando and K.Nishio, Convexity properties of operator radii associated with unitary ρ-dilations, Michigan Math. J., 20(1973), 303–307.

2. T.Ando and K.Takahashi, On operators with unitary ρ-dilalions, Ann. Pol. Math. LXVI(1997), 11–14.

3. J.A.R.Holbrook, On the power-bounded operators of Sz.-Nagy and Foias, Acta Sci. Math. (Szeged) 29(1968), 299–310.

4. C.K.Li, N.K. Tsing and F.Uhling, Numerical ranges of operator on an indefinite inner product space, Electronic J.Lin Alg., 1(1996), 1–17.

5. R.Mathias and K.Okubo, The induced norm of the Schur multiplication operator with respect to the operator radius, Linear and Multilinear Algebra 37(1994), 114-124.

6. B.Sz.-Nagy and C.Foias, On certain classes of power bounded operators in Hilbert space, Acta Sci. Math. 27(1966), 17–25.

7. B.Sz.-Nagy and C.Foias, Harmonic Analysis of Operators on Hilbert Space, North-Holland, Amsterdam, 1970.

8. H.Watanabe, Operators characterized by certain Cauchy-Schwarz type inequalities, Publ. Res. Inst. Math. Sci. 30(1994), 249–259.

Department of Mathematics
Faculty of Science
Hokkaido University
Sapporo 060, Japan

Mathematics Laboratory
Sapporo College
Hokkaido University of Education
Sapporo 002, Japan