



HOKKAIDO UNIVERSITY

Title	Perturbations of Caustics and fronts
Author(s)	Ishikawa, G.
Citation	Hokkaido University Preprint Series in Mathematics, 566, 1-17
Issue Date	2002-10
DOI	https://doi.org/10.14943/83711
Doc URL	https://hdl.handle.net/2115/69315
Type	departmental bulletin paper
File Information	re566.pdf



PERTURBATIONS OF CAUSTICS AND FRONTS

Goo Ishikawa

Series #566, October 2002

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #540 H. Ishii and T. Mikami, Motion of a graph by R -curvature, 28 pages. 2001.
- #541 M. Jinzenji and T. Sasaki, $\mathcal{N} = 4$ supersymmetric Yang-Mills theory on orbifold- $T^4/\mathbf{Z}_2$: higher rank case, 17 pages. 2001.
- #542 T. Nakazi, The Nevanlinna counting functions for Rudin's orthogonal functions, 7 pages. 2001.
- #543 K. Sugano, On H-separable extensions of QF-3 rings, 7 pages. 2001.
- #544 A. Arai, Non-relativistic limit of a Dirac-Maxwell operator in relativistic quantum electrodynamics, 27 pages. 2001.
- #545 O. Sawada, On time-local solvability of the Navier-Stokes equations in Besov spaces, 30 pages. 2001.
- #546 C. M. Elliott, Y. Giga, and S. Goto, Dynamic boundary conditions for Hamilton-Jacobi equations, 27 pages. 2001.
- #547 Y. Nakano, Minimizing coherent risk measures of shortfall in discrete-time models with cone constraints, 22 pages. 2002.
- #548 K. Tachizawa, A generalization of the Lieb-Thirring inequalities in low dimensions, 13 pages. 2002.
- #549 T. Nakazi, Absolute values and real parts for functions in the Smirnov class, 8 pages. 2002.
- #550 T. Nakazi and T. Watanabe, Properties of a Rubin's orthogonal function which is a linear combination of two inner functions, 9 pages. 2002.
- #551 T. Ohtsuka, A level set method for spiral crystal growth, 24 pages. 2002.
- #552 M.-H. Giga and Y. Giga, Minimal vertical singular diffusion preventing overturning for the Burgers equation, 18 pages. 2002.
- #553 Y. Giga and P. Rybka, Berg's effect, 12 pages. 2002.
- #554 Y. Tonegawa, Domain dependent monotonicity formula for a singular perturbation problem, 14 pages. 2002.
- #555 S. Izumiya and N. Takeuchi, New special curves and developable surfaces, 10 pages. 2002.
- #556 S. Izumiya, K. Saji and N. Takeuchi, Singularities of line congruences, 17 pages. 2002.
- #557 S. Izumiya, D. Pei and M. C. Romero Fuster, The lightcone Gauss map of a spacelike surface in Minkowski 4-space, 21 pages. 2002.
- #558 S. Izumiya, D. Pei and M. C. Romero Fuster, Umbilicity of spacelike submanifolds of Minkowski space, 14 pages. 2002.
- #559 S. Izumiya, D. Pei and M. Takahashi, Curves and surfaces in Hyperbolic space, 16 pages. 2002.
- #560 J. Kato, On the uniqueness of nondecaying solutions for the Navier-Stokes equations, 19 pages. 2002.
- #561 M. Jinzenji and T. Sasaki, An Approach to $\mathcal{N} = 4$ ADE gauge Theory on $K3$, 29 pages. 2002.
- #562 T. Nakazi and T. Yamamoto, Norms of some singular integral operators on weighted L^2 spaces, 27 pages. 2002.
- #563 A. Harris and Y. Tonegawa, A $\bar{\partial}\partial$ -poincaré lemma for forms near an isolated complex singularity, 8 pages. 2002.
- #564 M. Takahashi, Bifurcations of ordinary differential equations of Clairaut type, 23 pages. 2002.
- #565 G. Ishikawa, Classifying singular Legendre curves by contactomorphisms, 17 pages. 2002.

PERTURBATIONS OF CAUSTICS AND FRONTS

GOO ISHIKAWA

*Department of Mathematics, Hokkaido University,
Sapporo 060-0810, Japan.*

e-mail : ishikawa@math.sci.hokudai.ac.jp

Dedicated to Professor Tatsuo Suwa for his 60th birthday.

1. Introduction. A degenerate singularity of a plane caustic bifurcates into several cusps. Then we may ask:

How many cusps do there appear after a stable perturbation?

In this article, we review the algebraic formula for the number of complex cusps, thus the upper estimate of the number of real cusps, and, gradually, we show their several possible, known and unknown, generalizations.

We start with the ordinary singularity theory. Let $f : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ be an unstable holomorphic map-germ. We consider the problem of counting the number of cusps after a stable perturbation. We note that the cusps appear, for the perturbed f , at a point on the singular locus $J = 0$, defined by the Jacobian $J = J(f_1, f_2)$, where the extended mapping $(J, f_1, f_2) : \mathbf{C}^2 \rightarrow \mathbf{C}^3$ is not immersive. So we count the intersection number at 0 of the jet section $j^2 f : \mathbf{C}^2 \rightarrow J^2(\mathbf{C}^2, \mathbf{C}^2)$ with the variety in $J^2(\mathbf{C}^2, \mathbf{C}^2)$ defined by J and 2-minors of the Jacobi matrix of (J, f_1, f_2) . Since $j^2 f$ is an immersion, setting the ideal

$$\mathcal{J} = \mathcal{J}_{1,1} = \langle J(f_1, f_2), J(f_1, J(f_1, f_2)), J(f_2, J(f_1, f_2)) \rangle_{\mathcal{O}_2},$$

we have:

THEOREM 1.1. ([3], see also [8]) *The number κ of cusps for a stable perturbation $\tilde{f}$ from $f : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$ is equal to the dimension over $\mathbf{C}$ of the quotient algebra (doubly iterated Jacobian algebra) $Q = \mathcal{O}_2 / \mathcal{J}$:*

$$\kappa = \dim_{\mathbf{C}} \mathcal{O}_2 / \langle J(f_1, f_2), J(f_1, J(f_1, f_2)), J(f_2, J(f_1, f_2)) \rangle_{\mathcal{O}_2},$$

1991 *Mathematics Subject Classification*: Primary 58K60; Secondary 13C14, 53D12.
Financial support from Grant-in-Aid for Scientific Research, No. 14340020.

provided $\dim_{\mathbf{C}} Q < \infty$.

Second, we consider the problem on caustics. Let $f : (\mathbf{C}^2, 0) \rightarrow T^*\mathbf{C}^2 = \mathbf{C}^4$ be a Lagrangian immersion. Recall the cotangent bundle $T^*\mathbf{C}^2$ with the canonical coordinates $(p_1, p_2; q_1, q_2)$ has the symplectic 2-form $\omega = dp_1 \wedge dq_1 + dp_2 \wedge dq_2$ and the canonical Lagrangian projection $\pi : T^*\mathbf{C}^2 \rightarrow \mathbf{C}^2$, $\pi(p, q) = q$. We suppose the immersion f satisfies $f^*\omega = 0$. Then $\pi \circ f$ is called the Lagrangian mapping and its critical value set is called *caustics* of f . Now assume f is not necessarily Lagrange stable with respect to π , and ask:

How many cusps do there appear after a Lagrange stable perturbation ?

We denote by $\kappa = \kappa(\pi \circ \tilde{f})$ the number of cusps near the origin, under the projection π , of a Lagrange stable perturbation $\tilde{f}$ of f . Since $\pi \circ \tilde{f}$ is also a stable perturbation of $\pi \circ f$, we can calculate the number of cusps by the above theorem from $\pi \circ f : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$.

In Lagrangian singularity theory, Lagrangian immersions and their caustics are analyzed via their generating families [1]. Thus we are led to also the following question:

What is the algebraic formula for κ in term of the generating family of f ?

Let $F : (\mathbf{C}^r \times \mathbf{C}^2, 0) \rightarrow \mathbf{C}$ be a Morse family. (It is not assumed to be $\mathcal{R}^+$ -stable) [1]. That F is a Morse family means that the *catastroph set*

$$C(F) := \left\{ (x, \lambda) \in \mathbf{C}^r \times \mathbf{C}^2 \mid \frac{\partial F}{\partial x_1} = 0, \dots, \frac{\partial F}{\partial x_r} = 0 \right\}$$

is regularly defined in $\mathbf{C}^r \times \mathbf{C}^2$, namely, $\left(\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_r} \right) : (\mathbf{C}^r \times \mathbf{C}^2, 0) \rightarrow \mathbf{C}^r$ is submersive along $C(F)$. Then the mapping $f = L(F) : C(F) \rightarrow T^*\mathbf{C}^2$ defined by

$$L(F)(x, \lambda) = \left(\frac{\partial F}{\partial \lambda_1}, \frac{\partial F}{\partial \lambda_2}, \lambda_1, \lambda_2 \right),$$

is a Lagrangian immersion. Conversely any Lagrangian immersion can be constructed by this method, up to parametrization. Thus we call F the *generating family* of the Lagrangian immersion f . The singularities of caustics of f coincides with the singularities of the projection $\Pi|_{C(F)} : C(F) \rightarrow \mathbf{C}^2$, where $\Pi : \mathbf{C}^r \times \mathbf{C}^2 \rightarrow \mathbf{C}^2$ is the projection $(x, \lambda) \mapsto \lambda$. The critical value set of $\Pi|_{C(F)}$ is called the *bifurcation set* of F , which agrees with the caustic of the Lagrangian immersion $L(F)$.

Now we set the Hessian of F by

$$J(\nabla F) := J \left(\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_r} \right) = \frac{\partial \left(\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_r} \right)}{\partial (x_1, \dots, x_r)},$$

denoting by $J(a_1, \dots, a_r)$ the Jacobian $\det \left(\frac{\partial a_i}{\partial x_j} \right)$ of $a_1, \dots, a_r$ by $x_1, \dots, x_r$, and the “secondary Hessian” of F by

$$J_i^{(2)}(\nabla F) := J \left(\frac{\partial F}{\partial x_1}, \dots, J^{(i)}(\nabla F), \dots, \frac{\partial F}{\partial x_r} \right),$$

replacing $\frac{\partial F}{\partial x_i}$ by $J(\nabla F)$ for $i = 1, \dots, r$. Then we set

$$\mathcal{J}_{r,1,1} := \left\langle \frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_r}, J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F) \right\rangle_{\mathcal{O}_{r+2}}.$$

Then we have:

THEOREM 1.2. *The number κ of cusps of a Lagrange stable perturbation of $L(F)$ is equal to the dimension $\dim_{\mathbf{C}} Q$ of the doubly iterated Jacobian algebra*

$$Q := \mathcal{O}_{r+2} / \mathcal{J}_{r,1,1},$$

provided $\dim_{\mathbf{C}} Q < \infty$.

Note that, if $\dim_{\mathbf{C}} Q < \infty$, then $\dim_{\mathbf{C}} \mathcal{O}_r / \langle \partial \phi / \partial x_1, \dots, \partial \phi / \partial x_r \rangle < \infty$ for $\phi = F(x, 0)$. Thus ϕ has an $\mathcal{R}^+$ -versal deformation and F is induced from it.

EXAMPLE 1.3. Set $F(x_1, \lambda_1, \lambda_2) = x_1^5 + \lambda_1 x_1^2 + \lambda_2 x_1$, with $r = 1$. Then we have $\frac{\partial F}{\partial x_1} = 5x_1^4 + 2\lambda_1 x_1 + \lambda_2$, $J(\nabla F) = \frac{\partial^2 F}{\partial x_1^2} = 20x_1^3 + 2\lambda_1$, and $J_1^{(2)}(\nabla F) = \frac{\partial^3 F}{\partial x_1^3} = 60x_1^2$. Thus $Q = \mathcal{O}_3 / \langle x_1^2, \lambda_1, \lambda_2 \rangle$. Therefore we have $\kappa = \dim_{\mathbf{C}} Q = 2$.

We give here the proof of Theorem 1.2 shortly, since the proof provides the prototype for the proofs of possible generalizations.

Proof of Theorem 1.2: $C(F)$ is defined by $\nabla F = \left(\frac{\partial F}{\partial x_1}, \dots, \frac{\partial F}{\partial x_r} \right) = 0$ and the singular locus of $\Pi|_{C(F)}$ coincides with the singular locus of $(\nabla F, \lambda_1, \lambda_2) : \mathbf{C}^{r+2} \rightarrow \mathbf{C}^{r+2}$ on $C(F)$, and it is defined by $J(\nabla F) = 0$ on $C(F)$. Moreover the cusp locus coincides with the singular locus of $(\nabla F, J(\nabla F), \lambda_1, \lambda_2) : \mathbf{C}^{r+2} \rightarrow \mathbf{C}^{r+3}$. The $(r+2)$ -minors of its Jacobi matrix are

$$J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F).$$

Now consider the jet space $J^3(\mathbf{C}^{r+2}, \mathbf{C})$. Then jets of Morse families form an open subspace $M \subset J^3(\mathbf{C}^{r+2}, \mathbf{C})$. We set $C := \{j^3 F(x_0) \in M \mid \nabla F(x_0) = 0\}$. Then C is a submanifold of M of codimension r . Now the $(r+2)$ -minors of Jacobi matrix of $(\nabla F, J(\nabla F), \lambda_1, \lambda_2)$, which is considered as an $(r+2) \times (r+3)$ matrix with entries in $\mathcal{O}_C$, defines a determinantal variety $\Sigma \subset C$ of codimension 2. Then Σ is Cohen-Macaulay. Moreover the defining ideal of Σ is reduced over the regular locus of Σ . Then the intersection index of $j^2 F : \mathbf{C}^{r+2} \rightarrow M$ with Σ gives the number of cusps after an $\mathcal{R}^+$ -stable perturbation of F , and it is given by $\dim_{\mathbf{C}} Q$ via $(j^3 F)^* : \mathcal{O}_{r+2} \leftarrow \mathcal{O}_M$. $\square$

Here we have two points for the idea of the proof: First we note that, for instance, $J_i^{(2)}(\nabla F)$ is expressed by a unique (universal) polynomial on partial derivatives of order ≤ 3 of F , for any F . In fact, we use throughout this paper the following well-known fact:

LEMMA 1.4. *Let h be a polynomial function on $J^r(\mathbf{C}^m, \mathbf{C}^p)$. Then, for each i ($1 \leq i \leq m$), there exists a unique polynomial $\frac{d}{dx_i} h$ over $J^{r+1}(\mathbf{C}^m, \mathbf{C}^p)$ such that $\left(\frac{d}{dx_i} h \right) \circ j^{r+1} f = \frac{\partial}{\partial x_i} (h \circ j^r f)$, for any $f : (\mathbf{C}^m, x_0) \rightarrow (\mathbf{C}^p, y_0)$, $x_0 \in \mathbf{C}^m, y_0 \in \mathbf{C}^p$. The operator $h \mapsto \frac{d}{dx_i} h$ is characterised by the properties $\frac{d}{dx_i} (hk) = \left(\frac{d}{dx_i} h \right) k + h \left(\frac{d}{dx_i} k \right)$, and $\frac{d}{dx_i} x_j = \delta_{ij}$,*

$\frac{d}{dx_i}(y_{\alpha,j}) = y_{\alpha+e_i,j}$, where $y_{\alpha,j}(j^r f(x_0)) = \frac{\partial^{|\alpha|} y_j \circ f}{\partial x^\alpha}(x_0)$, and $e_i = (0, \dots, 1, \dots, 0)$ (i -th component).

Second, recall that a local ring A is called *Cohen-Macaulay* if there exists a regular sequence $a_1, \dots, a_n$, ($n = \dim(A)$) belonging to the maximal ideal; a_1 is a non-zero-divisor in A , a_2 is a non-zero-divisor in $A/\langle a_1 \rangle_A$, and so on ([21], Ch. 6). In general the length of a regular sequence does not exceed the dimension of the ring. Therefore a Cohen-Macaulay ring possesses the possibly longest regular sequence, and it is known by this property the intersection theory, or the theory of multiplicity, works very well ([7] Prop. 7.1). A regular local ring, for example $\mathcal{O}_n$, is Cohen-Macaulay. In fact, in $\mathcal{O}_n$, the coordinate functions $x_1, \dots, x_n$ form a regular sequence. If $\mathcal{O}_n/\mathcal{J}$ is Cohen-Macaulay for an ideal $\mathcal{J} \subset \mathcal{O}_n$, then we call the zero locus $Z(\mathcal{J})$ of $\mathcal{J}$ with the defining ideal $\mathcal{J}$ *Cohen-Macaulay variety* in $\mathbf{C}^n$. Non-singular submanifolds are surely Cohen-Macaulay varieties. Another class of important examples for Cohen-Macaulay varieties is given by “determinantal varieties”. In fact we have used the following fundamental algebraic fact:

PROPOSITION 1.5. ([13]. See also [4]) *Let A be a Cohen-Macaulay local ring. Then, for the ideal $\mathcal{J}$ generated by ℓ -minors of an $n \times m$ -matrix with entries in A , the quotient ring $A/\mathcal{J}$ is Cohen-Macaulay, provided $\dim(A) - \dim(A/\mathcal{J}) = (n+1-\ell)(m+1-\ell)$, the “ideal” codimension.*

We have applied Proposition 1.5 in the above proof of Theorem 1.2 to $A = \mathcal{O}_C$, $n = r+2$, $m = r+3$ and $\ell = r+2$. Also we can apply Proposition 1.5 to show Theorem 1.1.

The usage of the Cohen-Macaulay property for the problem of counting perturbed isolated singularities is first considered in [3], along the general framework of intersection theory. Later the original idea is re-explained in [8] for obtaining the new result on the number of double folds. Note that the idea itself of counting perturbed singularities is considered through the deep investigations of germs $\mathbf{C}^2 \rightarrow \mathbf{C}^3$ due to Mond (cf. [22][23][24]). See also [30][28][29][19][20] for the related studies. After [3][8], there appears a sequence of investigations on the problem of counting general isolated Thom-Boardman singularities [26][4][5][6]. Then we observe that there are two main points for obtaining right formulae: one is seeking the appropriate defining ideals of treating singularities on jet spaces and second is seeing their Cohen-Macaulay property. For complicated singularities, both points cause nontrivial problems. In the following sections we collect, around the problems of caustics and wavefronts, several situations where we can overcome these points.

The singularities of generating families of functions can be studied from the viewpoint of singularities of mappings: Let us consider again a Morse family $F : (\mathbf{C}^r \times \mathbf{C}^2, 0) \rightarrow \mathbf{C}$ for plane caustic. Then we consider $G := (F, \Pi) : (\mathbf{C}^r \times \mathbf{C}^2, 0) \rightarrow \mathbf{C} \times \mathbf{C}^2$, $G(x, \lambda) = (F(x, \lambda), \lambda)$. Then the critical locus of G coincides with the catastrophe set $C(F)$ of F . Moreover the cusp (namely, $\Sigma^{r,1,0}$) singular points of G are exactly the fold (namely, $\Sigma^{1,0}$) singular points of $\Pi|_{C(F)}$, while the swallowtail (namely, $\Sigma^{r,1,1,0}$) singular points of G are exactly the cusp (namely, $\Sigma^{1,1,0}$) singular points of $\Pi|_{C(F)}$. The $\mathcal{R}^+$ -stable deformation

of F induces a stable deformation of G . Therefore Theorem 1.2 follows, for instance, from the following (Corollary 4.4 (3) of [4]):

THEOREM 1.6. (Fukui, Nuño Ballesteros, Saia) *The number of $\Sigma^{n-2,1,1,0}$ -points for a stable perturbation of a $\mathcal{K}$ -finite map-germ $G : (\mathbf{C}^n, 0) \rightarrow (\mathbf{C}^3, 0)$ with corank 1 is equal to $\dim_{\mathbf{C}} \mathcal{O}_n / \mathcal{J}_{n-2,1,1}(G)$, where $\mathcal{J}_{n-2,1,1}(G)$ is generated by 3-minors $M_1, \dots$, of Jacobi matrix $\text{Jac}(G)$ of G , $(n-2)$ -minors $N_1, \dots$, of $\text{Jac}(G, M)$ and $(n-2)$ -minors $L_1, \dots$, of $\text{Jac}(G, M, N)$.*

Note that, if $G = (F, \Pi)$, the ideal $\mathcal{J}_{n-2,1,1}(G)$ defined in the above theorem coincides with the ideal treated in Theorem 1.2. Thus we have the second proof of Theorem 1.2.

The problem of counting the number of cusps of caustics is closely related to the problem of counting the number of swallowtails of wave fronts. In fact, $G|_{C(F)} : C(F) \rightarrow \mathbf{C} \times \mathbf{C}^2$ is the wave front of the Legendrian lifting $\tilde{L}(F) : C(F) \rightarrow \mathbf{C} \times T^*\mathbf{C}^2$ of the Lagrangian immersion $L(F) : C(F) \rightarrow T^*\mathbf{C}^2$. In fact $F|_{C(F)} : C(F) \rightarrow \mathbf{C}$ is a generating function of $L(F)$ since $dF = \frac{\partial F}{\partial \lambda_1} d\lambda_1 + \frac{\partial F}{\partial \lambda_2} d\lambda_2$ on $C(F) = \left\{ \frac{\partial F}{\partial x_1} = 0, \dots, \frac{\partial F}{\partial x_r} = 0 \right\}$. Thus actually the formula of Theorem 1.2 can be interpreted, from the singularity theory of mappings, as the formula for the number of swallowtails of perturbed wave front. Of course, since swallowtails project to cusps, we have the same formula.

In Legendrian singularity theory, the singularities of wave fronts are analysed via generating families of hypersurfaces [1]. So we give the counting formula for swallowtails of wave fronts in the language of generating families of hypersurfaces.

Let $H : (\mathbf{C}^r \times \mathbf{C}^3, 0) \rightarrow (\mathbf{C}, 0)$ be a function family. Assume the equation $H = 0$ defines a Morse family of hypersurfaces in $\mathbf{C}^r$. This means that the *big singular set*

$$\tilde{C}(H) := \left\{ (x, \lambda) \mid H = 0, \frac{\partial H}{\partial x_1} = 0, \dots, \frac{\partial H}{\partial x_r} = 0 \right\},$$

is non-singular. The image $\Pi(\tilde{C}(H))$ of $\tilde{C}(H)$ under the projection $\Pi : \mathbf{C}^r \times \mathbf{C}^3 \rightarrow \mathbf{C}^3$ is called the *discriminant* of H . This coincides with the front of the Legendrian immersion defined by H . In fact the mapping $\tilde{L}(H) : \tilde{C}(H) \rightarrow PT^*\mathbf{C}^3$ defined by

$$\tilde{L}(H)(x, \lambda) = \left(\left[\frac{\partial H}{\partial \lambda_1}, \frac{\partial H}{\partial \lambda_2}, \frac{\partial H}{\partial \lambda_3} \right], \lambda_1, \lambda_2, \lambda_3 \right)$$

is a Legendre immersion in the projective cotangent bundle $PT^*\mathbf{C}^3$ over $\mathbf{C}^3$. For the Legendre fibration $\pi : PT^*\mathbf{C}^3 \rightarrow \mathbf{C}^3$, the front of $\tilde{L}(H)$, that is the image of $\pi \circ \tilde{L}(H)$, coincides with the discriminant $\Pi(\tilde{C}(H))$. Any Legendre immersion to $PT^*\mathbf{C}^3$ is obtained by this process, up to parametrization.

THEOREM 1.7. *The number of swallowtails in the fronts of a stable perturbation of $\tilde{L}(H)$ is equal to the dimension $\dim_{\mathbf{C}} \tilde{Q}$ of the algebra*

$$\tilde{Q} := \mathcal{O}_{r+3} / \left\langle H, \frac{\partial H}{\partial x_1}, \dots, \frac{\partial H}{\partial x_r}, J(\nabla H), J_1^{(2)}(\nabla H), \dots, J_r^{(2)}(\nabla H) \right\rangle_{\mathcal{O}_{r+3}},$$

provided $\dim_{\mathbf{C}} \tilde{Q} < \infty$.

Proof: Up to $\mathcal{K}$ -equivalence, we may assume $H = F - \lambda_3$ for a Morse family $F(x, \lambda_1, \lambda_2)$. Then we apply Theorem 1.6 to $G : (\mathbf{C}^r \times \mathbf{C}^2, 0) \rightarrow \mathbf{C} \times \mathbf{C}^2$, where $G(x, \lambda_1, \lambda_2) = (F(x, \lambda), \lambda_1, \lambda_2)$. $\square$

Note that it is known ([27][18][11][12]) the more geometric formula on the number of cusps appering in a generic plane section of the discsriminant by means of Milnor number. However it seems to be open yet to relate them to the formula in term of the iterated Jacobians.

In the following sections we consider several generalizations of the above results including singular cases and try to understand them clearly: In §2, we introduce the counting formulae for the numbers of A_4 and D_4 singularities of caustics due to Fukui and Weyman. In §3, we count the numbers of simplest singularities of singular Lagrangian immersions: open Whitney umbrellas. In §4, we count isolated singularities of singular Lagrangian immersions composed with Lagrangian projections. Actually, as the result obtained firstly in this paper, we give the formulae for the numbers of singularities S_3, S_4, S_5 and T_5 introduced in [2].

In the last section, we give open questions. We do not treat, in this paper, the interesting problem on the algebraic equations for the number of real cusps and other isolated singularities over the real. Also we do not treat the topological invariance of the complex or real numbers of isolated singularities and Euler characteristics of non-isolated singularities of stable perturbations.

The present paper is based partly on my talk under the same title at Yokohama, Japan, on November 2000, in the occasion of the celebration for the 60th birthdays of Professor Fukuda and of Professor Izumi. I would like to thank Professor Toshizumi Fukui and Professor Jerzy Weyman for their encouragement to my writing of this note by their asking the question on perturbations of caustics. I would like to thank Professor D. Siersma for his reminding me the other geometric formulae on the numbers of cusps. I would like to thank organizers of the conference Caustics'02, for their encouragement to my completing of the present paper.

2. Space caustics. So far we have investigated the number of A_3 singularities of caustics on the plane. Then it is natural to ask about the number of A_4 and D_4 singularities of caustics in the three space [1].

REMARK 2.1. For the number of A_2 -singularities of a generating family $F : (\mathbf{C}^r \times \mathbf{C}, 0) \rightarrow \mathbf{C}$, $F = F(x, \lambda)$ is given by $\dim_{\mathbf{C}} \mathcal{O}_{r+1} / \mathcal{J}_{r,1}$, $\mathcal{J}_{r,1} = \langle \nabla F, J(\nabla F) \rangle$.

Let $F : (\mathbf{C}^r \times \mathbf{C}^3, 0) \rightarrow \mathbf{C}$ be a generating Morse family of a Lagrangian immersion $f : (\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$. The fold (A_2) locus of $\Pi|_{C(F)}$ is defined by $\nabla F = 0$ and $J(\nabla F) = 0$. The cusp (A_3) locus of $\Pi|_{C(F)}$ is defined by $\nabla F, J(\nabla F)$ and $J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F)$. Then, for the description of swallowtail (A_4) locus of $\Pi|_{C(F)}$, we intend to consider the

mapping

$$(\nabla F, J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F), \lambda_1, \lambda_2, \lambda_3) : (\mathbf{C}^{r+3}, 0) \rightarrow (\mathbf{C}^{2r+4}, 0),$$

as in the proof of Theorem 1.2. However actually there is no reason to expect that its singular locus coincides with the swallowtail locus of $\Pi|_{C(F)}$, because

$$\nabla F, J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F)$$

are far from independent. In fact, in [4][5], it is observed that the singular locus includes the locus of D_4 as well as the locus of A_4 .

Anyway we define the ideal $\mathcal{J}_{r,1,1,1}$ in $\mathcal{O}_{r+3}$ generated by

$$\nabla F, J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F)$$

and Jacobians of r elements from $\nabla F, J(\nabla F), J_1^{(2)}(\nabla F), \dots, J_r^{(2)}(\nabla F)$ with respect to $x_1, \dots, x_r$. Moreover we define the ideal $\mathcal{J}_{r,2}$ in $\mathcal{O}_{r+3}$ generated by ∇F and $(r-1)$ -minors of the Jacobi matrix $\text{Jac}(\nabla F)$ of ∇F . Then the following result is a rewrite of the remarkable Theorem 3.2 of [5] in the language of generating family.

THEOREM 2.2. (Fukui-Weyman [5]) *Let $F : (\mathbf{C}^r \times \mathbf{C}^3, 0) \rightarrow \mathbf{C}$ be a Morse family. Then we have the formulae for the number $\#D_4$ of D_4 -singular points and the number $\#A_4$ of A_4 -singular points, respectively, after a Lagrange stable perturbation:*

$$\#D_4 = \dim_{\mathbf{C}} \mathcal{O}_{r+3} / \mathcal{J}_{r,2},$$

and

$$\#A_4 = \dim_{\mathbf{C}} \mathcal{O}_{r+3} / \mathcal{J}_{r,1,1,1} - 4 \dim_{\mathbf{C}} \mathcal{O}_{r+3} / \mathcal{J}_{r,2},$$

provided $\dim_{\mathbf{C}} \mathcal{O}_{r+3} / \mathcal{J}_{r,2} < \infty$ and $\dim_{\mathbf{C}} \mathcal{O}_{r+3} / \mathcal{J}_{r,1,1,1} < \infty$.

EXAMPLE 2.3. Consider the generating family of type D_4 :

$$F(x_1, x_2, \lambda_1, \lambda_2, \lambda_3) = x_1^2 x_2 + x_2^3 + \lambda_1 x_2^2 + \lambda_2 x_1 + \lambda_3 x_2,$$

($r = 2$). Then

$$\dim_{\mathbf{C}} \mathcal{O}_5 / \mathcal{J}_{2,2} = \dim_{\mathbf{C}} \mathcal{O}_5 / \left\langle \frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial x_2}, \frac{\partial^2 F}{\partial x_1^2}, \frac{\partial^2 F}{\partial x_1 \partial x_2}, \frac{\partial^2 F}{\partial x_2^2} \right\rangle_{\mathcal{O}_5} = 1.$$

Moreover we have

$$\mathcal{J}_{2,1,1,1} = \langle x_1^2, x_1 x_2, x_2^2, \lambda_1 x_1, \lambda_1 x_2, \lambda_1^2, \lambda_2, \lambda_3 \rangle_{\mathcal{O}_5}.$$

Thus we see $\mathcal{O}_5 / \mathcal{J}_{2,1,1,1}$ has a basis $\{1, x_2, x_3, \lambda_1\}$ and

$$\dim_{\mathbf{C}} \mathcal{O}_5 / \mathcal{J}_{2,1,1,1} = 4,$$

for D_4 . On the other hand, for the generating family of type A_4 :

$$F(x_1, x_2, \lambda_1, \lambda_2, \lambda_3) = x_1^5 + x_2^2 + \lambda_1 x_1^3 + \lambda_2 x_1^2 + \lambda_3 x_1,$$

($r = 2$), we have $\dim_{\mathbf{C}} \mathcal{O}_5 / \mathcal{J}_{2,2} = 0$ and $\mathcal{O}_5 / \mathcal{J}_{2,1,1,1} = 1$. $\square$

Let $f : (\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$ be a Lagrangian immersion and $\pi : T^*\mathbf{C}^3 \rightarrow \mathbf{C}^3$ the canonical projection. We can re-formulate Theorem 2.2, without using the term of generating family, as follows: Considering the composition $\pi \circ f = (q_1 \circ f, q_2 \circ f, q_3 \circ f)$, we define $\mathcal{J}_2$ as the ideal generated by the 2-minors of the Jacobi matrix $\text{Jac}(q_1 \circ f, q_2 \circ f, q_3 \circ f)$ and define

$\mathcal{J}_{1,1,1}$ as the ideal generated by the Jacobian $J = J(q_1 \circ f, q_2 \circ f, q_3 \circ f)$, the secondary Jacobians

$$J_1^{(2)} = (J, q_2 \circ f, q_3 \circ f), J_2^{(2)} = J(q_1 \circ f, J, q_3 \circ f), J_3^{(2)} = J(q_1 \circ f, q_2 \circ f, J)$$

and the ternary Jacobians, namely, 3-minors of

$$\text{Jac}(q_1 \circ f, q_2 \circ f, q_3 \circ f, J, J_1^{(2)}, J_2^{(2)}, J_3^{(2)}).$$

Then we have:

COROLLARY 2.4. *Let $f : (\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$ be a Lagrangian immersion. Then the number $\#D_4$ of D_4 -singularities and $\#A_4$ of A_4 -singularities respectively appearing in a Lagrange stable perturbation of f is given by*

$$\#D_4 = \dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_2, \quad \#A_4 = \dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_{1,1,1} - 4 \dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_2,$$

provided $\dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_2 < \infty$ and $\dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_{1,1,1} < \infty$.

Proof: Let $F : (\mathbf{C}^r \times \mathbf{C}^3, 0) \rightarrow \mathbf{C}$ be a generating family of f . We may suppose $\varphi := \pi_r|_{C(F)} : (C(F), 0) \rightarrow (\mathbf{C}^r, 0)$, $C(F)$ beeing the catastroph set-germ, is a complex analytic isomorphism. Also consider the inclusion $i : (C(F), 0) \rightarrow (\mathbf{C}^r \times \mathbf{C}^3, 0)$. Then the composition of $i^* : \mathcal{O}_{r+3} \rightarrow \mathcal{O}_{C(F)}$ with the inverse of $\varphi^* : \mathcal{O}_r \rightarrow \mathcal{O}_{C(F)}$ induces an isomorphisms

$$\mathcal{O}_{r+3}/\mathcal{J}_{r,2} \cong \mathcal{O}_3/\mathcal{J}_2, \quad \text{and} \quad \mathcal{O}_{r+3}/\mathcal{J}_{r,1,1,1} \cong \mathcal{O}_3/\mathcal{J}_{1,1,1}.$$

□

Here we also give the result on the numbers of A_4 and D_4 for generating families $H(x_1, \dots, x_r, \lambda_0, \lambda_1, \dots, \lambda_3) = 0$ on $(\mathbf{C}^r \times \mathbf{C}^4, 0)$ of Legendre immersions $(\mathbf{C}^3, 0) \rightarrow PT^*\mathbf{C}^4$. We define, for H , similarly as above, ideals $\tilde{\mathcal{J}}_{r,2}$ and $\tilde{\mathcal{J}}_{r,1,1,1}$ in $\mathcal{O}_{r+4}$, but adding H to the generators of $\mathcal{J}_{r,2}$ and $\mathcal{J}_{r,1,1,1}$ respectively. Then we have

COROLLARY 2.5. *Let $H : (\mathbf{C}^r \times \mathbf{C}^4, 0) \rightarrow (\mathbf{C}, 0)$ be a Morse denenerating family of hypersurfaces for a Legendrian immersion $\tilde{L}(H) : \tilde{C}(H) \rightarrow PT^*\mathbf{C}^4$. Then the numbers $\#D_4$ and $\#A_4$ of D_4 and A_4 singularities respectively of a Legendrian perturbation of $\tilde{L}(H)$ are given by:*

$$\#D_4 = \dim_{\mathbf{C}} \mathcal{O}_{r+4}/\tilde{\mathcal{J}}_{r,2},$$

and

$$\#A_4 = \dim_{\mathbf{C}} \mathcal{O}_{r+4}/\tilde{\mathcal{J}}_{r,1,1,1} - 4 \dim_{\mathbf{C}} \mathcal{O}_{r+4}/\tilde{\mathcal{J}}_{r,2},$$

provided $\dim_{\mathbf{C}} \mathcal{O}_{r+4}/\tilde{\mathcal{J}}_{r,2} < \infty$ and $\dim_{\mathbf{C}} \mathcal{O}_{r+4}/\tilde{\mathcal{J}}_{r,1,1,1} < \infty$.

REMARK 2.6. The counting of isolated singularities of caustics in the four space has several difficulties: The ideals asociated to Thom-Boardman singularites are actually defined in general ([25][26][4][5]): They are called *Morin ideals* of Thom-Boardman singularites. However, for A_5 -singularities or $\Sigma^{r,1,1,1,1}$ -singularities, the Morin ideal does not define a Cahen-Macaulay variety (Theorem 3.1 of [5]). Moreover we do not know the coefficients of necessary correcting terms for the counting of the number of $\Sigma^{r,1,1,1,1}$ -singularities. Furthermore, we note that D_5 -singularity of caustics is no longer a Thom-

Boardman singularity. So we need, in general, the method to find the defining ideals for closures of $\mathcal{K}$ -orbits in jet spaces.

3. Open Whitney umbrellas. Let $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ be an isotropic mapping; $f^*\omega = 0$. In this section and the next section, we do not assume f is an immersion (i.e. of corank 0). Instead we do assume f is of corank ≤ 1 .

In this section we do not treat singularities of caustics nor singularities of Lagrangian projections. Instead we do treat the singularities of f up to the symplectic equivalence [16]. Actually we consider Thom-Boardman singularities of type $\Sigma^{1,1,\dots}$ for isotropic mappings.

In this paper, by the isotropic jet space, we mean the space of isotropic jets of corank ≤ 1 :

$$J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n) := \{j^r f(x) \mid x \in \mathbf{C}^n, f : (\mathbf{C}^n, x) \rightarrow T^*\mathbf{C}^n, f^*\omega = 0, \text{ corank}(f) \leq 1\}.$$

Moreover we set

$$J_I^r(n, 2n) := \{j^r f(0) \mid f : (\mathbf{C}^n, 0) \rightarrow (T^*\mathbf{C}^n, 0), f^*\omega = 0, \text{ corank}(f) \leq 1\}.$$

Recall that the Thom-Boardman singularity $\Sigma^{1_k} = \Sigma^{1,1,\dots}$ (k -times) is defined in the ordinary jet space $J^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ ($k \leq r$); $\Sigma^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n) \subseteq J^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ is a submanifold of codimension $(n+1)k$. Then simply we set

$$\Sigma_I^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n) := \Sigma^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n) \cap J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n).$$

Similarly, for $\Sigma^{1_k}(n, 2n) \subseteq J^r(n, 2n)$, we set

$$\Sigma_I^{1_k}(n, 2n) := \Sigma^{1_k}(n, 2n) \cap J_I^r(n, 2n).$$

Then, via the trivialization

$$J^r(\mathbf{C}^n, T^*\mathbf{C}^n) \cong \mathbf{C}^n \times T^*\mathbf{C}^n \times J^r(n, 2n),$$

we have

$$J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n) \cong \mathbf{C}^n \times T^*\mathbf{C}^n \times J_I^r(n, 2n)$$

and

$$\Sigma_I^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n) \cong \mathbf{C}^n \times T^*\mathbf{C}^n \times \Sigma_I^{1_k}(n, 2n).$$

We abbreviate $J^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ by J^r , and similarly we abbreviate $J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ (resp. $\Sigma_I^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n)$) by J_I^r (resp. $\Sigma_I^{1_k}$).

It is known that the isotropic map-germs of corank ≤ 1 is essentially described by $J^r(1, 2)$ up to symplectic equivalence ([14][15][16][17]). In particular we have:

LEMMA 3.1. ([14]) $\Sigma_I^{1_k}(n, 2n)$ is a submanifold of $J_I^r(n, 2n)$ of codimension $2k$.

$\Sigma_I^{1_k} = \Sigma_I^{1_k}(\mathbf{C}^n, T^*\mathbf{C}^n)$ is a submanifold of $J_I^r = J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ of codimension $2k$.

Generic isotropic mappings of corank ≤ 1 are classified into *open Whitney umbrellas* of type k ($0 \leq k \leq [n/2]$) ([14]). They are characterised by the symplectic stability [16]. Among them, the classification coincides with the classification by Thom-Boardman singularities: An open Whitney umbrella $f : (\mathbf{C}^n, x) \rightarrow (T^*\mathbf{C}^n, f(x))$ is of type k if and only if $j^r f(x) \in \Sigma_I^{1_k, 0} (= \Sigma_I^{1_k} \setminus \Sigma_I^{1_{k+1}})$. We abbreviate an open Whitney umbrella of type k by OWU_k .

For an isotropic mapping $f : (\mathbf{C}^{2n}, 0) \rightarrow T^*\mathbf{C}^{2n}$, open Whitney umbrellas of type n appear isolatedly by Lemma 3.1 after a symplectically stable perturbation. Then we naturally have the question:

How many open Whitney umbrellas of isolated type do there appear after a symplectically stable perturbation?

We define, in general, for a map-germ $f : (\mathbf{C}^m, 0) \rightarrow (\mathbf{C}^p, 0)$ with $m \leq p$, the ideal $\mathcal{J}_{1_k}$ of $\mathcal{O}_m$ generated by m -minors $M_1^{(1)}, M_2^{(1)}, \dots$, of the Jacobi matrix (an $m \times p$ -matrix) $\text{Jac}(f)$ of f , m -minors $M_1^{(2)}, M_2^{(2)}, \dots$, of the Jacobi matrix (an $m \times p + {}_pC_m$ -matrix) $\text{Jac}(f, M^{(1)})$, m -minors $M_1^{(3)}, M_2^{(3)}, \dots$, of $\text{Jac}(f, M^{(1)}, M^{(2)})$, $\dots$, and lastly m -minors of $\text{Jac}(f, M^{(1)}, M^{(2)}, \dots, M^{(k-1)})$.

Then we have:

PROPOSITION 3.2. *Let $f : (\mathbf{C}^{2n}, 0) \rightarrow T^*\mathbf{C}^{2n}$ be an isotropic map-germ of corank ≤ 1 . Then the number $\#\text{OWU}_n$ of open Whitney umbrellas of type n after a symplectically stable perturbation of f is given by*

$$\#\text{OWU}_n = \dim_{\mathbf{C}} \mathcal{O}_{2n} / \mathcal{J}_{1_n},$$

provided $\dim_{\mathbf{C}} \mathcal{O}_{2n} / \mathcal{J}_{1_n} < \infty$.

Proof: The corresponding ideal to $\mathcal{J}_{1_n}$ in $\mathcal{O}_{J^n}$ is a regular defining ideal of Σ^{1_n} . Moreover it gives a regular defining ideal in $\mathcal{O}_{J^n}$ of $\Sigma_I^{1_n}$, which is non-singular of codimension $2n$. This means that, at each point of $\Sigma_I^{1_n}$, there are $2n$ elements from the ideal which give a submersion at the point. The number of open Whitney umbrellas of type n after a symplectically stable perturbation is equal to the intersection number of $J^n f(\mathbf{C}^{2n})$ and $\Sigma_I^{1_n}$ in $J^n = J^n(\mathbf{C}^{2n}, T^*\mathbf{C}^{2n})$. This implies the result. $\square$

In particular, setting $n = 1$ in Proposition 3.2, we have the following result.

COROLLARY 3.3. *Let $f : (\mathbf{C}^2, 0) \rightarrow T^*\mathbf{C}^2$ be an isotropic map-germ of corank ≤ 1 . Denote by u the number of open Whitney umbrellas of symplectically stable perturbations of f . Then*

$$u = \dim_{\mathbf{C}} \mathcal{O}_2 / \mathcal{J}_1,$$

where $\mathcal{J}_1$ is the ideal generated by 2-minors of Jacobi matrix $\text{Jac}(f)$ of f .

We apply the above result to the problem of counting the number of folded umbrellas of fronts.

A map-germ $g : (\mathbf{C}^n, 0) \rightarrow (\mathbf{C}^{n+1}, 0)$ with dense immersion locus is called a *frontal map-germ* if, for any $t \in (\mathbf{C}^n, 0)$, there exists unique limit of tangent spaces through immersive points $\lim_{s \rightarrow t} g_*(T_s \mathbf{C}^n) =: T_t$ depending smoothly on t .

For example, the swallowtail $(x_1, x_2) \mapsto (x_1, x_2^3 + x_1 x_2, \frac{3}{4}x_2^4 + \frac{1}{2}x_1 x_2^2)$ and the folded umbrella $(x_1, x_2) \mapsto (x_1, x_2^2, x_1 x_2^3)$ are frontal, while the Whitney umbrella $(x_1, x_2) \mapsto (x_1, x_2^2, x_1 x_2)$ are not frontal, when they are regarded as map-germs $(\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^3, 0)$.

Let $g : (\mathbf{C}^n, 0) \rightarrow (\mathbf{C}^{n+1}, 0)$ be frontal. Then the map-germ to the Grassmannian

$$G : (\mathbf{C}^n, 0) \rightarrow \text{Gr}(n, T\mathbf{C}^{n+1}) = PT^*\mathbf{C}^{n+1}$$

defined by $G(t) = T_t$ is an *integral* map-germ: If we consider the contact form $\alpha = dr - \sum_{i=1}^n p_i dq_i$ on $\mathbf{C}^{2n+1} = J^1(\mathbf{C}^n, \mathbf{C}) \subset PT^*\mathbf{C}^{n+1}$, then $G^*\alpha = 0$ provided $G(0) \in J^1(\mathbf{C}^n, \mathbf{C})$. For integral mappings see [1]. Then we have the following:

PROPOSITION 3.4. *Let $g : (\mathbf{C}^{2n}, 0) \rightarrow (\mathbf{C}^{2n+1}, 0)$ be a frontal mapping with the integral lifting $G : (\mathbf{C}^{2n}, 0) \rightarrow J^1(\mathbf{C}^{2n}, \mathbf{C}) = \mathbf{C}^{4n+1}$ of corank ≤ 1 . Set*

$$G = (g_1, \dots, g_{2n}, e; a_1, \dots, a_{2n})$$

with $de = a_1 df_1 + \dots + a_{2n} df_{2n}$. Then the number $\#FU_n$ of folded umbrellas of type n after a frontal stable perturbation of g is equal to $\#OWU_n$ for the isotropic map-germ $f = (g_1, \dots, g_{2n}, a_1, \dots, a_{2n}) : (\mathbf{C}^{2n}, 0) \rightarrow T^\mathbf{C}^{2n}$. Therefore*

$$\#FU_n = \dim_{\mathbf{C}} \mathcal{O}_{2n} / \mathcal{I}_n(f).$$

□

In particular consider the case $n = 2$. It is known that any integral map $G : (\mathbf{C}^2, 0) \rightarrow \text{Gr}(2, T\mathbf{C}^2)$ of corank ≤ 1 is approximated by an integral map $\tilde{G}$ such that $\tilde{g} = \pi \circ \tilde{G}$ has only swallowtails and folded umbrellas as singularities (cf. [9]). Denote by s (resp. u) the number of swallowtails (resp. folded umbrellas) for perturbations $\tilde{g} = \pi \circ \tilde{G}$ from g . Then we have:

COROLLARY 3.5. *Let $g : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^3, 0)$ be a frontal map-germ with the integral lifting $G : (\mathbf{C}^2, 0) \rightarrow J^1(\mathbf{C}^2, \mathbf{C}) = \mathbf{C}^5$ of corank ≤ 1 . Set $G = (g_1, g_2, e; a_1, a_2)$ with $de = a_1 dg_1 + a_2 dg_2$. Then we have $s = \dim_{\mathbf{C}} Q$ with*

$$Q := \mathcal{O}_2 / \langle J(g_1, g_2), J(g_1, J(g_1, g_2)), J(J(g_1, g_2), g_2) \rangle_{\mathcal{O}_2}.$$

Moreover we have $u = \dim_{\mathbf{C}} Q'$ with

$$Q' := \mathcal{O}_2 / \left\langle \begin{array}{ccc} J(g_1, g_2), & J(g_1, a_1), & J(g_1, a_2) \\ J(g_2, a_1), & J(g_2, a_2), & J(a_1, a_2) \end{array} \right\rangle_{\mathcal{O}_2}.$$

REMARK 3.6. Lowering more the dimension, consider a plane curve $g : (\mathbf{C}, 0) \rightarrow (\mathbf{C}^2, 0)$ as front. Then the number κ of cusps of frontal perturbations of g is equal to the order of the plane curve-germ g : $\kappa = \text{ord}(g)$.

4. Supersingularities of caustics. In this section we count isolated singularities of isotropic mappings relatively to the Lagrangian projections, namely singularities of composed isotropic mapping $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ with the Lagrangian fibration $\pi : T^*\mathbf{C}^n \rightarrow \mathbf{C}^n$. If f is an immersion, then f is a Lagrangian immersion, and the composed mapping $\pi \circ f$ is exactly the Lagrangian mapping of f . In general, the singularity of composed mapping is not necessarily reflected to the singularity of the caustic; the set of critical values of the composed mapping $\pi \circ f$. This is the reason to use the term “supersingularity”.

We denote by $\pi : T^*\mathbf{C}^n \rightarrow \mathbf{C}^n$ the canonical projection. We define Thom-Boardman singularity $\Sigma_{i_1, i_2, \dots, I}^{1_k}$ for isotropic mappings of corank ≤ 1 by

$$\Sigma_I^{1_k} \cap \pi_r^{-1}(\Sigma^{i_1, i_2, \dots}) \subset J_I^r = J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n).$$

Here we use the letter “ I ” for “isotropic”, and denote by

$$\pi_r : J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n) \rightarrow J^r(\mathbf{C}^n, \mathbf{C}^n)$$

the projection induced by $\pi : T^*\mathbf{C}^n \rightarrow \mathbf{C}^n$. Note that the closure $\overline{\Sigma_{i_1, i_2, \dots, I}^{1_k}}$ in $J_I^r = J_I^r(\mathbf{C}^n, T^*\mathbf{C}^n)$ is equal to

$$\Sigma_I^{1_k} \cap \pi_r^{-1}(\overline{\Sigma_{i_1, i_2, \dots}^{1_k}}).$$

In [2], we classify simple Lagrangian projections of open Whitney umbrellas of type 1. Moreover we know the “nice range” for isotropic mappings $\mathbf{C}^n \rightarrow T^*\mathbf{C}^n$ of corank ≤ 1 under the Lagrangian equivalence is $\{n \in \mathbf{N} \mid n \leq 4\}$. This means that any isotropic map-germ $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ of corank ≤ 1 is perturbed into an isotropic mapping with only Lagrange stable germs at any point in a sufficiently small neighbourhood of 0, provided $n \leq 4$. Note that a generic isotropic map-germ $(\mathbf{C}^4, 0) \rightarrow T^*\mathbf{C}^4$ of type $\Sigma^{1,1}$ has unique Lagrange equivalence class (Theorem 3 of [14]). Also note that the nice range for Lagrangian immersions is $\{n \in \mathbf{N} \mid n \leq 5\}$ ([1][17]).

Now we consider the supersingularities of caustics with $n \leq 4$. From the classification in [2], we have:

THEOREM 4.1. *Let $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ be an isotropic map-germs of corank ≤ 1 with $n = 1, 2, 3$ or 4. Then f has a Lagrange stable perturbation with singularities only in the following list:*

- (1) $n = 1$. $A_2 = \Sigma_{1,0,I}^0$ the fold.
- (2) $n = 2$. In addition, $A_3 = \Sigma_{1,1,0,I}^0$ the cusp, and $S_3 = \Sigma_{1,0,I}^{1,0}$ the (original) open Whitney umbrella.
- (3) $n = 3$. In addition, $A_4 = \Sigma_{1,1,1,0,I}^0$ the swallowtail, $D_4 = \Sigma_{2,0,I}^0$ the umbilic, and $S_4 = \Sigma_{1,1,0,I}^{1,0}$.
- (4) $n = 4$. In addition, $A_5 = \Sigma_{1,1,1,0,I}^0$, D_5 , $S_5 = \Sigma_{1,1,1,0,I}^{1,0}$, $T_5 = \Sigma_{2,0,I}^{1,0}$ and $W_5 = \Sigma_{1,1,0,I}^{1,1,0}$.

REMARK 4.2. Note that, in the case $n = 4$, as a generic isotropic mapping, the open Whitney umbrellas of type 2 appear as well as that of type 1. Thus we must put, in addition to the list from Theorem 1 of [2], the Lagrangian equivalence classes of generic open Whitney umbrellas of type 2. However generic projections of open Whitney umbrellas of type 2 has unique Lagrangian equivalence class by Theorem 3 of [14]. In this paper we denote the unique class by W_5 . It was denoted in [14] by $f_{4,2}$. The singularity T_5 , found by Ilya Bogaevski([2]) is given by a stable projection of corank two at the singular point of the open Whitney umbrella of type 1 in the smallest dimension.

Now naturally we ask:

What is the algebraic formula for the number of isolated supersingularities after a Lagrange stable perturbation of an isotropic map-germ of corank one?

We give the formulae for the numbers of $A_2, A_3, S_3, S_4, S_5, W_5$ and T_5 for isotropic map-germs of corank one. For S_5 , yet we have an inequality (Proposition 4.9). We observe naturally a possible similarity between the pair (D_4, A_4) and the pair (T_5, S_5) . Thus we conjecture that the equality for S_5 in Proposition 4.9 holds.

Now we start with A_2 . The following fact is very easy to see:

PROPOSITION 4.3. *Let $f : (\mathbf{C}, 0) \rightarrow T^*\mathbf{C}$ be a map-germ. Then the number of A_2 -singular points appearing in a Lagrange stable perturbation of f is given by $\#A_2 = \dim \mathcal{O}_1/\mathcal{J}_1$ provided $\dim \mathcal{O}_1/\mathcal{J}_1 < \infty$, where $\mathcal{J}_1 := \langle J(\pi \circ f) \rangle$.*

Next we turn to A_3 . The formula of A_3 for a Lagrangian immersion is given in §1. Also for isotropic map-germ of corank one, we have the following:

PROPOSITION 4.4. *Let $f : (\mathbf{C}^2, 0) \rightarrow T^*\mathbf{C}^2$ be an isotropic map-germ. Then the number of A_3 -singular points appearing in a Lagrange stable perturbation of f is given by*

$$\#A_3 = \dim \mathcal{O}_2/\mathcal{J}_{1,1}$$

provided $\dim \mathcal{O}_2/\mathcal{J}_{1,1} < \infty$, where we set as before

$$\mathcal{J}_{1,1} := \langle J(f_1, f_2), J(f_1, J(f_1, f_2)), J(J(f_1, f_2), f_2) \rangle,$$

setting $\pi \circ f = (f_1, f_2) : (\mathbf{C}^2, 0) \rightarrow (\mathbf{C}^2, 0)$.

Proof: Since a Lagrange stable perturbation of f induces a stable perturbation of $\pi \circ f$, in this case, Proposition 4.4 follows from Theorem 1.1. $\square$

Next we proceed to count supersingularities. First we note that $\#S_3 = \#\text{OWU}_1$ (resp. $\#W_5 = \#\text{OWU}_2$) for a Lagrange stable perturbation of $f : (\mathbf{C}^2, 0) \rightarrow T^*\mathbf{C}^2$ (resp. $f : (\mathbf{C}^4, 0) \rightarrow T^*\mathbf{C}^4$). Thus already we have given the formulae for S_3 and W_5 in §3.

Now we study on S_4 , namely, $\Sigma_{1,1,0,I}^{1,0}$ singularity.

Let $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ be an isotropic map-germ. Consider the ideal $\mathcal{J}_{1,1}^1$ in $\mathcal{O}_n$ generated by n -minors of the Jacobi matrix $\text{Jac}(f)$ and n -minors of $\text{Jac}(J(\pi \circ f), \pi \circ f)$.

LEMMA 4.5. *The ideal $\tilde{\mathcal{J}}_{1,1}^1$ in $\mathcal{O}_{J_I^2}$ induced naturally from $\mathcal{J}_{1,1}^1$ defines the Cohen-Macaulay varieties $\overline{\Sigma_{1,1,I}^1}$ of codimension 3 in J_I^2 . $\hat{\mathcal{J}}$ is regular on the submanifold $\Sigma_{1,1,I}^1$ of codimension 3 in J_I^2 . Moreover $\overline{\Sigma_{1,1,I}^1} \setminus \Sigma_{1,1,I}^1$ has codimension > 3 in J_I^2 .*

Proof: Since $\hat{\mathcal{J}}_{1,1}^1$ defines $\overline{\Sigma_{1,1,I}^1}$, which a hypersurface in the submanifold Σ_I^1 , we see $\overline{\Sigma_{1,1,I}^1}$ with the ideal $\hat{\mathcal{J}}_{1,1}^1$ is a Cohen-Macaulay variety. Note that $\text{codim}(\Sigma_{1,1,I}^1) = 3$. For the typical singularity S_4 , $f : (\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$, we have $\dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_{1,1}^1 = 1$ by a simple calculation. This shows that $\hat{\mathcal{J}}_{1,1}^1$ is regular on $\Sigma_{1,1,I}^1$. Since

$$\overline{\Sigma_{1,1,I}^1} \setminus \Sigma_{1,1,I}^1 = \overline{\Sigma_{2,I}^1}$$

is of codimension 4, we have the result. $\square$

In particular we have the following formula, by Lemma 4.5, and by counting the intersection number of $j^2 f(\mathbf{C}^3)$ and $\overline{\Sigma_{1,1,I}^1}(\mathbf{C}^3, T^*\mathbf{C}^3)$ in $J_I^2 = J_I^2(\mathbf{C}^3, T^*\mathbf{C}^3)$:

THEOREM 4.6. *Let $f : (\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$ be an isotropic map-germ of corank 1. Then the number of S_4 -singularities of a Lagrangian stable perturbation of f is given by*

$$\#S_4 = \dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_{1,1}^1,$$

provided $\dim_{\mathbf{C}} \mathcal{O}_3/\mathcal{J}_{1,1}^1 < \infty$. □

Lastly we turn to T_5 and S_5 . Actually we study $\Sigma_{1,1,1,0,I}^{1,0}$ and $\Sigma_{2,0,I}^{1,0}$ singularities.

Let $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ be an isotropic map-germ of corank ≤ 1 . Consider the ideal $\mathcal{J}_{1,1,1}^1$ in $\mathcal{O}_n$ generated by n -minors of the Jacobi matrix $\text{Jac}(f)$, n -minors $J_1^{(2)}, \dots, J_n^{(2)}$ (and $J(\pi \circ f)$) of $\text{Jac}(J(\pi \circ f), \pi \circ f)$, and n -minors of $\text{Jac}(J_1^{(2)}, \dots, J_n^{(2)}, J(\pi \circ f), \pi \circ f)$.

Moreover consider the ideal $\mathcal{J}_2^1$ in $\mathcal{O}_n$ generated by n -minors of the Jacobi matrix $\text{Jac}(f)$ and $(n-1)$ -minors of $\text{Jac}(\pi \circ f)$.

Then we have:

LEMMA 4.7.

(1) The ideal $\widehat{\mathcal{J}}_2^1$ in $\mathcal{O}_{J_I^1}$ induced naturally from $\mathcal{J}_2^1$ defines the Cohen-Macaulay varieties $\overline{\Sigma_{2,I}^1}$ of codimension 4 in J_I^1 . $\widehat{\mathcal{J}}_2^1$ is reduced on the submanifold $\Sigma_{2,I}^1$ of codimension 4 in J_I^1 . Moreover $\overline{\Sigma_{2,I}^1} \setminus \Sigma_{2,I}^1$ has codimension > 4 in J_I^1 .

(2) The ideal $\widehat{\mathcal{J}}_{1,1,1}^1$ in $\mathcal{O}_{J_I^3}$ induced naturally from $\mathcal{J}_{1,1,1}^1$ defines the variety

$$\overline{\Sigma_{1,1,1,I}^1 \cup \Sigma_{2,I}^1}$$

of codimension 4 in J_I^3 . $\widehat{\mathcal{J}}_{1,1,1}^1$ is reduced on the submanifold $\Sigma_{1,1,1,I}^{1,0}$ of codimension 4 in J_I^3 , but of multiplicity 4 on the submanifold $\Sigma_{2,0,I}^{1,0}$. Moreover

$$\overline{\Sigma_{1,1,1,I}^1 \cup \Sigma_{2,I}^1} \setminus (\Sigma_{1,1,1,I}^{1,0} \cup \Sigma_{2,0,I}^{1,0})$$

has codimension > 5 in J_I^3 .

Proof: (1) Set $X := \overline{\Sigma_{2,I}^1} \subset J_I^1$, and $Y := \Sigma_I^1 \subset J_I^1$. Then X has codimension 4 in J_I^1 ([15]). Moreover $X \subset Y \subset J_I^1$ and Y is a regular submanifold of codimension 2 in J_I^1 (§3). Now let $z \in Y$. Then the $2n$ -column vectors of the $n \times 2n$ -matrix z generates a $(n-1)$ -dimensional vector space $V = V(z) \subset \mathbf{C}^n$ since $z \in \Sigma^1$. Take a local frame $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$ near z of V . Let $\mathbf{q}_1, \dots, \mathbf{q}_n$ be the column vectors of $\text{Jac}(\pi \circ z)$. Set

$$(\mathbf{q}_1, \dots, \mathbf{q}_n) = (\mathbf{e}_1, \dots, \mathbf{e}_{n-1})A(z).$$

Then $A(z)$ is a matrix of size $(n-1) \times n$ with entries in $\mathcal{O}_Y$. Then $z \in \overline{\pi^{-1}\Sigma_2}$ if and only if $\text{rank} A(z) \leq n-2$. We see in $\mathcal{O}_Y$, $\widehat{\mathcal{J}}_2^1$ agrees with the ideal generated by $(n-1)$ -minors of $A(z)$. Moreover X is of codimension 2 in Y . Thus X is a determinantal variety in Y . Therefore X is Cohen-Macaulay. For T_5 singularity, by a simple calculation, we have $\dim_{\mathbf{C}} \mathcal{O}_4/\mathcal{J}_2^1 = 1$.

(2) First note that the zero locus $Z = Z(\widehat{\mathcal{J}}_{1,1,1}^1)$ is included in Σ_I^1 . Since $\widehat{\mathcal{J}}_{1,1,1}^1$ defines a hypersurface in Σ_I^1 , we see there exists an element $J^{(2)} \in \mathcal{O}_{J_I^3}$ such that $\widehat{\mathcal{J}}_{1,1,1}^1 = \widehat{\mathcal{J}}^1 + \langle J^{(2)} \rangle_{\mathcal{O}_{J_I^3}}$. Let $z \in \Sigma_I^1$. Then n -column vectors of $\text{Jac}(f)$ generates an $(n-1)$ -dimensional space $V = V(z)$. We take a local frame $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}$ of $V(z)$ over $\Sigma_I^1 \subset J_I^3$. We add $\mathbf{e}_n$ to get a local frame $\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \mathbf{e}_n$ of $\mathbf{C}^n$. We set

$$\text{Jac}(\pi \circ f, J(\pi \circ f), J^{(2)}) = (\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \mathbf{e}_n)C(z),$$

for an $n \times (n+2)$ -times matrix $C(z)$. Then $\widehat{\mathcal{J}}_{1,1,1}^1$ is generated by n -minors of $C(z)$ over Σ_I^1 . Thus we see

$$Z(\widehat{\mathcal{J}}_{1,1,1}^1) = \overline{\Sigma_{1,1,1,I}^1 \cup \Sigma_{2,I}^1}.$$

and it is of codimension 2 in Σ_I^1 .

By a simple calculation, when $n = 4$, we have $\dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_{1,1,1}^1 = 1$, for a S_5 singularity, and $\dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_{1,1,1}^1 = 4$, for a T_5 singularity. Here we present the calculation for T_5 . For an isotropic map-germ $f : (\mathbf{C}^n, 0) \rightarrow T^*\mathbf{C}^n$ ($n \geq 4$) of type T_5 , up to Lagrange equivalence, we have

$$\pi \circ f(t, x_2, x_3, \dots, x_{n-1}) = \left(\frac{1}{2}t^2 - x_2x_3, -\frac{1}{3}t^3 - 3x_2^2 - x_2x_3t - 2x_2x_4, x_3, \dots, x_{n-1} \right),$$

and moreover, up to a symplectomorphism of $(T^*\mathbf{C}^n, f(0))$, we have

$$f(t, x_2, x_3, \dots, x_{n-1}) = (x_2t, \frac{1}{3}t^3, 0, \dots, 0, \frac{1}{2}t^2, x_2, x_3, \dots, x_{n-1}).$$

Note that a diffeomorphism of $(T^*\mathbf{C}^n, f(0))$ preserves the ideal $\mathcal{J}^1$ generated by n -minor of $\text{Jac}(f)$. Then we have

$$\mathcal{J}_{1,1,1}^1 = \langle t, x_2, x_3x_4, x_4^2, x_3^2 \rangle_{\mathcal{O}_n},$$

which defines 4-fold T_5 -locus $\{t = 0, x_2 = 0, x_3 = 0, x_4 = 0\}$. Thus we have the required results. $\square$

By Lemma 4.7, counting the intersection number of $j^3 f(\mathbf{C}^4)$ with the zero locus $Z(\widehat{\mathcal{J}}_2^1)$ and $Z(\widehat{\mathcal{J}}_{1,1,1}^1)$ in $J_I^3(\mathbf{C}^4, T^*\mathbf{C}^4)$, we have:

THEOREM 4.8. *Let $f : (\mathbf{C}^4, 0) \rightarrow T^*\mathbf{C}^4$ be an isotropic map-germ of corank ≤ 1 . Then the number $\#T_5$ of T_5 -singularities of a Lagrangian stable perturbation of f is given by the following formula:*

$$\#T_5 = \dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_2^1,$$

provided $\dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_2^1 < \infty$.

PROPOSITION 4.9. *Let $f : (\mathbf{C}^4, 0) \rightarrow T^*\mathbf{C}^4$ be an isotropic map-germ of corank ≤ 1 . Then the number $\#S_5$ of S_5 -singularities is estimated by*

$$\#S_5 \leq \dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_{1,1,1}^1 - 4 \dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_2^1,$$

provided $\dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_2^1 < \infty$ and $\dim_{\mathbf{C}} \mathcal{O}_4 / \mathcal{J}_{1,1,1}^1 < \infty$.

5. Open questions. We collect here several practical questions related to the results in this paper.

(Q1) Recall that the formulae for the numbers of A_4 and of D_4 for Lagrangian immersions are given in §2. Thus we ask: *Do the same formulae hold on the numbers of A_4 and D_4 , not only for a Lagrange immersion, but also for an isotropic map-germ of corank one $(\mathbf{C}^3, 0) \rightarrow T^*\mathbf{C}^3$?*

(Q2) We have observed a similarity between the pair (D_4, A_4) and the pair (T_5, S_5) in §4. *Does the similarity have any reason?*

(Q3) *What are the formulae for the number of A_5 and D_5 of a Lagrangian immersion and of an isotropic map-germ of corank one?* See Remark 2.6.

(Q4) We can formulate the following conjecture:

Let $f : (\mathbf{C}^n, 0) \rightarrow T\mathbf{C}^n$ be an isotropic map-germ of corank at most one. Then the number $\#\Sigma_{i_1, i_2, \dots, I}^{1_k}$ of isolated singularities, with an isotropic Thom-Boardman symbol

$\Sigma_{i_1, i_2, \dots, l}^{1_k}$, appearing in a Lagrange stable perturbation of f is equal to the dimension of the quotient $\mathcal{O}_n / \mathcal{J}_{i_1, i_2, \dots}^{1_k}$ by the Morin ideal $\mathcal{J}_{i_1, i_2, \dots}^{1_k}$ for the corresponding (non-isotropic) Thom-Boardman singularity $\Sigma_{i_1, i_2, \dots}^{1_k}$ of composed mappings.

Is it true the conjecture in general? We have shown, in this paper, it is true for the cases A_2, A_3, S_3, S_4, T_5 and for W_5 , in the nice range $n \leq 4$, except for the cases A_4, D_4, A_5 and S_5 . Remark that D_5 -singularity is not described as a Thom-Boardman singularity (Remark 2.6). See §§2, 3 and §4 for the details.

References

- [1] V.I. Arnol'd, S.M. Gusein-Zade, A.N. Varchenko, *Singularities of Differentiable Maps I*, Birkhäuser, 1985.
- [2] I. Bogaevskii, G. Ishikawa, *Lagrange mappings of the first open Whitney umbrella*, Pacific Journal of Math., **203**–1 (2002), 115–138.
- [3] T. Fukuda, G. Ishikawa, *On the number of cusps of stable perturbations of a plane-to-plane singularity*, Tokyo Journal of Math., **10**–2 (1987), 375–384.
- [4] T. Fukui, J.J. Nuño Ballesteros, M.J. Saia, *On the number of singularities in generic deformations of map germs*, J. London Math. Soc., **58** (1998), 141–152.
- [5] T. Fukui and J. Weyman, *Cohen-Macaulay properties of Thom-Boardman strata I: Morin's ideal*, Proc of London Math. Soc., **80**–2 (2000), 257–303.
- [6] T. Fukui and J. Weyman, *Cohen-Macaulay properties of Thom-Boardman strata II: The defining ideals of $\Sigma^{i,j}$* , Preprint.
- [7] W. Fulton, *Intersection Theory*, Ergebnisse der Mathematik, **3**–2, Springer-Verlag, 1984.
- [8] T. Gaffney, D.M.Q. Mond, *Cusps and double folds of germs of analytic maps $\mathbf{C}^2 \rightarrow \mathbf{C}^2$* , J. London Math. Soc., **43** (1991), 185–192.
- [9] A.B. Givental', *Lagrangian imbeddings of surfaces and open Whitney umbrella*, Funkt. Anal. Prilozhen, **20**–3 (1986), 35–41. English transl., Funct. Anal. Appl., **20** (1986), 197–203.
- [10] A.B. Givental', *Singular Lagrangian manifolds and their Lagrangian mappings*, Itogi Nauki Tekh., Ser. Sovrem. Prob. Mat., (Contemporary Problems of Mathematics) **33**, VINITI, 1988, pp. 55–112. English transl., J. Soviet Math. **52** (1990), no. 4, 3246–3278.
- [11] G.M. Greuel, *Die Zahl der Spitzen und die Jacobi-Algebra einer isolierten Hyperflähensingularität*, Manuscripta Math., **21**–3 (1977), 227–241.
- [12] G.M. Greuel, *Ergänzung und Berichtigung zu: "Die Zahl der Spitzen und die Jacobi-Algebra einer isolierten Hyperflähensingularität"*, Manuscripta Math., **25**–2 (1978), 205–208.
- [13] M. Hochster, J.A. Eagon, *Cohen-Macaulay rings, invariant theory, and the generic perfection of determinantal loci*, Amer. J. Math., **93** (1971), 1020–1058.
- [14] G. Ishikawa, *The local model of an isotropic map-germ arising from one-dimensional symplectic reduction*, Math. Proc. Camb. Phil. Soc., **111** (1992), 103–112.
- [15] G. Ishikawa, *Transversalities for Lagrange singularities of isotropic mappings of corank one*, in Singularities and Differential Equations, Banach Center Publications, **33** (1996), 93–104.
- [16] G. Ishikawa, *Symplectic and Lagrange stabilities of open Whitney umbrellas*, Invent. math., **126**–2 (1996), 215–234.
- [17] G. Ishikawa, *Determinacy, transversality and Lagrange stability*, in Geometry and Topology of Caustics – Caustics' 98, Banach Center Publications, **50** (1999), 123–135.

- [18] B. Iversen, Lê Dung Tráng, *Calcul du nombre de cusps dans la déformation semi-universelle d'une singularité isolée d'hypersurface complexe*, Bull. Soc. Math. France, **102** (1974), 99–107.
- [19] W.L. Marar, D. Mond, *Multiple point schemes for corank 1 maps*, J. London Math. Soc., **39-3** (1989), 553–567.
- [20] W.L. Marar, J. A. Montaldi, M. A. S.Ruas, *Multiplicities of zero-schemes in quasihomogeneous corank-1 singularities $\mathbf{C}^n \rightarrow \mathbf{C}^n$* , Singularity theory (Liverpool, 1996), 353–367, London Math. Soc. Lecture Note Ser., **263**, Cambridge Univ. Press, Cambridge, 1999.
- [21] H. Matsumura, *Commutative Algebra*, Mathematical Lecture Note Series, **56**, Benjamin, 1980.
- [22] D. Mond, *Some remarks on the geometry and classification of germs of maps from surfaces to 3-space*, Topology **26-3** (1987), 361–383.
- [23] D. Mond, *Vanishing cycles for analytic maps*, in Singularity Theory and its Applications, I: Geometric Aspects of Singularities, Lecture Notes in Math., **1462**, Springer-Verlag, (1991).
- [24] D. Mond, *The number of vanishing cycles for a quasihomogeneous mapping from $\mathbf{C}^2$ to $\mathbf{C}^3$* , Quart. J. Math. Oxford Ser. **42-167** (1991), 335–345.
- [25] B. Morin, *Calcul jacobien*, Ann. Sci. École Norm. Sup., **8** (1975), 1–98.
- [26] J.J. Nuño Ballesteros, M.J. Saia, *Multiplicity of Boardman strata and deformations of map germs*, Glasgow Math. J., **40** (1998), 21–32.
- [27] F. Pham, *Courbes discriminantes des singularités planes d'ordre 3*, Singularités à Cargèse, Rencontre Singularités Géom. Anal., Inst. Études Sci., Cargèse, 1972, pp. 363–391, Asterisque, **7/8**, Soc. Math. France, Paris, 1973.
- [28] J.H. Rieger, *Families of maps from the plane to the plane*, J. London Math. Soc., **36-2** (1987), 351–369.
- [29] J.H. Rieger, *Versal topological stratification and the bifurcation geometry of map-germs of the plane*, Math. Proc. Cambridge Philos. Soc., **107-1** (1990), 127–147.
- [30] V. M. Zakalyukin, *Reconstructions of fronts and caustics depending on a parameter, and versality of mappings*, Current problems in mathematics, Vol. **22**, pp. 56–93, Itogi Nauki i Tekhniki, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1983. English transl., Journal of Soviet Math., **27-3** (1984), 2713–2735.
- [31] V. M. Zakalyukin, *Generating ideals of Lagrangian varietes*, Theory of Singularities and its Applications, ed. by V.I. Arnold, Advances in Soviet Math., vol.1, Amer. Math. Soc., (1990), pp. 201–210.