



HOKKAIDO UNIVERSITY

Title	Spectra of Toeplitz Operators and Uniform Algebras
Author(s)	Nakazi, T.; Osawa, T.
Citation	Hokkaido University Preprint Series in Mathematics, 581, 1-9
Issue Date	2003-01
DOI	https://doi.org/10.14943/83726
Doc URL	https://hdl.handle.net/2115/69330
Type	departmental bulletin paper
File Information	re581.pdf



**Spectra Of Toeplitz Operators
And Uniform Algebras**

Takahiko Nakazi and Tomoko Osawa

Series #581. January 2003

HOKKAIDO UNIVERSITY
PREPRINT SERIES IN MATHEMATICS

- #555 S. Izumiya and N. Takeuchi, New special curves and developable surfaces, 10 pages. 2002.
- #556 S. Izumiya, K. Saji and N. Takeuchi, Singularities of line congruences, 17 pages. 2002.
- #557 S. Izumiya, D. Pei and M. C. Romero Fuster, The lightcone Gauss map of a spacelike surface in Minkowski 4-space, 21 pages. 2002.
- #558 S. Izumiya, D. Pei and M. C. Romero Fuster, Umbilicity of spacelike submanifolds of Minkowski space, 14 pages. 2002.
- #559 S. Izumiya, D. Pei and M. Takahashi, Curves and surfaces in Hyperbolic space, 16 pages. 2002.
- #560 J. Kato, On the uniqueness of nondecaying solutions for the Navier-Stokes equations, 19 pages. 2002.
- #561 M. Jinzenji and T. Sasaki, An Approach to $\mathcal{N} = 4$ *ADE* gauge Theory on $K3$, 29 pages. 2002.
- #562 T. Nakazi and T. Yamamoto, Norms of some singular integral operators on weighted L^2 spaces, 27 pages. 2002.
- #563 A. Harris and Y. Tonegawa, A $\bar{\partial}\partial$ -poincaré lemma for forms near an isolated complex singularity, 8 pages. 2002.
- #564 M. Takahashi, Bifurcations of ordinary differential equations of Clairaut type, 23 pages. 2002.
- #565 G. Ishikawa, Classifying singular Legendre curves by contactomorphisms, 17 pages. 2002.
- #566 G. Ishikawa, Perturbations of Caustics and fronts, 17 pages. 2002.
- #567 Y. Giga and O. Sawada, On regularizing-decay rate estimates for solutions to the Navier-Stokes initial value problem, 12 pages. 2002.
- #568 T. Miyao, Strongly supercommuting self-adjoint operators, 34 pages. 2002.
- #569 Jun-Muk Hwang and K. Yamaguchi, Characterization of Hermitian symmetric spaces by fundamental forms, 10 pages. 2002.
- #570 H. Ishii and T. Mikami, Convexified Gauss curvature flow of bounded open sets in an anisotropic external field: a stochastic approximation and PDE, 37 pages. 2002.
- #571 Y. Nakano, Minimization of shortfall risk in a jump-diffusion model, 10 pages. 2002.
- #572 K. Izuchi and T. Nakazi, Backward Shift Invariant Subspaces in the Bidisc, 8 pages. 2002.
- #573 S. Izumiya, D. Pei and M. C. Romero-Fuster, The horospherical geometry of surfaces in Hyperbolic 4-space, 17 pages. 2002.
- #574 S. Izumiya and M. C. Romero-Fuster, The hyperbolic Gauss-Bonnet type theorem, 10 pages. 2002.
- #575 S. Izumiya and S. Janeczko, A symplectic framework for multiplane gravitational lensing, 19 pages. 2002.
- #576 S. Izumiya, M. Kossowski, D. Pei and M. C. Romero-Fuster, Singularities of C^∞ -lightlike hypersurfaces in Minkowski 4-space, 18 pages. 2002.
- #577 S. Izumiya, D. Pei and M. Takahashi, Evolutes of hypersurfaces in Hyperbolic space, 21 pages. 2002.
- #578 Y. Giga, S. Matsui and S. Sasayama, Blow up rate for semilinear heat equation with subcritical nonlinearity, 29 pages. 2002.
- #579 M. Tsujii, Physical measures for partially hyperbolic surface endomorphisms, 71 pages. 2003.
- #580 Y. Giga and K. Yamada, On viscous Burgers-like equations with linearly growing initial data, 19 pages. 2003.

Spectra Of Toeplitz Operators And Uniform Algebras

by

Takahiko Nakazi*

and

Tomoko Osawa*

*This research was partially supported by Grant-in-Aid for Scientific Research,
Ministry of Education

Mathematics Subject Classification 2000 : 46 J 10, 46 J 15, 47 A 10, 47 B 35

Key words and phrases : Toeplitz operator, uniform algebra, spectrum, convex-hull, index

Abstract. Let A be a uniform algebra on X and $\mathcal{P}$ a set of all probability measures on X . For each μ in $\mathcal{P}$, $H^2(\mu)$ is the closure of A in $L^2(\mu)$ and T_ϕ^μ is a Toeplitz operator on $H^2(\mu)$ for a continuous function ϕ on X . In this paper we study the invertibility and the spectrum of $\mathbf{T}_\phi = \sum_{\mu \in \mathcal{P}} \oplus T_\phi^\mu$. We show that if $\mathbf{T}_\phi$ is invertible then the index of ϕ is zero and if the converse is true for an arbitrary continuous function ϕ then A is a Dirichlet algebra on X . Moreover we study the spectrum of $\mathbf{T}_\phi$.

§1. Introduction

Let X be a compact Hausdorff space, $C = C(X)$ an algebra of all continuous complex-valued functions on X , and A a uniform algebra on X . $\mathcal{P}$ denotes the set of all positive Borel measures on X with total mass 1. For each μ , $H^2(\mu)$ is defined as the closure of A in $L^2(\mu)$. Let P^μ be the orthogonal projection from $L^2(\mu)$ onto $H^2(\mu)$. For ϕ in C and f in $H^2(\mu)$, put

$$T_\phi^\mu f = P^\mu(\phi f).$$

In this paper, we study Toeplitz operators $\mathbf{T}_\phi$ on the Hardy space $\mathbf{H}^2$ for ϕ in C , where

$$\mathbf{T}_\phi = \sum_{\mu \in \mathcal{P}} \oplus T_\phi^\mu \quad \text{and} \quad \mathbf{H}^2 = \sum_{\mu \in \mathcal{P}} \oplus H^2(\mu).$$

$\sigma(\mathbf{T}_\phi)$ denotes the spectrum of $\mathbf{T}_\phi$ and then $\sigma(\mathbf{T}_\phi) \supseteq \bigcup_{\mu \in \mathcal{P}} \sigma(T_\phi^\mu)$. For ϕ, ψ in C $T_\phi^\mu T_\psi^\mu = P^\mu \mathbf{T}_\phi \mathbf{T}_\psi |H^2(\mu)$ and so $\mathbf{T}_\phi$ is a power dilation of T_ϕ^μ . $\|\mathbf{T}_\phi\| = \sup_{x \in X} |\phi(x)|$ and $\|T_\phi^\mu\| = \mu - \text{esssup}_{x \in X} |\phi(x)|$. T_ϕ^μ is the local part of $\mathbf{T}_\phi$ and $\mathbf{T}_\phi$ is more strongly related with A than T_ϕ^μ . The local part of $\mathbf{T}_\phi$ has been studied in [7] for arbitrary uniform algebra.

When X is the unit circle ∂D , A is the disc algebra $\mathcal{A}$ and μ is the normalized Lebesgue measure $d\theta/2\pi$, $H^2 = H^2(\mu)$ is the classical Hardy space and $T_\phi = T_\phi^\mu$ is the usual Toeplitz operator on H^2 . When μ is a finite positive Borel measure on ∂D , $H^2(\mu)$ is called a weighted Hardy space. In this classical case, our result shows that $\sigma(\mathbf{T}_\phi) = \sigma(T_\phi)$ for arbitrary ϕ in C . When X is the closed unit disc $\bar{D}$, A is the disc algebra $\mathcal{A}$ and μ is the normalized area measure $\text{rdrd}\theta/\pi$, $L_a^2 = H^2(\mu)$ is the Bergman space and $T'_\phi = T_\phi^\mu$ is the usual Toeplitz operator on L_a^2 . In this case, $\sigma(\mathbf{T}_\phi) \neq \sigma(T'_\phi)$. By definition, $\mathbf{T}_\phi$ is strongly related with the uniform algebra, that is, $\mathcal{A}|\partial D$ or $\mathcal{A}|\bar{D}$ but T_ϕ or T'_ϕ is related with $d\theta/2\pi$ or $\text{rdrd}\theta/\pi$. Suppose the classical Hardy space H^∞ is the weak $*$ closure of $\mathcal{A}$ in $L^\infty = L^\infty(\mu)$ with $\mu = d\theta/2\pi$. When X is the maximal ideal space of L^∞ , A is the Gelfand transform of H^∞ on X and μ is a representing measure on X for the origin, $H^2(\mu)$ is also the classical Hardy space H^2 and T_ϕ^μ is the usual Toeplitz operator T_ϕ . Our result implies that $\sigma(\mathbf{T}_\phi) = \sigma(T_\phi)$ for arbitrary ϕ in $C =$ the Gelfand transform of L^∞ .

Let A be an arbitrary uniform algebra on X and $M(A)$ the maximal ideal space of A . For a function ϕ in C , we say that the index of ϕ is zero if there exists a nonvanishing function g in $C(M(A))$ such that ϕg has a continuous logarithm on X . By the Arens-Royden theorem [3, p89], we can choose g as an element in A^{-1} . When A is a natural uniform algebra on the complex plane, our definition of the index of ϕ is same to the classical case (see [4, p281]).

In Section 2, we show that $\mathbf{T}_\phi$ is invertible if and only if there exist a positive constant δ and a function g in A^{-1} such that

$$\text{Re} \phi g \geq \delta > 0 \quad \text{on } X.$$

This result is similar as that of the classical Toeplitz operator by Widom and Devinatz (see [2]). That is, T_ϕ^μ is invertible if and only if there exist a positive constant δ and a

function g in $H^\infty(\mu)^{-1}$ such that $\operatorname{Re}\phi g \geq \delta > 0$ a.e. on ∂D , when ϕ is in $L^\infty(\mu)$ and $\mu = d\theta/2\pi$. When C is a commutative C^* -subalgebra of $L^\infty(\mu)$, our theorem implies that if ϕ is a function in C and $\mathbf{T}_\phi$ is invertible then there exist a positive constant δ and a function g in $H^\infty(\mu)^{-1} \cap C$ such that $\operatorname{Re}\phi g \geq \delta > 0$ on ∂D . In the general setting, our result shows that if $\mathbf{T}_\phi$ is invertible then the index of ϕ is zero. When A is a Dirichlet algebra on X , if the index of ϕ is zero then $\mathbf{T}_\phi$ is invertible. We show that the converse is also true. If $\mathbf{T}_\phi$ is always invertible for an arbitrary function ϕ in C with index $\phi = 0$, then A is a Dirichlet algebra on X . In Section 3, we show that $\phi(X) \subseteq \{\lambda \in \mathcal{C} ; \text{index } (\phi - \lambda) \neq 0\} \subseteq \sigma(\mathbf{T}_\phi) \subseteq \text{the convex hull of } \phi(X)$. Moreover when ϕ is in A or ϕ is real-valued in C , $\sigma(\mathbf{T}_\phi)$ is completely described.

In this paper, $A^\perp$ denotes the set of all annihilating measures on X and $M(A)$ denotes the maximal ideal space of A . $M(A)$ is called simply connected when the first Čech cohomology group of $M(A)$ with integer coefficients is zero. $\mathcal{R}(\phi)$ is the range $\phi(X)$ of ϕ in X . $\langle f, g \rangle_\mu = \int_X f \bar{g} d\mu$ is the inner product in $L^2(\mu)$ and $\|f\|_\mu = (\langle f, f \rangle_\mu)^{1/2}$. $\|F\|$ is the norm of F in $\mathbf{H}^2$. $\|f\|_X = \sup_{x \in X} |f(x)|$ and $\|f + A\|_X = \inf_{g \in A} \|f + g\|_X$. $\|\mathbf{T}_\phi\|$ denotes the norm of the operator $\mathbf{T}_\phi$ on $\mathbf{H}^2$ and $|T_\phi^\mu|$ denotes the norm of the operator T_ϕ^μ on $H^2(\mu)$.

§2. Invertibility of $\mathbf{T}_\phi$

For each μ in $\mathcal{P}$, let I^μ be the identity operator on $L^2(\mu)$. For ϕ in C and f in $H^2(\mu)$, put $H_\phi^\mu f = (I^\mu - P^\mu)(\phi f)$ and $\mathbf{H}_\phi = \sum_{\mu \in \mathcal{P}} \oplus H_\phi^\mu$. Lemma 1 is similar to a theorem of Nehari [8]. Lemma 2 is similar to a result of Nakazi [6] which was proved by Widom and Devinatz [2] when ϕ is unimodular. Theorem 1 is an analogue of a theorem of Widom and Devinatz [2] in the classical case.

Lemma 1. *If ϕ is a function in C , then $\|\mathbf{H}_\phi\| = \|\phi + A\|_X$.*

Proof. It is clear that $\|\mathbf{H}_\phi\| \leq \|\phi + A\|_X$. Fix $\phi \in C$ with $\phi \notin A$. By the Hahn-Banach theorem and the Riesz representation theorem, there exists a finite Borel measure $\nu \in A^\perp$ with $\|\nu\| = 1$ such that

$$\|\phi + A\|_X = \int \phi d\nu.$$

Put $F = d|\nu|/d\nu$ and $\mu = |\nu|$, then $F \in L^2(\mu) \cap A^\perp$. Hence

$$\|\phi + A\|_X = \int \phi \cdot 1 \cdot \bar{F} d\mu = \langle H_\phi^\mu 1, F \rangle_\mu \leq |H_\phi^\mu| \leq \|\mathbf{H}_\phi\|.$$

Lemma 2. *Suppose ϕ is a function in C . $\mathbf{T}_\phi$ is left invertible if and only if there exists a positive constant ε and a function g in A such that*

$$|\phi + g|^2 \leq |\phi|^2 - \varepsilon \text{ on } X.$$

Proof. The ‘if’ part is easy. In fact,

$$\mathbf{H}_\phi^* \mathbf{H}_\phi + \mathbf{T}_\phi^* \mathbf{T}_\phi = \mathbf{T}_{|\phi|^2}$$

because $(H_\phi^\mu)^* H_\phi^\mu + (T_\phi^\mu)^* T_\phi^\mu = T_{|\phi|^2}^\mu$ for all $\mu \in \mathcal{P}$. Hence

$$\mathbf{H}_\phi^* \mathbf{H}_\phi = \mathbf{H}_{\phi+g}^* \mathbf{H}_{\phi+g} \leq \mathbf{T}_{|\phi+g|^2} \leq \mathbf{T}_{|\phi|^2-\varepsilon}$$

and so

$$\mathbf{T}_\phi^* \mathbf{T}_\phi = \mathbf{T}_{|\phi|^2} - \mathbf{H}_\phi^* \mathbf{H}_\phi \geq \mathbf{T}_\varepsilon.$$

Now we will show the ‘only if’ part. If $\mathbf{T}_\phi$ is left invertible, then there exists $\varepsilon > 0$ such that $\mathbf{T}_\phi^* \mathbf{T}_\phi \geq \mathbf{T}_{\sqrt{2\varepsilon}}$. Hence $(T_\phi^\mu)^* T_\phi^\mu \geq T_{\sqrt{2\varepsilon}}^\mu$ for all $\mu \in \mathcal{P}$ and so

$$\int |\phi f|^2 d\mu \geq 2\varepsilon \int |f|^2 d\mu \quad (f \in A)$$

for all $\mu \in \mathcal{P}$. Therefore $|\phi|^2 \geq 2\varepsilon > 0$. Hence $\mathbf{H}_\phi^* \mathbf{H}_\phi \leq \mathbf{T}_{|\phi|^2-2\varepsilon} \leq \mathbf{T}_{|\phi|^2-\varepsilon}$ and so $\|\mathbf{H}_\phi F\|^2 \leq \|(\mathbf{T}_{|\phi|^2-\varepsilon})^{1/2} F\|^2$ for all $F \in \mathbf{H}^2$. Therefore for any $\mu \in \mathcal{P}$, $f \in A$ and $g \in L^2(d\mu) \cap A^\perp$,

$$\begin{aligned} \left| \int \phi f \bar{g} d\mu \right|^2 &= |\langle H_\phi^\mu f, g \rangle_\mu|^2 \leq \|H_\phi^\mu f\|_\mu^2 \|g\|_\mu^2 \\ &\leq \langle T_{|\phi|^2-\varepsilon}^\mu f, f \rangle \|g\|_\mu^2 \leq \int (|\phi|^2 - \varepsilon) |f|^2 d\mu \int |g|^2 d\mu \end{aligned}$$

because $\|H_\phi^\mu f\|_\mu^2 \leq \|(T_{|\phi|^2-\varepsilon}^\mu)^{1/2} f\|_\mu^2$ for all $f \in A$. If $v^2 d\mu \in \mathcal{P}$ and v is an invertible function in C , then for any $f \in A$ and $G \in L^2(v^2 d\mu) \cap A^\perp$

$$\left| \int \phi f \bar{G} v^2 d\mu \right|^2 \leq \int (|\phi|^2 - \varepsilon) |f|^2 v^2 d\mu \int |G|^2 v^2 d\mu.$$

It is easy to see that $v^2(L^2(v^2 d\mu) \cap A^\perp) = L^2(d\mu) \cap A^\perp$. Hence for any $f \in A$ and $g \in L^2(d\mu) \cap A^\perp$

$$\left| \int \phi f \bar{g} d\mu \right|^2 \leq \int (|\phi|^2 - \varepsilon) |f|^2 v^2 d\mu \int |g|^2 v^{-2} d\mu$$

because $G = g/v^2$ belongs to $L^2(v^2 d\mu) \cap A^\perp$. Put $a^{-1} = \int (|\phi|^2 - \varepsilon)^{-1/2} d\mu$ and $v^2 = a(|\phi|^2 - \varepsilon)^{-1/2}$, then $\int v^2 d\mu = 1$ and so $v^2 d\mu \in \mathcal{P}$ by the definition of $\mathcal{P}$ in Introduction. Then v^2 is an invertible function in C because $|\phi|^2 \geq 2\varepsilon > 0$. Hence for any $f \in A$ and $g \in L^2(d\mu) \cap A^\perp$

$$\left| \int \phi f \bar{g} d\mu \right|^2 \leq \int (|\phi|^2 - \varepsilon)^{1/2} a |f|^2 d\mu \int (|\phi|^2 - \varepsilon)^{1/2} a^{-1} |g|^2 d\mu. \quad (\star)$$

Put $u = (|\phi|^2 - \varepsilon)^{1/2}$, then $(A \cdot u^{-1})^\perp \cap (C \cdot u^{-1})^* = \{u d\lambda ; \lambda \in A^\perp\}$ where $*$ denotes the dual. By the Hahn-Banach theorem,

$$\inf \{ \|(\phi + h)u^{-1}\|_X ; h \in A \} = \sup \left\{ \left| \int \phi d\lambda \right| ; \lambda \in A^\perp \text{ and } \int u d|\lambda| \leq 1 \right\}.$$

If $\phi \notin A$ then there exists a nonzero $\nu \in A^\perp$ with $\int u d|\nu| = 1$ such that

$$\inf \|(\phi + h)u^{-1}\|_X = \int \phi d\nu.$$

Put $F = d|\nu|/d\nu$, then $F \in L^2(d\nu) \cap A^\perp$ and $1 \in A$, hence by $(\star)$

$$\int \phi d\nu = \int \phi \cdot 1 \cdot \bar{F} d|\nu| \leq \int (|\phi|^2 - \varepsilon)^{1/2} d|\nu| = \int u d|\nu| = 1.$$

Thus $\inf \|(\phi + h)u^{-1}\|_X \leq 1$. If $\varepsilon_1 < \varepsilon$, then for some $\delta > 0$ $|\phi|^2 - \varepsilon_1 \geq (1 + \delta)(|\phi|^2 - \varepsilon)$ and hence there exists a function g in A

$$|\phi|^2 - \varepsilon_1 \geq (1 + \delta)(|\phi|^2 - \varepsilon) \geq |\phi + g|^2.$$

Lemma 3. *Suppose ϕ is a function in C . $\mathbf{T}_\phi$ is left invertible if and only if there exists a positive constant δ and a function g in A such that*

$$\operatorname{Re} \bar{\phi} g \geq \delta > 0 \quad \text{on } X.$$

Proof. If $\mathbf{T}_\phi$ is left invertible, then by Lemma 2 there exists g in A such that $|\phi|^2 \geq \varepsilon^2 + |\phi - g|^2$ for some constant $\varepsilon > 0$. Then, $|\phi|^2 \geq \varepsilon^2$, and so $1 \geq \varepsilon^2/|\phi|^2 \geq \varepsilon_1 > 0$ for some constant ε_1 . Hence

$$1 - \varepsilon_1 \geq 1 - \frac{\varepsilon^2}{|\phi|^2} \geq \left| 1 - \frac{g}{\phi} \right|^2.$$

Therefore there exist constants $\varepsilon_2, \varepsilon_3$ such that $\operatorname{Re} \frac{g}{\phi} \geq \varepsilon_2 > 0$ and so $\operatorname{Re} \bar{\phi} g \geq \varepsilon_2 |\phi|^2 \geq \varepsilon_2 \varepsilon^2 > 0$. Conversely if $\operatorname{Re} \bar{\phi} g \geq \delta > 0$, then $0 < \varepsilon_3 \leq |\phi| \leq \gamma < \infty$ and so $\operatorname{Re} \frac{g}{\phi} \geq \frac{\delta}{\gamma} > 0$.

Hence there exist two positive constant ε_4 and ε_5 such that $\left| \varepsilon_4 \frac{g}{\phi} - 1 \right|^2 \leq 1 - \varepsilon_5$. Hence $|\varepsilon_4 g - \phi|^2 \leq (1 - \varepsilon_5) |\phi|^2$ and so

$$|\varepsilon_4 g - \phi|^2 + \varepsilon_5 \varepsilon^2 \leq |\varepsilon_4 g - \phi|^2 + \varepsilon_5 |\phi|^2 \leq |\phi|^2.$$

Lemma 4. *Suppose ϕ is a function in C . $\mathbf{T}_\phi$ is invertible if and only if $\mathbf{T}_{\frac{\phi}{|\phi|}}$ and $\mathbf{T}_{|\phi|}$ are invertible.*

Proof. If $\mathbf{T}_\phi$ is invertible, and $\mathbf{T}_\phi^* = \mathbf{T}_{\bar{\phi}}$ is also invertible and so by Lemma 3 there exists a constant δ_1 and a function g in A such that

$$\operatorname{Re} \phi g \geq \delta_1 > 0.$$

Then ϕ is invertible in C and so there exists a constant δ_2 such that

$$\operatorname{Re} \frac{\phi}{|\phi|} g \geq \delta_2 > 0.$$

Hence by Lemma 3 $\mathbf{T}_{\phi/|\phi|}$ and $\mathbf{T}_{|\phi|}$ are invertible. The converse can be proved similarly by Lemma 3.

Theorem 1. *Suppose ϕ is a function in C . Then, $\mathbf{T}_\phi$ is invertible if and only if there exist a constant δ and a function g in A^{-1} such that*

$$\operatorname{Re} \phi g \geq \delta > 0 \text{ on } X.$$

Proof. By Lemma 4, we may assume that ϕ is unimodular. If $\mathbf{T}_\phi$ is invertible, then by Lemma 2 there exists a function g in A such that $\|\phi + g\| < 1$ and so $\|1 + \bar{\phi}g\| < 1$. Hence $\mathbf{T}_{\bar{\phi}}\mathbf{T}_g$ is invertible and so $\mathbf{T}_g$ is invertible because $\mathbf{T}_\phi^* = \mathbf{T}_{\bar{\phi}}$ is invertible. Therefore $gH^2(\mu) = H^2(\mu)$ for any $\mu \in \mathcal{P}$ and so g^{-1} belongs to $H^2(\mu)$. Lemma 4 implies that $g^{-1} \in C$. We will show that g^{-1} belongs to A . Then the proof of Lemma 3 implies the theorem. If $g^{-1} \notin A$, there exists a finite measure λ in $A^\perp$ such that $\|\lambda\| = 1$ and $\int g^{-1} d\lambda \neq 0$. Let $F = d\lambda/d|\lambda|$, then $\bar{F}$ is orthogonal to $H^2(|\lambda|)$. Since $\int g^{-1} F d|\lambda| \neq 0$, $g^{-1} \notin H^2(|\lambda|)$. This contradiction implies that $g^{-1} \in A$. Conversely if there exist a constant δ and a function g in A^{-1} such that $\operatorname{Re} \phi g \geq \delta > 0$, then there exists a constant δ' such that $\operatorname{Re} \phi \frac{g}{|g|} \geq \delta' > 0$. Hence $\operatorname{Re} \bar{\phi} \frac{f}{|f|} \geq \delta' > 0$ where $f = g^{-1}$. By lemma 3, $\mathbf{T}_\phi$ is invertible.

Corollary 1. *Let ϕ be a function in C . If $\mathbf{T}_\phi$ is invertible, then there exists a function g in A^{-1} such that ϕg has a continuous logarithm on X , that is, the index of ϕ is zero.*

The converse of Corollary 1 is not true as Corollary 2 shows.

Corollary 2. *If $\mathbf{T}_\phi$ is always invertible for an arbitrary function ϕ in C with index $\phi = 0$, then A is a Dirichlet algebra on X .*

Proof. If $v \in C$ is real-valued and $\phi = e^{iv}$, then index $\phi = 0$. By hypothesis, $\mathbf{T}_\phi$ is invertible and so by Theorem 1 there exists a function g in A such that $\operatorname{Re} \bar{\phi} g > 0$ on X . Corollary 4.7 in [5] shows that A is a Dirichlet algebra.

§3. Spectrum of $\mathbf{T}_\phi$

Let B be a subset of C^{-1} . We define a generalization of the convex hull of the range $\mathcal{R}(\phi)$ of ϕ in C . That is, $\operatorname{Hull} \{\mathcal{R}(\phi), B\}$ is a set of all complex numbers λ which satisfy the following : There do not exist a constant δ and a function g in B such that

$\operatorname{Re}(\phi - \lambda)g \geq \delta > 0$ on X . $\operatorname{Hull} \{\mathcal{R}(\phi), B\}$ need not be determined by $\mathcal{R}(\phi)$ and B in general. $\operatorname{Hull} \{\mathcal{R}(\phi), B\}$ contains $\mathcal{R}(\phi)$. If $B = \{\lambda \in \mathcal{C} ; \lambda \neq 0\}$, then $\operatorname{Hull} \{\mathcal{R}(\phi), B\}$ is the convex hull of $\mathcal{R}(\phi)$. If $B = C^{-1}$, then $\operatorname{Hull} \{\mathcal{R}(\phi), B\} = \mathcal{R}(\phi)$. If $B = \exp C$, then $\operatorname{Hull} \{\mathcal{R}(\phi), B\} = \{\lambda \in \mathcal{C} ; \phi - \lambda \text{ does not have a continuous logarithm on } X\}$. In this section, we are interested in $\operatorname{Hull} \{\mathcal{R}(\phi), \exp A\}$ and $\operatorname{Hull} \{\mathcal{R}(\phi), A^{-1}\}$. If ϕ is in A then $\operatorname{Hull} \{\mathcal{R}(\phi), A^{-1}\} = \hat{\phi}(M(A))$. This is proved in the proof of (4) of Theorem 2. (2) and (4) of Theorem 2, and Corollaries 3 and 4 are similar to theorems in [1, Chapter 7]. For example, (2) of Theorem 2 is an analogue of a theorem of Hartman-Wintner (cf. [1, 7.20 Theorem]).

Theorem 2. *Let ϕ be a function in C .*

- (1) $\sigma(\mathbf{T}_\phi) = \operatorname{Hull} \{\mathcal{R}(\phi), A^{-1}\}$.
- (2) $\mathcal{R}(\phi) \subseteq \sigma(\mathbf{T}_\phi) \subseteq \text{the convex hull of } \mathcal{R}(\phi)$.
- (3) $\sigma(\mathbf{T}_\phi) \supseteq \{\lambda \in \mathcal{C} ; \operatorname{index}(\phi - \lambda) \neq 0\}$.
- (4) *If ϕ is in A , then $\sigma(\mathbf{T}_\phi) = \hat{\phi}(M(A))$.*
- (5) *If ϕ is real-valued, $a = \min \phi$ and $b = \max \phi$, then $\sigma(\mathbf{T}_\phi) \supseteq \mathcal{R}(\phi) \ni a, b$ and $\sigma(\mathbf{T}_\phi) \subseteq [a, b]$. Moreover $\lambda \in [a, b] \setminus \sigma(\mathbf{T}_\phi)$ if and only if χ_{E_λ} belongs to A where $E_\lambda = \{x \in X ; \phi(x) - \lambda > 0\}$.*

Proof. (1) is clear by Theorem 1. (2) is a result of (1). (3) is a result of (1) and the definition of the index. (4) If $\lambda \notin \sigma(\mathbf{T}_\phi)$, then by Theorem 1 there exist a positive constant δ and $g \in A^{-1}$ such that $\operatorname{Re}(\phi - \lambda)g \geq \delta > 0$. Hence there exists a function $f \in A$ such that $(\phi - \lambda)g = e^f$ and so $\phi - \lambda \in A^{-1}$. Therefore $\lambda \notin \hat{\phi}(M(A))$. Conversely if $\lambda \notin \hat{\phi}(M(A))$, then $\phi - \lambda \in A^{-1}$. Put $g = (\phi - \lambda)^{-1}$ and $\delta = 1$, then $\operatorname{Re}(\phi - \lambda)g = \delta > 0$. Hence Theorem 1 implies that $\lambda \notin \sigma(\mathbf{T}_\phi)$. (5) (1) implies that $\sigma(\mathbf{T}_\phi) \supseteq \mathcal{R}(\phi) \ni a, b$ and $\sigma(\mathbf{T}_\phi) \subseteq [a, b]$. If $\lambda \in [a, b] \setminus \sigma(\mathbf{T}_\phi)$, then by (1) there exist δ and $g \in A^{-1}$ such that $\operatorname{Re}(\phi - \lambda)g \geq \delta > 0$. Since $(\phi - \lambda)/|\phi - \lambda| = 2\chi_{E_\lambda} - 1$, $\operatorname{Re} g \geq \delta/|\phi - \lambda|$ on E_λ and $\operatorname{Re} g \leq -\delta/|\phi - \lambda|$ off E_λ . A theorem of Runge implies that χ_{E_λ} belongs to A . Conversely if $\chi_{E_\lambda} \in A$, then $g = 2\chi_{E_\lambda} - 1$ belongs to A^{-1} and so $\operatorname{Re}(\phi - \lambda)g \geq \delta > 0$. (1) implies that $\lambda \in [a, b] \setminus \sigma(\mathbf{T}_\phi)$.

Corollary 3. *Suppose χ_E is a characteristic function of a subset E in X , and a and b are real numbers with $a < b$. If $\phi = a\chi_E + b\chi_{E^c}$ is in C , then $\sigma(\mathbf{T}_\phi) = \{a, b\}$ when χ_E is in A and $\sigma(\mathbf{T}_\phi) = [a, b]$ when χ_E is not in A .*

Corollary 4. *Suppose A is antisymmetric. If ϕ is a real-valued function in C with $a = \min \phi$ and $b = \max \phi$, then $\sigma(\mathbf{T}_\phi) = [a, b]$.*

Corollary 5. *Suppose A is a Dirichlet algebra on X and ϕ is a function in C .*

- (1) $\sigma(\mathbf{T}_\phi) \subseteq \{\lambda \in \mathcal{C} ; \phi - \lambda \text{ does not have a continuous logarithm on } X\}$.
- (2) *If $M(A)$ is simply connected, then $\sigma(\mathbf{T}_\phi) = \{\lambda \in \mathcal{C} ; \phi - \lambda \text{ does not have the continuous logarithm on } X\}$.*

Proof. (1) If $\log(\phi - \lambda) \in C$, then there exist two real-valued functions u, v

such that $\phi - \lambda = e^{u+iv}$. Since A is a Dirichlet algebra, there exists $f \in A$ such that $\|v - \operatorname{Im} f\| < \pi/2$. Put $g = e^{-f}$ then $\operatorname{Re}(\phi - \lambda)g \geq \delta > 0$ for some constant δ and $g \in A^{-1}$. By Theorem 2, λ belongs to $\sigma(\mathbf{T}_\phi)^c$. (2) Since $M(A)$ is simply connected, by the Arens-Royden theorem $A^{-1} = \exp A$. Hence (1) of Theorem 2 imply (2).

References

1. R. G. Douglas, Banach Algebra Techniques in Operator Theory, Academic Press, New York and London, 1972.
2. A. Devinatz, Toeplitz operators on H^2 spaces, Trans.Amer.Math.Soc., 112(1964), 304-317.
3. T. Gamelin, Uniform Algebras, Prentice-Hall, Englewood Cliffs, New Jersey, 1969
4. T. Gamelin, J. Garnett, L. Rubel and A. Shields, On badly approximable functions, J. Approx.Th. 17(1976), 280-296.
5. D. H. Luecking, On badly approximable functions and uniform algebras, J. Approx.Th. 22(1978), 161-176.
6. T. Nakazi, Absolute values of Toeplitz operators and Hankel operators, Canad.Math.Bull., 34(1991), 249-253.
7. T. Nakazi and M. Yamada, Invertible Toeplitz operators and uniform algebras, Acta Sci.Math.(Szeged) 59(1994), 173-185.
8. Z. Nehari, On bounded bilinear forms, Ann. of Math. 65(1957), 153-162.

Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan
e-mail : nakazi@math.sci.hokudai.ac.jp

and

Mathematical and Scientific Subjects
Asahikawa National College of Technology
Asahikawa 071-8142, Japan