



HOKKAIDO UNIVERSITY

Title	Interpolation Of Weighted L^q Sequences By H^p Functions
Author(s)	Nakazi, Takahiko
Citation	Hokkaido University Preprint Series in Mathematics, 617, 1-11
Issue Date	2003-11-29
DOI	https://doi.org/10.14943/83762
Doc URL	https://hdl.handle.net/2115/69366
Type	departmental bulletin paper
File Information	re617.pdf



Interpolation Of Weighted ℓ^q Sequences By H^p Functions

by

Takahiko Nakazi*

* This research was partially supported by Grant-in-Aid for Scientific Research, Ministry of Education.

Mathematics Subject Classification 2000 : 30 D 55, 30 E 05

Keywords and phrases : weighted Hardy space, weighted sequence space, interpolation

Abstract. Let (z_n) be a sequence of points in the open unit disc D and $\rho_n = \prod_{m \neq n} |(z_n - z_m)(1 - \bar{z}_m z_n)^{-1}| > 0$. Let $a = (a_j)_{j=1}^\infty$ be a sequence of positive numbers and $\ell^s(a) = \{(w_j) ; (a_j w_j) \in \ell^s\}$ where $1 \leq s \leq \infty$. When $1 \leq p \leq \infty$ and $1/p + 1/q = 1$, we show that $\{(f(z_n)) ; f \in H^p\} \supset \ell^s(a)$ if and only if there exists a finite positive constant γ such that $\left\{ \sum_{n=1}^\infty (a_n \rho_n)^{-t} (1 - |z_n|^2)^t |f(z_n)|^t \right\}^{1/t} \leq \gamma \|f\|_q$ ($f \in H^q$), where $1/s + 1/t = 1$. As results, we show that $\{(f(z_j)) ; f \in H^p\} \supset \ell^1(a)$ if and only if $\sup_n (a_n \rho_n)^{-1} (1 - |z_n|^2)^{1/p} < \infty$, and $\{(f(z_n)) ; f \in H^1\} \supset \ell^\infty(a)$ if and only if $\sum_n (a_n \rho_n)^{-1} (1 - |z_n|^2) \delta_{z_n}$ is finite measure on D . These are also proved in the case of weighted Hardy spaces.

§1. Introduction

H^p ($0 < p \leq \infty$) denotes the usual Hardy space in the open unit disc D . In this paper, we assume that a sequence (z_j) in D satisfies that $\sum_{j=1}^{\infty} (1 - |z_j|) < \infty$, that is, there exists a Blaschke product

$$B(z) = \prod_{j=1}^{\infty} -\frac{\bar{z}_j}{|z_j|} \frac{z - z_j}{1 - \bar{z}_j z}.$$

Let

$$\begin{aligned} \rho_{k,n} &= \prod_{\substack{j=1 \\ j \neq k}}^n \left| \frac{z_k - z_j}{1 - \bar{z}_j z_k} \right|, \quad 1 \leq k \leq n, \\ \rho_k &= \prod_{\substack{j=1 \\ j \neq k}}^{\infty} \left| \frac{z_k - z_j}{1 - \bar{z}_j z_k} \right|. \end{aligned}$$

Then $\rho_{k,n} \geq \rho_{k,n+1}$ and $\lim_{n \rightarrow \infty} \rho_{k,n} = \rho_k$ for $k \geq 1$. We assume that $\rho_k > 0$ for $k = 1, 2, \dots$.

For a positive sequence $a = (a_j)$, $\ell^s(a)$ denotes $\{(w_j) ; w_j \in \mathcal{C} \text{ and } \sum_{j=1}^{\infty} (a_j |w_j|)^s < \infty\}$ and $\ell^\infty(a)$ denotes $\{(w_j) ; w_j \in \mathcal{C} \text{ and } \sup_{1 \leq j < \infty} a_j |w_j| < \infty\}$. In this paper, we study the following problem : Find a necessary and sufficient condition on (z_j) so that $\{(f(z_j)) ; f \in H^p\} \supset \ell^s(a)$ where $1 \leq p \leq \infty$ and $1 \leq s \leq \infty$.

Suppose $a_j = 1$ for all $j \geq 1$. When $p = s = \infty$, this was solved by L. Carleson [1]. That is, $\{(f(z_j)) ; f \in H^\infty\} \supset \ell^\infty$ if and only if $\inf_j \rho_j > 0$. (z_j) is called a uniformly separated sequence when $\inf_j \rho_j > 0$. When $p = \infty$ and $1 \leq s < \infty$, A. K. Snyder [13] (cf. [7],[11]) proved that $\{(f(z_j)) ; f \in H^\infty\} \supset \ell^s$ if and only if $\inf_j \rho_j > 0$. A. K. Snyder [13] and P. L. Duren and H. S. Shapiro [3] showed that there exists a sequence (z_j) which is not uniformly separated, that is, $\inf_j \rho_j = 0$ and has the property : $\{(f(z_j)) ; f \in H^p\} \supset \ell^\infty$ when $p \neq \infty$. B. A. Taylor and D. L. Williams [14] showed that for $1 \leq p \leq \infty$ $\{(f(z_j)) ; f \in H^p\} \supset \ell^\infty$ if and only if there exists a positive finite constant γ such that $\sum_{j=1}^{\infty} \frac{1}{\rho_j} (1 - |z_j|^2) |g(z_j)| \leq \gamma \|g\|_q$ for all g in H^q and $1/p + 1/q = 1$.

Suppose $1 \leq p = s \leq \infty$. When $a_j = (1 - |z_j|^2)^{1/p}$ for all $j \geq 1$, this was solved by H. S. Shapiro and A. L. Shields [11]. That is, $\{(f(z_j)) ; f \in H^p\} \supset \ell^p(a)$ if and only if $\inf_j \rho_j > 0$. When $a_j = \rho_j^2$ for all $j \geq 1$, J. P. Earl [4] showed that $\{(f(z_j)) ; f \in H^\infty\}$ contains $\ell^\infty(a)$ always. This was pointed out by A. M. Gleason (see [6]). On the other hand, when $a_j = \rho_j$ for all $j \geq 1$, T. Nakazi [10] showed that $\{(f(z_j)) ; f \in H^\infty\} \supset \ell^\infty(a)$ if and only if (z_j) is the union of a finite number of uniformly separated sequences. For a general weight $a = (a_j)$, J. D. McPhail [9] gave a necessary and sufficient condition about

(z_j) that $\{(f(z_j)) ; f \in H^p\} \supset \ell^p(a)$. In fact, he studied such a problem in weighted Hardy spaces.

In §2, we give a necessary and sufficient condition about (z_j) for that $\{(f(z_j)) ; f \in H^p\} \supset \ell^s(a)$ where $1 \leq p \leq \infty$, $1 \leq s \leq \infty$ and $a = (a_j)$ is arbitrary weight. As a result, we show that $\{(f(z_j)) ; f \in H^1\} \supset \ell^s(a)$ if and only if $\sum_{j=1}^{\infty} (a_j \rho_j)^{-t} (1 - |z_j|^2)^t < \infty$

where $1/s + 1/t = 1$. Moreover, when $1 < p \leq \infty$ and $a = (\rho_j^{-1})$, we show that $\{(f(z_j)) ; f \in H^p\} \supset \ell^p(a)$ if and only if (z_j) is a finite sum of uniformly separated sequences. This is a generalization of a result in [10] for $p = \infty$.

In §3, when $1 \leq p \leq \infty$, we show that $\{(f(z_j)) ; f \in H^p\} \supset \ell^1(a)$ if and only if $\sup_j (a_j \rho_j)^{-1} (1 - |z_j|^2)^{1/p} < \infty$. As a result, a theorem of A. K. Snyder [13] follows, that is, $\{(f(z_j)) ; f \in H^{\infty}\} \supset \ell^s$ if and only if $\inf_j \rho_j > 0$.

In §4, we give a necessary and sufficient condition about (z_j) for that $\{(f(z_j)) ; f \in H^p\} \supset \ell^{\infty}(a)$. Put $\mu = \sum_{j=1}^{\infty} (a_j \rho_j)^{-1} (1 - |z_j|^2) \delta_{z_j}$. Then $\{(f(z_j)) ; f \in H^1\} \supset \ell^{\infty}(a)$ if and only if μ is a finite measure on D , and $\{(f(z_j)) ; f \in H^{\infty}\} \supset \ell^{\infty}(a)$ if and only if μ is a Carleson measure on D .

In §5, we give a necessary and sufficient condition about (z_j) for that $\{(s(z_j)f(z_j)) ; f \in H^p(W)\} \supset \ell^p$, where $H^p(W)$ is a weighted Hardy space and $s(z_j) = \inf\{\int |f|^p W d\theta / 2\pi ; f(z_j) = 1\}$. We assume only that $\log W$ is in L^1 . J.D.McPhail [9] studied such a problem when W satisfies the (A_p) -condition of Muckenhoupt.

Our interests in this paper are in the differences between interpolations for $\ell^1(a)$ and $\ell^{\infty}(a)$ and in the interpolation problems for weighted Hardy spaces. For example, it is very easy to prove that $\{(f(z_j)) ; f \in H^{\infty}\} \supset \ell^1$ if and only if $\{z_j\}$ is uniformly separated.

§2. General results

In this section, we obtain a general result for interpolation problems for $\ell^s(a)$ ($1 \leq s \leq \infty$) by H^p ($1 \leq p \leq \infty$). For $1 \leq j \leq n$, let

$$B_n(z) = \prod_{j=1}^n \frac{z - z_j}{1 - \bar{z}_j z} \quad \text{and} \quad B_{nj}(z) = B_n(z) \frac{1 - \bar{z}_j z}{z - z_j}.$$

If we put $b_{nj} = B_{nj}(z_j)$, then

$$\rho_{j,n} = |b_{nj}| \quad (1 \leq j \leq n).$$

Suppose for $n = 1, 2, \dots$

$$f_n(z) = \sum_{j=1}^n b_{nj}^{-1} w_j B_{nj}(z).$$

Then f_n is in H^∞ and $f_n(z_j) = w_j$ ($1 \leq j \leq n$). Lemma 1 is essentially known.

Lemma 1. *Let $1 \leq p \leq \infty$ and $1/p + 1/q = 1$. Suppose w_j is a complex number for $j = 1, 2, \dots$. There exists a function f in H^p such that $f(z_j) = w_j$ for $j = 1, 2, \dots$ if and only if there exists a positive finite constant γ such that for any $n \geq 1$ and for all g in H^q ,*

$$\left| \sum_{j=1}^n \frac{w_j}{b_{nj}} (1 - |z_j|^2) g(z_j) \right| \leq \gamma \|g\|_q.$$

Proof. Put for $n \geq 1$

$$m_{p,n}(w) = \inf \{ \|f_n + B_n h\|_p ; h \in H^p \}.$$

Then by [2, p142],

$$m_{p,n}(w) = \sup \left\{ \left| \sum_{j=1}^n \frac{w_j}{b_{nj}} (1 - |z_j|^2) g(z_j) \right| ; g \in H^q \text{ and } \|g\|_q \leq 1 \right\}.$$

There exists a function f in H^p such that $f(z_j) = w_j$ for $j = 1, 2, \dots$ if and only if $\sup_n m_{p,n}(w) < \infty$ because the unit ball of H^p is compact in the weak topology or the weak $*$ topology. This implies the lemma.

Theorem 1. *Let $1 \leq p \leq \infty$ and $1 \leq s \leq \infty$. $\{(f(z_n)) ; f \in H^p\} \supset \ell^s(a)$ if and only if there exists a finite positive constant γ such that*

$$\left\{ \sum_{n=1}^{\infty} (a_n \rho_n)^{-t} (1 - |z_n|^2)^t |f(z_n)|^t \right\}^{1/t} \leq \gamma \|f\|_q$$

for f in H^q , where $1/p + 1/q = 1$ and $1/s + 1/t = 1$.

Proof. For the ‘only if’ part, since $\{(f(z_j)) ; f \in H^p\} \supset \ell^s(a)$, by Lemma 1 there exists a positive finite constant γ such that for any $n \geq 1$

$$\sup_{\substack{w \in \ell^s(a) \\ \|w\| \leq 1}} \left| \sum_{j=1}^n \frac{w_j}{b_{nj}} (1 - |z_j|^2) g(z_j) \right| \leq \gamma \|g\|_q \quad (g \in H^q)$$

where $w = (w_j)$ and $\|w\| = \left(\sum_{j=1}^{\infty} |w_j a_j|^s \right)^{1/s}$. Hence for any $n \geq 1$

$$\left\{ \sum_{j=1}^{\infty} (a_j \rho_{n,j})^{-t} (1 - |z_j|^2)^t |g(z_j)|^t \right\}^{1/t} \leq \gamma \|g\|_q \quad (g \in H^q).$$

Assuming $\|g\|_q = 1$,

$$\sum_{j=1}^{\infty} (a_j \rho_{n,j})^{-t} (1 - |z_j|^2)^t |g(z_j)|^t \leq \gamma^t.$$

For any $\varepsilon > 0$, there exists a positive integer n_j for each j such that for all $n \geq n_j$

$$(a_j \rho_{n,j})^{-t} (1 - |z_j|^2)^t |g(z_j)|^t - \frac{\varepsilon}{2^j} \leq (a_j \rho_{n,j})^{-t} (1 - |z_j|^2)^t |g(z_j)|^t$$

because $\rho_{j,n} \geq \rho_{j,n+1}$ and $\lim_{n \rightarrow \infty} \rho_{j,n} = \rho_j$. Thus, $\{(f(z_j)) ; f \in H^p\} \supset \ell^s(a)$ if and only if for any $\varepsilon > 0$ and any $n \geq \max(n_1, \dots, n_n)$

$$\sum_{j=1}^n (a_j \rho_j)^{-t} (1 - |z_j|^2)^t |g(z_j)|^t - \varepsilon \leq \sum_{j=1}^n (a_j \rho_j)^{-t} (1 - |z_j|^2)^t |g(z_j)|^t \leq \gamma^t$$

This implies the ‘only if’ part.

For the ‘if’ part, by Lemma 1 it is sufficient to show that there exists a finite positive constant γ such that for all $n \geq 1$

$$\sup_{\substack{w \in \ell^s(a) \\ \|w\| \leq 1}} \sup_{\|g\|_q \leq 1} \left| \sum_{j=1}^n \frac{w_j}{b_{nj}} (1 - |z_j|^2) g(z_j) \right| \leq \gamma < \infty.$$

In fact, for all $n \geq 1$

$$\begin{aligned} & \sup_{\substack{w \in \ell^s(a) \\ \|w\| \leq 1}} \sup_{\|g\|_q \leq 1} \left| \sum_{j=1}^n \frac{w_j}{b_{nj}} (1 - |z_j|^2) g(z_j) \right| \\ & \leq \sup_{\|g\|_q \leq 1} \left\{ \sum_{j=1}^n (a_j \rho_{j,n})^{-t} (1 - |z_j|^2)^t |g(z_j)|^t \right\}^{1/t} \\ & \leq \sup_{\|g\|_q \leq 1} \left\{ \sum_{j=1}^{\infty} (a_j \rho_j)^{-t} (1 - |z_j|^2)^t |g(z_j)|^t \right\}^{1/t} < \infty \end{aligned}$$

Corollary 1. Let $1 \leq s \leq \infty$. $\{(f(z_n)) ; f \in H^1\} \supset \ell^s(a)$ if and only if

$$\sum_{n=1}^{\infty} (a_n \rho_n)^{-t} (1 - |z_n|^2)^t < \infty$$

where $1/s + 1/t = 1$. Hence, when $a = (a_n) = (\rho_n^{-1})$ it is always true that $\{(f(z_n)) ; f \in H^1\} \supset \ell^s(a)$.

Proof. The first part is clear by Theorem 1. When $a = (\rho_n^{-1})$, $\{(f(z_n)) ; f \in H^1\} \supset \ell^s(a)$ if and only if $\sum_{n=1}^{\infty} (1 - |z_n|^2)^t < \infty$. This implies the second part.

Corollary 2. Let $1 \leq p \leq \infty$, $1 \leq s \leq \infty$ and $a = (\rho_n^{-1})$. $\{(f(z_n)) ; f \in H^p\} \supset \ell^s(a)$ if and only if there exists a finite positive constant γ such that

$$\left\{ \sum_{n=1}^{\infty} (1 - |z_n|^2)^t |f(z_n)|^t \right\}^{1/t} \leq \gamma \|f\|_q$$

for f in H^q , where $1/p + 1/q = 1$ and $1/s + 1/t = 1$. When $1 < p \leq \infty$, $\{(f(z_n)) ; f \in H^p\} \supset \ell^p(a)$ if and only if (z_n) is a finite sum of uniformly separated sequences.

Proof. The first part is clear by Theorem 1. The second part follows from the first one and [8].

In Corollary 2, when $1 < p \leq \infty$ and $1 < s \leq \infty$ and $s > p$, if $\{(f(z_n)) ; f \in H^p\} \supset \ell^s(a)$ then (z_n) is a finite sum of uniformly separated sequences but the converse is not true. When $s < p$, if (z_n) is a finite sum of uniformly separated sequences then $\{(f(z_n)) ; f \in H^p\} \supset \ell^s(a)$ but the converse is not true.

§3. Interpolations for $\ell^1(a)$

$\ell^1(a)$ is the smallest sequence space among $\ell^p(a)$ ($1 \leq p \leq \infty$) for the same $a = \{a_j\}$. Then the interpolations for $\ell^1(a)$ are very special as the following shows.

The case of $p = \infty$ in Corollary 3 was proved by A.Snyder [13] (see [7], [11]). Corollary 4 is due to O. Hatori [7].

Theorem 2. Let $1 \leq p \leq \infty$. $\{(f(z_n)) ; f \in H^p\} \supset \ell^1(a)$ if and only if

$$\sup_n (a_n \rho_n)^{-1} (1 - |z_n|^2)^{1/p} < \infty.$$

Proof. By Theorem 1, $\{(f(z_n)) ; f \in H^p\} \supset \ell^1(a)$ if and only if there exists a finite positive constant γ such that

$$\sup_n (a_n \rho_n)^{-1} (1 - |z_n|^2) |f(z_n)| \leq \gamma \|f\|_q$$

for all f in H^q . For each n , $\sup_{\|f\|_q=1} |f(z_n)| = (1 - |z_n|^2)^{-1/q}$ by [2, p144] and so the theorem follows.

Corollary 3. Let $1 \leq p \leq \infty$. $\{(f(z_n)) ; f \in H^p\} \supset \ell^1$ if and only if $\sup_n \frac{1}{\rho_n} (1 - |z_n|^2)^{1/p} < \infty$. Hence if $p = \infty$, $\{(f(z_n)) ; f \in H^\infty\} \supset \ell^1$ if and only if $\inf_n \rho_n > 0$.

Corollary 4. Let $1 \leq p \leq \infty$. $\{((1 - |z_n|^2)^{1/p} f(z_n)) ; f \in H^p\} \supset \ell^1$ if and only if $\inf_n \rho_n > 0$.

Proof. Note that $\{((1 - |z_n|^2)^{1/p} f(z_n)) ; f \in H^p\} \supset \ell^1$ if and only if $\{(f(z_n)) ; f \in H^p\} \supset \ell^1(a)$ and $a = ((1 - |z_n|^2)^{1/p})$.

Corollary 5. Let $1 \leq p \leq \infty$. For any (z_n) , $\{(f(z_n)) ; f \in H^p\} \supset \ell^1(a)$ where $a = (\rho_n^{-1})$.

Let (b_j) be a uniformly separated sequence in D such that $0 < \operatorname{Re} b_j \nearrow 1$ and $\operatorname{Im} b_j \searrow 0$. For $j \geq 1$, put $z_{2j-1} = b_j$ and $z_{2j} = \bar{b}_j$. Let B be the Blaschke product associated with $\{z_n\}$. Then for each j

$$B = \frac{z - b_j}{1 - \bar{b}_j z} \frac{z - \bar{b}_j}{1 - b_j z} B_{1j} B_{2j}$$

where B_{1j} (or B_{2j}) is a Blaschke product with zeros $\{b_\ell\}_{\ell \neq j}$ (or $\{\bar{b}_\ell\}_{\ell \neq j}$). Then

$$\rho_{2j-1} = \left| \frac{b_j - \bar{b}_j}{1 - \bar{b}_j b_j} \right| \prod_{\ell \neq j} \left| \frac{b_j - b_\ell}{1 - \bar{b}_\ell b_j} \right| \prod_{\ell \neq j} \left| \frac{b_j - \bar{b}_\ell}{1 - b_\ell b_j} \right|$$

and

$$\rho_{2j} = \left| \frac{\bar{b}_j - b_j}{1 - \bar{b}_j \bar{b}_j} \right| \prod_{\ell \neq j} \left| \frac{\bar{b}_j - b_\ell}{1 - \bar{b}_\ell \bar{b}_j} \right| \prod_{\ell \neq j} \left| \frac{\bar{b}_j - \bar{b}_\ell}{1 - b_\ell \bar{b}_j} \right|.$$

Hence $\rho_{2j-1} = \rho_{2j}$ for $j \geq 1$ and

$$\delta^2 \frac{|\bar{b}_j - b_j|}{1 - |b_j|^2} \leq \rho_{2j} = \rho_{2j-1} \leq \frac{|\bar{b}_j - b_j|}{1 - |b_j|^2} \quad (j \geq 1)$$

where

$$0 < \delta = \min \left\{ \inf_j \prod_{\ell \neq j} \left| \frac{b_j - b_\ell}{1 - \bar{b}_\ell b_j} \right|, \inf_j \prod_{\ell \neq j} \left| \frac{b_j - \bar{b}_\ell}{1 - b_\ell b_j} \right| \right\}.$$

Hence

$$\frac{(1 - |z_n|^2)^{1+1/p}}{|z_n - \bar{z}_n|} \leq \frac{(1 - |z_n|^2)^{1/p}}{\rho_n} \leq \delta^{-2} \frac{(1 - |z_n|^2)^{1+1/p}}{|z_n - \bar{z}_n|}.$$

Thus $\{(f(z_n)) ; f \in H^p\} \supset \ell^1$ if and only if $\sup_n (1 - |z_n|^2)^{1+1/p} / |z_n - \bar{z}_n| < \infty$.

§4. Interpolations for $\ell^\infty(a)$

$\ell^\infty(a)$ is the largest sequence space among $\ell^p(a)$ ($1 \leq p \leq \infty$) for the same $a = (a_j)$. Then the interpolations for $\ell^\infty(a)$ are special as the following shows. The case of $p = \infty$ of Corollary 6 is known in [10].

Theorem 3. *Let $1 \leq p \leq \infty$ and $1/p + 1/q = 1$, $\{(f(z_n)) ; f \in H^p\} \supset \ell^\infty(a)$ if and only if there exists a finite positive constant γ such that*

$$\sum_n (a_n \rho_n)^{-1} (1 - |z_n|^2) |f(z_n)| \leq \gamma \|f\|_q$$

for all f in H^q . When $p = 1$, $\{(f(z_n)) ; f \in H^1\} \supset \ell^\infty(a)$ if and only if $\mu = \sum_n (a_n \rho_n)^{-1} (1 - |z_n|^2) \delta_{z_n}$ is a finite measure on D . When $p = \infty$, $\{(f(z_n)) ; f \in H^\infty\} \supset \ell^\infty(a)$ if and only if $\mu = \sum_n (a_n \rho_n)^{-1} (1 - |z_n|^2) \delta_{z_n}$ is a Carleson measure on D .

Corollary 6. *Let $1 \leq p \leq \infty$ and $1/p + 1/q = 1$ and $a = (\rho_n^{-1})$. $\{(f(z_n)) ; f \in H^p\} \supset \ell^\infty(a)$ if and only if there exists a finite positive constant γ such that*

$$\sum_n (1 - |z_n|^2) |f(z_n)| \leq \gamma \|f\|_q$$

for all f in H^q .

- (1) *When $p = 1$, for any (z_n) , $\{(f(z_n)) ; f \in H^1\} \supset \ell^\infty(a)$.*
- (2) *When $p = \infty$, $\{(f(z_n)) ; f \in H^\infty\} \supset \ell^\infty(a)$ if and only if (z_n) is a finite union of uniformly separated sequences.*
- (3) *When $1 < p < \infty$, there exists a sequence (z_n) in D such that $\{(f(z_n)) ; f \in H^p\} \supset \ell^\infty(a)$ and (z_n) is not a union of finitely many uniformly separated sequences. If $\sum_{n=1}^{\infty} (1 - |z_n|^2)^{1/p} < \infty$, then $\{(f(z_n)) ; f \in H^p\} \supset \ell^\infty(a)$.*

Suppose that (z_n) is the sequence in D which was used in the end of Section 3, and $1 \leq p < \infty$. If $0 < \gamma_1 \leq \frac{(1 - |z_n|^2)^{1+1/p-\varepsilon}}{|z_n - \bar{z}_n|} \leq \gamma_2 < \infty$ for some $0 < \varepsilon < 1/p$, then $\{(f(z_n)) ; f \in H^p\} \supset \ell^\infty$. This was proved by B. A. Taylor and D. L. Williams [14].

§5. Weighted Hardy space

Let W be a nonnegative function in L^1 with $\log W \in L^1$ and $1 \leq p < \infty$. $H^p(W)$ denotes the closure of the set of all analytic polynomials in $L^p(W) = L^p(W d\theta/2\pi)$. $H^p(W)$ is called a weighted Hardy space. For $b \in D$, put

$$s(b) = s(b, p, W) = \inf \left\{ \int |f|^p W d\theta/2\pi ; f(b) = 1 \right\}.$$

Let h be an outer function in H^p such that $|h|^p = W$.

Lemma 2. *For $1 \leq p < \infty$ and $b \in D$,*

$$\begin{aligned} s(b, p, W) &= (1 - |b|^2) \exp(\log W)^\sim(b) \\ &= (1 - |b|^2) |h(b)|^p, \end{aligned}$$

where $(\log W)^\sim(b)$ denotes the Poisson integral of $\log W$ at b .

Proof. It is well known (cf. [5, p136]) that $s(0, p, W) = \exp \int_0^{2\pi} \log W d\theta/2\pi$. It is easy to show the lemma using $f(b) = f \circ \phi_b(0)$, where $\phi_b(z) = (z + b)/(1 + \bar{b}z)$.

Lemma 3. *Suppose (z_j) is a sequence of points in D . For $1 \leq p < \infty$ and $1 \leq s < \infty$, $\{(s(z_j, p, W)^{1/p} f(z_j)) ; f \in H^p(W)\} \supset \ell^s$ if and only if $\{(F(z_j)) ; F \in H^p\} \supset \ell^s(a)$, where $a = (a_j)$ and $a_j = s(z_j)^{1/p}/h(z_j)$.*

Proof. Since $H^p(W) = h^{-1}H^p$, $f \in H^p(W)$ if and only if $f = h^{-1}F$ and $F \in H^p$. For each j , $s(z_j)^{1/p}f(z_j) = w_j$ if and only if $F(z_j) = h(z_j)w_j/s(z_j)^{1/p}$ if and only if $F(z_j) = \zeta_j$, $w_j = a_j\zeta_j$. $(w_j) \in \ell^p$ if and only if $(\zeta_j) \in \ell^s(a)$. Now the lemma follows.

Theorem 4. *Let $1 \leq p < \infty$, $1 \leq s \leq \infty$, and $1/p + 1/q = 1/s + 1/t = 1$. Then, $\{(s(z_n, p, W)^{1/p}f(z_n)) ; f \in H^p(W)\} \supset \ell^s$ if and only if*

$$\left\{ \sum_{n=1}^{\infty} \frac{1}{\rho_n^t} s(z_n)^{t/q} |g(z_n)|^t \right\}^{1/t} \leq \gamma \|g\|_{H^q(W)}$$

for g in $H^q(W)$.

Proof. By Lemma 3, $\{(s(z_n)^{1/p}f(z_n)) ; f \in H^p(W)\} \supset \ell^s$ if and only if $\{(F(z_n)) ; F \in H^p\} \supset \ell^s(a)$, where $a_n = s(z_n)^{1/p}/|h(z_n)|$. By Theorem 1, this is equivalent to saying that there exists a finite positive constant γ such that

$$\left\{ \sum_{n=1}^{\infty} \frac{1}{\rho_n^t} \frac{1}{a_n^t} (1 - |z_n|^2)^t |G(z_n)|^t \right\}^{1/t} \leq \gamma \|G\|_q$$

for $G \in H^q$. Since $H^q(W) = h^{-p/q}H^q$, $g \in H^q(W)$ if and only if $g = h^{-p/q}G$ and $G \in H^q$. Hence $\|g\|_{H^q(W)} = \|G\|_{H^q}$ and for each $n \geq 1$

$$\begin{aligned} & a_n^{-t} (1 - |z_n|^2)^t |G(z_n)|^t \\ &= s(z_n)^{-(t/p)} |h(z_n)|^t (1 - |z_n|^2)^t |h(z_n)|^{pt/q} |g(z_n)|^t \\ &= s(z_n)^{-(t/p)} (1 - |z_n|^2)^t |h(z_n)|^{t(q+p)/q} |g(z_n)|^t \\ &= s(z_n)^{-(t/p)} s(z_n)^t |g(z_n)|^t \\ &= s(z_n)^{t/q} |g(z_n)|^t. \end{aligned}$$

This implies the theorem.

Corollary 7. *Let $1 < p < \infty$ and $1/p + 1/q = 1$. Then $\{(s(z_n, p, W)^{1/p}f(z_n)) ; f \in H^p(W)\} \supset \ell^1$ if and only if $\inf_n \rho_n > 0$.*

Proof. By Theorem 4, $\{(s(z_n, p, W)^{1/p}f(z_n)) ; f \in H^p(W)\} \supset \ell^1$ if and only if

$$\sup_n \frac{1}{\rho_n} s(z_n, p, W)^{1/p} s(z_n, q, W)^{-1/q} < \infty.$$

Now Lemma 2 implies the corollary.

References

1. L. Carleson, An interpolation problem for bounded analytic functions, Amer.J.Math. 80(1958), 921-930.
2. P. L. Duren, Theory of H^p Spaces, Academic Press, New York, 1970.

3. P. L. Duren and H. S. Shapiro, Interpolation in H^p spaces, Proc.Amer.Math.Soc. 31(1972), 162-164.
4. J. P. Earl, On the interpolation of bounded sequences by bounded functions, J.London Math.Soc (2)2(1970), 544-548.
5. T.W.Gamelin, Uniform Algebras, Chelsea Publishing Company, 1969.
6. J. Garnett, Two remarks on interpolation by bounded analytic functions, Banach Spaces of Analytic Functions (Baker et al., eds)(Lecture Notes in Math. Vol.604, 32-40), Springer-Verlag, Berlin.
7. O. Hatori, The Shapiro-Shields theorem on finite connected domains, Surikaiseikikenkyūsho Kōkyūroku 1049(1998), 21-29 (in Japanese).
8. S. McDonald and C. Sundberg, Toeplitz operators on the disc, Indiana Univ.Math. J. 28(1979), 595-611.
9. J. D. McPhail, A weighted interpolation problem for analytic functions, Studia Math.T.XCVI(1990), 105-106.
10. T. Nakazi, Notes on interpolation by bounded analytic functions, Canad.Math.Bull. 31(1988), 404-408.
11. T. Nakazi, Interpolation problem for ℓ^1 and a uniform algebra, J.Austral.Math.Soc. 72(2002), 1-11.
12. H. S. Shapiro and A.L.Shields, On some interpolation problems for analytic functions, Amer.J.Math. 83(1961), 513-532.
13. A. K. Snyder, Sequence spaces and interpolation problems for analytic functions, Studia Math. 39(1971), 137-153.
14. B. A. Taylor and D. L. Williams, Interpolation of ℓ^q sequences by H^p functions, Proc.Amer.Math.Soc. 34(1972), 181-186.
15. H. Turku and S. V. Švedenko, Interpolation of ℓ^p sequences by H^2 functions over a half-plane, Math.Balkanica 6 : 43(1976), 273-280.

Department of Mathematics
Hokkaido University
Sapporo 060-0810, Japan

nakazi@math.sci.hokudai.ac.jp