



HOKKAIDO UNIVERSITY

Title	LECTURES ON NONLINEAR DISPERSIVE EQUATIONS II
Author(s)	Ozawa, T.; Tsutsumi, Y.
Description	Program N. Tzvetkov Ill-posedness issues for nonlinear dispersive equations H. Koch Dispersive estimates and applications
Citation	Hokkaido University technical report series in mathematics, 86, 1
Issue Date	2004-01-01
DOI	https://doi.org/10.14943/645
Doc URL	https://hdl.handle.net/2115/703
Type	departmental bulletin paper
File Information	tn86.pdf



LECTURES ON NONLINEAR DISPERSIVE EQUATIONS II

Edited by
T. Ozawa and Y. Tsutsumi

Sapporo, 2004

Partially supported by JSPS

- Grant-in-Aid for formation of COE
“Mathematics of Nonlinear Structure via Singularities”
- Grant-in-Aid for Scientific Research
S(2) #16104002 (T. Ozawa)
A(2) #15204008 (Y. Tsutsumi)

PREFACE

This volume, together with the last, is intended as the proceedings of expository lectures in Special Months “Nonlinear Dispersive Equations. ”

Nonlinear dispersive equations, such as nonlinear Schrödinger equations, KdV equation, and Benjamin-Ono equation, are of mathematical and physical importance. Expository courses in September 2004 are intended to cover a broad spectrum of the issues, especially the Cauchy problem and related topics.

We wish to express our sincere thanks to

- J. Bona, H. Koch, F. Planchon, P. Raphaël, and N. Tzvetkov for excellent lectures.
- M. Ikawa and A. Ogino for efficient arrangements.

T. Ozawa and Y. Tsutsumi

CONTENTS

Program

N. Tzvetkov

Ill-posedness issues for nonlinear dispersive equations

H. Koch

Dispersive estimates and applications

COE Program Special Months

Lectures on Nonlinear Dispersive Equations II

Organizers: T. Ozawa (Hokkaido Univ.)
Y. Tsutsumi (Kyoto Univ.)

Period :

- September 15 - 17, 2004
- September 27 - 29, 2004

Venue :

Department of Mathematics, Hokkaido University
Science Building #8 Room 309

Program :

	10:30–12:00	13:30–15:00	15:30–17:00
Sep. 15 (Wed)	–	–	Tzvetkov (1)
16 (Thu)	–	Tzvetkov (2)	Tzvetkov (3)
17 (Fri)	–	Tzvetkov (4)	Tzvetkov (5)
Sep. 27 (Mon)	–	Koch (1)	Koch (2)
28 (Tue)	–	Koch (3)	Koch (4)
29 (Wed)	Koch (5)	–	–

- N. Tzvetkov (Université Lille I)
Ill-posedness issues for nonlinear dispersive equations
- H. Koch (Universität Dortmund)
Dispersive estimates and applications

– – *COE Program Special Months Homepage* – –

<http://coe.math.sci.hokudai.ac.jp/news/special/index02.html.en>

Secretariat: Atsuko Ogino TEL: 011-706-4671 FAX: 011-706-4672

ILL-POSEDNESS ISSUES FOR NONLINEAR DISPERSIVE EQUATIONS

N. TZVETKOV

ABSTRACT. These notes are devoted to the notion of well-posedness of the Cauchy problem for nonlinear dispersive equations. We present recent methods for proving ill-posedness type results for dispersive PDE's. The common feature in the analysis is that the proof of such results requires the construction of high frequency approximate solutions on small time intervals (possibly depending on the frequency). The classical notion of well-posedness, going back to Hadamard, requires the existence, the uniqueness and the continuity of the flow map on the spaces where the existence is established. It turns out that in many cases a stronger form of well-posedness holds. Namely, the flow map enjoys better continuity properties as for example being Lipschitz continuous on bounded sets. In such a situation we say that the corresponding problem is semi-linearly well-posed in the corresponding functional setting. Our main message is that for dispersive PDE's, contrary to the case of hyperbolic PDE's, the verification whether an equation in hand is semi-linearly well-posed in a given functional framework requires a considerable care. Our examples are KdV type equations and non linear Schrödinger equations.

1. INTRODUCTION

We will discuss here the Cauchy problem for nonlinear PDE's which can be written in the form

$$(1.1) \quad \dot{u}(t) = Lu(t) + F(u(t)), \quad u(0) = u_0,$$

where $u(t)$, $t \in \mathbb{R}$ is a function defined on a Riemannian manifold (M, g) with values either in \mathbb{R} or in \mathbb{C} . In (1.1), L is a linear map acting as an isometry on the Sobolev spaces $H^s(M)$ while $F(u(t))$ represents the nonlinear interaction. The initial data u_0 is supposed to belong to $H^s(M)$. This choice is natural because, for the models we are interested in, the equation (1.1) enjoys conservation laws providing a uniform control on (low regularity) Sobolev norms of the solutions of (1.1). An important aspect of the analysis of the Cauchy problem (1.1) is to understand the impact of

1991 *Mathematics Subject Classification.* 35Q53, 35A07, 35B30 .

Key words and phrases. well-posedness, ill-posedness, dispersive equations.

PRELIMINARY VERSION

the interplay between L and F on the behavior on the solutions of (1.1). Here we will study this issue only for *small times* t . As far as the Sobolev spaces $H^s(M)$ are chosen for phase spaces, the local in time behavior of the solutions is naturally linked to the notion of well-posedness of the Cauchy problem (1.1) that we recall now.

Definition 1.1. *We say that the Cauchy problem (1.1) is well-posed for data in $H^s(M)$, if for every bounded set B of $H^s(M)$ there exist $T > 0$ and a Banach space \mathcal{X}_T continuously embedded in $C([-T, T]; H^s(M))$ such that if $u_0 \in B$ then there exists a unique solution u of (1.1) on $[-T, T]$ in the class \mathcal{X}_T . Moreover :*

1. *The flow map $u_0 \mapsto u$ is continuous from B to $C([-T, T]; H^s(M))$.*
2. *Higher smoothness is propagated by the flow. More precisely, if $u_0 \in H^\sigma(M)$, $\sigma \geq s$ then $u \in C([-T, T]; H^\sigma(M))$.*

Let us notice that in the above definition, the time of existence T depends only on the bounded set B , i.e. on an H^s bound of the initial data. There are several important examples of the so called critical problems when the time of existence is depending in a more complicated way on the initial data. It is worth noticing that “usually”, if a problem in hand is critical for data in H^s then it is well-posed in the sense of Definition 1.1 for data in H^σ , $\sigma > s$. It is also “usual” that the well-posedness in H^s , implies the well-posedness in $H^{s'}$, $s' \geq s$.

A very common way to prove the well-posedness of (1.1) is to solve by a contraction principle an equivalent integral equation, exactly as we do in the proof of the Cauchy-Lipschitz theorem in the theory of the ordinary differential equations. More precisely, the problem (1.1) can be rewritten, at least formally, as an integral equation (Duhamel formula)

$$(1.2) \quad u(t) = \exp(tL)u_0 + \int_0^t \exp((t-t')L)F(u(t'))dt'.$$

The well-posedness of (1.1) is reduced to finding a functional spaces \mathcal{X}_τ , $\tau > 0$ continuously embedded in $C([- \tau, \tau]; H^s(M))$ such that for every bounded set B of $H^s(M)$ there exists $T > 0$ such that for every $u_0 \in B$ the right hand-side of (1.2) is a contraction in a suitable ball of \mathcal{X}_T . In some cases, the space $C([- \tau, \tau]; H^s(M))$ can give the contraction properties. However, in these cases the assumption on s is quite restrictive. In order to include a larger possible values of s , the whole difficulty in making work the above approach is to find functional spaces \mathcal{X}_τ , $\tau > 0$ which are adapted in the best way to the equation in hand. This problematic has now a long history and remains an active research field. Once the existence and the uniqueness in \mathcal{X}_T is established, it is natural to look for a larger uniqueness class,

for instance one may ask whether the uniqueness holds in $C([-T, T]; H^s(M))$ (cf. e.g. [42]).

It turns out that if we are able to show the well-posedness of (1.1) by the above procedure then the flow map enjoys better continuity properties, for example it is Lipschitz continuous on B , and, in the case of polynomial nonlinearities it is a C^∞ map from $H^s(M)$ to $C([-T, T]; H^s(M))$. These properties seem to be related to what we call a semi-linearly well-posed problem. The following definition seems to be natural (cf. e.g. [5, 6, 12] ...).

Definition 1.2. *We say that the Cauchy problem (1.1) is semi-linearly well-posed for data in $H^s(M)$, if it is well-posed in the sense of Definition 1.1, and, in addition the flow map $u_0 \mapsto u$ is **uniformly continuous** from B to $C([-T, T]; H^s(M))$.*

The notion of well-posedness of Definition 1.1 is invariant under changes of variables in the phase space which are continuous on H^s . Similarly the notion of semi-linear well-posedness is invariant under uniformly continuous changes of variables. Therefore, it is not excluded that, by a change of variables (gauge transform)

$$u(t) \longrightarrow v(t)$$

in (1.1) which is continuous on H^s but not uniformly continuous, the equation for $v(t)$ to be semi-linearly well-posed even if the equation for $u(t)$ is not semi-linearly well-posed.

Another and quite different way to solve (1.1) is to apply a compactness argument. Roughly speaking, it means to solve the equation by passing to a (weak) limit in a family of approximate solutions. Usually this method can provide the well-posedness of (1.1), but it does not give directly the semi-linear well-posedness as the contraction method does. A natural question is whether there exists PDE's which are well-posed but not semi-linearly well-posed in $H^s(M)$. Probably the simplest example of such a PDE is the Burgers equation

$$(1.3) \quad u_t + uu_x = 0,$$

posed on $H^s(\mathbb{R})$ for real valued u (if u is not real valued the situation is quite different, as it is shown in [27]). It turns out that (1.3) is well-posed in $H^s(\mathbb{R})$, $s > 3/2$ but not semi-linearly well-posed in this same space. Let us explain how we prove the well-posedness of (1.3) for data in $H^s(\mathbb{R})$, $s > 3/2$. Let u be a smooth solution of (1.3) which belong to all H^σ , $\sigma \in \mathbb{R}$. Our purpose is to establish a priori bounds for u . Denote by D^s the Fourier multiplier with symbol $(1 + \xi^2)^{s/2}$, i.e.

$$\widehat{D^s u}(\xi) = (1 + \xi^2)^{s/2} \widehat{u}(\xi),$$

where the Fourier transform is defined as follows

$$\widehat{u}(\xi) = \int_{-\infty}^{\infty} e^{-ix\xi} u(x) dx .$$

Notice that $\|u\|_{H^s} = \|D^s u\|_{L^2}$. Applying D^s to (1.3), multiplying it with $D^s u$ and an integration by parts gives,

$$\frac{d}{dt} \|u(t, \cdot)\|_{H^s}^2 = \int_{-\infty}^{\infty} u_x(t, x) (D^s u(t, x))^2 dx - 2 \int_{-\infty}^{\infty} ([D^s, u]u_x)(t, x) D^s u(t, x) dx .$$

Using the Kato-Ponce (cf. [43]) commutator estimate

$$(1.4) \quad \|[D^s, f]g\|_{L^2} \leq C(\|f_x\|_{L^\infty} \|D^{s-1}g\|_{L^2} + \|D^s f\|_{L^2} \|g\|_{L^\infty})$$

with $f = u$ and $g = u_x$, we obtain that

$$\frac{d}{dt} \|u(t, \cdot)\|_{H^s}^2 \leq C \|u_x(t, \cdot)\|_{L^\infty} \|u(t, \cdot)\|_{H^s}^2 .$$

Thus the Gronwall lemma yields that for every $0 \leq t \leq T$,

$$(1.5) \quad \|u(t, \cdot)\|_{H^s} \leq \|u(0, \cdot)\|_{H^s} \exp(C \|u_x\|_{L^1([0, T]; L^\infty)}) .$$

If $s > 3/2$, the Sobolev embedding gives,

$$(1.6) \quad \|u_x\|_{L^1([0, T]; L^\infty)} \leq CT \|u\|_{L^\infty([0, T]; H^s)} .$$

Combining (1.5) and (1.6), using a continuity argument, we deduce that there exist $c > 0$ and $C > 0$ such that if

$$T \leq c(1 + \|u_0\|_{H^s})^{-1}$$

then

$$(1.7) \quad \|u_x\|_{L^1([0, T]; L^\infty)} \leq C$$

and

$$(1.8) \quad \|u\|_{L^\infty([0, T]; H^s)} \leq C \|u(0, \cdot)\|_{H^s} .$$

The priori estimate (1.8) is the key to perform a classical compactness argument (cf. e.g. [55]) which provides the existence. The uniqueness is easily ensured by the Gronwall lemma. The propagation of the higher Sobolev regularity readily follows from (1.7) and (1.8).

The continuous dependence is a slightly more delicate issue and can be obtained for instance by the Bona-Smith argument [7] (cf. also [40]). Let us briefly recall this argument. Fix a bump function $\rho \in \mathcal{S}(\mathbb{R})$ such that $\widehat{\rho} \in C_0^\infty(\mathbb{R})$ and $\widehat{\rho}(\xi) = 1$ for ξ in a neighborhood of 0. For $\varepsilon > 0$, we set $\rho_\varepsilon(x) := \varepsilon^{-1} \rho(x/\varepsilon)$. Let u be a solution of (1.3) with data $u(0) \in H^s(\mathbb{R})$, $s > 3/2$ which belongs to a *fixed bounded set* of

$H^s(\mathbb{R})$. Denote by u^ε the solution of the Burgers equation (1.3) with initial data $\rho_\varepsilon \star u(0)$. One can easily check that

$$\|\rho_\varepsilon \star u(0)\|_{H^s} \leq C\|u_0\|_{H^s}, \quad \varepsilon \in]0, 1]$$

and therefore we can assume that u^ε enjoys the bounds (1.7) and (1.8) on the time of existence of u . For $\varepsilon > \varepsilon' > 0$, we set $v := u^\varepsilon - u^{\varepsilon'}$. Then v is a solution of the equation

$$(1.9) \quad 2v_t + (u_x^\varepsilon + u_x^{\varepsilon'})v + (u^\varepsilon + u^{\varepsilon'})v_x = 0.$$

It is easy to check that

$$(1.10) \quad \|v(0)\|_{H^s} = o(1), \quad \|v(0)\|_{L^2} = \mathcal{O}(\varepsilon^s)$$

as $\varepsilon \rightarrow 0$. Multiplying (1.9) with v and applying (1.7) (with u^ε and $u^{\varepsilon'}$ instead of u) gives the bound

$$(1.11) \quad \|v(t, \cdot)\|_{L^2} \leq C\varepsilon^s$$

for t in the time of existence of u . Applying D^s to (1.9), multiplying it with $D^s v$ and using the Kato-Ponce estimate (1.4) yields the estimate

$$(1.12) \quad \begin{aligned} \frac{d}{dt} \|v(t, \cdot)\|_{H^s}^2 &\leq C(\|u^\varepsilon(t, \cdot)\|_{H^s} + \|u^{\varepsilon'}(t, \cdot)\|_{H^s}) \|v(t, \cdot)\|_{H^s}^2 + \\ &+ C(\|u^\varepsilon(t, \cdot)\|_{H^{s+1}} + \|u^{\varepsilon'}(t, \cdot)\|_{H^{s+1}}) \|v(t, \cdot)\|_{H^{s-1}} \|v(t, \cdot)\|_{H^s}. \end{aligned}$$

Using (1.8) gives

$$(1.13) \quad \|u^\varepsilon(t, \cdot)\|_{H^{s+1}} \leq C\|u^\varepsilon(0, \cdot)\|_{H^{s+1}} \leq C\varepsilon^{-1}.$$

On the other hand, thanks to (1.11),

$$(1.14) \quad \|v(t, \cdot)\|_{H^{s-1}} \leq \|v(t, \cdot)\|_{L^2}^{\frac{1}{s}} \|v(t, \cdot)\|_{H^s}^{1-\frac{1}{s}} \leq C\varepsilon \|v(t, \cdot)\|_{H^s}^{1-\frac{1}{s}}$$

Using (1.12), (1.13), (1.14), a variant of the Gronwall lemma, and (1.10) gives

$$(1.15) \quad \|u^\varepsilon(t, \cdot) - u^{\varepsilon'}(t, \cdot)\|_{H^s} = \|v(t, \cdot)\|_{H^s} = o(1)$$

as $\varepsilon \rightarrow 0$. We can now easily obtain the continuity of the flow map. Indeed, let $(u_{0,n})$ be a sequence converging to u_0 in $H^s(\mathbb{R})$ with corresponding solutions (u_n) . Then $\rho_\varepsilon \star u_{0,n} \rightarrow u_{0,n}$ as $\varepsilon \rightarrow 0$ in $H^s(\mathbb{R})$, *uniformly in n* , and, exactly as above, we can show

$$(1.16) \quad \|u_n^\varepsilon(t, \cdot) - u^\varepsilon(t, \cdot)\|_{H^s} \leq C\|u_{0,n} - u_0\|_{H^s} + o(1),$$

where $o(1) \rightarrow 0$ as $\varepsilon \rightarrow 0$. It is now a routine procedure to show that (1.15) and (1.16) imply the continuity on $H^s(\mathbb{R})$, $s > 3/2$ of the flow map of the Burgers equation (1.3).

At this point, it is worth to notice that the argument based on a priori estimates for proving the well-posedness that we have just presented is less perturbative (“more nonlinear”) than the contraction method explained after Definition 1.1. It has the advantage to have a larger scope of applicability compared to the contraction method, but, at the present moment, to make it work one should require considerably more regularity on the initial data.

Let us next describe an argument providing the lack of semi-linear well-posedness of (1.3). We first observe that if u solves (1.3) then so does

$$(1.17) \quad v(t, x) = u(t, x - \omega t) + \omega, \quad \omega \in \mathbb{R}.$$

The shift in the spatial variable in (1.17) is “responsible” for the failure of uniform continuity of the flow map. The constant ω in (1.17) can be replaced by a function which is zero at infinity, thanks to the finite propagation speed of the Burgers equation. More precisely, inspired by (1.17), we look for an approximate solution of the Burgers equation of the form

$$(1.18) \quad u_{ap}^{\omega, \lambda}(t, x) = \omega \lambda^{-1} \tilde{\varphi}(x/\lambda^\delta) + \lambda^{-\delta/2-s} \varphi(x/\lambda^\delta) \cos(\lambda x - \omega t),$$

where $s > 3/2$, $\lambda \in [-1, 1]$, $\omega \in \mathbb{R}$, $1 < \delta < 2$ and $\varphi, \tilde{\varphi}$ are non zero $C_0^\infty(\mathbb{R})$ functions such that $\tilde{\varphi}$ is equal to one on the support of φ . We can then show that there exists $\varepsilon > 0$ such that

$$(1.19) \quad \|\partial_t u_{ap}^{\omega, \lambda} + u_{ap}^{\omega, \lambda} \partial_x u_{ap}^{\omega, \lambda}\|_{L^2(\mathbb{R})} \leq C \lambda^{-\varepsilon-s}.$$

Thanks to (1.19) and the well-posedness analysis in H^s , $s > 3/2$, we obtain that (1.18) is indeed a good approximate solution, in H^s , of the Burgers equation. Considering the sequences $(u_{ap}^{1, \lambda})$ and $(u_{ap}^{-1, \lambda})$, $\lambda \gg 1$ gives the failure of uniform continuity on $H^s(\mathbb{R})$, $s > 3/2$ of the flow map of the Burgers equation.

However, the Burgers equation (1.3) does not fit in the class of dispersive PDE’s and one may think that the above described property of (1.3) is only related to its the hyperbolic nature. It turns out that the Benjamin-Ono equation

$$(1.20) \quad u_t + H u_{xx} + u u_x = 0,$$

posed on $H^s(\mathbb{R})$ (in (1.20) H denotes the Hilbert transform which is a “zero order” operator) is well-posed in $H^s(\mathbb{R})$, $s > 3/2$ but not semi-linearly well-posed in this same space. The equation (1.20) fits in the class of the dispersive equations because

of the presence of the term Hu_{xx} . We use the term “dispersive equation” since any solution of

$$(1.21) \quad u_t + Hu_{xx} = 0,$$

issued from $L^1(\mathbb{R})$ initial data disperses as $t \rightarrow \infty$, more precisely,

$$\lim_{t \rightarrow \infty} \|u(t, \cdot)\|_{L^\infty(\mathbb{R})} = 0.$$

However this property is for large times, and, since we are concerned with a small time analysis a more relevant property related to the dispersive nature of the equation (1.21) is the (small time) Strichartz inequality (cf. e.g. [60]). More precisely, there exists $C > 0$ such that for every $T > 0$, every $u_0 \in L^2(\mathbb{R})$ the solution u of (1.21) with data u_0 satisfies,

$$(1.22) \quad \|u\|_{L^p([0,T]; L^q(\mathbb{R}))} \leq C \|u_0\|_{L^2(\mathbb{R})}, \quad \frac{2}{p} + \frac{1}{q} = \frac{1}{2}, \quad p \geq 4.$$

Estimates of type (1.22) are usually very useful to apply the contraction strategy but in the case of (1.20) they are not sufficient to make it work.

We next consider the KdV equation

$$(1.23) \quad u_t + u_{xxx} + uu_x = 0,$$

posed on $H^s(\mathbb{R})$, which have a higher order dispersion compared to (1.20). It turns out that, in sharp contrast with the Burgers and the Benjamin-Ono equations, the KdV equation (1.23) is semi-linearly well-posed for data in $H^s(\mathbb{R})$, $s > -3/4$. Therefore, the notion of semi-linear well-posedness makes a natural classification in the class of KdV type models, i.e. equation (1.1) with $F(u) = uu_x$ and $L = |D_x|^\alpha \partial_x$, depending on the order of dispersion $\alpha > 0$.

Another set of models where the notion of semi-linear well-posedness is naturally involved (but less understood) are the nonlinear Schrödinger equations (NLS). Let (M, g) be a compact smooth boundaryless Riemannian manifold of dimensions $d = 2, 3$. Denote by Δ the Laplace-Beltrami operator associated to the metric g . We consider the nonlinear Schrödinger equation

$$(1.24) \quad iu_t + \Delta u + |u|^2 u = 0,$$

posed on M . In (1.24), u is complex valued function on M . Let us first consider the case $d = 2$, i.e. the case when M is a surface. If M is the flat torus \mathbb{T}^2 then the Cauchy problem associated to (1.24) is semi-linearly well-posed for data in $H^s(\mathbb{T}^2)$, provided $s > 0$. This result is essentially sharp, since for $s < 0$ the problem is not semi-linearly well-posed for data in $H^s(M)$ for an arbitrary (M, g) . On the

other hand if M is the standard sphere S^2 then the Cauchy problem for (1.24) is not semi-linearly well-posed for data in $H^s(S^2)$, $s < 1/4$, in sharp contrast with the case of the torus \mathbb{T}^2 . Hence the same equation (1.24) behaves quite differently with respect to the semi-linear well-posedness depending on the geometry of the spatial domain. It is an interesting open problem whether (1.24) posed on S^2 might be well-posed for some $s \in [0, 1/4]$. Let us next consider the case $d = 3$. If M is the torus \mathbb{T}^3 or the sphere S^3 then the Cauchy problem for (1.24) is semi-linearly well-posed for data in H^s , $s > 1/2$. It turns out that this result is essentially sharp even regarding the classical notion of well-posedness. More precisely, for $0 < s < 1/2$, the Cauchy problem for (1.24) posed on an arbitrary M is not well-posed for data in H^s .

One may ask for the critical threshold in the scale of H^s for the well-posedness of (1.1). It means to find a real number s_c such that for $s < s_c$ (1.1) is not well-posed for data in $H^s(M)$, while for $s > s_c$ (1.1) is well-posed for data in $H^s(M)$. Similarly, one can define a critical threshold for the semi-linear well-posedness. In this context, the discussion around (1.24) above simply affirms that, for $d = 3$ and $M = \mathbb{T}^3$, the value $s_c = \frac{1}{2}$ is the critical threshold for both the well-posedness and the semi-linear well-posedness, as far as positive values of the Sobolev regularity s are considered. It is a natural question whether the critical threshold for the well-posedness and the semi-linear well-posedness may be different. The answer of this question is positive as shows the following example. Consider the following version of the modified KdV equation

$$(1.25) \quad u_t + u_{xxx} + \left(u^2 - \int_{\mathbb{T}} u^2(t, x) dx\right) u_x = 0,$$

posed on the torus $\mathbb{T} = \mathbb{R}/\mathbb{Z}$. The equation (1.25) can be obtained from the modified KdV equation

$$(1.26) \quad v_t + v_{xxx} + v^2 v_x = 0,$$

by the *gauge transformation* $u \rightarrow v$ defined as

$$v(t, x) = u\left(t, x - \int_0^t \int_{\mathbb{T}} u^2(\tau, y) dy d\tau\right).$$

The Cauchy problem for (1.25) is semi-linearly well-posed for data in $H^s(\mathbb{T})$, $s > 1/2$ (cf. [10]), it is not semi-linearly well-posed for data in $H^s(\mathbb{T})$, $3/8 < s < 1/2$ (cf. [62]), but ... it is still well-posed for data in $H^s(\mathbb{T})$, $s \in [3/8, 1/2]$ (cf. [62, 38, 39]). Hence the critical threshold for the well-posedness and the semi-linear well-posedness can be different.

Let us complete this introduction by noticing that, in the last years, gauge transformations were an important tool in the study of dispersive PDE's, cf. e.g. [10, 34, 56, 62, 64, 65, 66] ...

2. KdV TYPE PROBLEMS

Consider the Cauchy problem for the Korteweg de Vries (KdV) equation

$$(2.1) \quad u_t + u_{xxx} + uu_x = 0, \quad u(0) = u_0.$$

The best known result regarding the well-posedness of (2.1) is due to Kenig-Ponce-Vega.

Theorem 2.1 (cf. [46]). *For $s > -3/4$ the Cauchy problem (2.1) is semi-linearly well-posed for data in $H^s(\mathbb{R})$.*

To prove Theorem 2.1 one uses the contraction method as explained after Definition 1.1 of the previous section. The spaces X_T where one performs the argument are the Fourier transform restriction spaces introduced by Bourgain [8, 9, 10], equipped with the norm

$$\|u\|_{X_T} = \inf \{ \|w\|_X, \quad w \in X \quad \text{with} \quad w|_{[-T,T]} = u \},$$

where

$$\|w\|_X^2 = \int_{\mathbb{R}^2} (1 + |\tau - \xi^3|^2)^b (1 + |\xi|^2)^s |\widehat{w}(\tau, \xi)|^2 d\tau d\xi$$

with $b > 1/2$ sufficiently close to $1/2$. The spaces of Bourgain are very useful to recover the derivative loss in the nonlinearity. We refer to [13, 32] for an introduction to the Fourier transform restriction method of Bourgain. There has been a number of works preceding Theorem 2.1 where the well-posedness for bigger values of s were established (cf. e.g. [61, 44, 45, 10]). A particularly important step was done in [45], where it is realized for the first time that the KdV equation can be semi-linearly well-posed. The value $s = -3/4$ in Theorem 2.1 is optimal, as far as the semi-linear well-posedness is concerned (cf. [24]). But it is a priori not excluded (2.1) to be well-posed for some $s < -3/4$.

Next we consider the Cauchy problem for the Benjamin-Ono (BO) equation (cf. [3])

$$(2.2) \quad u_t + H u_{xx} + u u_x = 0, \quad u(0) = u_0.$$

In (2.2), H denotes the Hilbert transform, namely,

$$(Hf)(x) := 2 \lim_{\varepsilon \rightarrow 0} \int_{|x-y| \geq \varepsilon} \frac{f(y)}{x-y} dy.$$

It is easy to check that

$$\widehat{Hf}(\xi) = -i \operatorname{sign}(\xi) \widehat{f}(\xi).$$

Therefore the Hilbert transform is acting essentially as a zero order operator. The presence of H in (2.2) is important to establish some monotonicity properties of the local mass of the solutions of (2.2), but it will not play an essential role in our discussion here.

There has been many works regarding the well-posedness of (2.2) (cf. [61, 1, 36, 60, 38, 48, 66]). The best result in the present moment is due to Tao.

Theorem 2.2 (cf. [66]). *For $s \geq 1$ the Cauchy problem (2.2) is well-posed for data in $H^s(\mathbb{R})$.*

One may ask whether, similarly to the KdV case, we also have the semi-linear well-posedness in Theorem 2.2. It turns out that the answer is negative.

Theorem 2.3 (cf. [50]). *In Theorem 2.2, one can not replace the well-posedness with semi-linear well-posedness.*

Therefore in the well-posedness analysis of (2.2), it is not a question to find a suitable space to perform the contraction method, simply this method for proving the well-posedness does not work, as far as the classical Sobolev spaces H^s are considered as a space for the initial data. This fact was first detected in [58].

A related to Theorem 2.3 result is obtained in [4] where it is shown the lack of semi-linear well-posedness for (2.2) with data in $H^s(\mathbb{R})$, $s < -1/2$.

Interestingly, the modified Benjamin-Ono equation

$$u_t + H u_{xx} + u^2 u_x = 0, \quad u(0) = u_0$$

turns out to be semi-linearly well-posed for data in $H^s(\mathbb{R})$, $s > 1/2$ (cf. [56]). Hence, even if the dispersion is the same, the semi-linear well-posedness may also be sensitive to the “degree” of the nonlinearity¹.

It is clear that Theorem 2.3 is a consequence of the following statement.

Theorem 2.4 (cf. [50]). *Let $s > 0$. There exist two positive constants c and C and two sequences (u_n) and (\tilde{u}_n) of solutions of the Benjamin-Ono equation such that for every $t \in [0, 1]$,*

$$\sup_n \|u_n(t, \cdot)\|_{H^s(\mathbb{R})} + \sup_n \|\tilde{u}_n(t, \cdot)\|_{H^s(\mathbb{R})} \leq C,$$

¹The example of KdV and Benjamin-Ono equations is an instance when the the semi-linear well-posedness depends on the degree of the dispersion.

(u_n) and (\tilde{u}_n) satisfy initially

$$\lim_{n \rightarrow \infty} \|u_n(0, \cdot) - \tilde{u}_n(0, \cdot)\|_{H^s(\mathbb{R})} = 0,$$

but, for every $t \in [0, 1]$,

$$\liminf_{n \rightarrow \infty} \|u_n(t, \cdot) - \tilde{u}_n(t, \cdot)\|_{H^s(\mathbb{R})} \geq c \sin t.$$

In the proof of Theorem 2.4, we will make use of the following well-posedness result for (2.2).

Proposition 2.5. *Fix $s \geq \sigma > 3/2$. Then for every $u_0 \in H^s(\mathbb{R})$ there exists a unique global solution $u \in C(\mathbb{R}; H^s(\mathbb{R}))$ of (2.2). Moreover*

$$\|u(t, \cdot)\|_{H^s(\mathbb{R})} \leq C \|u_0\|_{H^s(\mathbb{R})},$$

provided $|t| \leq c \|u_0\|_{H^\sigma(\mathbb{R})}^{-4}$.

Sketch of the proof. The proof of Proposition 2.5 is based on the compactness argument explained in the introduction in the context of the Burgers equation (1.3). One first proves the result for $s = \sigma$. The nature of the restriction $|t| \leq c \|u_0\|_{H^\sigma(\mathbb{R})}^{-4}$ is related to the scaling of (2.2). It turns out that one can reduce the matters to the problem of existence on the time interval $[0, 1]$ with initial data with small norm in $H^\sigma(\mathbb{R})$. Suppose that there exists a positive constant γ such that if the initial data of the Benjamin-Ono equation satisfies $\|u_0\|_{H^\sigma} \leq \gamma$ then we can find a unique solution on the time interval $[0, 1]$. We now prove that for $u_0 \in H^\sigma$ of arbitrary size we can solve (2.2) for time of order $\|u_0\|_{H^\sigma}^{-4}$. Indeed, given $u_0 \in H^\sigma$ we choose $\lambda \ll 1$ such that

$$(2.3) \quad 0 < \lambda^{\frac{1}{2}}(1 + \lambda^\sigma) \|u_0\|_{H^\sigma} \ll \gamma.$$

Set $\tilde{u}_0(x) := \lambda u_0(\lambda x)$. Then due to (2.3), $\|\tilde{u}_0\|_{H^\sigma} \leq \gamma$ and we can apply our assumption to \tilde{u}_0 . Let $\tilde{u}(t, x)$ be the solution of the Benjamin-Ono equation with data \tilde{u}_0 up to time one. Then one can easily verify that $u(t, x) := \lambda^{-1} \tilde{u}(\lambda^{-2}t, \lambda^{-1}x)$ is a solution of the Benjamin-Ono equation up to time λ^2 which in view of (2.3) is of order $\|u_0\|_{H^\sigma}^{-4}$. Hence we may reduce the matters to the existence on $[0, 1]$ for small data.

Let u be a sufficiently smooth in the scale of the Sobolev spaces solution of the Benjamin-Ono equation (2.2). Then, as in the case of the Burgers equation, one gets the bound

$$(2.4) \quad \|D^\sigma u\|_{L^\infty([0, T]; L^2)} \leq \|u(0)\|_{H^\sigma} \exp(C \|u_x\|_{L^1([0, T]; L^\infty)}).$$

Notice that the key quantity $\int_0^T \|u_x(t)\|_{L^\infty} dt$ is invariant with respect to the scaling of the equation. More precisely if $u(t, x) = \lambda^{-1} \tilde{u}(\lambda^{-2}t, \lambda^{-1}x)$ then

$$\int_0^{\lambda^2} \|u_x(t)\|_{L^\infty} dt = \int_0^1 \|\tilde{u}_x(t)\|_{L^\infty} dt.$$

With

$$F(T) := \|u_x\|_{L_T^1 L^\infty} + \|D^\sigma u\|_{L_T^\infty L^2}, \quad T \in [0, 1]$$

we can deduce from (2.4) and the Sobolev inequality (here we use that $\sigma > 3/2$) that

$$F(T) \leq C \|u(0)\|_{H^\sigma} \exp(cF(T)).$$

Now a straightforward continuity argument shows that there exist positive constants γ and C such that if $\|u(0)\|_{H^\sigma} \leq \gamma$ (and hence $F(0) \leq \gamma$) then $F(1) \leq C$, and in particular

$$(2.5) \quad \int_0^1 \|u_x(t)\|_{L^\infty} dt \leq C$$

Using (2.5) and (2.4) (with $T = 1$) we obtain that if u is a smooth solution of the Benjamin-Ono equation, then

$$(2.6) \quad \|D^\sigma u\|_{L^\infty([0,1]; L^2(\mathbb{R}))} \leq C \|u(0)\|_{H^s},$$

provided $\|u(0)\|_{H^s} \leq \gamma$. Moreover the solution satisfies (2.5). The bounds (2.5) and (2.6) enable one to perform a standard compactness argument for the proof of the existence. As for the Burgers equation, the uniqueness follows from the Gronwall lemma, the assumption $\sigma > 3/2$ and the Sobolev embedding. Let us next show the bound for the higher Sobolev norms. Let $s > \sigma$. Then we clearly have an analog of (2.4) on the H^s level. Namely,

$$\|D^s u\|_{L^\infty([0,1]; L^2)} \leq \|u(0)\|_{H^s} \exp\left(c \int_0^1 \|u_x(t)\|_{L^\infty} dt\right) \leq C \|u(0)\|_{H^s},$$

where in the last inequality, we used (2.5). Finally the global well-posedness follows from the conservation laws enjoyed by the solutions of the (2.2). Indeed one has controls (cf. e.g. [1]) on $\|u(t, \cdot)\|_{H^{k/2}}$ for $k = 0, 1, 2, \dots$. Hence the assertion of global existence is straightforward for $s \geq 2$. For $s < 2$ one may use the $H^{3/2}$ well-posedness result of Theorem 2.2 and the $H^{3/2}$ control. This completes the discussion on the proof of Proposition 2.5. \square

Next, we pick a usual bump function $\varphi \in C_0^\infty(\mathbb{R})$ such that $\varphi(x) = 1$ for $|x| < 1$ and $\varphi(x) = 0$ for $|x| > 2$. Let $\tilde{\varphi} \in C_0^\infty(\mathbb{R})$ be equal to one on the support of φ . Notice

that $\varphi\tilde{\varphi} = \varphi$. For $0 < \delta < 1$, we set

$$(2.7) \quad \varphi_\lambda(x) := \varphi\left(\frac{x}{\lambda^{1+\delta}}\right), \quad \tilde{\varphi}_\lambda(x) := \tilde{\varphi}\left(\frac{x}{\lambda^{1+\delta}}\right).$$

The assertion of Theorem 2.4 is a corollary of the following statement.

Theorem 2.6. *Let $\max(1-s, 0) < \delta < 1$ and $|\omega| \ll \lambda^{\frac{1-\delta}{2}}$. Let $u_{\omega,\lambda}$ be the unique global solution of the Benjamin-Ono equation subject to initial data*

$$u_{\omega,\lambda}(0, x) = -\omega \lambda^{-1} \tilde{\varphi}_\lambda(x) - \lambda^{-\frac{1}{2}-\frac{\delta}{2}-s} \varphi_\lambda(x) \cos \lambda x.$$

Then the identity

$$u_{\omega,\lambda}(t, x) = -\lambda^{-\frac{1}{2}-\frac{\delta}{2}-s} \varphi_\lambda(x) \cos(-\lambda^2 t + \lambda x + \omega t) + \mathcal{O}\left(\lambda^{-\frac{\min\{\delta, 1-\delta\}}{4(s+2)}} + |\omega| \lambda^{-\frac{1-\delta}{2}}\right)$$

holds in $H_x^s(\mathbb{R})$, uniformly in $t \in [0, 1]$.

Let us notice that if $\omega = 0$, the solution propagates as a high frequency linear Benjamin-Ono wave while when $\omega \neq 0$, the solution propagates as a high frequency linear dispersive wave with modified propagation speed which is the crucial nonlinear effect.

Let us now show why Theorem 2.6 implies Theorem 2.4. Apply Theorem 2.6 with $\omega = \pm 1$ and $\lambda = 1, 2, \dots$. We thus obtain two families $(u_{1,\lambda})$ and $(u_{-1,\lambda})$ of solutions to the Benjamin-Ono equation. Notice that

$$\|u_{1,\lambda}(0, \cdot) - u_{-1,\lambda}(0, \cdot)\|_{H^s} \leq C \lambda^{-\frac{1-\delta}{2}}$$

and moreover due to Theorem 2.6, setting $\kappa = -\lambda^2 t + \lambda x$, we arrive at

$$\|u_{1,\lambda}(t, \cdot) - u_{-1,\lambda}(t, \cdot)\|_{H^s} = \|\lambda^{-(\frac{1+\delta}{2}+s)} \varphi_\lambda(x) (\cos(\kappa + t) - \cos(\kappa - t))\|_{H_x^s} + o(1),$$

if $t \in [0, 1]$ and where $o(1) \rightarrow 0$ as $\lambda \rightarrow \infty$. At this point we need the following elementary lemma whose proof will be omitted.

Lemma 2.7. *Fix $s \geq 0$, $0 < \delta < 1$, $\alpha \in \mathbb{R}$ and $\varphi \in C_0^\infty(\mathbb{R})$. Then*

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\frac{1+\delta}{2}-s} \|\varphi_\lambda(x) \cos(\lambda x + \alpha)\|_{H_x^s} = \frac{1}{\sqrt{2}} \|\varphi\|_{L^2},$$

where φ_λ is defined by (2.7).

Using Lemma 2.7, we get

$$\lim_{\lambda \rightarrow \infty} \|\lambda^{-(\frac{1+\delta}{2}+s)} \varphi_\lambda(x) (\cos(\kappa + t) - \cos(\kappa - t))\|_{H_x^s} = \sqrt{2} |\sin t| \|\varphi\|_{L^2}.$$

Therefore Theorem 2.6 implies Theorem 2.4.

Proof of Theorem 2.6. Let $u_{low}(t, x)$ be the solution of (2.2) with initial data

$$(2.8) \quad u_{low}(0, x) = -\omega \lambda^{-1} \tilde{\varphi}_\lambda(x), \quad 0 < \delta < 1, \quad \omega \in \mathbb{R}.$$

In the next lemma, we collect several bounds for $u_{low}(t, x)$.

Lemma 2.8. *Let $k \geq 0$. Then the following estimates hold :*

$$(2.9) \quad \|\partial_x^k u_{low}(t, \cdot)\|_{L^2(\mathbb{R})} \leq C|\omega| \lambda^{-\frac{1-\delta}{2}-k(1+\delta)},$$

$$(2.10) \quad \|\partial_x u_{low}(t, \cdot)\|_{L^\infty(\mathbb{R})} \leq C|\omega| \lambda^{-2-\delta},$$

$$(2.11) \quad \|u_{low}(t, \cdot) - u_{low}(0, \cdot)\|_{L^2(\mathbb{R})} \leq C|\omega| \lambda^{-2-\delta},$$

if $|t| \leq 1$ and $|\omega| \ll \lambda^{\frac{1-\delta}{2}}$.

Proof of Lemma 2.8. Rescale by setting

$$(2.12) \quad v(t, x) := \lambda^{1+\delta} u_{low}(\lambda^{2+2\delta} t, \lambda^{1+\delta} x).$$

Then v is again a solution of the Benjamin-Ono equation. Since $v(0, x) = -\omega \lambda^\delta \tilde{\varphi}(x)$ we obtain

$$\|v(0, \cdot)\|_{H^s} = |\omega| \lambda^\delta \|\tilde{\varphi}\|_{H^s}$$

and therefore by Proposition 2.5

$$(2.13) \quad \|v(t, \cdot)\|_{H^s} \leq C|\omega| \lambda^\delta,$$

if $|t| \lesssim |\omega|^{-4} \lambda^{-4\delta}$ and $s > 3/2$. But since the right hand-side of (2.13) does not depend on s , we conclude that (2.13) is valid for any real s . The Sobolev embedding and (2.13) now give

$$(2.14) \quad \|v_x(t, \cdot)\|_{L^\infty} \leq C|\omega| \lambda^\delta,$$

if $|t| \lesssim |\omega|^{-4} \lambda^{-4\delta}$.

Using (2.12) and the restriction on $|\omega|$, we deduce from (2.14) by scaling back that

$$(2.15) \quad \|\partial_x u_{low}(t, \cdot)\|_{L^\infty} \leq C|\omega| \lambda^{-2-\delta},$$

if $|t| \leq 1$ which is (2.10).

We now turn to the proof of (2.9) and (2.11). Differentiating (2.12) and using (2.13) (with $s = k$) yields

$$(2.16) \quad \|\partial_x^k u_{low}(t, \cdot)\|_{L^2} \leq C|\omega| \lambda^{-\frac{1-\delta}{2}-k(1+\delta)}, \quad k = 0, 1, 2, \dots$$

if $|t| \leq 1$. Estimate (2.16) is indeed (2.9). Next, using (2.15), (2.16) and the equation satisfied by u_{low} gives

$$\|\partial_t u_{low}(t, \cdot)\|_{L^2} \leq C(\|\partial_x^2 u_{low}(t, \cdot)\|_{L^2} + \|\partial_x u_{low}(t, \cdot)\|_{L^\infty} \|u_{low}(t, \cdot)\|_{L^2}) \leq C|\omega| \lambda^{-2-\delta},$$

if $|t| \leq 1$. We now observe that (2.11) can be deduced from the above bound via the fundamental theorem of calculus, applied to u_{low} in the time variable. This completes the proof of Lemma 2.8. \square

We now set for $\lambda \geq 1$, $0 < \delta < 1$ and $|\omega| \ll \lambda^{\frac{1-\delta}{2}}$,

$$(2.17) \quad u_{ap}(t, x) := u_{low}(t, x) - \lambda^{-\frac{1}{2}-\frac{\delta}{2}-s} \varphi_\lambda(x) \cos(-\lambda^2 t + \lambda x - \lambda t u_{low}(0, x)).$$

The above function is an approximate solution of (2.2) for $\lambda \gg 1$ and $s > 0$ as shows the next statement.

Lemma 2.9. *Let $s > 0$, $0 < \delta < 1$, $|\omega| \ll \lambda^{\frac{1-\delta}{2}}$ and $|t| \leq 1$. Set*

$$F := (\partial_t + H\partial_x^2)u_{ap} + u_{ap} \partial_x u_{ap}.$$

Then there exist positive constants C and λ_0 such that for $\lambda \geq \lambda_0$ one has

$$\|F(t, \cdot)\|_{L^2(\mathbb{R})} \leq C \left(\lambda^{-\delta-s} + \lambda^{\frac{1-\delta}{2}-2s} \right).$$

Proof. Set $\Phi := -\lambda^2 t + \lambda x + \omega t$. We observe that

$$u_{ap}(t, x) = u_{low}(t, x) - \lambda^{-\frac{1}{2}-\frac{\delta}{2}-s} \varphi_\lambda(x) \cos \Phi.$$

Furthermore, we define the high frequency part of u_{ap} by setting

$$u_h(t, x) := -\lambda^{-\frac{1}{2}-\frac{\delta}{2}-s} \varphi_\lambda(x) \cos \Phi.$$

Next, we can write

$$(\partial_t + H\partial_x^2)u_{ap} + u_{ap} \partial_x u_{ap} = F_1 + F_2 + F_3 + F_4 + F_5,$$

where

$$\begin{aligned} F_1 &:= (\partial_t + H\partial_x^2)u_{low} + u_{low} \partial_x u_{low} \\ F_2 &:= -\lambda^{-\frac{1+\delta}{2}-s} \cos \Phi \partial_x \left(u_{low} \varphi_\lambda \right) \\ F_3 &:= u_h \partial_x u_h \\ F_4 &:= -\lambda^{-\frac{1+\delta}{2}-s} \left[H\partial_x^2, \varphi_\lambda \right] \cos \Phi \\ F_5 &:= -\lambda^{-\frac{1+\delta}{2}-s} \varphi_\lambda (\partial_t + H\partial_x^2 + u_{low} \partial_x) \cos \Phi. \end{aligned}$$

Since u_{low} is a solution of the Benjamin-Ono equation, we deduce that $F_1 = 0$. Using that $\varphi_\lambda \tilde{\varphi}_\lambda = \varphi_\lambda$, we readily obtain that

$$F_5 = \lambda^{\frac{1-\delta}{2}-s} (u_{low}(t, x) - u_{low}(0, x)) \varphi_\lambda(x) \sin \Phi.$$

Using Lemma 2.8, we get

$$(2.18) \quad \|F_5(t, \cdot)\|_{L^2} \leq C \lambda^{\frac{1-\delta}{2}-s} |\omega| \lambda^{-2-\delta} \leq C \lambda^{-1-2\delta-s}.$$

It remains to bound F_2 , F_3 and F_4 . Let us expand F_4 as

$$(2.19) \quad F_4 = \lambda^{\frac{3-\delta}{2}-s} [H, \varphi_\lambda] \cos \Phi + 2\lambda^{-\frac{1+\delta}{2}-s-\delta} H((\varphi')_\lambda \sin \Phi) - \\ - \lambda^{-\frac{5}{2}-\frac{5\delta}{2}-s} H((\varphi'')_\lambda \cos \Phi),$$

where

$$(\varphi')_\lambda = \varphi' \left(\frac{x}{\lambda^{1+\delta}} \right), \quad (\varphi'')_\lambda = \varphi'' \left(\frac{x}{\lambda^{1+\delta}} \right).$$

The first term in the right hand-side of (2.19) is controlled in L^2 by the estimate

$$\|[H, \varphi_\lambda] \cos \Phi\|_{L^2} \leq C_N \lambda^{-N}$$

which follows easily from the definition of the Hilbert transform. The L^2 norm of the other terms in the right hand-side of (2.19) are readily estimated by $c\lambda^{-\delta-s}$. Therefore

$$(2.20) \quad \|F_4(t, \cdot)\|_{L^2} \leq C \lambda^{-\delta-s}.$$

Expanding $\partial_x u_h$, the L^2 norm of F_3 is controlled as follows

$$(2.21) \quad \|F_3(t, \cdot)\|_{L^2} \leq C \lambda^{-\frac{3}{2}-\frac{3\delta}{2}-2s} + C \lambda^{\frac{1}{2}-\frac{\delta}{2}-2s} \leq C \lambda^{\frac{1-\delta}{2}-2s}.$$

Next, using Lemma 2.8 and the assumption on $|\omega|$, we obtain

$$(2.22) \quad \|F_2(t, \cdot)\|_{L^2} \leq C(\lambda^{-s} \|\partial_x u_{low}(t, \cdot)\|_{L^\infty} + \lambda^{-\frac{3+3\delta}{2}-s} \|u_{low}(t, \cdot)\|_{L^2}) \leq \\ \leq C \lambda^{-\frac{3+3\delta}{2}-s}.$$

Collecting (2.18), (2.20), (2.21) and (2.22) completes the proof of Lemma 2.9. \square

Let us now finish the proof of Theorem 2.6. The first step is to bound $u_{\omega, \lambda}$ in high Sobolev norms. We distinguish two cases : $s > 3/2$ and $0 < s \leq 3/2$. In the second case we will need to exploit the higher conservation laws for the Benjamin-Ono equation while in the first case we use Proposition 2.5 instead.

Let $s > 3/2$. Observe that for $3/2 < \sigma < s$

$$\|u_{\omega, \lambda}(0, \cdot)\|_{H^\sigma} \leq C(\lambda^{\sigma-s} + |\omega| \lambda^{-\frac{1-\delta}{2}}).$$

Therefore for $k \geq s$, it follows from Proposition 2.5 that

$$(2.23) \quad \|u_{\omega, \lambda}(t, \cdot)\|_{H^k} \leq C \|u_{\omega, \lambda}(0, \cdot)\|_{H^k} \leq C \lambda^{k-s}, \quad |t| \leq 1.$$

Let $0 < s \leq \frac{3}{2}$. Using the conservation laws associated to the Benjamin-Ono equation (cf. [1, Lemma 3.3.2]), we get the following bound uniformly in $t \in \mathbb{R}$

$$(2.24) \quad \|u_{\omega, \lambda}(t, \cdot)\|_{H^2} \leq C(\|u_{\omega, \lambda}(0, \cdot)\|_{H^2} + \|u_{\omega, \lambda}(0, \cdot)\|_{L^2}^5) \leq C(1 + \lambda^{2-s}),$$

and therefore we obtain

$$(2.25) \quad \|u_{\omega,\lambda}(t, \cdot)\|_{H^2} \leq C\lambda^{2-s}, \quad t \in \mathbb{R}, \quad \lambda \geq 1.$$

Let u_{ap} be as in (2.17). Set

$$v_{\omega,\lambda} := u_{\omega,\lambda} - u_{ap}.$$

The aim is to show that $v_{\omega,\lambda}$ is small comparing to u_{ap} in the H^s norm.

Due to Lemma 2.8, we get

$$\|u_{low}(t, \cdot)\|_{H^s} \leq C|\omega|\lambda^{-\frac{1-\delta}{2}},$$

if $|t| \leq 1$. Next, using Lemma 2.7, we obtain the bound

$$\|u_{ap}(t, \cdot)\|_{H^k(\mathbb{R})} \leq C\lambda^{k-s},$$

if $|t| \leq 1$ and $k \geq s$.

Therefore using (2.23) and (2.25), we get the bounds for the high Sobolev norms

$$(2.26) \quad \|v_{\omega,\lambda}(t, \cdot)\|_{H^k} \leq C\lambda^{k-s},$$

if $|t| \leq 1$ and $3/2 < s < k$, and

$$(2.27) \quad \|v_{\omega,\lambda}(t, \cdot)\|_{H^2} \leq C\lambda^{2-s},$$

if $t \in \mathbb{R}$ and $0 < s < 3/2$.

Further, we prove a good bound of the L^2 norm of $v_{\omega,\lambda}$. Clearly

$$(2.28) \quad (\partial_t + H\partial_x^2)v_{\omega,\lambda} + v_{\omega,\lambda}\partial_x v_{\omega,\lambda} + \partial_x(u_{ap}v_{\omega,\lambda}) + F = 0, \quad v_{\omega,\lambda}(0, x) = 0$$

with

$$F = (\partial_t + H\partial_x^2)u_{ap} + u_{ap}\partial_x u_{ap},$$

which satisfies

$$\|F(t, \cdot)\|_{L^2} \leq C\lambda^{-\frac{\min\{\delta, 1-\delta\}}{2}-s}$$

by Lemma 2.9 and the assumption $1 - s < \delta < 1$.

The second endpoint in the bounds for $v_{\omega,\lambda}$ is the L^2 estimate

$$(2.29) \quad \|v_{\omega,\lambda}(t, \cdot)\|_{L^2} \leq C\lambda^{-\frac{\min\{\delta, 1-\delta\}}{2}-s}, \quad |t| \leq 1.$$

To prove (2.29), we multiply (2.28) by $v_{\omega,\lambda}$ and we integrate in x

$$\frac{d}{dt}\|v_{\omega,\lambda}(t, \cdot)\|_{L^2}^2 \leq C(\|\partial_x u_{ap}(t, \cdot)\|_{L^\infty}\|v_{\omega,\lambda}(t, \cdot)\|_{L^2}^2 + \|v_{\omega,\lambda}(t, \cdot)\|_{L^2}\|F(t, \cdot)\|_{L^2}).$$

Hence, since we have for $1 - s < \delta < 1$ and $\lambda \gg 1$,

$$\|\partial_x u_{ap}(t, \cdot)\|_{L^\infty} \leq C\|\partial_x u_{low}(t, \cdot)\|_{L^\infty} + C\lambda^{\frac{1-\delta}{2}-s} \leq C|\omega|\lambda^{-2-\delta} + C\lambda^{\frac{1-\delta}{2}-s} \ll 1,$$

we readily get the bound (2.29).

We now complete the proof by an interpolation argument. Let first $s > 3/2$. Choose $k \in [s + \frac{1}{2}, s + 2]$ and interpolate between (2.26) and (2.29) as follows

$$\|v_{\omega,\lambda}(t, \cdot)\|_{H^s} \leq \|v_{\omega,\lambda}(t, \cdot)\|_{L^2}^{\frac{k-s}{k}} \|v_{\omega,\lambda}(t, \cdot)\|_{H^k}^{\frac{s}{k}} \leq C\lambda^{-\frac{\min\{\delta, 1-\delta\}}{4(s+2)}}.$$

If $s \leq \frac{3}{2}$ we obtain the same estimate by using $k = 2$ in the interpolation and (2.27) instead of (2.26). This completes the proof of Theorem 2.6. \square

We end this section by a series of remarks.

The method of proof of Theorem 2.6 can be generalized to many other equations. For example the corresponding to Theorem 2.6 result in the context of the KdV equation provides a family of essentially linear KdV waves ($\omega \rightarrow 0$) as approximate solutions and thus no instability property of the flow is displayed.

The proof of Theorem 2.2 is based on a gauge transform reducing (2.2) to a problem which, despite the lack of semi-linear well-posedness displayed by Theorem 2.3, shares many features with a semi-linearly well-posed problem.

One may consider the higher dispersion versions of the (2.2)

$$(2.30) \quad u_t - Lu_x + uu_x = 0, \quad u(0) = u_0,$$

where L is Fourier multiplier with symbol $|\xi|^\gamma$, $1 \leq \gamma \leq 2$. The KdV equation corresponds to $\gamma = 2$, and, thanks to Theorem 2.1 in this case (2.30) is semi-linearly well-posed in $H^s(\mathbb{R})$, $s > -3/4$. On the other hand, in view of the result of [58], it seems reasonable to conjecture that for $1 < \gamma < 2$, the Cauchy problem (2.30) is not semi-linearly well-posed in *all* $H^s(\mathbb{R})$.

Another instance when the notion of semi-linear well-posedness is naturally involved is the analysis of the Cauchy problem for the Kadomtsev-Petviashvili equations. The Kadomtsev-Petviashvili (KP) equations are natural two dimensional generalizations of the KdV equation (cf. [37]). There are two KP equations, the KP-I equation

$$(2.31) \quad (u_t + u_{xxx} + uu_x)_x - u_{yy} = 0,$$

and the KP-II equation

$$(2.32) \quad (u_t + u_{xxx} + uu_x)_x + u_{yy} = 0.$$

It is known that the Cauchy problem for the KP-II equation (2.32) is semi-linearly well-posed in $H^s(\mathbb{R}^2)$, $s \geq 0$ and even in Sobolev type spaces of negative indices

(spaces of distributions), cf. [11, 63, 35]. On the other hand, in view of the result of [59], it seems reasonable to conjecture that the Cauchy problem for the KP-I equation (2.31) is not semi-linearly well-posed in all $H^s(\mathbb{R}^2)$.

3. NONLINEAR SCHRÖDINGER EQUATIONS (NLS)

3.1. Nonlinear Schrödinger equations on \mathbb{R}^d . Consider the Cauchy problem for the nonlinear Schrödinger equation

$$(3.1) \quad iu_t + \Delta u + |u|^2 u = 0, \quad u(0) = u_0,$$

posed on the Euclidean space \mathbb{R}^d , $d \geq 1$. Equation (3.1) is a focusing model. The defocusing model

$$(3.2) \quad iu_t + \Delta u - |u|^2 u = 0$$

is also of interest. The long time dynamics of (3.1) and (3.2) are quite different. But for our discussion here (small time analysis) it will be relevant to concentrate only on (3.1). There has been a large number of articles studying (3.1), (3.2) and their generalizations, when $|u|^2 u$ (which is the term involved in many applications) is replaced by a more general nonlinear term $f(|u|^2)u$ (cf. [29, 30, 31, 41, 42, 23, 67, 68] ...). The equation (3.1) is an infinite dimensional Hamiltonian equation with canonical coordinates (u, \bar{u}) and Hamiltonian

$$H(u, \bar{u}) = \frac{1}{2} \int_{\mathbb{R}^d} |\nabla u|^2 - \frac{1}{4} \int_{\mathbb{R}^d} |u|^4.$$

The Hamiltonian is formally preserved by the flow of (3.1). So is the L^2 norm of u . Therefore the space $H^1(\mathbb{R}^d)$ is a natural phase space² for (3.1) and (3.2) at least for $d \leq 4$ when the second term of the Hamiltonian is dominated by the first one and the L^2 norm of u . Fortunately, we can achieve this regularity for $d \leq 3$ in the context of the well-posedness theory of (3.1). More precisely, we have the following result regarding the well-posedness of (3.1).

Theorem 3.1 (cf. [23]). *Let $s > \frac{d-2}{2}$, $d \geq 2$. Then the Cauchy problem (3.1) is semi-linearly well-posed for data in $H^s(\mathbb{R}^d)$.*

Proof. We will give the proof because it is “typical” for a semi-linearly well-posed problem. It is worth noticing that such a proof is indeed quite different from the reasoning in the proof of Proposition 2.5 above. To simplify a little the notations we will only consider the case $d = 2$, the proof in higher dimensions being very similar. The proof is based on the following Strichartz inequality for the free evolution.

²In this space the well-posedness of (3.2) is actually global in time. For (3.1) the well-posedness is global as far as the data is small in a suitable sense, for large data solutions developing singularities in finite time appear.

Proposition 3.2. *Let (p, q) such that, $\frac{1}{p} + \frac{1}{q} = \frac{1}{2}$, $p > 2$. Then there exists a constant $C > 0$ such that for every $T > 0$, every $u_0 \in L^2(\mathbb{R}^2)$,*

$$\|e^{it\Delta}u_0\|_{L^p([0, T]; L^q(\mathbb{R}^2))} \leq C\|u_0\|_{L^2(\mathbb{R}^2)}.$$

Proof. The proof of Proposition 3.2 can be found in [22]. \square

Let us now show how Proposition 3.2 implies Theorem 3.1. Consider the integral equation corresponding to (3.1)

$$(3.3) \quad u(t) = e^{it\Delta}u_0 + i \int_0^t e^{i(t-\tau)\Delta}(|u(\tau)|^2u(\tau))d\tau.$$

Let us fix a real number σ satisfying

$$0 < \sigma < \min \left\{ s, \frac{1}{2} \right\}.$$

The value of σ being fixed, we define $q \in [2, 4[$ by the identity

$$\frac{1}{q} - \frac{1}{4} = \frac{\sigma}{2}.$$

Next, we define p such that

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{2}.$$

Set

$$X_T = L^\infty([0, T]; H^s(\mathbb{R}^2)) \cap Y_T \cap Z_T,$$

where Y_T, Z_T are equipped with the norms

$$\|u\|_{Y_T}^2 = \sum_{N\text{-dyadic}} N^{2s} \|\Delta_N(u)\|_{L_T^p L^q(\mathbb{R}^2)}^2, \quad \|u\|_{Z_T}^2 = \sum_{N\text{-dyadic}} N^{2s} \|\Delta_N(u)\|_{L_T^4 L^4(\mathbb{R}^2)}^2.$$

The sums over N are running over all dyadic values of N , i.e $N = 2^n$, $n \geq 0$ and

$$u = \sum_N \Delta_N(u)$$

is a Littlewood-Paley decomposition³ of u . More precisely, Δ_N are the Fourier multipliers defined by

$$\widehat{\Delta_N(u)}(\xi) = \varphi(N^{-1}\xi)\hat{u}(\xi), \quad N = 2^n, \quad n \geq 1$$

and

$$\widehat{\Delta_1(u)}(\xi) = \psi(\xi)\hat{u}(\xi)$$

³Actually we do not need to use the precise form of Δ_N here, cut-off projectors would also make work the argument.

with $\varphi \in C_0^\infty([1, 2[)$ and

$$\psi(\xi) + \sum_{n \geq 1} \varphi(2^{-n}\xi) = 1.$$

Notice that if $u \in \mathcal{S}'(\mathbb{R}^2)$ then $\Delta_N(u)$ is localized at frequencies of order N . It is also useful to see the norm in Y_T as $\|N^s \Delta_N(u)\|_{l_N^2 L_T^p L^q}$.

Next, using Proposition 3.2, we get

$$\|e^{it\Delta} \Delta_N(u_0)\|_{L_T^p L^q(\mathbb{R}^2)} + \|e^{it\Delta} \Delta_N(u_0)\|_{L_T^4 L^4(\mathbb{R}^2)} \leq C \|\Delta_N(u_0)\|_{L^2(\mathbb{R}^2)}$$

and therefore

$$(3.4) \quad \|e^{it\Delta} u_0\|_{X_T} \leq C \|u_0\|_{H^s}.$$

Similarly, using the Minkowski inequality, we get the bound,

$$(3.5) \quad \left\| \int_0^t e^{i(t-\tau)\Delta} (|u(\tau)|^2 u(\tau)) d\tau \right\|_{X_T} \leq C \int_0^T \left\| e^{-i\tau\Delta} (|u(\tau)|^2 u(\tau)) \right\|_{H^s} d\tau.$$

In order to bound the right hand-side of (3.5), we will show that there exists $C > 0$ such that for every u_1, u_2, u_3 in X_T ,

$$(3.6) \quad \int_0^T \left\| e^{-i\tau\Delta} (u_1(\tau) \overline{u_2(\tau)} u_3(\tau)) \right\|_{H^s} d\tau \leq CT^\sigma \prod_{j=1}^3 \|u_j\|_{X_T}.$$

By duality, to show (3.6), it suffices to obtain that for every $w \in H^{-s}(\mathbb{R}^2)$,

$$(3.7) \quad \int_0^T \int_{\mathbb{R}^2} (u_1(\tau) \overline{u_2(\tau)} u_3(\tau)) (e^{i\tau\Delta} \overline{w}) d\tau \leq CT^\sigma \|w\|_{H^{-s}} \prod_{j=1}^3 \|u_j\|_{X_T}.$$

Notice that if $u_1(\tau), \overline{u_2(\tau)}, u_3(\tau)$ are localized at frequencies N_1, N_2, N_3 respectively then only frequencies of order $\leq C(N_1 + N_2 + N_3)$ of $e^{i\tau\Delta} \overline{w}$ contribute to the left hand-side of (3.7). Writing down the Littlewood-Paley decompositions of $u_1(\tau), \overline{u_2(\tau)}, u_3(\tau)$ et w , using the Hölder inequality and Proposition 3.2 to bound $e^{i\tau\Delta} \overline{w}$, we deduce that we can bound the left hand-side of (3.7) by

$$(3.8) \quad \sum_{N \leq C(N_1 + N_2 + N_3)} \|\Delta_{N_1}(u_1)\|_{L_T^4 L^4} \|\Delta_{N_2}(u_2)\|_{L_T^4 L^4} \|\Delta_{N_3}(u_3)\|_{L_T^4 L^4} \|\Delta_N(w)\|_{L^2}$$

By symmetry, we can suppose that in (3.8) the summation is restricted to

$$N_1 \geq N_2 \geq N_3.$$

Next, using the Sobolev embedding $W^{\sigma,q}(\mathbb{R}^2) \subset L^4(\mathbb{R}^2)$, and, the Hölder inequality in the time variable, we get the bound

$$\|\Delta_{N_2}(u_2)\|_{L_T^4 L^4} \|\Delta_{N_3}(u_2)\|_{L_T^4 L^4} \leq CT^\sigma (N_2 N_3)^\sigma \|\Delta_{N_2}(u_2)\|_{L_T^p L^q} \|\Delta_{N_3}(u_3)\|_{L_T^p L^q}.$$

Set

$$c_1(N) = N^s \|\Delta_{N_1}(u_1)\|_{L_T^4 L^4}, \quad c_j(N) = N^s \|\Delta_N(u_j)\|_{L_T^p L^q}, \quad j = 2, 3,$$

and $d(N) = N^{-s} \|\Delta_N(w)\|_{L^2}$. We obtain that (3.8) is bounded by

$$(3.9) \quad \sum_{N \leq CN_1} \sum_{N_2, N_3} \left(\frac{N}{N_1}\right)^s \frac{CT^\sigma}{(N_2 N_3)^{s-\sigma}} c_1(N_1) c_2(N_2) c_3(N_3) d(N).$$

Summing geometric series in N_2, N_3 , we obtain that (3.9) is bounded by

$$(3.10) \quad CT^\sigma \|u_2\|_{X_T} \|u_3\|_{X_T} \sum_{N \leq CN_1} \left(\frac{N}{N_1}\right)^s c_1(N_1) d(N).$$

To bound (3.10), we use the following lemma.

Lemma 3.3. *For every $\Lambda > 0$, every $s > 0$ there exists $C > 0$ such that if (c_{N_0}) and (d_{N_1}) are two sequences of nonnegative numbers indexed by the dyadic integers, then,*

$$\sum_{N_0 \leq \Lambda N_1} \left(\frac{N}{N_1}\right)^s c_{N_0} d_{N_1} \leq C \left(\sum_{N_0} c_{N_0}^2\right)^{\frac{1}{2}} \left(\sum_{N_1} d_{N_1}^2\right)^{\frac{1}{2}}.$$

Proof. Let us set

$$K(N_0, N_1) := \mathbf{1}_{N_0 \leq \Lambda N_1} \frac{N_0^s}{N_1^s}.$$

Summing geometric series imply that there exists $C > 0$ such that

$$\sup_{N_0} \sum_{N_1} K(N_0, N_1) + \sup_{N_1} \sum_{N_0} K(N_0, N_1) \leq C.$$

Therefore the Schur lemma implies the boundedness on $l_{N_0}^2 \times l_{N_1}^2$ of the bilinear form with kernel $K(N_0, N_1)$. This completes the proof of Lemma 3.3. \square

Using Lemma 3.3, we bound (3.10) by the right hand-side of (3.7) which completes the proof of (3.6).

Estimates (3.4), (3.5) et (3.6) yield that for every bounded set B of $H^s(\mathbb{R}^2)$ there exists $T > 0$ such that for every $u_0 \in B$ the right hand-side of (3.3) is a contraction in a suitable ball of X_T .

Let us finally explain how we obtain the propagation of regularity property for data in $H^{\tilde{s}}$, $\tilde{s} \geq s$. Denote by $X_T^{\tilde{s}}$ the space X_T use above associated to the $H^{\tilde{s}}$

regularity. It is easy to observe that the preceding analysis also gives the tame estimate

$$\left\| \int_0^t e^{i(t-\tau)\Delta} (|u(\tau)|^2 u(\tau)) d\tau \right\|_{X_T^{\frac{3}{2}}} \leq CT^\sigma \|u\|_{X_T^s}^2 \|u\|_{X_T^{\frac{3}{2}}}$$

which implies the propagation of the $H^{\frac{3}{2}}$ regularity in a straightforward way. This ends the discussion on the proof of Theorem 3.1. \square

The indice $\frac{d-2}{2}$ appeared in Theorem 3.1 is closely related to the scaling of the equation (3.1). More precisely if $u(t, x)$ solves (3.1) then so does

$$u_\lambda(t, x) := \lambda u(\lambda^2 t, \lambda x).$$

The norm of u_λ in the homogeneous Sobolev \dot{H}^s is independent of λ only for $s = \frac{d-2}{2}$.

At this point, it is worth noticing that the scaling invariance is responsible for the existence of solutions of (3.1) which concentrate in a point. Such kind of concentrations may give ill-posedness results only below the scaling norm. As we will see later concentration on higher dimensional objects as curves are responsible for ill-posedness above the scaling exponent.

It turns out that the result of Theorem 3.1 is essentially sharp, i.e. the point concentration coming from of the scaling invariance are the worst possible.

Theorem 3.4 (cf. [25]). *Let $d \geq 2$. Then :*

1. *For $d = 2$, the Cauchy problem (3.1) is not semi-linearly well-posed for data in $H^s(\mathbb{R}^d)$, $s < 0$ ($= \frac{d-2}{2}$). Moreover, it is not well-posed for data in $H^s(\mathbb{R}^d)$, $s \leq -1$ ($= -\frac{d}{2}$).*
2. *For $d \geq 3$, the Cauchy problem (3.1) is not well-posed for data in $H^s(\mathbb{R}^d)$, $0 < s < \frac{d-2}{2}$ or $s \leq -\frac{d}{2}$. Moreover, it is not semi-linearly well-posed for data in $H^s(\mathbb{R}^d)$, $-\frac{d}{2} \leq s \leq 0$.*

Proof. In order to simplify the exposition, we will give the proof of Theorem 3.4, for $d \geq 5$ and s a positive integer. This will cover the most interesting case $s = 1$, i.e. the ill-posedness of (3.1) in the “energy space” $H^1(\mathbb{R}^d)$, $d \geq 5$. The proof of Theorem 3.4 in the other cases has a very similar flavor.

Let us first observe that it suffices to prove the following statement.

Proposition 3.5. *Let $d \geq 5$ and $s \in]0, \frac{d-2}{2}[$ be a positive integer. Then there exist a sequence (t_n) of positive numbers tending to zero and a sequence $(u_n(t))$ of $C^\infty(\mathbb{R}^d) \cap H^s(\mathbb{R}^d)$ functions defined for $t \in [0, t_n]$, such that*

$$(i\partial_t + \Delta)u_n + |u_n|^2 u_n = 0$$

with

$$\lim_{n \rightarrow \infty} \|u_n(0, \cdot)\|_{H^s(\mathbb{R}^d)} = 0,$$

and

$$\lim_{n \rightarrow \infty} \|u_n(t_n, \cdot)\|_{H^s(\mathbb{R}^d)} = \infty.$$

Proof. Let us consider an initial data concentrating in the point $x = 0$

$$u_n(0, x) := \kappa_n n^{\frac{d}{2}-s} \varphi(nx), \quad n \gg 1,$$

where φ is a non identically zero smooth compactly supported function and

$$\kappa_n = \log^{-\delta_1}(n)$$

with $\delta_1 > 0$ to be fixed later. Remark that

$$\|u_n(0, \cdot)\|_{H^s(\mathbb{R}^d)} \sim \kappa_n.$$

The function

$$v_n(t, x) = \kappa_n n^{\frac{d}{2}-s} \varphi(nx) e^{it[\kappa_n n^{\frac{d}{2}-s} \varphi(nx)]^2}$$

is the solution of the equation

$$(3.11) \quad i\partial_t v_n + |v_n|^2 v_n = 0, \quad v_n(0, x) = u_n(0, x).$$

It turns out that for very small times v_n is near the actual solutions⁴ of (3.1).

Next, for a fixed integer $l > d/2$, we define quantity,

$$E_n(u) := \left(n^{2s} \|u\|_{L^2(\mathbb{R}^d)}^2 + n^{-2(l-s)} \|u\|_{H^l(\mathbb{R}^d)}^2 \right)^{\frac{1}{2}}$$

which can be seen as a semi-classical energy of u . Notice that, uniformly in n ,

$$(3.12) \quad \|u\|_{H^s} \leq C E_n(u).$$

The main point in the proof of Proposition 3.5 is the next statement.

Lemma 3.6. *Fix $\delta_2 \in \mathbb{R}$ such that*

$$0 < \delta_2 < \frac{1}{l+1}.$$

Then the solution u_n of (3.1) with initial data

$$u_0(x) = \kappa_n n^{\frac{d}{2}-s} \varphi(nx)$$

exists for $0 \leq t \leq t_n$, where

$$t_n = \log^{\delta_2}(n) n^{-2(\frac{d}{2}-s)}.$$

⁴A similar idea was used, in a different context, by Kuksin [51].

Moreover, there exists $\varepsilon > 0$ such that for $t \in [0, t_n]$,

$$E_n(u_n(t) - v_n(t)) \leq Cn^{-\varepsilon}.$$

Proof. Since the initial data are in H^l , $l > d/2$, we know that $u_n(t)$ exist on small time interval $[0, \tilde{t}_n]$. Consequently, to prove Lemma 3.6, we simply prove the *a priori* estimates which ensure, by a classical bootstrap argument, both the existence and the control on $E_n(u_n(t) - v_n(t))$ for $t \in [0, t_n]$. Let us set

$$w_n := u_n - v_n.$$

The *a priori* estimates involved in the proof are energy inequalities in the equation satisfied by w_n ,

$$\begin{aligned} (i\partial_t + \Delta)w_n &= -\Delta v_n - v_n^2 \bar{w}_n - 2|v_n|^2 w_n - 2v_n |w_n|^2 - \bar{v}_n w_n^2 - |w_n|^2 w_n \\ &:= -\Delta v_n + \Lambda(v_n, w_n). \end{aligned}$$

Using the explicit formula for v_n , we have that for $0 \leq t \leq t_n$,

$$(3.13) \quad \|v_n(t, \cdot)\|_{L^\infty(\mathbb{R}^d)} \leq Cn^{\frac{d}{2}-s}$$

and for $\sigma \geq 0$, $0 \leq t \leq t_n$,

$$(3.14) \quad \|v_n(t, \cdot)\|_{H^\sigma(\mathbb{R}^d)} \leq Cn^{\sigma-s} \log^{\delta_{2\sigma}}(n).$$

Let us now estimate $E_n(\Lambda(v_n(t), w_n(t)))$ for $0 \leq t \leq t_n$. Using the Gagliardo-Nirenberg inequality

$$\|f\|_{L^\infty(\mathbb{R}^d)} \leq C\|f\|_{L^2(\mathbb{R}^d)}^{1-\frac{d}{2l}}\|f\|_{H^l(\mathbb{R}^d)}^{\frac{d}{2l}}, \quad l > d/2,$$

we infer that

$$(3.15) \quad \|f\|_{L^\infty(\mathbb{R}^d)} \leq Cn^{\frac{d}{2}-s} E_n(f).$$

Coming back to the expression for $\Lambda(v_n, w_n)$, we get

$$\|\Lambda(v_n, w_n)\|_{L^2} \leq C(\|v_n\|_{L^\infty}^2 + \|w_n\|_{L^\infty}^2)\|w_n\|_{L^2}.$$

Using (3.13) and (3.15), we obtain that for $0 \leq t \leq t_n$,

$$n^s \|\Lambda(v_n(t), w_n(t))\|_{L^2} \leq Cn^{2(\frac{d}{2}-s)} (E_n(w_n(t)) + E_n^3(w_n(t))).$$

Next, using several times the classical bilinear inequality

$$\|fg\|_{H^l} \leq C(\|f\|_{L^\infty}\|g\|_{H^l} + \|g\|_{L^\infty}\|f\|_{H^l}),$$

we deduce that

$$\begin{aligned} \|\Lambda(v_n, w_n)\|_{H^l} &\leq C(\|v_n\|_{L^\infty}^2\|w_n\|_{H^l} + \|v_n\|_{L^\infty}\|v_n\|_{H^l}\|w_n\|_{L^\infty} + \\ &\quad + \|v_n\|_{H^l}\|w_n\|_{L^\infty}^2 + \|v_n\|_{L^\infty}\|w_n\|_{L^\infty}\|w_n\|_{H^l} + \|w_n\|_{L^\infty}^2\|w_n\|_{H^l}). \end{aligned}$$

Using (3.13), (3.14) and (3.15), we infer that for $0 \leq t \leq t_n$,

$$n^{-(l-s)} \|\Lambda(v_n(t), w_n(t))\|_{H^l} \leq C n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n) (E_n(w_n(t)) + E_n^3(w_n(t))).$$

Summarizing the above discussion yields that for $0 \leq t \leq t_n$,

$$E_n(\Lambda(v_n(t), w_n(t))) \leq C n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n) (E_n(w_n(t)) + E_n^3(w_n(t))).$$

Next, we estimate the source term $-\Delta v_n$. Using (3.14), we obtain that for $0 \leq t \leq t_n$,

$$\|\Delta v_n(t, \cdot)\|_{L^2} \leq C \|v_n(t, \cdot)\|_{H^2} \leq C n^{2-s} \log^{2\delta_2}(n).$$

Similarly,

$$\|\Delta v_n(t, \cdot)\|_{H^l} \leq C \|v_n(t, \cdot)\|_{H^{l+2}} \leq C n^{l+2-s} \log^{(l+2)\delta_2}(n).$$

Therefore, for $0 \leq t \leq t_n$,

$$E_n(\Delta v_n(t)) \leq C n^2 \log^{(l+2)\delta_2}(n).$$

Coming back to the equation solved by w_n , we deduce that for $0 \leq t \leq t_n$,

$$\begin{aligned} \frac{d}{dt} E_n^2(w_n(t)) &\leq C n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n) (E_n^2(w_n(t)) + E_n^4(w_n(t))) + \\ &\quad + C n^2 \log^{(l+2)\delta_2}(n) E_n(w_n(t)). \end{aligned}$$

Suppose first that $E_n(w_n(t)) \leq 1$ which is clearly the case at least for $t \ll 1$ since $w_n(0, x) = 0$. Using the elementary inequality

$$2n^2 \log^{(l+2)\delta_2}(n) E_n(w_n) \leq n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n) E_n^2(w_n) + n^{4-2(\frac{d}{2}-s)} \log^{(l+4)\delta_2}(n)$$

we obtain that for $0 \leq t \leq t_n$,

$$\frac{d}{dt} \left[e^{-C t n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n)} E_n^2(w_n(t)) \right] \leq C n^{4-2(\frac{d}{2}-s)} \log^{(l+4)\delta_2}(n) e^{-C t n^{2(\frac{d}{2}-s)} \log^{\delta_2 l}(n)}.$$

Integrating between 0 and t yields,

$$E_n(w_n(t)) \leq C n^{2-2(\frac{d}{2}-s)} \log^{2\delta_2}(n) e^{C \log^{(l+1)\delta_2}(n)}.$$

The assumption $s < (d-2)/2$ implies $2 - 2(\frac{d}{2} - s) < 0$. Moreover δ_2 is such that $(l+1)\delta_2 < 1$. Therefore there exists $\varepsilon > 0$ such that

$$E_n(w_n(t)) \leq C n^{-\varepsilon}.$$

Finally, a bootstrap argument allows to drop the assumption

$$E_n(w_n(t)) \leq 1.$$

This completes the proof of Lemma 3.6. \square

Let us now finish the proof of Proposition 3.5. We need to make a proper choice of the number δ_1 involved in the definition of κ_n . Using the explicit formula for v_n , we easily obtain that for $s \in \mathbb{Z}^+$,

$$\|v_n(t_n, \cdot)\|_{H^s} \geq C\kappa_n (t_n [\kappa_n n^{\frac{d}{2}-s}]^2)^s,$$

provided $t_n [\kappa_n n^{\frac{d}{2}-s}]^2 \gg 1$, i.e. $\log^{\delta_2 - 2\delta_1}(n) \gg 1$. The first assumption on δ_1 is thus

$$0 < \delta_1 < \frac{\delta_2}{2}.$$

Therefore, using Lemma 3.6 and (3.12), we obtain that for $n \gg 1$,

$$\begin{aligned} \|u_n(t_n, \cdot)\|_{H^s} &\geq C\|v_n(t_n, \cdot)\|_{H^s} - Cn^{-\varepsilon} \geq C\kappa_n (t_n [\kappa_n n^{\frac{d}{2}-s}]^2)^s - Cn^{-\varepsilon} = \\ &= C\log^{s\delta_2 - (1+2s)\delta_1}(n) - Cn^{-\varepsilon}. \end{aligned}$$

The proof of Proposition 3.5 is completed by choosing $\delta_1 \in \mathbb{R}$ such that

$$0 < \delta_1 < \frac{s\delta_2}{1+2s}.$$

□

The ansatz with v_n as an approximate solution still holds for $-\frac{d}{2} < s \leq 0$ and very small times of order $\sim \log^\delta(n)n^{-2(\frac{d}{2}-s)}$ with a suitable $\delta > 0$. Unfortunately, we can no longer bound from below $\|v_n(t_n, \cdot)\|_{H^s}$ as above. For that reason one can not get the failure of well-posedness for data in $H^s(\mathbb{R}^d)$, $-\frac{d}{2} < s \leq 0$. One can however still obtain the lack of semi-linear well-posedness, by an argument very similar to the one that we presented in the context of the Benjamin-Ono equation. This ends the discussion on the proof of Theorem 3.4. □

Remark 3.7. *The approach to Theorem 3.4 we present here may seem more involved than in [25] but it has the advantage to avoid the scaling considerations of [25]. In particular it works for variable coefficients second order operators instead of Δ or for (3.1) posed on a curved space.*

Let us next consider (2.1) posed on the real line \mathbb{R} . In this case the critical threshold for the semi-linear well-posedness is shifted with respect to the scaling regularity.

Theorem 3.8 (cf. [23, 24, 67]). *For $d = 1$ the Cauchy problem (3.1) is semi-linearly well-posed for data in $H^s(\mathbb{R})$, $s \geq 0$, and, it is not semi-linearly well-posed for data in $H^s(\mathbb{R})$, $s < 0$.*

It is worth noticing that in the proof of the lack of semi-linear well-posedness for $s < 0$, one uses a family of solutions which concentrate on the line $\{t = 0\}$ of the space time (t, x) . This family of solutions is related to the Galilean invariance

of (3.1).

Notice that for $d \leq 4$, the space H^1 is essentially covered by Theorem 3.1. For $d \geq 5$, the result of Theorem 3.1 is far from the regularity H^1 , and moreover as we have shown, for $d \geq 5$ the Cauchy problem (3.1) is not well-posed for data in $H^1(\mathbb{R}^d)$.

In order to have an H^1 theory in dimensions $d \geq 5$, it is reasonable to replace (3.1) with the equation

$$(3.16) \quad (i\partial_t + \Delta)u = F(u),$$

where the nonlinear interaction F is supposed to satisfy $F(0) = 0$ and is supposed of the form $F = \frac{\partial V}{\partial \bar{z}}$ with a *positive* $V \in C^\infty(\mathbb{C}; \mathbb{R})$ satisfying $V(e^{i\theta}z) = V(z)$, $\theta \in \mathbb{R}$, $z \in \mathbb{C}$, and, for some $\alpha > 1$,

$$|\partial_z^{k_1} \partial_{\bar{z}}^{k_2} V(z)| \leq C_{k_1, k_2} (1 + |z|)^{1+\alpha-k_1-k_2}.$$

The number α involved in the second condition on V corresponds to the degree of the nonlinearity $F(u)$. The Hamiltonian associated to (3.16) is

$$(3.17) \quad \int_{\mathbb{R}^d} |\nabla u|^2 + \int_{\mathbb{R}^d} V(u).$$

which controls the \dot{H}^1 norm. If $\alpha < 1 + \frac{4}{d-2}$, the second term in (3.17) is controlled by the first one and the L^2 norm of u . It is therefore reasonable to expect that for $\alpha < 1 + \frac{2}{d-2}$ the Cauchy problem for (3.16) is well-posed for data in H^1 . It turns out to be the case at least for $d = 5, 6$.

Theorem 3.9 (cf. [29, 40, 18]). *Let $\alpha < 1 + \frac{4}{d-2}$ and $d \leq 6$. Then the Cauchy problem associated to (3.16) is semi-linearly well-posed for data in $H^1(\mathbb{R}^d)$.*

Remark 3.10. *The proof of Theorem 3.9 relies crucially on the Strichartz inequalities for the non-homogeneous linear problem (recall that in the proof of Theorem 3.1 we only used the Strichartz inequalities for the free evolution). For $d \geq 7$ one can still prove the existence and the uniqueness. Since the nonlinearity in (3.16) is not polynomial, the propagation of the regularity is a nontrivial problem in the analysis of (3.16) (cf. e.g. [33]). In addition, even if one can prove the regularity propagation, the semi-linear well-posedness of (3.16) for $d \geq 7$ remains a non trivial issue.*

3.2. Nonlinear Schrödinger equations on compact manifolds. Let (M, g) be a compact smooth boundaryless Riemannian manifold of dimension $d \geq 2$. Denote by Δ_g the Laplace operator associated to the metric g . In this section we consider

the analog of (3.1) on (M, g)

$$(3.18) \quad iu_t + \Delta_g u + |u|^2 u = 0, \quad u(0) = u_0.$$

We have the following well-posedness result for (3.18)

Theorem 3.11 (cf. [15]). *The Cauchy problem (3.18) is semi-linearly well-posed for data in $H^s(M)$, $s > \frac{d-1}{2}$.*

The proof of Theorem 3.11 is based on the following Strichartz inequality with derivative losses for the free evolution

$$\|\exp(it\Delta_g)u_0\|_{L^p([0,T]; L^q(M))} \leq C_T \|u_0\|_{H^{\frac{1}{p}}(M)},$$

where

$$(3.19) \quad \frac{2}{p} + \frac{d}{q} = \frac{d}{2}, \quad p \geq 2, \quad (p, q) \neq (2, \infty).$$

Then the contraction argument is performed in the space

$$X_T = L^\infty([0, T]; H^s(M)) \cap L^p([0, T]; W^{s-\frac{1}{p}, q}(M)),$$

with suitable (p, q) satisfying (3.19).

Notice that there is a gap between the regularity $\frac{d-1}{2}$ of Theorem 3.11 and the regularity $\frac{d-2}{2}$ of Theorem 3.1. It is natural question what happens for data in H^s , $s \in [\frac{d-2}{2}, \frac{d-1}{2}]$. It turns out that in the case of the flat torus \mathbb{T}^d we can recover the result of the case \mathbb{R}^d .

Theorem 3.12 (cf. [8]). *Let M be the flat torus. Then (3.18) is semi-linearly well-posed for data in $H^s(\mathbb{T}^d)$, $s > \frac{d-2}{2}$.*

As in the case of the KdV equation of the previous section, the proof of Theorem 3.12 uses the Fourier transform restrictions spaces. An important additional element in the analysis is the use of bilinear improvements of the Strichartz inequalities.

The spaces of Bourgain and bilinear Strichartz estimates can also be used in the case of the sphere to get the following result.

Theorem 3.13 (cf. [19, 20]). *Let M be the standard sphere S^d . Then :*

1. *If $d = 2$ then (3.18) is semi-linearly well-posed for data in $H^s(S^2)$, $s > \frac{1}{4}$.*
2. *If $d \geq 3$ then (3.18) is semi-linearly well-posed for data in $H^s(S^d)$, $s > \frac{d-2}{2}$.*

It is worth noticing that the ill-posedness result of Theorem 3.4 still applies in the setting of the Riemannian manifolds (cf. Remark 3.7). Therefore, for $d \geq 3$, the indice $s = \frac{d-2}{2}$, turns out to be the critical one for both the well-posedness and

the semi-linear well-posedness of (3.18) posed on the flat torus or on the standard sphere.

We observe that in the case $d = 2$ the assumption $s > 1/4$ in Theorem 3.13 is more restrictive than in the case of the torus \mathbb{T}^2 . It turns out that this assumption is sharp, as far as the semi-linear well-posedness is concerned.

Theorem 3.14 (cf. [16]). *Let M be the standard sphere S^2 . Then (3.18) is not semi-linearly well-posed for data in $H^s(S^2)$, $s < \frac{1}{4}$.*

Proof. The assertion of Theorem 3.14 is a consequence of the description of the evolution by the flow of (3.18) of the highest weight spherical harmonics.

Proposition 3.15. *Let $T > 0$, $s \in]\frac{3}{20}, \frac{1}{4}[$, $\kappa \in]0, 1[$. Take $M = S^2$ with the canonical metric in (3.18). For $n \in \mathbb{N}$, we denote by $\psi_n : S^2 \rightarrow \mathbb{C}$ the restriction to S^2 of the harmonic polynomial $(x_1 + ix_2)^n$. Then there exists n_0 such that for $n \geq n_0$, the solution $u_n(t)$ of (3.18) with initial data $\kappa\varphi_n$, where $\varphi_n = n^{\frac{1}{4}-s}\psi_n$ is globally defined, and, for $t \in [0, T]$ it can be represented as*

$$(3.20) \quad u_n(t) = \kappa e^{-it(n(n+1)-\kappa^2\omega_n)} (\varphi_n + r_n(t)),$$

where $\omega_n \approx n^{\frac{1}{2}-2s}$ and $r_n(t)$ satisfies

$$\|r_n(t)\|_{H^s(S^2)} \leq C_T n^{-\delta}$$

where $\delta > 0$ and C_T depends on T but not on n . Moreover there exists $C > 0$, independent of T and n such that

$$(3.21) \quad \|u_n\|_{L^\infty(\mathbb{R}; H^s(S^2))} \leq C\kappa.$$

Proof of Proposition 3.15. Recall that ψ_n is an eigenfunction of $-\Delta_g$ associated to an eigenvalue $n^2 + n$. An easy computation shows that

$$\|\psi_n\|_{L^p} \approx n^{-\frac{1}{2p}}, \quad n \gg 1.$$

Therefore $\|\varphi_n\|_{H^s} \approx 1$ and $\|\varphi_n\|_{L^2} \leq Cn^{-s}$. Similarly to the Euclidean case, the solutions of (3.18) enjoy the conservation laws

$$(3.22) \quad \int_M |u(t, x)|^2 dx = \text{Const}$$

and

$$(3.23) \quad \int_M |\nabla u(t, x)|^2 dx - \frac{1}{2} \int_M |u(t, x)|^4 dx = \text{Const}.$$

The well-posedness result of Theorem 3.11 applies for the initial data $u_0 = \kappa\varphi_n$ and we obtain a local solution $u_n(t)$. But since $\|\varphi_n\|_{L^2} \leq Cn^{-s}$, using the conservation laws (3.22), (3.23) and the Gagliardo-Nirenberg inequalities, we deduce that for

$n \gg 1$ the H^1 norm of $u_n(t)$ is bounded uniformly with respect to t . Therefore, we can reiterate the well-posedness results and to obtain that for $n \gg 1$, the solutions $u_n(t)$ are globally defined.

For every $\alpha \in \mathbb{R}$, we denote by R_α the rotation of \mathbb{R}^3 defined by

$$R_\alpha(x_1, x_2, x_3) = (\cos \alpha x_1 - \sin \alpha x_2, \sin \alpha x_1 + \cos \alpha x_2, x_3)$$

and by R_α^* the associated unitary operator of $L^2(S^2)$,

$$R_\alpha^* u(x) = u(R_\alpha(x)).$$

Observe that $R_\alpha^* \psi_n = \exp(in\alpha) \psi_n$ for every $\alpha \in \mathbb{R}$. The following elementary lemma will be useful in the sequel.

Lemma 3.16. *Let $n \in \mathbb{Z}^+$ and $u \in L^2(S^2)$ be such that for every $\alpha \in \mathbb{R}$,*

$$(3.24) \quad R_\alpha^* u = \exp(in\alpha) u.$$

Then the decomposition of u in spherical harmonics reads

$$u = c\psi_n + \sum_j g_j$$

where $c \in \mathbb{C}$ and each g_j is a spherical harmonic of degree $> n$.

Proof. Since the family $(R_\alpha^*)_{\alpha \in \mathbb{R}}$ is a one-parameter group of unitary operators leaving invariant the space of spherical harmonics of degree l , one can find an orthonormal basis (h_k) of $L^2(S^2)$ such that, for every k , h_k is a spherical harmonic satisfying, for some $n_k \in \mathbb{Z}$, for every $\alpha \in \mathbb{R}$,

$$(3.25) \quad R_\alpha^* h_k = \exp(in_k \alpha) h_k.$$

Comparing (3.24) and (3.25), the decomposition of u in the basis (h_k) reads

$$(3.26) \quad u = \sum_{k: n_k = n} c_k h_k.$$

Let h be a spherical harmonic of degree l satisfying property (3.24) for every $\alpha \in \mathbb{R}$. Denote by P the l -homogeneous polynomial on \mathbb{R}^3 such that $h = P|_{S^2}$. Then (3.24) is equivalent to

$$(3.27) \quad \forall x \in \mathbb{R}^3, \quad P(R_\alpha(x)) = \exp(in\alpha) P(x).$$

Let us decompose P according to the powers of $z = x_1 + ix_2$ and $\bar{z} = x_1 - ix_2$

$$(3.28) \quad P(x_1, x_2, x_3) = \sum_{p+q \leq l} a_{pq} z^p \bar{z}^q x_3^{l-p-q}$$

where $a_{pq} \in \mathbb{C}$. In view of (3.27), (3.28) and

$$(3.29) \quad P(R_\alpha(x)) = \sum_{p+q \leq l} a_{pq} \exp(i(p-q)\alpha) z^p \bar{z}^q x_3^{l-p-q},$$

we conclude that $a_{pq} = 0$ unless $p - q = n$. As a consequence,

$$l \geq p + q \geq p - q = n$$

and, if $l = n$, then $p = n$ and $q = 0$, so that $P = cz^n$, i.e. $h = c\psi_n$ for some $c \in \mathbb{C}$. Coming back to decomposition (3.26) completes the proof of Lemma 3.16. \square

Using Lemma 3.16, we can write

$$|\varphi_n|^2 \varphi_n = \omega_n \varphi_n + r_n,$$

where r_n contains only spherical harmonics of degree $> n$ in its spectral decomposition and

$$\omega_n = \frac{\|\varphi_n\|_{L^4}^4}{\|\varphi_n\|_{L^2}^2} \approx n^{\frac{1}{2}-2s}.$$

Observe that that $R_\alpha^* u_n$ is a solution of (3.18) with data $u_0 = \kappa e^{in\alpha} \varphi_n$. On the other hand $e^{in\alpha} u_n$ is also a solution of (3.18) with the same initial data. Therefore, using the uniqueness assertion of Theorem 2.6 (in spaces invariant under the action of R_α^*) for the Cauchy problem (3.18), we obtain

$$R_\alpha^* u_n = e^{in\alpha} u_n.$$

Using Lemma 3.16, we deduce that $u_n(t)$ is a linear combination of ψ_n and spherical harmonics of degree $> n$.

Let us give the heuristic argument which permits us to find an ansatz for $u_n(t)$. In view of the above discussion, we may hope that $u_n(t)$ can be written as

$$u_n(t) = \kappa c_n(t) \varphi_n + \text{“small error”}.$$

Substituting this in the equation (3.18), neglecting the “small error” and projection on φ_n yields the equation

$$i\kappa \dot{c}_n - n(n+1)\kappa c_n + \kappa^3 \omega_n |c_n|^2 c_n = 0, \quad c_n(0) = 0$$

which gives

$$c_n(t) = e^{-it(n(n+1)-\kappa^2 \omega_n)}.$$

In order to make the above formal discussion rigorous, we set

$$u_n(t) = \kappa e^{-it(n(n+1)-\kappa^2 \omega_n)} ((1 + z_n(t)) \varphi_n + q_n(t)),$$

where $z_n(0) = 0$, $q_n(0) = 0$ and $q_n(t)$ contains only spherical harmonics of degree $> n$ in its spectral decomposition. Proposition 3.15 is clearly a consequence of the following statements.

Lemma 3.17. *There exists a constant $C > 0$, independent of T and n such that*

$$\|q_n(t)\|_{H^s} \leq Cn^{-\frac{1}{4}-s}.$$

Lemma 3.18. *There exists a constant $C_T > 0$, which depends on T but not on n such that*

$$\sup_{t \in [0, T]} |z_n(t)| \leq C_T n^{\frac{1}{4}-3s}.$$

Proof of Lemma 3.17. Let us first rewrite the conservation laws (3.22), (3.23) in terms of $z_n(t)$ and $q_n(t)$. Since φ_n is orthogonal to $q_n(t)$ in $L^2(S^2)$ as well as $\nabla\varphi_n$ to $\nabla q_n(t)$, we can rewrite (3.22) and (3.23) as

$$(3.30) \quad |1 + z_n(t)|^2 \|\varphi_n\|_{L^2}^2 + \|q_n(t)\|_{L^2}^2 = \|\varphi_n\|_{L^2}^2,$$

$$(3.31) \quad |1 + z_n(t)|^2 \|\nabla\varphi_n\|_{L^2}^2 + \|\nabla q_n(t)\|_{L^2}^2 - \frac{1}{2\kappa^2} \|u_n(t)\|_{L^4}^4 = \|\nabla\varphi_n\|_{L^2}^2 - \frac{\kappa^2}{2} \|\varphi_n\|_{L^4}^4.$$

Observe that

$$\|\nabla\varphi_n\|_{L^2}^2 = \langle \varphi_n, -\Delta\varphi_n \rangle = n(n+1)\|\varphi_n\|_{L^2}^2,$$

where $\langle \cdot, \cdot \rangle$ denotes the $L^2(S^2)$ scalar product. Therefore multiplying (3.30) with $-n(n+1)$ and adding it to (3.31) gives

$$(3.32) \quad \|\nabla q_n(t)\|_{L^2}^2 - n(n+1)\|q_n(t)\|_{L^2}^2 = \frac{1}{2\kappa^2} \|u_n(t)\|_{L^4}^4 - \frac{\kappa^2}{2} \|\varphi_n\|_{L^4}^4 \leq \frac{1}{2\kappa^2} \|u_n(t)\|_{L^4}^4.$$

Let us bound the right hand-side of (3.32) as follows

$$\begin{aligned} \|u_n(t)\|_{L^4}^4 &\leq C(\|\varphi_n\|_{L^4}^4 + \|q_n(t)\|_{L^4}^4) \leq \\ &\leq C(n^{\frac{1}{2}-4s} + \|q_n(t)\|_{L^4}^4) \leq C(n^{\frac{1}{2}-4s} + \|q_n(t)\|_{H^{\frac{1}{2}}}^4), \end{aligned}$$

where we used the Sobolev inequality in the last estimate. We can decompose

$$q_n(t) = \sum_{l \geq n+1} q_{n,l}(t)$$

where $q_{n,l} \in \text{Ker}(\Delta_{S^2} + l(l+1))$. Hence

$$\|\nabla q_n(t)\|_{L^2}^2 - n(n+1)\|q_n(t)\|_{L^2}^2 = \sum_{l \geq n+1} (l(l+1) - n(n+1))\|q_{n,l}(t)\|_{L^2}^2.$$

If $l \geq n+1$, we have $l(l+1) - n(n+1) \geq l$ and therefore

$$\|\nabla q_n(t)\|_{L^2}^2 - n(n+1)\|q_n(t)\|_{L^2}^2 \geq \|q_n(t)\|_{H^{\frac{1}{2}}}^2.$$

Coming back to (3.32),

$$\|q_n(t)\|_{H^{\frac{1}{2}}}^2 \leq C(n^{\frac{1}{2}-4s} + \|q_n(t)\|_{H^{\frac{1}{2}}}^4).$$

Since $q_n(0) = 0$ by a bootstrap argument,

$$(3.33) \quad \|q_n(t)\|_{H^{\frac{1}{2}}} \leq Cn^{\frac{1}{4}-2s}.$$

On the other hand, we also have that if $l \geq n+1$ then $l(l+1) - n(n+1) \geq n$ and thus

$$n\|q_n(t)\|_{L^2}^2 \leq C(n^{\frac{1}{2}-4s} + n^{1-8s}).$$

Since $s > 3/20$, we get the bound

$$(3.34) \quad \|q_n(t)\|_{L^2} \leq Cn^{-\frac{1}{4}-2s}.$$

Using (3.33) and (3.34), we finally arrive at

$$\|q_n(t)\|_{H^s} \leq \|q_n(t)\|_{L^2}^{1-2s} \|q_n(t)\|_{H^{\frac{1}{2}}}^{2s} \leq Cn^{-\frac{1}{4}-s}.$$

This completes the proof of Lemma 3.17 \square

Proof of Lemma 3.18. Let us set $w_n(t) := z_n(t)\varphi_n + q_n(t)$. By projecting the equation

$$(i\partial_t + \Delta)u_n + |u_n|^2 u_n = 0$$

on the mode φ_n , we get that z_n solves the equation

$$\begin{aligned} i\dot{z}_n - \kappa^2 \omega_n z_n &= -\frac{\kappa^2}{\|\varphi_n\|_{L^2}^2} \left(\langle |\varphi_n + w_n|^2 (\varphi_n + w_n), \varphi_n \rangle - \langle |\varphi_n|^2 \varphi_n, \varphi_n \rangle \right) = \\ &= -\frac{\kappa^2}{\|\varphi_n\|_{L^2}^2} \left(\int (2|\varphi_n|^2 w_n + \varphi_n^2 \bar{w}_n) \bar{\varphi}_n + (2\text{Re}(\bar{\varphi}_n w_n) w_n \bar{\varphi}_n + |w_n|^2 |\varphi_n|^2 + |w_n|^2 w_n \bar{\varphi}_n) \right). \end{aligned}$$

Next, the equation for $z_n(t)$ can be rewritten as

$$i\dot{z}_n - \kappa^2 \omega_n z_n = -2\omega_n \kappa^2 z_n - \omega_n \kappa^2 \bar{z}_n + \frac{\kappa^2}{\|\varphi_n\|_{L^2}^2} \mathcal{O}(|z_n|^2 \int |\varphi_n|^4 + |z_n|^3 \int |\varphi_n|^4 + \int |q_n|^3 |\varphi_n| + \int |q_n|^2 |\varphi_n|^2 + |\langle q_n, r_n \rangle|).$$

Let us estimate the source terms. Write using Lemma 3.17

$$\begin{aligned} \frac{\int |q_n|^3 |\varphi_n|}{\|\varphi_n\|_{L^2}^2} &\leq Cn^{2s} \|q_n\|_{L^3}^3 \|\varphi_n\|_{L^\infty} \leq Cn^{2s} \|q_n\|_{L^2} \|q_n\|_{H^{\frac{1}{2}}}^2 \|\varphi_n\|_{L^\infty} \leq \\ &\leq Cn^{2s} n^{-\frac{1}{4}-2s} n^{\frac{1}{2}-4s} n^{\frac{1}{4}-s} = Cn^{\frac{1}{2}-5s}. \end{aligned}$$

Further we have

$$\frac{\int |q_n|^2 |\varphi_n|^2}{\|\varphi_n\|_{L^2}^2} \leq Cn^{2s} \|q_n\|_{L^2}^2 \|\varphi_n\|_{L^\infty}^2 \leq Cn^{2s} n^{-\frac{1}{2}-4s} n^{\frac{1}{2}-2s} = Cn^{-4s}.$$

and

$$\frac{|\langle q_n, r_n \rangle|}{\|\varphi_n\|_{L^2}^2} \leq Cn^{2s} \|q_n\|_{L^2} \|r_n\|_{L^2} \leq Cn^{2s} n^{-\frac{1}{4}-2s} n^{\frac{1}{2}-3s} = Cn^{\frac{1}{4}-3s}.$$

Therefore, if $s > \frac{1}{8}$, the equation for $z_n(t)$ can be written as

$$(3.35) \quad i\partial_t z_n = -2\omega_n \kappa^2 \operatorname{Re}(z_n) + \mathcal{O}(\omega_n |z_n|^2 + \omega_n |z_n|^3 + n^{\frac{1}{4}-3s})$$

with $z_n(0) = 0$. Moreover using once again the L^2 conservation law (3.30), we have

$$1 - |1 + z_n|^2 = \frac{\|q_n(t)\|_{L^2}^2}{\|\varphi_n\|_{L^2}^2} = \mathcal{O}(n^{-\frac{1}{2}-2s}).$$

Therefore

$$|2\operatorname{Re}(z_n) + |z_n|^2| = \mathcal{O}(n^{-\frac{1}{2}-2s})$$

and the equation (3.35) takes the form

$$i\partial_t z_n = \mathcal{O}(\omega_n |z_n|^2 + \omega_n |z_n|^3 + n^{\frac{1}{4}-3s}),$$

with $\omega_n = \mathcal{O}(n^{\frac{1}{2}-2s})$. Hence if we set

$$M_n(T) = \sup_{0 \leq t \leq T} |z_n(t)|,$$

we obtain

$$(3.36) \quad M_n(T) \leq CT(n^{\frac{1}{2}-2s} [M_n(T)]^2 + n^{\frac{1}{2}-2s} [M_n(T)]^3 + n^{\frac{1}{4}-3s}).$$

In view of (3.36), we set

$$\tilde{M}_n(T) = n^{3s-\frac{1}{4}} M_n(T)$$

and therefore (3.36) yields

$$\tilde{M}_n(T) \leq CT(1 + n^{\frac{3}{4}-5s}[\tilde{M}_n(T)]^2 + n^{1-8s}[\tilde{M}_n(T)]^3).$$

Since $\tilde{M}_n(0) = 0$ and $s > \frac{3}{20}$, we obtain that $\tilde{M}_n(T) \leq CT$ uniformly with respect to n . This completes the proof of Lemma 3.18. \square

This proof of Proposition 3.15 is now completed. \square

Notice that the assertion of Theorem 3.14 is new only for $0 \leq s < 1/4$. Indeed for $s < 0$, we can apply Theorem 3.4.

Let us now show how Proposition 3.15 implies Theorem 3.14 for $3/20 < s < 1/4$. The main point is that for $s < 1/4$ we have $\omega_n \rightarrow \infty$ as $n \rightarrow \infty$. Let us fix $T > 0$, $\kappa \in]0, 1]$ and let (κ_n) be a sequence of positive numbers such that

$$(\kappa^2 - \kappa_n^2)\omega_n = n^\beta, \quad 0 < \beta < 1.$$

Since $s < 1/4$ and $\omega_n \approx n^{\frac{1}{2}-2s}$, we have that for $\beta \ll 1$, $\kappa_n \rightarrow \kappa$. Let $(u_{\kappa,n})$ and $(u_{\kappa_n,n})$ be the solutions of (3.18) with data $\kappa\varphi_n$ and $\kappa_n\varphi_n$ respectively. Then

$$\|u_{\kappa,n}(0, \cdot) - u_{\kappa_n,n}(0, \cdot)\|_{H^s} \leq C|\kappa - \kappa_n| \rightarrow 0$$

but thanks to Proposition 3.15, for $t \in [0, T]$,

$$\|u_{\kappa,n}(t, \cdot) - u_{\kappa_n,n}(t, \cdot)\|_{H^s} \geq c|e^{itn^\beta} - 1| - CTn^{-\delta}$$

with $\delta > 0$. The proof of Theorem 3.14 for $3/20 < s < 1/4$ is completed by observing that for all $n \gg 1$,

$$\sup_{0 \leq t \leq T} |e^{itn^\beta} - 1| = 2.$$

When $0 \leq s \leq 3/20$, we need to perform a slight modification of the argument. Indeed, in this case it suffices to remark that in fact we need to justify the ansatz only on a *small interval*⁵ $[0, T_n]$ with T_n satisfying

$$\lim_{n \rightarrow \infty} n^{\frac{1}{2}-2s}T_n = \infty.$$

The bound of Lemma 3.17 is uniform in time. We only need to slightly modify the proof of Lemma 3.18. In the case $0 \leq s \leq 3/20$, we define

$$\tilde{M}_n(T) := T_n^{-1}n^{3s-\frac{1}{4}}M_n(T), \quad 0 \leq T \leq T_n,$$

and, the argument of Lemma 3.18 yields the bound

$$\tilde{M}_n(T) \leq C(1 + T_n^2n^{\frac{3}{4}-5s}[\tilde{M}_n(T)]^2 + T_n^3n^{1-8s}[\tilde{M}_n(T)]^3).$$

⁵Recall that a similar idea was used in the discussion around Theorem 3.4.

If we set

$$T_n := n^{\frac{5s}{2} - \frac{3}{8} - \varepsilon}$$

with $0 < \varepsilon \ll 1$, a bootstrap argument gives

$$\tilde{M}_n(T) \leq C$$

and therefore

$$|z_n(t)| \leq C n^{\frac{1}{4} - 3s} n^{\frac{5s}{2} - \frac{3}{8} - \varepsilon} = C n^{-\frac{1}{8} - \frac{s}{2} - \varepsilon}, \quad 0 \leq t \leq T_n.$$

Therefore the ansatz is valid on $[0, T_n]$, which is a sufficiently large small interval to get the instability property of the flow map. Indeed

$$n^{\frac{1}{2} - 2s} T_n = n^{\frac{1}{8} + \frac{s}{2} - \varepsilon}$$

which gives the big oscillations needed to assure the lack of uniform continuity of the flow map. Let us finally mention that in the case $s = 0$, to assure the global existence one also has to impose a smallness assumption (independent of n) on the parameter κ . This completes the proof of Theorem 3.14. \square

We end this section by several remarks.

The result of Theorem 3.14 is another instance when we see that the critical indice for the semi-linear well-posedness is shifted from the scaling one because of concentration on a curve (a closed geodesic). It would be interesting to develop a notion of critical exponent associated to curve similarly to the one associated to a point via the scaling invariance.

It would be interesting to decide whether for some $0 \leq s \leq 1/4$, the Cauchy problem (3.18) is well-posed (probably after a suitable gauge transform) for data in $H^s(S^2)$. Recall that such a phenomenon is not excluded as shows the experience with the modified KdV equation.

We do not know for an analog of Theorem 3.9 in the setting of compact manifolds. Moreover, it is known that in case of the sphere S^6 the assertion of Theorem 3.9 fails. More precisely the Cauchy problem

$$iu_t + \Delta_g u = (1 + |u|^2)^{\frac{\alpha}{2}} u, \quad u(0) = u_0, \quad 0 < \alpha \leq 1,$$

posed on S^6 is not semi-linearly well-posed for data in $H^1(S^6)$ (cf. [16, 18])

We refer to the work of Banica [2], where the ansatz used in the proof of Theorem 3.14 is justified up to time one for $0 < s \leq 3/20$. Let us also mention the work [17] where the approach of Theorem 3.14 is extended to (3.18) posed on the unit

disc of \mathbb{R}^2 with Dirichlet boundary conditions. We also refer to [26] for ill-posedness results for the cubic NLS posed on the circle S^1 .

4. FINAL REMARKS

There has been a number of works, closely related to the discussion in these notes for nonlinear wave equations (cf. [14, 25, 28, 54, 52, 53] ...). In the context of the nonlinear wave equations, again, *families* of solutions concentrating at a point contradict the well-posedness (or semi-linear well-posedness) below the scaling exponent. The finite propagation speed of the wave equation is exploited in [53] to construct a *single* solution, concentrating in an infinite number of points, which stays bounded in H^s (for some suitable s) and becomes instantaneously very large in H^σ , $\sigma > s$. It would be interesting to prove the analogue of Lebeau's result in the context of the NLS. Despite the lack of the finite propagation speed for the Schrödinger operator, the reasoning in the proof of Theorem 3.4 above is of a semi-classical nature (cf. also [15]) and thus finite propagation speed considerations could be employed. It is worth noticing that, again, in the case of nonlinear wave equations ill-posedness above the scaling is closely related to concentrations on curves, for instance the Lorenz invariance provides families of solutions concentrating on light rays.

The problematic discussed in these notes fits naturally in the context of parabolic PDE's. There has been some first results in that direction (cf. [21, 57] ...), and, we believe there is further progress to come.

REFERENCES

- [1] L. Abdelouhab, J. Bona, M. Felland, J.C. Saut, *Nonlocal models for nonlinear, dispersive waves*, Phys. D 40 (1989) 360–392.
- [2] V. Banica, *On the nonlinear Schrödinger dynamics on S^2* J. Math. Pure Appl. 83 (2004) 77–98.
- [3] T.B. Benjamin, *Internal waves of permanent form in fluids of great depth*, J. Fluid Mech. 29 (1967) 559–592.
- [4] H. Biagioni, F. Linares, *Ill-posedness for the derivative Schrödinger and generalized Benjamin-Ono equations*, Trans. Amer. Math. Soc. 353 (2001), 3649–3659.
- [5] B. Birnir, C. Kenig, G. Ponce, N. Svanstedt, L. Vega, *On the ill-posedness of the IVP for the generalized Korteweg-de Vries and nonlinear Schrödinger equations*, Annales I.H.P., Analyse non linéaire 13 (1996) 529–535.
- [6] B. Birnir, G. Ponce, N. Svanstedt, *The local ill-posedness of the modified KdV equation*, J. London Math. Soc. 53 (1996) 551–559.
- [7] J.L. Bona, R. Smith, *The initial-value problem for the Korteweg-de Vries equation*, Philos. Trans. Roy. Soc. London Ser. A 278 (1975), 1287, 555–601.
- [8] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear evolution equations I. Schrödinger equations*, Geom. Functional Analysis 3 (1993) 107–156.

- [9] J. Bourgain, *Exponential sums and nonlinear Schrödinger equations*, Geom. Functional Analysis 3 (1993) 157-178.
- [10] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and application to nonlinear evolution equations II. The KdV equation*, Geom. Functional Analysis 3 (1993) 209-262.
- [11] J. Bourgain, *On the Cauchy problem for the Kadomtsev-Petviashvili equation*, Geom. Functional Analysis 3 (1993) 315-341.
- [12] J. Bourgain, *Periodic Korteweg de Vries equation with measures as initial data*, Sel. Math. New. Ser. 3 (1997) 115-159.
- [13] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, Colloq. Publications, American Math. Soc., 1999.
- [14] P. Brenner, P. Kumlin, *On wave equations with supercritical nonlinearities*, Archiv der Mathematik 74 (200) 129-147.
- [15] N. Burq, P. Gérard, N. Tzvetkov, *Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds*, Amer. J. Math. 126 (2004) 569-605.
- [16] N. Burq, P. Gérard, N. Tzvetkov, *An instability property of the nonlinear Schrödinger equation on S^d* , Math. Research Letters, 9 (2002) 323-335.
- [17] N. Burq, P. Gérard, N. Tzvetkov, *Two singular dynamics of the nonlinear Schrödinger equation on a plane domain*, Geom. Functional Analysis, 13 (2003) 1-19.
- [18] N. Burq, P. Gérard, N. Tzvetkov, *The Cauchy problem for the nonlinear Schrödinger equation on compact manifold*, J. Nonlinear Math. Physics, 10 (2003) 12-27.
- [19] N. Burq, P. Gérard, N. Tzvetkov, *Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces*, Invent. Math. (to appear).
- [20] N. Burq, P. Gérard, N. Tzvetkov, *Multilinear eigenfunction estimates and global existence for the three dimensional nonlinear Schrödinger equations*, Preprint July 2004.
- [21] M. Canone, G. Karch, *On the validity of Picard algorithm for nonlinear parabolic equation*, Preprint 2003.
- [22] T. Cazenave, *An introduction to nonlinear Schrödinger equations, second edition*, Textos de Metodes Mathematicos 26, 1993.
- [23] T. Cazenave, F. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear. Anal. TMA 14 (1990) 807-836.
- [24] M. Christ, J. Colliander, T. Tao, *Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations*, Amer. J. Math. 125 (2003) 1235-1293.
- [25] M. Christ, J. Colliander, T. Tao, *Ill-posedness for nonlinear Schrödinger and wave equations*, Preprint 2003.
- [26] M. Christ, J. Colliander, T. Tao, *Instability of the periodic nonlinear Schrödinger equation*, Preprint 2003.
- [27] M. Christ, *Illposedness of a Schrödinger equation with derivative nonlinearity*, Preprint 2003.
- [28] P. D'Ancona, V. Georgiev, *On the continuity of the solution operator to the wave map system*, Comm. Pure Appl. Math. 57 (2004) 357-383.
- [29] J. Ginibre, G. Velo, *On a class of nonlinear Schrödinger equations*, J. Func. Anal. 32 (1979) 1-71.
- [30] J. Ginibre, G. Velo, *The global Cauchy problem for the nonlinear Schrödinger equation*, Ann. Inst. H. Poincare (Analyse non linéaire) 2 (1985) 309-327.
- [31] J. Ginibre, G. Velo, *Smoothing properties and retarded estimates for some dispersive evolution equations*, Comm. Math. Phys. 144 (1992) 163-188.

- [32] J. Ginibre, *Le problème de Cauchy pour des EDP semi-linéaires périodiques en variables d'espace (d'après Bourgain)* Séminaire Bourbaki 796, Astérisque (1995) 163–187.
- [33] J. Ginibre, *Introduction aux équations de Schrödinger non linéaires*, Cours de DEA, Université de Paris Sud 1994-1995. Edition Paris Onze.
- [34] N. Hayashi, T. Ozawa, *Remarks on Schrödinger equations in one space dimension*, Diff. Int. Equations 7 (1994) 453-461.
- [35] P. Isaza, J. Mejia, *Local and global Cauchy problems for the Kadomtsev-Petviashvili (KP-II) equation in Sobolev spaces of negative indices*, Comm. PDE 26 (2001) 1027-1054.
- [36] J.R. Iorio, *On the Cauchy problem for the Benjamin-Ono equation*, Comm. PDE 11 (1986) 1031-1081.
- [37] B.B. Kadomtsev, V.I. Petviashvili, *On the stability of solitary waves in weakly dispersive media*, Soviet Phys. Dokl. 15 (1970) 539-541.
- [38] T. Kappeler, P. Topalov, *Global well-posedness of KdV in $H^{-1}(\mathbb{T}, \mathbb{R})$* , Preprint 2003.
- [39] T. Kappeler, P. Topalov, *Global well-posedness of the mKdV in $L^2(\mathbb{T}, \mathbb{R})$* , Preprint 2003.
- [40] T. Kato, *Quasilinear equations of evolution, with applications to partial differential equations*, Springer Lecture Notes in Math. 448 (1975), 27-50.
- [41] T. Kato, *On nonlinear Schrödinger equations*, Ann. Inst. H. Poincaré Phys. Theor. 46 (1986) 113-129.
- [42] T. Kato, *On nonlinear Schrödinger equations II. H^s solutions and unconditional well-posedness*, J. Anal. Math. 67 (1995), 281-306.
- [43] T. Kato, G. Ponce, *Commutator estimates and the Euler and Navier-Stokes equations*, Comm. Pure Appl. Math. 41 (1988), 891-907.
- [44] C. Kenig, G. Ponce, L. Vega, *On the (generalized) Korteweg- de Vries equation*, Duke Math. J. 59 (1989), 585-610.
- [45] C. Kenig, G. Ponce, L. Vega, *Well-posedness and scattering results for the generalized Korteweg de Vries equation via the contraction principle*, Comm. Pure Appl. Math, 46 (1993) 527-620.
- [46] C. Kenig, G. Ponce, L. Vega, *A bilinear estimate with application to the KdV equation*, J. Amer. Math. Soc. 9 (1996) 527-620.
- [47] C. Kenig, G. Ponce, L. Vega, *On the ill-posedness of some canonical dispersive equations*, Duke Math. J. 106 (2001) 617-633.
- [48] C. Kenig, K. Koehnig, *On the local well-posedness of the Benjamin-Ono and modified Benjamin-Ono equations*, Math. Res. Letters 10 (2003) 879-895.
- [49] H. Koch, N. Tzvetkov, *On the local well-posedness of the Benjamin-Ono equation in $H^s(\mathbb{R})$* , IMRN, 26 (2003) 1449-1464.
- [50] H. Koch, N. Tzvetkov, *Nonlinear wave interactions for the Benjamin-Ono equation*, Preprint 2003.
- [51] S. Kuksin, *On squeezing and of energy for nonlinear wave equations*, Geom. Functional Analysis, 5 (1995) 668-701.
- [52] G. Lebeau, *Non linear optic and supercritical wave equation*, Bull. Soc. Roy. Sci. Liège 70 (2002) 267–306.
- [53] G. Lebeau, *Perte de régularité pour les équations d'ondes sur-critiques* Preprint 2002.
- [54] H. Lindblad, *A sharp counterexample to the local existence of low-regularity solutions to nonlinear wave equations*, Duke Math. J. 72 (1993) 503-539.
- [55] J.L. Lions *Quelques méthodes de résolution des équations aux dérivées partielles non linéaires*, Dunod, Paris, 1969.

- [56] L. Molinet, F. Ribaud, *Well-posedness results for the generalized Benjamin-Ono equation with arbitrary large initial data*, Preprint 2004.
- [57] L. Molinet, F. Ribaud, A. Youssfi, *Ill-posedness issues for a class of parabolic equations* Proc. Roy. Soc. Edinburg Sect. A 132 (2002) 1407-1416.
- [58] L. Molinet, J.C. Saut, N. Tzvetkov, *Ill-posedness issues for the Benjamin-Ono and related equations*, SIAM J. Math. Anal. 33 (2001), 982-988.
- [59] L. Molinet, J.C. Saut, N. Tzvetkov, *Well-posedness and ill-posedness results for the Kadomtsev-Petviashvili-I equation*, Duke Math. J. 115 (2002) 353-384.
- [60] G. Ponce, *On the global well-posedness of the Benjamin-Ono equation*, Diff. Integral Eq. 4 (1991), 527-542.
- [61] J.C. Saut, *Sur quelques généralisations de l'équation de Korteweg- de Vries*, J. Math. Pures Appl. 58 (1979) 21-61.
- [62] H. Takaoka, Y. Tsutsumi, *Well-posedness of the Cauchy problem for the modified KdV equation with periodic boundary conditions*, Preprint 2003.
- [63] H. Takaoka, N. Tzvetkov, *On the local regularity of Kadomtsev-Petviashvili-II equation*, IMRN, 8 (2001) 77-114.
- [64] T. Tao, *Global regularity of wave maps I. Small critical Sobolev norm in high dimension*, IMRN 7 (2001), 299-328.
- [65] T. Tao, *Global regularity of wave maps II. Small energy in two dimensions*, Comm. Math. Phys. 224 (2001) 443-544.
- [66] T. Tao, *Global well-posedness of the Benjamin-Ono equation in $H^1(R)$* , J. Hyperbolic Diff. Eq. 1 (2004) 27-49.
- [67] Y. Tsutsumi, *L^2 - solutions for nonlinear Schrödinger equations and nonlinear groups*, Funkcial. Ekvak. 30 (1987) 115-125.
- [68] K. Yajima, *Existence of solutions for Schrödinger evolution equations*, Commun. Math. Phys. 110 (1987) 415-426.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ LILLE I, 59 655 VILLENEUVE D'ASCQ CEDEX, FRANCE

Dispersive estimates and applications

Herbert Koch, Dortmund

Strichartz estimates are one of several basic tools in the study of linear and nonlinear dispersive equations. They quantify the limitations of the extent to which waves can be focussed over a certain time interval. Analytically they extend estimates which are valid for elliptic operators to certain non elliptic operators.

The lectures series will focus on Strichartz estimates for equations with rough coefficients, which are motivated by strong wave interactions for nonlinear equations.

The first talk will give a survey on Strichartz estimates for equations with rough coefficients and their applications. The second lecture will describe the technique of wave packet decomposition leading to a calculus based on the FBI transform.

This wave packet decomposition is applied in the third lecture to linear dispersive equations, leading to a representation of the fundamental solution. This wave packet decomposition implies dispersive estimates.

Finally these techniques are applied to nonlinear dispersive equations and to spectral projections.

Most of the lecture series is based on the joint paper with D. Tataru:

Dispersive estimates for principally normal pseudodifferential operators,
aiXiv:math.AP/0401234

and on joint work with N. Tzvetkov:

On the local well-posedness of the Benjamin-Ono equation in $H^s(\mathbb{R})$, Int. Math. Res. Not. (26), 1449-1464(2003).