HOKKAIDO UNIVERSITY

Title	Immunoglobulin G (IgG)-Based Imaging Probe Accumulates in M1 Macrophage-Infiltrated Atherosclerotic Plaques Independent of IgG Target Molecule Expression
Author(s)	Shimizu, Yoichi; Hanzawa, Hiroko; Zhao, Yan et al.
Citation	Molecular imaging and biology, 19(4), 531-539 https://doi.org/10.1007/s11307-016-1036-8
Issue Date	2017-08
Doc URL	https://hdl.handle.net/2115/71143
Rights	The original publication is available at www.springerlink.com
Type	journal article
File Information	11307_2016_1036_MOESM1_ESM.pdf, Supplementary material

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Electronic Supplementary Materials

Immunoglobulin-G (IgG)-Based Imaging Probe Accumulates in M1 Macrophage-Infiltrated Atherosclerotic Plaques Independent of IgG Target Molecule Expression

Journal: Molecular Imaging and Biology

Yoichi Shimizu[1,2*], Hiroko Hanzawa[3], Yan Zhao[4], Sagiri Fukura[4], Ken-ichi Nishijima[2,4], Takeshi Sakamoto[5], Songji Zhao[4], Nagara Tamaki[4], Mikako Ogawa[1], Yuji Kuge[2,4]

[1] Faculty of Pharmaceutical Sciences, Hokkaido University, Kita 12 Nishi 6, Kita-ku, 060-0812 Sapporo, Japan


[2] Central Institute of Isotope Science, Hokkaido University, Kita 15 Nishi 7, Kita-ku, 060-0815 Sapporo, Japan

[3] Center for Exploratory Research, Research & Development Group, Hitachi, Ltd., 350-0395 Hatoyama, Japan

[4] Hokkaido University Graduate School of Medicine, Kita 15 Nishi 7, Kita-ku, 060-8638 Sapporo, Japan

[5] Center for Technology Innovation − Healthcare, Research & Development Group, Hitachi, Ltd., 185-8601 Kokubunji, Japan

Supplemental Figure



Supplemental Figure 1. Size exclusion HPLC analysis of [^{99m}Tc]IgG monitored using a UV detector at 280 nm (a) and a radiodetector (b).