



HOKKAIDO UNIVERSITY

Title	Substrate recognition of the catalytic α -subunit of glucosidase II from <i>Schizosaccharomyces pombe</i>
Author(s)	Okuyama, Masayuki; Miyamoto, Masashi; Matsuo, Ichiro et al.
Citation	Bioscience biotechnology and biochemistry, 81(8), 1503-1511 https://doi.org/10.1080/09168451.2017.1320520
Issue Date	2017-08
Doc URL	https://hdl.handle.net/2115/71180
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis in Bioscience biotechnology and biochemistry on 2017 Aug, available online: http://www.tandfonline.com/10.1080/09168451.2017.1320520 .
Type	journal article
File Information	text_170411.pdf



Regular paper

Title

Substrate recognition of the catalytic α -subunit of glucosidase II from *Schizosaccharomyces pombe*

Authors

Masayuki Okuyama^{1,†,*}, Masashi Miyamoto^{1,†}, Ichiro Matsuo², Shogo Iwamoto², Ryo Serizawa¹, Masanari Tanuma¹, Min Ma¹, Patcharapa Klahan¹, Yuya Kumagai¹, Takayoshi Tagami¹ and Atsuo Kimura^{1,*}

¹ *Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan.* ² *Graduate School of Science and Technology, Gunma University, Kiryu 376-8515, Japan.*

Email addresses

MO, okuyama@abs.agr.hokudai.ac.jp; MM, masashi.m1229@gmail.com; IM, matsuo@gunma-u.ac.jp; SI, t11802104@gunma-u.ac.jp; RS, ryo.serizawa@frontier.hokudai.ac.jp; MT, tanuma.marbo.0415@gmail.com; MM, mamin@abs.agr.hokudai.ac.jp; PK, patchakl@abs.agr.hokudai.ac.jp; YK, ykumagai@abs.agr.hokudai.ac.jp; TT, tagami@abs.agr.hokudai.ac.jp; AK, kimura@abs.agr.hokudai.ac.jp

Running title

Substrate recognition of catalytic subunit of ER GII

*Corresponding authors: E-mail, kimura@abs.agr.hokudai.ac.jp. (AK);

Email: okuyama@abs.agr.hokudai.ac.jp (MO)

[†]The first two authors contributed equally to this work.

Funding

This work was supported by JSPS KAKENHI Grant Numbers 22780082 and 26450114.

Abstract

The recombinant catalytic α -subunit of *N*-glycan processing glucosidase II from *Schizosaccharomyces pombe* (SpGII α) was produced in *Escherichia coli*. The recombinant SpGII α exhibited quite low stability, with a reduction in activity to < 40% after 2-days preservation at 4°C, but the presence of 10% (v/v) glycerol prevented this loss of activity. SpGII α , a member of the glycoside hydrolase family 31 (GH31), displayed the typical substrate specificity of GH31 α -glucosidases. The enzyme hydrolyzed not only α -(1 $\rightarrow$ 3)- but also α -(1 $\rightarrow$ 2)-, α -(1 $\rightarrow$ 4)- and α -(1 $\rightarrow$ 6)-glucosidic linkages, and *p*-nitrophenyl α -glucoside. SpGII α displayed most catalytic properties of glucosidase II. Hydrolytic activity of the terminal α -glucosidic residue of Glc₂Man₃-Dansyl was faster than that of Glc₁Man₃-Dansyl. This catalytic α -subunit also removed terminal glucose residues from native *N*-glycans (Glc₂Man₉GlcNAc₂ and Glc₁Man₉GlcNAc₂) although the activity was low.

Key words: catalytic α -subunit of ER glucosidase II; heterologous expression; glycoside hydrolase family 31; substrate specificity

Abbreviations

cleavage-1, conversion of G2M9 to G1M9; cleavage-2, conversion of G1M9 to M9; BSA, bovine serum albumin; ER, endoplasmic reticulum; GH31, glycoside hydrolase family 31; GI, glucosidase I; GII, glucosidase II; GII α , α -subunit of glucosidase II; GII β , β -subunit of glucosidase II; G1M3, Glc₁Man₃; G2M3, Glc₂Man₃; G1M9, Glc₁Man₉GlcNAc₂; G1M9-PA, pyridylaminated G1M9; G2M9, Glc₂Man₉GlcNAc₂; G3M9, Glc₃Man₉GlcNAc₂; G3M9-PA, pyridylaminated G1M9; HPLC, high-performance liquid chromatography; IPTG, isopropyl β -thiogalactopyranoside; M9, Man₉GlcNAc₂; pNPG, *p*-nitrophenyl α -glucopyranoside; ScGI, glucosidase I of *Saccharomyces cerevisiae*; SpGII α , α -subunit of glucosidase II of *Schizosaccharomyces pombe*.

Introduction

Secretory proteins of eukaryotic cells are *N*-glycosylated in the endoplasmic reticulum (ER). *N*-Glycosylation is initiated with the transfer of the precursor glycan, Glc₃Man₉GlcNAc₂ (G3M9), to an Asn residue in the motif Asn-Xaa-Ser/Thr of nascent polypeptides. Mannoses in G3M9 are organized in a 3'-trimannosyl branch (arm A) and a 6'-petamannosyl branch (composed of arms B and C), with the former type capped by three glucose residues. While still in ER, terminal glucoses in arm A and mannose in arm B are removed by glycosidases. The removal of glucose residues is catalyzed by glucosidases I and II (GI and GII, respectively), and the removal of the mannose residue is catalyzed by ER mannosidase. GI specifically trims the outermost α -(1→2)-linked glucose residue of G3M9 to produce Glc₂Man₉GlcNAc₂ (G2M9). Subsequently, GII removes a further two α -(1→3)-linked glucose residues in succession: cleavage-1, conversion from G2M9 to Glc₁Man₉GlcNAc₂ (G1M9); and cleavage-2, conversion from G1M9 to Man₉GlcNAc₂ (M9). These cleavage processes by GII have important implications for folding and quality control of nascent glycoproteins.¹⁾

GII is a heterodimeric protein composed of tightly bound α - and β -subunits (GII α and GII β , respectively). GII α is the catalytic subunit and displays significant amino acid sequence similarity to glycoside hydrolase family 31 (GH31) α -glucosidases. GH31 α -glucosidases are widespread in many organisms and are believed to be involved in the degradation of starch and α -glucooligosaccharides (e.g., maltooligosaccharides and α -glucobioses). The catalytic domain of GH31 α -glucosidase adopts the (β / α)₈-barrel fold and the active site is located in a pocket at the C-terminus of the inner β -barrel.²⁻⁶⁾ Recently, the three-dimensional structures of GII α s from *Chaetomium thermophilum* and murine were determined.^{7,8)} The structures are almost identical to other GH31 α -glucosidases, except for an N-terminal segment. The structure of murine GII α contains a portion of the GII β .⁸⁾ This structure reveals that several

salt bridges and hydrogen bonds are associated with the formation of the GII heterodimer. GII β is necessary for solubility, stability, activity and localization of GII.^{9,10)} GII β is involved in the identification of the substrate because it recognizes the arms B and C by its mannose 6-phosphate receptor homolog domain (MRH domain).^{11,12)} Structural analysis of the MRH domain of *Schizosaccharomyces pombe* GII β indicates that a tyrosine residue is closely associated with the binding of the mannose residue.¹³⁾

GII α without GII β exhibits hydrolysis activity toward chromogenic substrates such as *p*-nitrophenyl α -glucopyranoside (pNPG), but not the glucosidic residues in the physiological glycans, G2M9 and G1M9. Moreover, there are a few reports that GII α can trim glucose residues in native glycans without GII β . GII α from *Saccharomyces cerevisiae* can catalyze cleavage-1, but not cleavage-2 without GII β .¹⁴⁾ *S. pombe* GII α (SpGII α) exhibits limited cleavage-1 activity *in vivo*, but not *in vitro*.¹²⁾ *Bombyx mori* GII α , a purified recombinant protein, shows weak cleavage-2 activity.¹⁵⁾ However, the detailed catalytic specificity of GII α toward the natural *N*-glycan remains unresolved.

In this study, we revealed the substrate recognition of SpGII α for the first time by successfully obtaining soluble recombinant SpGII α , and the stability of recombinant SpGII α was improved by coexistence with glycerol. Substrates used in this study were: i) substrates (maltooligosaccharides, α -glucobioses, and pNPG) of GH31 α -glucosidases, which share the common ancestral protein with GII α ; ii) dansyl substrates (G2M3-Dansyl and G1M3-Dansyl) mimicking arm A of *N*-glycan; and iii) natural *N*-glycan substrates (G2M9 and G1M9). SpGII α hydrolyzes the substrates of GH31 α -glucosidase with broad specificity. Dansyl substrates and natural *N*-glycan substrates are also cleaved even without GII β . However, the hydrolysis rate on *N*-glycan substrates is slow and this observation suggests that the assistance of GII β is required for proper activity.

Material and methods

Materials. Bacto yeast extract, bacto peptone and bacto yeast nitrogen base without amino acids and ammonium sulfate were purchased from BD biosciences (Sparks, MD, USA). α,α -Trehalose, pNPG and glycerol were from Nacalai Tesque (Kyoto, Japan). The maltose used was manufactured by Nihon Shokuhin Kako, Tokyo, Japan. Kojibiose, nigerose, maltotriose, maltotetraose, maltopentaose, maltohexaose and maltoheptaose were purchased from Wako Pure Chemical Industries (Osaka, Japan). Isomaltose was from Tokyo Chemical Industry (Tokyo, Japan). Pyridylaminated G1M9 and G3M9 (G1M9-PA and G3M9-PA) were purchased from Masuda Chemical industries Co., Ltd. (Kagawa, Japan). All chemicals used were of analytical grade unless otherwise noted.

Strains and vectors. *Escherichia coli* (*E. coli*) strain DH5 α was used for cloning, whereas BL21-CodonPlus (DE3)-RIL and -RP (Stratagene, La Jolla, CA, USA), and RosettaTM (DE3) (Novagen-Merck Millipore, Billerica, MA, USA) were used for recombinant protein expression. *E. coli* expression vectors, pCold I DNA and pET vectors (pET23d and pET41a), were purchased from Takara Bio (Otsu, Japan) and Novagen-Merck Millipore, respectively. *S. cerevisiae* NBRC 1136 (S288C) was from the National Institute of Technology and Evaluation Biological Resource Center, Kisarazu, Japan, and *Pichia pastoris* GS115 and the pPICZ α A plasmid vector were from Invitrogen (Thermo Fisher Scientific, Carlsbad, CA, USA). *S. pombe* AHU 3179 was kindly supplied by the Applied Microbiology Laboratory, Research Faculty of Agriculture, Hokkaido University, Sapporo, Japan.

Cloning of the gls2 gene and construction of its expression plasmid. The *gls2* gene (SPAC1002.03c) encoding SpGII α was amplified by PCR using genomic *S. pombe* DNA as the template using the two primers (sense_1 of 5'-TGTAACCTTCTCCCGGGAAGATTCC-

3' and antisense_1 of 5'-TTTCTCCATAAACGTTAAATATTGG-3') and thermostable KOD-plus-DNA polymerase (Toyobo, Osaka, Japan). The PCR product was cloned into pBluescript II SK(+) (Stratagene) to obtain SK-gls2. The recombinant SpGII α was produced as a fusion protein with a His₆-affinity tag at the C-terminus as follows. DNA encoding the predicted mature SpGII α (Ala26 to Val923) was amplified by PCR using SK-gls2 as the template with primers: sense_2 (5'-TGCCCCATGGCATTTTCGACATCAATTTAAA-3', the *Nco*I site is underlined) and antisense_2 (5'-AATCTCGAGAACCACAAAAAAGTTGTGGATT-3', the *Xho*I site is underlined). The presence and location of the signal peptide cleavage site was predicted by the Signal P server (<http://www.cbs.dtu.dk/services/SignalP/>).¹⁶⁾ The PCR product was digested with *Nco*I and *Xho*I, and then introduced into the *Nco*I–*Xho*I sites of pET23d to generate the pET23d-gls2 plasmid for recombinant SpGII α production.

Cloning of the CWH41 gene and construction of its expression plasmid. The expression vector of the CWH41 gene (YGL027C) encoding *S. cerevisiae* processing glucosidase I (ScGI) was constructed according to previous reports.^{17,18)} The enzyme was a truncated form (Glu33 to Phe833) with the N-terminal *S. cerevisiae* α -factor secretion signal and a C-terminal His₆-affinity tag. ScGI was produced in *P. pastoris* using pPICZ α A as follows. The CWH41 gene was amplified by PCR using genomic *S. cerevisiae* DNA as the template with primers: CWH41_F (5'-AAGTAGTGGATAATAACGGTTCAGG-3') and CWH41_R (5'-CTTACTAGTAAGCGTCCAAGGATGTTGAC-3', the *Spe*I site is underlined). The insertion of *Spe*I resulted in a substitution of an amino acid residue, Phe833→Leu. Primestart Max DNA polymerase (Takara Bio) was used as the thermostable DNA polymerase. The PCR product was cloned into pBluescript II SK(+) to obtain SK-CWH41. SK-CWH41 was digested by *Eco*RI and *Spe*I, cloned into the *Eco*RI–*Xho*I sites of pPICZ α A, and designated as pPICZ α A-CWH41 for the production of ScGI.

Production and purification of SpGIIa. *E. coli* BL21-CodonPlus (DE3)-RIL cells were transformed by pET23d-gls2. Transformants were selected on an LB agar plate (5 g/L NaCl, 10 g/L bacto tryptone, 5 g/L bacto yeast extract and 15 g/L agar), supplemented with 50 mg/L ampicillin and 30 mg/L chloramphenicol. A transformant was grown in 40 mL of LB liquid medium with 50 mg/L ampicillin and 30 mg/L chloramphenicol at 30°C overnight. The resultant 30 mL of overnight LB-culture was inoculated into TB medium (3 L; 12 g/L bacto tryptone, 24 g/L bacto yeast extract, 0.4% (v/v) glycerol, 170 mM KH₂PO₄ and 720 mM K₂HPO₄) supplemented with 50 mg/L ampicillin and cell culturing was continued at 37°C. When the OD₆₀₀ reached 0.7, the culture broth was cooled on ice for 30 min. Induction of the culture was achieved at 12°C for 48 h without the addition of isopropyl β-thiogalactopyranoside (IPTG). Bacterial cells, harvested by centrifugation (11,600 × *g*, 4°C, 10 min), were suspended in 150 mL of 20 mM sodium phosphate buffer (pH 7.5) containing 10% (v/v) glycerol (buffer-A) and disrupted by sonication. The cell-free extract, obtained by centrifugation (9,600 × *g*, 4°C, 10 min), was applied to a Co-chelating Sepharose Fast Flow column (1.5 cm I.D. × 4 cm, GE Healthcare, Buckinghamshire, United Kingdom) equilibrated with buffer-A containing 0.5 M NaCl (buffer-B). After thorough washing with buffer-B and 10 mM imidazole-containing buffer-B in this order, the adsorbed protein was eluted with a linear gradient of imidazole from 10 to 200 mM in buffer-B. The active fractions were loaded on to a Bio-Gel P6 fine column (3 cm I.D. × 28 cm, Bio-Rad, Richmond, CA) equilibrated with buffer-A. The desalted sample was then loaded onto a DEAE-Sepharose Fast Flow column (2.6 cm I.D. × 8.5 cm, GE Healthcare) equilibrated with buffer-A. After washing with buffer-A, the adsorbed protein was eluted with a linear gradient of NaCl from 0 to 1.0 M in buffer-A. The active fractions, desalted by the aforementioned procedure using Bio-Gel P6, were subjected to a second DEAE-Sepharose Fast Flow column chromatography (1.5 cm I.D.

× 6 cm) step. The adsorbed protein was eluted isocratically with 65 mM NaCl in buffer-A. The active fractions were pooled and desalted by Bio-Gel P6 using buffer-A. The purity of the protein was analyzed by SDS-PAGE. The concentration of the purified protein was estimated by amino acid analysis of the protein hydrolysate (6 M HCl, 110°C, 24 h) using JLC-500/V (JOEL, Tokyo, Japan) equipped with a ninhydrin-detection system. The extinction coefficient of 1 mg/mL purified SpGIIα at 280 nm was 2.27.

Production and purification of ScGI. Transformation of *P. pastoris* GS115 was performed as described by Lin-Cereghino *et al.*¹⁹⁾ pPICZαA-CWH41A was linearized with *SacI* and introduced into *P. pastoris* cells by electroporation. Transformants were selected on YPDSZ agar plates (10 g/L bacto yeast extract, 20 g/L bacto peptone, 2 g/L glucose, 1 M sorbitol, 100 mg/L Zeocin and 15 g/L agar). The selected transformant was grown in BMGY medium [3L; 10 g/L bacto yeast extract, 20 g/L bacto peptone, 3.4 g/L yeast nitrogen base without amino acids and ammonium sulfate, 10 g/L ammonium sulfate, 0.1 M potassium phosphate buffer (pH 6.0), 10 g/L glycerol and 0.4 mg/L biotin] at 30°C overnight. The cells collected by centrifugation at 4°C for 10 min (3,000 × *g*) were resuspended in 500 mL of BMMY medium that contains 1% (v/v) methanol instead of the glycerol of BMGY and incubated at 22°C for 96 h under vigorous shaking. One hundredth of the culture volume of methanol was added every 24 h. The supernatant was removed by centrifugation (11,600 × *g*, 4°C, 10 min) and its pH was adjusted to 7.5 by addition of 1 M NaOH with stirring. Debris were removed by centrifugation (11,600 × *g*, 4°C, 10 min) and the recovered supernatant was loaded onto a Ni-chelating Sepharose Fast Flow column (1.5 cm I.D. × 5 cm) equilibrated with 50 mM sodium phosphate buffer (pH 7.5) containing 0.5 M NaCl (buffer-C). After washing with buffer-C and then 5 mM imidazole-containing buffer-C, the adsorbed protein was eluted with 200 mM imidazole in buffer-C. The fractions containing purified ScGI, which was determined by SDS-

PAGE, were dialyzed against 20 mM potassium phosphate buffer (pH 6.8) containing 0.1 M NaCl and concentrated using Centriprep YM-50 centrifugal filter units (Novagen-Merck Millipore).

Standard enzyme assay. The activity of SpGII α was determined by measuring the increase of *p*-nitrophenol during hydrolysis of pNPG. A reaction mixture (50 μ L) consisting of an appropriate concentration of SpGII α , 2 mM pNPG, 40 mM MES-NaOH buffer (pH 6.5), 2% (v/v) glycerol and 0.2 mg/mL bovine serum albumin (BSA) was incubated at 30°C for 10 min. The reaction was stopped by mixing with two volumes of 1 M sodium carbonate. The amount of *p*-nitrophenol released was measured by absorption at 400 nm in a 1-cm cuvette, using a molar extinction coefficient of 5,560 M⁻¹ cm⁻¹. One unit of SpGII α activity was defined as the amount of enzyme that produced 1 μ mol *p*-nitrophenol per min under these conditions.

Optimum pH and stability to pH and heat. The optimum pH of the catalytic reaction was determined by measuring the hydrolytic rate at various pH values. The SpGII α concentration and buffer used were 8.2 μ g/mL and 80 mM Britton–Robinson buffer (pH 2.7–11.5), respectively. The other reaction conditions were the same as those of the standard assay method. For measurement of pH stability, SpGII α (82 μ g/mL) was incubated in 10-fold-diluted Britton–Robinson buffer (pH 3.3 to 10.5) containing 0.1% BSA and 10% (v/v) glycerol at 4°C for 24 h, followed by measurement of the residual activity under the standard assay conditions. The stable region was defined as the pH range exhibiting residual activity of > 90%. For measurement of thermal stability, SpGII α (2.7 μ g/mL) in 67 mM MES-NaOH buffer (pH 6.5) containing 0.1% BSA and 10% (v/v) glycerol was kept at 26 to 55°C for 15 min, followed by measurement of their residual activities under the standard assay conditions.

The stable region was defined as the temperature range exhibiting residual activity of > 90%.

Effect of additives on enzyme stability. SpGII α in 20 mM HEPES-NaOH (pH 7.5) was incubated with NaCl (0.05, 0.2, or 0.5 M), CaCl₂ (0.05 or 0.2 M), EDTA·2Na (0.1, 0.5, or 1 mM), glycerol [2, 10, or 20% (v/v)], or Triton X-100 [0.5 or 1% (v/v)] at 4°C. At the indicated time, an aliquot of the mixture was taken and diluted with 40 mM MES-NaOH (pH 6.5) containing 0.1% BSA and 10% (v/v) glycerol. The residual activity was evaluated by hydrolysis of 0.2% maltose at 30°C. The hydrolysis reaction was terminated by mixing with two volumes of 2 M Tris-HCl buffer (pH 7.0) and liberated glucose was measured with the Glucose C II-Test Wako (Wako Pure Chemical Industries).

Effect of various salts on the enzyme reaction. The enzyme reactions were performed in 40 mM MES-NaOH (pH 6.5) containing 1.6 μ g/mL SpGII α , 0.2% maltose, 2% (v/v) glycerol and 0.2 mg/mL BSA at 30°C for 10 min by adding salt (0–40 mM KCl, MgCl₂, or various sodium salts). The concentration of liberated glucose was measured, as described above.

Kinetic parameters for hydrolysis of various substrates. Kinetic parameters for hydrolysis of pNPG, nigerose, kojibiose and a series of maltooligosaccharides were calculated from the initial rates at various substrate concentrations by fitting to the Michaelis–Menten equation using KaleidaGraph 3.6J (Synergy Software, Reading, PA, USA). Substrate concentrations were as follows: 0.4–15 mM for pNPG, 0.5–10 mM for nigerose, 1–80 mM for kojibiose, 3.2–80 mM for maltose, 2.0–64 mM for maltotriose, 2.0–80 mM for maltotetraose and maltopentaose, 2.0–40 mM for maltohexaose and 0.8–40 mM for maltoheptaose. Since the K_m value for isomaltose was so large, its k_{cat}/K_m value was determined from the slope of Lineweaver-Burk plots at concentrations from 16 to 80 mM.

The amount of *p*-nitrophenol released from pNPG was measured by absorption at 400 nm, as described above. For other substrates, the liberated glucose was measured as described above.

Measurement of the hydrolytic rates of G2M3-Dansyl and G1M3-Dansyl. The hydrolytic rates for G2M3-Dansyl and G1M3-Dansyl, both of which were prepared as described previously,^{20,21)} and nigerose were determined by measuring the liberated glucose concentration. A reaction mixture (180 μ L) consisting of SpGII α (0.35 μ g/mL for nigerose and G2M3-Dansyl, 1.8 μ g/mL for G1M3-Dansyl), 0.4 mM substrate, 40 mM MES-NaOH buffer (pH 6.5), 2% (v/v) glycerol and 0.2 mg/mL BSA was incubated at 30°C. At 5, 10, 15 and 20 min, an aliquot of the reaction mixture was taken and heated at 100°C for 3 min to terminate the reaction. The glucose concentration was measured by high-performance anion exchange chromatography [Dionex ICS-3000 system with pulsed amperometric detection (Dionex/Thermo Fisher Scientific, Idstein, Germany)]. The analytical column (CarboPac PA1, 4 mm I.D. $\times$ 250 mm, Dionex/Thermo Fisher Scientific) was equilibrated by 100 mM NaOH at a flow speed of 0.8 mL min⁻¹. Separations were performed with the 6 min-linear gradient of 100–640 mM NaOH for nigerose and with the 42 min-linear gradient of 0–600 mM sodium acetate in 100 mM NaOH for dansyl oligosaccharides. The NaOH solution was prepared from super special grade 50% NaOH (Wako Pure Chemical Industries). Standards and samples contained 0.1 mM fructose as an internal reference. Peak areas and retention times of eluted carbohydrates were evaluated with the Chromeleon software (Dionex/Thermo Fisher Scientific). The concentration of glucose in each sample was calculated from its peak area using a calibration curve prepared from standard glucose and internal fructose.

Detection of deglycosylation of pyridylaminated oligosaccharides. A reaction mixture (100 μ L) consisting of SpGII α (1.04 μ g/mL), 0.4 μ M substrate (G1M9-PA or G3M9-PA), 40

mM MES-NaOH buffer (pH 6.5), 2% (v/v) glycerol and 0.2 mg/mL BSA was incubated at 30°C. For ScGI-catalyzed hydrolysis of the outermost glucose residue of G3M9-PA, 2.1 µg/mL ScGI was used under the same reaction conditions as SpGIIα. The reaction was terminated by incubation at 95°C for 5 min. The oligosaccharides were separated by high-performance liquid chromatography (HPLC) using an Asahipack NH2-50-4E column (4.6 mm I.D. × 250 mm, Shodex, Tokyo, Japan) at 40°C. The elution was done at a flow speed of 0.8 mL min⁻¹ with a linear gradient of acetonitrile from 68 to 34% in 0.3% ammonium acetate buffer (pH 7.0). Pyridylaminated oligosaccharides were monitored by the fluorescence signal (excitation wave length, 310 nm; emission wavelength, 380 nm) using a fluorescence detector FP-2020 Plus (Jasco, Tokyo, Japan).

Results and discussion

Production and characterization of SpGIIα

The expression conditions for obtaining soluble recombinant SpGIIα in *E. coli* were examined. The DNA coding for the mature SpGIIα was in-frame inserted into different expression plasmids (pET23d, pCold I, or pET41a) and enzyme production was tested using three different *E. coli* strains [Rosetta (DE3), BL21-CodonPlus (DE3)-RIL, or -RP]. In most expression systems tested, recombinant SpGIIα was produced as an insoluble form; however, the BL21-CodonPlus (DE3)-RIL strain transformed with the pET23d was the best combination for soluble protein production. Furthermore, cultivation in TB medium supplemented with 50 µg/mL ampicillin at 12°C without IPTG induction was also effective for production of soluble SpGIIα.

The induced cells cultivated under the optimum conditions for 48 h were disrupted in 20 mM sodium phosphate buffer (pH 7.5) including 10% glycerol, and the crude extract obtained contained 11.5 U per 1 L culture. The presence of glycerol in both the purification and storage

275 buffers was essential for preventing SpGII α from inactivation, as described below.
276 Recombinant SpGII α produced by *E. coli* was purified using cobalt-affinity and anion-
277 exchange chromatographies. The final desalting column step yielded 1.8 mg of
278 electrophoretically homogeneous SpGII α (Fig. 1) from 3 L of culture with specific activity
279 of 4.0 U/mg.

280 Figure 2 shows the effects of various additives on the stability of SpGII α . The residual
281 activity of SpGII α without any additive decreased to < 40% of the initial activity after 2 d
282 (Fig. 2A), and to < 1% after 23 d (Figs. 2B, C). Inhibitory effect was observed when the
283 protein was incubated with NaCl, CaCl₂ and Triton X-100 (Fig. 2A). In particular, addition
284 of CaCl₂ and Triton X-100 induced substantially activity loss. Both glycerol and EDTA as
285 additives improved the stability of recombinant SpGII α (Figs. 2B, C). It is hard to understand
286 how EDTA contributes to the stability of SpGII α . It is anticipated that EDTA removes
287 contaminating divalent ions such as the calcium ion, which adversely affects the SpGII α
288 stability. The addition of $\geq 10\%$ (v/v) glycerol was more effective in stabilizing SpGII α than
289 EDTA, with the enzyme maintaining > 90% activity after 92 d (Fig. 2C). Glycerol is known
290 to induce protein compaction, reduce protein flexibility, stabilize specific partially unfolded
291 intermediates and affect protein aggregation,²²⁾ resulting in its wide use as a protein stabilizer.
292 Based on the effect of glycerol on protein stability, Vangenende *et al.* proposed that glycerol
293 prevents protein aggregation thorough preferential interaction with hydrophobic surface
294 regions.²²⁾ In nature, GII β forms a heterodimer with GII α . Thus, glycerol was anticipated to
295 affect the exposed dimer interface of GII β lacking the GII α subunit, because generally an
296 oligomer interface is hydrophobic. However, the three-dimensional structure of murine GII α
297 complexed with an N-terminal portion of GII β showed that the interface area is not that
298 hydrophobic,⁸⁾ suggesting less hydrophobicity of the equivalent regions of SpGII α . Therefore,
299 it is possible that SpGII α has other surface exposed hydrophobic regions where glycerol

interacts to prevent instability of the protein. Nonetheless, stabilization conditions found in this study allow us to perform purification and characterization of SpGII α .

The effects of various salts on the activity were investigated by measuring the initial hydrolytic velocity on maltose using 2% (v/v) glycerol. NaCl, NaH₂PO₄, Na₂SO₄, NaNO₃, KCl, MgCl₂, or EDTA·2Na with concentrations up to 40 mM did not affect activity, while the addition of 40 mM CaCl₂ decreased to 63% of the original initial velocity. The calcium ion should be harmful to both activity and stability of SpGII α . As shown in Fig. 2A, 50 mM CaCl₂ causes its inactivation within 2 d, which shows a negative effect of this salt on its stability.

The effects of pH and temperature on the activity were examined in the presence of 10% (v/v) glycerol. The pH optimum was 6.5 and SpGII α was stable between pH 6.2 and 9.1. SpGII α was thermally stable to 40°C with complete inactivation observed at 55°C (15 min treatment). In the absence of glycerol, this thermal stability was reduced to < 30% of the initial activity observed when incubated in the presence of glycerol for 15 min at 40°C.

Substrate specificity of SpGII α : substrates for GH31 α -glucosidase

Substrate specificity of GII α is of interest, because this protein has a common ancestor with GH31 α -glucosidases, which can hydrolyze α -(1→2)-, α -(1→3)-, α -(1→4)- and α -(1→6)-glucosidic linkages.^{23–27} By primarily exhibiting specificity to α -(1→4)-glucosidic linkage, GH31 α -glucosidases are mostly linked with metabolic pathways of starch and maltooligosaccharides degradation.²⁸ While GII α plays a different role in biological processes when compared with that of GH31 α -glucosidases, both enzymes might display similar substrate specificities, because catalytic function of a protein is generally conserved during molecular evolution. Substrate specificity of SpGII α towards common substrates of α -glucosidases was evaluated by determining kinetic parameters (Table 1). SpGII α hydrolyzed all substrates tested, proving broader specificity than ER-resident processing GI, which

recognizes the outermost α -1,2-glucosy residue in the *N*-glycan so strictly that kojibiose acts as an inhibitor.^{29,30)} Among the disaccharides, nigerose was the best substrate with the highest k_{cat}/K_m value (Table 1). The K_m values for maltose and kojibiose were around 10 times as large as that for nigerose, and that for isomaltose could not be obtained due to unsaturated curve in $s - v$ plots up to 80 mM isomaltose, indicating that SpGII α displays weaker binding to α -(1 $\rightarrow$ 4)-, α -(1 $\rightarrow$ 2)- and α -(1 $\rightarrow$ 6)-glucosidic linkages. This specificity is similar to the GH31 α -1,3-glucosidase, which hydrolyzes α -(1 $\rightarrow$ 4)-, α -(1 $\rightarrow$ 2)- and α -(1 $\rightarrow$ 3)-glucosidic linkages together with weak specificity towards the α -(1 $\rightarrow$ 6)-glucosidic linkage.³¹⁻³³⁾ This enzyme has been found in some microorganisms and is hypothesized to be involved in a metabolic pathway rather than the trimming of sugar chains.³¹⁻³³⁾ Hydrolysis of pNPG was characterized by low k_{cat} and K_m values, which are common features of the GH31 α -glucosidases and the α -1,3-glucosidase.^{23-25,28,31,32)} These results demonstrate that SpGII α conserves substrate specificity while its localization and physiological role are different. The higher k_{cat}/K_m value for maltose than that for malto-triose, -tetraose and -pentaose (Table 1) indicates that SpGII α prefers maltose, and this observation is consistent with the structure of the active-site pocket of GII α being capable of accommodating only disaccharides.^{7,8)} The k_{cat}/K_m values for malto-hexaose and -heptaose increased 3.0 to 5.2 times over -triose, -tetraose and -pentaose. The GH31 sugar beet α -glucosidase possesses the machinery that accommodates long-chain substrates,³⁴⁾ but SpGII α has no equivalent element. Thus, it is difficult to explain the increase in specificity toward longer-chain substrates.

Substrate specificity of SpGII α : N-glycan-relating substrates

We measured the hydrolytic rates on 0.4 mM G2M3-Dansyl and G1M3-Dansyl, which mimic arm A of *N*-glycan, together with 0.4 mM nigerose for comparison. SpGII α was able to hydrolyze the α -glucosidic linkage in G2M3-Dansyl and G1M3-Dansyl with hydrolytic

rates of 1.23 and 0.274 $\mu\text{mol min}^{-1} \text{mg}^{-1}$, respectively, although these values were slower than that of nigerose (2.42 $\mu\text{mol min}^{-1} \text{mg}^{-1}$). The difference in hydrolytic rates between G2M3-Dansyl and G1M3-Dansyl is in agreement with the observation that GII catalyzes the hydrolysis of the $\alpha\text{-Glu-(1}\rightarrow\text{3)-Glc}$ linkage (cleavage-1) faster than that of the $\alpha\text{-Glu-(1}\rightarrow\text{3)-Man}$ linkage (cleavage-2).^{35,36)} The difference in the catalytic rate between both cleavages was accounted for by binding of GII β to mannose residues in arms B and C.^{35–37)} Alternatively, recent crystal structure analyses proposed that an interaction between OH-2 of a mannose residue and the carboxy group of the acid/base catalyst of GII α is responsible for the slow reaction rate of cleavage-2.^{7,8)} Our SpGII α , devoid of GII β , shows cleavage-2, corroborating the latter explanation.

GII α without the GII β is able to hydrolyze much smaller substrates, such as pNPG, but not able to deglycosylate G2M9 and G1M9. These distinct results are also reported as follows. Stigliano *et al.* demonstrated that SpGII α was able to degrade G2M9 and G1M9 *in vivo*.¹²⁾ GII α in *S. cerevisiae* catalyzes cleavage-1 but not cleavage-2.¹⁴⁾ GII α from *B. mori* weakly hydrolyzes G1M9 *in vitro*.¹⁵⁾ According to these observations, GII α does not seem to require GII β to hydrolyze the $\alpha\text{-glucosidic}$ linkages in *N*-glycan. We thus examined hydrolysis of G1M9-PA and G3M9-PA using the purified SpGII α to evaluate its hydrolytic ability toward *N*-glycan. SpGII α weakly but clearly catalyzed cleavage-2 to generate M9-PA from G1M9-PA (Fig. 3A). After 1 h of the reaction, the peak of M9-PA appeared, and then increased to the same level of G1M9-PA after the reaction had proceeded for 12 h. As shown in Fig. 3B, SpGII α cannot degrade G3M9-PA (Fig. 3B), which coincides with the general specificity of GII. Conversion from G3M9 to G2M9 is known to be catalyzed by GI, which removes the outermost $\alpha\text{-Glc-(1}\rightarrow\text{2)-Glc}$ linkage of G3M9. ScGI rapidly hydrolyzed G3M9-PA to produce G2M9-PA (Fig. 3C). Figure 3D shows the hydrolysis of G3M9-PA in the presence of SpGII α and ScGI, which demonstrates that G2M9-PA, generated by the rapid reaction of ScGI until

1 h, is degraded to G1M9-PA and M9-PA at 12 h. These results indicate that the purified SpGII α catalyzes the trimming of glucose residues from G2M9-PA and G1M9-PA without the assistance of GII β . Although the purified SpGII α certainly catalyzes cleavage-1 and -2, their reaction rates are not high (Figs. 3A, 3D). According to the result of Watanabe *et al.*,³⁷⁾ 50% conversion of 25 μ M G1M9 to M9 requires 45 min by a membrane fraction containing GII, even though the accurate concentration of GII was unknown. The present study provides a more quantitative analysis in which 9.6 nM SpGII α (1.0 μ g/mL SpGII α) and a 12-h reaction yield approximately 50% conversion of 0.4 μ M G1M9 (Fig. 3A). The low hydrolysis of G1M9 and G2M9 by SpGII α is probably because of insufficient binding energy for efficiently hydrolyzing these natural substrates, because SpGII α can only hold the small terminal moiety of the comparatively large structure of G1M9 or G2M9. The active-site pocket of GII α s has the gourd-shaped bilocular pocket, which can accommodate only disaccharides,^{7,8)} and GII β would be necessary to stabilize the enzyme-substrate complex. Olson *et al.* demonstrates an importance of the MRH domain of GII β to the hydrolytic activity of GII α through a recognition of arms B and C of the natural *N*-glycan.¹³⁾ We need to direct our future studies to understanding the effect of GII β on the catalytic properties including substrate recognition.

Author contributions

Conceived and designed the experiments: M. Okuyama. Performed the experiments: M. Okuyama, M. Miyamoto, I. Matsuo, S. Iwamoto, R. Serizawa and M. Tanuma. Contributed reagents/materials/analysis tools: M. Ma, P Klahan and Y. Kumagai. Wrote the paper: M. Okuyama, T. Tagami and A. Kimura.

Disclosure statement

No potential conflict of interest was reported by the authors.

400

401 **Acknowledgements**

402 We thank the staff of the Instrumental Analysis Division of the Creative Research
403 Institution at Hokkaido University for the amino acid analysis, and Applied Microbiology
404 Laboratory, Research Faculty of Agriculture, Hokkaido University for the supply of *S. pombe*
405 AHU 3179. We also thank Dr. S. Fukiya of the Laboratory of Microbial Physiology, Research
406 Faculty of Agriculture, Hokkaido University for the analysis of pyridylaminated
407 oligosaccharides. This work was supported by JSPS KAKENHI Grant Numbers 22780082
408 and 26450114.

409

References

1. Parodi AJ. Role of *N*-oligosaccharide endoplasmic reticulum processing reactions in glycoprotein folding and degradation. *Biochem. J.* 2000;348:1–13.
2. Ernst HA, Lo Leggio L, Willemoes M, et al. Structure of the *Sulfolobus solfataricus* α -glucosidase: Implications for domain conservation and substrate recognition in GH31. *J. Mol. Biol.* 2006;358:1106–1124.
3. Sim L, Quezada-Calvillo R, Sterchi EE, et al. Human intestinal maltase-glucoamylase: crystal structure of the N-terminal catalytic subunit and basis of inhibition and substrate specificity. *J. Mol. Biol.* 2008;375:782–792.
4. Tan K, Tesar C, Wilton R, et al. Novel α -glucosidase from human gut microbiome: substrate specificities and their switch. *FASEB J.* 2010;24:3939–3949.
5. Ren L, Qin X, Cao X, et al. Structural insight into substrate specificity of human intestinal maltase-glucoamylase. *Protein Cell.* 2011;2:827–836.
6. Tagami T, Yamashita K, Okuyama M, et al. Molecular basis for the recognition of long-chain substrates by plant α -glucosidases. *J. Biol. Chem.* 2013;288:19296–19303.
7. Satoh T, Toshimori T, Yan G, et al. Structural basis for two-step glucose trimming by glucosidase II involved in ER glycoprotein quality control. *Sci. Rep.* 2016;6:20575.
8. Caputo AT, Alonzi DS, Marti L, et al. Structures of mammalian ER α -glucosidase II capture the binding modes of broad-spectrum iminosugar antivirals. *Proc. Natl. Acad. Sci. USA.* 2016;113:E4630–E4638.
9. Pelletier MF, Marcil A, Sevigny G, et al. The heterodimeric structure of glucosidase II is required for its activity, solubility, and localization *in vivo*. *Glycobiology.* 2000;10:815–827.

- 435 10. Trembl K, Meimaroglou D, Hentges A, et al. The α - and β -subunits are required for
436 expression of catalytic activity in the hetero-dimeric glucosidase II complex from
437 human liver. *Glycobiology*. 2000;10:493–502.
- 438 11. Totani K, Ihara Y, Matsuo I, et al. Substrate specificity analysis of endoplasmic
439 reticulum glucosidase II using synthetic high mannose-type glycans. *J. Biol. Chem.*
440 2006;281:31502–31508. .
- 441 12. Stigliano ID, Caramelo JJ, Labriola CA, et al. Glucosidase II β subunit modulates *N*-
442 glycan trimming in fission yeasts and mammals. *Mol. Biol. Cell*. 2009;20:3974–
443 3984. .
- 444 13. Olson LJ, Orsi R, Peterson FC, et al. Crystal structure and functional analyses of the
445 lectin domain of glucosidase II: Insights into oligomannose recognition.
446 *Biochemistry*. 2015;54:4097–4111.
- 447 14. Wilkinson BM, Purswani J, Stirling CJ. Yeast GTB1 encodes a subunit of
448 glucosidase II required for glycoprotein processing in the endoplasmic reticulum. *J.*
449 *Biol. Chem.* 2006;281:6325–6333.
- 450 15. Watanabe S, Kakudo A, Ohta M, et al. Molecular cloning and characterization of the
451 α -glucosidase II from *Bombyx mori* and *Spodoptera frugiperda*. *Insect Biochem.*
452 *Mol. Biol.* 2013;43:319–327.
- 453 16. Petersen TN, Brunak S, von Heijne G, et al. SignalP 4.0: discriminating signal
454 peptides from transmembrane regions. *Nat. Methods*. 2011;8:785–786.
- 455 17. Faridmoayer A, Scaman C. Truncations and functional carboxylic acid residues of
456 yeast processing α -glucosidase I. *Glycoconj. J.* 2007;24:429–437
- 457 18. Barker MK, Wilkinson BL, Faridmoayer A, et al. Production and crystallization of
458 processing α -glucosidase I: *Pichia pastoris* expression and a two-step purification
459 toward structural determination. *Protein Expr. Purif.* 2011;79:96–101.

19. Lin-Cereghino J, Wong WW, Xiong S, et al. Condensed protocol for competent cell preparation and transformation of the methylotrophic yeast *Pichia pastoris*. Biotechniques. 2005;38:44–48.
20. Iino K, Iwamoto S, Kasahara Y, et al. Facile construction of 1,2-cis glucosidic linkage using sequential oxidation–reduction route for synthesis of an ER processing α -glucosidase I substrate, Tetrahedron Lett. 2012;53:4452–4456.
21. Iwamoto S, Kasahara Y, Kamei K, et al. Measurement of endo- α -mannosidase activity using a fluorescently labeled oligosaccharide derivative. Biosci. Biotechnol. Biochem. 2014;78:927–936.
22. Vagenende V, Yap MGS, Trout BL. Mechanisms of protein stabilization and prevention of protein aggregation by glycerol. Biochemistry. 2009;48:11084–11096.
23. Okuyama M, Tanimoto Y, Ito T, et al. Purification and characterization of the hyperglycosylated extracellular α -glucosidase from *Schizosaccharomyces pombe*. Enzyme Microb. Technol. 2005;37:472–480.
24. Nakai H, Ito T, Hayashi M, Kamiya K, et al. Multiple forms of α -glucosidase in rice seeds (*Oryza sativa* L., var Nipponbare). Biochimie. 2007;89:49–62.
25. Sato F, Okuyama M, Nakai H, et al. Glucoamylase originating from *Schwanniomyces occidentalis* is a typical α -glucosidase. Biosci. Biotechnol. Biochem. 2005;69:1905–1913.
26. Tagami T, Okuyama M, Nakai H, et al. Key aromatic residues at subsites + 2 and + 3 of glycoside hydrolase family 31 α -glucosidase contribute to recognition of long-chain substrates. Biochim. Biophys. Acta. 2013;1834:329–335.
27. Saburi W, Okuyama M, Kumagai Y, et al. Biochemical properties and substrate recognition mechanism of GH31 α -glucosidase from *Bacillus* sp. AHU 2001 with broad substrate specificity. Biochimie. 2015;108:140–148.

- 485 28. Frandsen TP, Svensson B. Plant α -glucosidases of the glycoside hydrolase family 31.
486 Molecular properties, substrate specificity, reaction mechanism, and comparison
487 with family members of different origin. *Plant. Mol. Biol.* 1998;37:1–13.
- 488 29. Dhanawansa R, Faridmoayer A, van der Merwe G, et al. Overexpression,
489 purification, and partial characterization of *Saccharomyces cerevisiae* processing α -
490 glucosidase I. *Glycobiology*. 2002;12:229–234. .
- 491 30. Miyazaki T, Matsumoto Y, Matsuda K, et al. Heterologous expression and
492 characterization of processing α -glucosidase I from *Aspergillus brasiliensis* ATCC
493 9642. *Glycoconj. J.* 2011;28:563–571.
- 494 31. Yamamoto T, Unno T, Watanabe Y, et al. Purification and characterization of
495 *Acremonium implicatum* α -glucosidase having regioselectivity for α -1,3-glucosidic
496 linkage. *Biochim. Biophys. Acta*. 2004;1700:189–198.
- 497 32. Kang M-S, Okuyama M, Mori H, et al. The first α -1,3-glucosidase from bacterial
498 origin belonging to glycoside hydrolase family 31. *Biochimie*. 2009;91:1434–1442.
- 499 33. Song KM, Okuyama M, Kobayashi K, et al. Characterization of a glycoside
500 hydrolase family 31 α -glucosidase involved in starch utilization in *Podospora*
501 *anserina*. *Biosci. Biotechnol. Biochem.* 2013;77:2117–2124.
- 502 34. Tagami T, Yamashita K, Okuyama M, et al. Structural advantage of sugar beet α -
503 glucosidase to stabilize the Michaelis complex with long-chain substrate. *J. Biol.*
504 *Chem.* 2015;290:1796–1803.
- 505 35. Kaushal GP, Pastuszak I, Hatanaka K, et al. Purification to homogeneity and
506 properties of glucosidase II from mung bean seedlings and suspension-cultured
507 soybean cells. *J. Biol. Chem.* 1990;265:16271–16279.
- 508 36. Totani K, Ihara Y, Matsuo I, et al. Effects of macromolecular crowding on
509 glycoprotein processing enzymes. *J. Am. Chem. Soc.* 2008;130:2101–2107.

- 510 37. Watanabe T, Totani K, Matsuo I, et al. Genetic analysis of glucosidase II β -subunit
511 in trimming of high-mannose-type glycans. *Glycobiology*. 2009;19:834–840.
- 512 38. Deprez P, Gautschi M, Helenius A. More than one glycan is needed for ER
513 glucosidase II to allow entry of glycoproteins into the calnexin/calreticulin cycle.
514 *Mol. Cell*. 2005;19:183–195.
515

Table

Table 1. Kinetic parameters of SpGII α hydrolysis of GH31 α -glucosidase substrates.

Substrate	K_m (mM)	k_{cat} (sec ⁻¹)	k_{cat}/K_m (sec ⁻¹ M ⁻¹)
<i>p</i> NPG	3.62 $\pm$ 0.02	0.0150 $\pm$ 0.0001	4.15
Nigerose	2.13 $\pm$ 0.06	0.102 $\pm$ 0.001	48.1
Kojibiose	21.0 $\pm$ 1.6	0.0575 $\pm$ 0.001	2.75
Isomaltose	N.D.	N.D.	0.014 $\pm$ 0.007
Maltose	18.0 $\pm$ 0.1	0.241 $\pm$ 0.007	13.4
Maltotriose	16.0 $\pm$ 0.3	0.0343 $\pm$ 0.0001	2.15
Maltotetraose	28.5 $\pm$ 2.5	0.0477 $\pm$ 0.0012	1.68
Maltopentaose	26.7 $\pm$ 0.2	0.0367 $\pm$ 0.0004	1.38
Maltohexaose	15.3 $\pm$ 0.6	0.109 $\pm$ 0.001	7.15
Maltoheptaose	15.0 $\pm$ 0.2	0.0965 $\pm$ 0.0007	6.43

N.D.: Values could not be determined because s-v plots were not saturated.

Figure captions

Fig. 1. Expression and purification of recombinant SpGII α .

Notes: Expression and purification of GII α were analyzed by 10% SDS–PAGE. Lane 1, IPTG-induced whole *E. coli* cells harboring pET23d-gls2; lane 2, supernatant from cell lysate of lane 1; lane 3, eluate from Co²⁺ affinity chromatography; lane 4, Bio-Gel P6 chromatography (1st); lane 5, anion exchange chromatography (1st); lane 6, Bio-Gel P6 chromatography (2nd); lane 7, anion exchange chromatography (2nd); lane 8, Bio-Gel P6 chromatography (3rd); lane M, marker proteins with molecular masses (kDa) shown at the side.

Fig. 2. Effect of additives on long-term stability of SpGII α .

Notes: (A) Residual activity of GII α was measured after incubation with various additives at 4 °C for 2 days. (B) Time course of residual activity of GII α in the presence of 0.1 mM (○), 0.5 mM (■) and 1 mM (□) EDTA·2Na. (C) Time course of residual activity of GII α in the presence of 2% (○), 10% (■), and 20% (□) glycerol. The residual activity without any additives is represented by closed circles (●) in both (B) and (C).

Fig. 3. HPLC analysis of the hydrolysates of (A) G1M9 by SpGII α , (B) G3M9 by SpGII α , (C) G3M9 by ScGI, and (D) G3M9 by ScGI and SpGII α .

Notes: The reaction mixture samples were taken at 0, 1 and 12 h and were analyzed by HPLC. Symbol used for the structure formulae: ▲, glucose; ○, mannose; and ■, *N*-acetylglucosamine.




Figure 1. Okuyama et al




Figure 2. Okuyama et al




Figure 3. Okuyama et al