



HOKKAIDO UNIVERSITY

Title	Association of perfluoroalkyl substances exposure in utero with reproductive hormone levels in cord blood in the Hokkaido Study on Environment and Children's Health
Author(s)	Itoh, Sachiko; Araki, Atsuko; Mitsui, Takahiko et al.
Citation	Environment International, 94, 51-59 https://doi.org/10.1016/j.envint.2016.05.011
Issue Date	2016-09
Doc URL	https://hdl.handle.net/2115/71600
Rights	© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	ENVINT-D-15-01711R2(3)(1)-2.pdf



**Association of perfluoroalkyl substances exposure in utero
with reproductive hormone levels in cord blood in the
Hokkaido Study on Environment and Children's Health**

Sachiko Itoh,^a Atsuko Araki,^a Takahiko Mitsui,^b Chihiro Miyashita,^a Houman
Goudarzi,^a Seiko Sasaki,^c Kazutoshi Cho,^d Hiroyuki Nakazawa,^e Yusuke Iwasaki,^e
Nobuo Shinohara,^f Katsuya Nonomura,^{f,g} Reiko Kishi,^{a*}

^a Center for Environmental Health and Sciences, Hokkaido University, Sapporo, Japan

^b Department of Urology, Hokkaido University Hospital, Sapporo, Japan

^c Department of Public Health, Hokkaido University Graduate School of Medicine,
Sapporo, Japan

^d Department of Obstetrics and Gynecology, Hokkaido University Hospital, Sapporo,
Japan

^e Department of Analytical Chemistry, Faculty of Pharmaceutical Sciences, Hoshi
University, Tokyo, Japan

^f Department of Renal and Genitourinary Surgery, Hokkaido University Graduate
School of Medicine, Sapporo, Japan

^g Kushiro Rosai Hospital, Kushiro, Japan

***Corresponding author:**

Reiko Kishi

Center for Environmental Health and Sciences, Hokkaido University

North 12 West 7, Kita-ku, Sapporo 060-0812, Japan

Tel: +81 11 706 4746

E-mail: rkishi@med.hokudai.ac.jp

Running title: PFASs and cord blood reproductive hormone levels

Acknowledgements

We thank all the mothers and children who participated in the study, and all staff at Sapporo Toho Hospital. This study was financially supported by a Grant-in Aid from the Japanese Ministry of Health, Labour and Welfare, Health and Labour Sciences Research Grants; Grants in Aid of Scientific Research from the Japan Society for the Promotion of Science, the Ministry of Education, Culture, Sports, Science and Technology (13307015, 16209022, 19209024, 22249021); and the Environment

Research and Technology Development Fund (5C-1252, 5-1554) from the Ministry of the Environment, Japan. The funders had no role in the study's design, data collection and analysis, decision to publish, or preparation of the manuscript.

Financial interests:

The authors declare they have no actual or potential competing financial interests.

Abbreviations

PFASs; perfluoroalkyl substances

PFOS; perfluorooctane sulfonate

PFOA; perfluorooctanoate

E2; estradiol

T; testosterone

P4; progesterone

LH; luteinizing hormone

FSH; follicle-stimulating hormone

SHBG; steroid hormone binding globulin

PRL; prolactin

INSL3; insulin-like factor 3

Abstract

BACKGROUND: Exposure to perfluoroalkyl substances (PFASs) may disrupt reproductive function in animals and humans. Although PFASs can cross the human placental barrier, few studies evaluated the effects of prenatal PFAS exposure on the fetus' reproductive hormones.

OBJECTIVE: To explore the associations of prenatal exposure to perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) with cord blood reproductive hormones.

METHODS: In the prospective birth cohort (Sapporo cohort of the Hokkaido study), we included 189 mother-infant pairs recruited in 2002-2005 with both prenatal maternal and cord blood samples. PFOS and PFOA levels in maternal blood after the second trimester were measured via liquid chromatography-tandem mass spectrometry. We also measured cord blood levels of the fetuses' reproductive hormones, including estradiol (E2), total testosterone (T), progesterone (P4), inhibin B, insulin-like factor 3 (INSL3), steroid hormone binding globulin (SHBG), follicle-stimulating hormone, and luteinizing hormone, and prolactin (PRL).

RESULTS: The median PFOS and PFOA levels in maternal serum were 5.2 ng/mL and 1.4 ng/mL, respectively. In the fully adjusted linear regression analyses of the male

infants, maternal PFOS levels were significantly associated with E2 and positively, and T/E2, P4, and inhibin B inversely; PFOA levels were positively associated with inhibin B levels. Among the female infants, there were significant inverse associations between PFOS levels and P4 and PRL levels, although there were no significant associations between PFOA levels and the female infants' reproductive hormone levels.

CONCLUSIONS: These results suggest that the fetal synthesis and secretion of reproductive hormones may be affected by in utero exposure to measurable levels of PFOS and PFOA.

Keywords

Perfluoroalkyl substances, reproductive hormones, cord blood, prenatal exposure, birth cohort

1. Introduction

Perfluoroalkyl substances (PFASs) are widely-used in industrial products and are commonly detected in the environment. Human exposure to PFASs mainly occurs orally, via the intake of contaminated food, water, and dust (Fromme et al., 2009). As perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) are the most commonly detected PFASs in the environment and in humans, their presence in human blood has been reported in several countries (Butenhoff et al., 2006; Calafat et al., 2007; Harada et al., 2007; Midasch et al., 2007). Furthermore, in 2009, PFOS was added to Annex B of the Stockholm Convention on Persistent Organic Pollutants. PFOA is now proposed to be listed on Stockholm Convention by EU. Although PFOS and PFOA are being voluntarily phased out by several industries, they are still present in older products and have long elimination half-lives in human serum (PFOS: 5.3 years, PFOA: 3.8 years) (Olsen et al., 2007). PFOS and PFOA can cross the placental barrier, and be transferred from mother to fetus in humans (Inoue et al., 2004; Midasch et al., 2007). Therefore, significant concern has been raised regarding the adverse effects of in utero exposure to PFOS and PFOA on the fetus.

Epidemiological studies have reported that prenatal exposures to PFOS and PFOA were inversely associated with birth size and neurodevelopment during

childhood (Apelberg et al., 2007; Chen et al., 2013). Our group, the Hokkaido Study on Environment and Children's Health, has also reported an inverse association between maternal PFOS serum levels and birth weight among female infants (Washino et al., 2009). Moreover, we found inverse associations of PFOS with essential fatty acid and triglyceride serum levels in pregnant mothers (Kishi et al., 2015), at levels of PFOS and PFOA that were lower than those found in other countries and areas in Japan (Okada et al., 2013).

Recent studies have suggested that PFASs can disrupt the endocrine system, such as levels of reproductive hormones, toxicity on reproductive cells and gene expression. For example, PFOS and PFOA treatment in adult mice caused decreases in serum testosterone (T) and epididymal sperm counts (Wan et al., 2011), and an increase in progesterone (P4) levels (Zhao Y et al., 2010). In addition, Zhao et al. (2014) have reported that PFOS exposure was associated with a reduced number of fetal Leydig cells, reduced steroidogenic enzyme gene expression, and lower T levels in pregnant rats. As for human populations, a cross-sectional study in children at 6-9 years of age in Mid-Ohio Valley reported that PFAS concentrations inversely associated with serum levels of T, estradiol (E2) and insulin-like growth factor-1 (IGF-1) (Lopez-Espinosa et al., 2016). A study among Taiwanese at 12-17 years of age also reported the

associations of PFAS with lowered T, follicle stimulating hormone (FSH) and sex-hormone binding globulin (SHBG) levels (Tsai et al., 2015). In adult populations, some cross-sectional reports have also revealed associations between serum PFAS concentrations and altered serum levels of E2, T, and luteinizing hormone (LH) (Joensen et al., 2009; Joensen et al., 2013; Knox et al., 2011; Raymer et al., 2012). Furthermore, two prospective studies have reported that PFOS and PFOA exposures in utero were positively associated with serum T levels among young girls (Maisonet et al., 2015), and that PFOA exposure in utero was positively associated with LH and FSH levels, and inversely associated with sperm concentrations and total sperm counts among adult men (Vested et al., 2013). Those studies appear to indicate that prenatal exposure to PFOS and PFOA disrupts reproductive functions by altering reproductive hormone secretions at a later age.

Prenatal period is very important for the growth of fetuses. Disrupted endocrine environment during the perinatal period is considered to influence on the reproductive health in adulthood (Lagiou et al., 2011). Although reproductive issues among adults may have fetal origins (Crain et al., 2008; Juul et al., 2014), there is limited numbers of studies investigating the PFAS impact on fetal reproductive health. It is necessary to clarify the effects of PFAS exposure during the fetal period. Therefore, the present study

aimed to investigate the effects of prenatal exposure to PFOS and PFOA on fetal reproductive hormones.

2. Materials and Methods

2.1 Participants

This prospective birth cohort study was based on the Hokkaido Study on Environment and Children's Health. Details regarding the study population, data collection, biological specimen sampling, and questionnaire's contents have been described elsewhere (Kishi et al., 2013; Kishi et al., 2011). In brief, native Japanese citizens who resided in Sapporo or its suburbs were recruited at 23–35 weeks of gestation between July 2002 and October 2005 at Sapporo Toho Hospital (Sapporo, Hokkaido, Japan). Of 1796 potentially eligible pregnant women, the following women were excluded: decided to participate in the Japanese cord blood bank (22% of those approached), delivered at another hospital (3% of those approached). Of the remaining eligible subjects, 514 (28.6% of those approached) pregnant women agreed to participate in this study. All participants provided written informed consent before the participation in the Hokkaido Study and the study protocol was approved by the ethics review board for epidemiological studies at Hokkaido University Graduate School of

Medicine and the Hokkaido University Center for Environmental and Health Sciences.

This study was performed in accordance with principles of the Declaration of Helsinki.

Among the 514 participants, we excluded women who experienced miscarriage, stillbirth, relocation, or voluntary withdrawal (n = 10), and women who delivered twins (n = 7). 447 maternal blood samples were available for the PFOS and PFOA measurements, and 295 cord blood samples were available for the reproductive hormone measurements. In addition, among 257 participants with paired maternal and cord blood samples, maternal blood samples from 68 women were obtained after their delivery due to anemia. Since these post-delivery blood samples exhibited significantly lower PFOS and PFOA (before delivery: PFOS = 5.2 and PFOA = 1.4 ng/mL; post-delivery: PFOS = 3.5 ng/mL and PFOA = 1.3 ng/mL), those 68 women were excluded. Finally, 189 mother-infant pairs were included in the statistical analysis.

2.2 Exposure assessment

We used the data of PFOS and PFOA concentrations in maternal serum measured by liquid chromatography-tandem mass spectrometry (LC/MS/MS) in our previous study. Detailed methods for the measurement of PFOS and PFOA have been described in our previous reports (Kishi et al., 2015; Nakata A, 2009). 6.9% of the samples exhibited PFOA levels that were below the limit of detection (LOD; 0.50

ng/mL). In those cases, we assigned the sample a value of 0.25 ng/mL (50% of the LOD).

2.3 Outcome measures

A blood sample (10–30 mL) was collected from the umbilical cord at the time of delivery and was stored at –80°C until analysis. We analyzed cord blood samples for their levels of E2, T, P4, LH, FSH, SHBG, prolactin (PRL), and inhibin B, as described in our previous report (Araki et al., 2014). The concentration of insulin-like factor 3 (INSL3) was measured for all male infants using an enzyme immunoassay (INSL3 / RLF [human] EIA kit; Phoenix Pharmaceuticals Inc., CA, USA). However, due to the low detection rate among female infants in previous studies (Bay et al., 2007; Szydlarska et al., 2012), we only tested 25 female infants for INSL3. All reproductive hormone measurements were performed at Asuka Pharma Medical Co. Ltd (Kanagawa, Japan). Samples that exhibited values below the LOD of the test were assigned a value that was 50% of that test’s LOD.

2.4 Questionnaire and medical records

All participants completed a self-administered questionnaire at enrollment regarding their maternal age, educational level, household income, smoking status and medical history. Maternal smoking status during pregnancy was classified as

non-smoker (never smoked or quit smoking during the first trimester) and smoker (had smoked after the first trimester). Medical records were obtained to collect information regarding body mass index (BMI) before pregnancy, pregnancy complications, gestational age, infant sex, parity, congenital anomalies (including hypospadias and cryptorchidism), and infant birth sizes.

2.5 Statistical analyses

The associations of PFOS and PFOA with the maternal and fetal characteristics were explored using Spearman's correlation test and the Mann-Whitney U test.

Correlations of maternal serum PFOS and PFOA levels with cord blood reproductive hormone levels were analyzed using Spearman's correlation test. We also performed two different analyses: multiple linear regression analysis and dose-response analysis.

In the linear regression analysis, levels of PFOS, PFOA, and reproductive hormones were converted using a log₁₀ scale to account for their skewed distribution

(Supplemental Figure 1, 2). In the dose-response analyses, PFOS and PFOA levels were categorized into four quartiles, and reproductive hormones levels were converted using a log₁₀ scale. The least square means of each reproductive hormone were calculated and back transformed. The p-value for trend was calculated using linear contrast coefficient -3, -1, +1, +3 assigned to quartile first, second, third and fourth, respective

(Goudarzi et al., 2016; Kishi et al., 2015). The first quartile was also compared to the second, third, and fourth quartiles of PFOS and PFOA, and the p-values were calculated using Hsu-Dunnett's test. In the fully adjusted model, potential confounders were selected based on the existing literature and the results of the present study ($p < 0.05$ in Table 1; infant factor: gestational age at birth [weeks] [continuous variable], maternal factors: age at delivery [years], parity [$0/\geq 1$], BMI before pregnancy [kg/m^2], annual income ($<5/\geq 5$ million yen], smoking during pregnancy [yes/no], caffeine consumption during pregnancy (mg/day), and gestational weeks of blood sampling for the PFOS and PFOA measurements [continuous variable]). We also performed sensitivity analyses and conducted the backward procedure to exclude some confounders, and compared to the final models. As for BMI, no correlation was found between maternal BMI and hormone levels in cord blood. However, we selected BMI as one of confounders according to the associations between obesity and hormone levels during pregnancy (Lassance et al., 2015). Because of association of steroid hormone levels in fetal serum to gestational age (Procianoy and de Oliveira-Filho, 1996), we included it as a confounder. We also included annual income as an indicator of socioeconomic status into the final models. All statistical analyses were performed using JMP Pro software (version 12; SAS Institute Inc., NC, USA).

3. Results

Among the 189 mother-infant pairs, the median maternal levels of PFOS and PFOA were 5.2 ng/mL (interquartile range [IQR]: 3.85–7.15 ng/mL) and 1.4 ng/mL (IQR: 0.90–2.20 ng/mL), respectively. Table 1 shows the maternal serum PFOS and PFOA levels in relation to the maternal and infants' characteristics; PFOS and PFOA levels were significantly higher among primiparous women. More than 20% of women had smoked during pregnancy, and PFOS level was significantly lower among smokers. The amount of caffeine intake was negatively associated with PFOA levels. The mean of the gestational weeks of blood sampling for the PFOS and PFOA measurement was 33.3 weeks. PFOS and PFOA levels were negatively correlated with increasing gestational weeks of pregnancy. Maternal PFOS was significantly correlated with the birth weight of female infants.

The cord blood reproductive hormone levels are shown in Table 2. Less than 30% of female infants exhibited detectable levels of LH, FSH, and inhibin B. Therefore, we omitted those hormones from further analyses in the female infants. Furthermore, because male infants exhibited significantly higher median levels of T, LH, FSH, inhibin B, and INSL3 (vs. female infants), we performed all further analyses after

stratifying the data according to infant sex.

Table 3 shows the correlations of maternal PFOS and PFOA levels with cord blood reproductive hormone levels. In the male infants, PFOS levels were positively correlated with E2, and were inversely correlated with the T/E2 ratio and inhibin B levels. Maternal PFOA levels were positively correlated with male P4 and inhibin B levels. In the female infants, maternal PFOS was inversely correlated with PRL levels.

The relationships between maternal PFOS and PFOA levels and reproductive hormone concentrations, according to our linear regression analyses for male and female infants are shown in Table 4 and Table 5, respectively. After fully adjusting for the potential confounders, maternal PFOS exhibited significant positive association with E2 in male infants ($\beta = 0.372$; 95% confidence interval [CI]: 0.057, 0.687; $p = 0.021$), and negative associations with the male infants' values for the T/E2 ratio ($\beta = -0.399$; 95% CI: $-0.643, -0.156$; $p = 0.008$), P4 ($\beta = -0.344$; 95% CI: $-0.678, -0.010$; $p = 0.043$), and inhibin B ($\beta = -0.439$; 95% CI: $-0.620, -0.257$; $p < 0.001$). Among the male infants, maternal PFOA was positively associated with inhibin B ($\beta = 0.197$; 95% CI: 0.009, 0.384; $p = 0.040$). Similarly, among the female infants, we observed negative associations between maternal PFOS and P4 ($\beta = -0.552$; 95% CI: $-0.894, -0.210$; $p = 0.002$), and PRL ($\beta = -0.491$; 95% CI: $-0.764, -0.218$; $p = 0.001$), although PFOA was

not significantly associated with any hormone levels.

We also investigated the dose-response relationships between the quartiles of maternal PFOS and PFOA levels and cord blood reproductive hormone levels (using factors in Table 4 and 5 with a p-value of <0.1). Figure 1 shows the hormones in male infants with a significant p-value for trend (<0.05). After fully adjusting for the confounders, these analyses revealed increasing trend for E2 (A), decreasing trends for the T/E2 ratio (B), and inhibin B levels (D) in relation to the PFOS quartiles. The analyses among female infants are shown in Figure 2. Significant decreasing trends were observed for P4 (A) and PRL levels (B) in relation to the PFOS quartiles.

4. Discussion

Reproductive hormones in prenatal period could have permanent effects on the differences in reproductive structure, behavior and cognition between males and females (Collaer and Hines, 1995). Therefore, it is important to examine the disruptive effects of PFAS exposure on reproductive hormone balances during fetal period. To our knowledge, this is the first study to report the associations between prenatal PFAS exposures and cord blood reproductive hormone levels. Our analysis of 8 hormones (including steroid, peptide, and pituitary hormones) and 1 binding protein allow us to

evaluate the PFAS effects on not only changes of each hormone level but also hormone balances. In the present study, we found that maternal PFOS levels were significantly and positively associated with E2 and inversely associated with T/E2, P4, and inhibin B in male's cord blood, and that maternal PFOA levels were positively associated with the male infants' levels of inhibin B. Among the female infants, we found that maternal PFOS levels were significantly and inversely associated with P4 and PRL levels, and that there were no significant associations between maternal PFOA levels and female infants' reproductive hormones levels.

There are some previous studies that evaluated the associations between in utero PFOS and PFOA exposure and adolescents' reproductive hormone levels. As for the exposure levels, a UK study reported median levels for PFOS and PFOA of 19.2 ng/mL and 3.6 mg/mL in the median pregnancy week 16 between 1991 and 1992, respectively (Maisonet et al., 2015), and a Danish study reported median levels of 21.2 ng/mL and 3.8 mg/mL in pregnancy week 30 in 1988-1989, respectively (Vested et al., 2013). In this study, maternal blood samples were obtained during 2002-2005 and the median gestational week of blood sampling for PFOS and PFOA was 33.3 weeks. Median levels of PFOS and PFOA are 5.2 ng/mL and 1.4 ng/mL, respectively, which were lower than in other areas of Japan (Harada et al., 2010) and in other previous

303 studies. Our results suggest that even low levels of PFOS and PFOA exposure can
304 disrupt reproductive hormone imbalance in the fetus. It is noted that the percentage of
305 smoking women in Hokkaido tend to be higher compared to other area in Japan
306 (Matsuzaki et al., 2014). Table 1 shows that 21.7 % of our participants continued
307 smoking during pregnancy, which was not significantly different from participants in
308 original cohort (n=514) (Supplemental Table 1). The level of PFOS among smoking
309 women was significantly lower than that of non-smoking women. Our result is
310 consistent with previous study.

311 There is few study that investigated the levels of reproductive hormone levels
312 in cord blood. Regarding the comparison of those levels in cord blood, our data in Table
313 2 did not differ dramatically from previous studies (Hollier et al., 2014; Kuijper et al.,
314 2013; Warembourg et al., 2016).

315 Among the male infants, we observed that maternal PFOS levels were
316 significantly and inversely associated with P4 levels and the T/E2 ratio, while
317 associated with increased E2 levels. We examined the T/E2 ratio as a marker of
318 aromatase activity, which convert T to E2 (Simpson et al., 1994). The fetal adrenal uses
319 the large amounts of P4 supplied by the placenta (Mastorakos and Ilias, 2003). One
320 reason of decreased P4 in this study can be explained by a recent study that has reported

that PFOS exposure suppressed the secretion of P4 by inducing placental cell apoptosis (Zhang DY et al., 2015). However, p-for trend in quartile was not significant (Figure 1). Our result indicates the there is little dose-response toxicity of PFOS on P4. The inverse association between PFOS levels and the T/E2 ratio is consistent with the results from a Danish cross-sectional study of adult men (Joensen et al., 2013). Zhang Q et al. (2010) have reported that the T/E2 ratio in the seminal plasma of infertile men was significantly lower than that in normospermic men, which suggests that a lower T/E2 may indicate a corresponding reduction in Leydig cell function and spermatogenesis. In regard to steroid synthesis, the association of maternal PFOS with increased E2 and decreased T/E2 in this study is not consistent with the results in previous study that PFOS did not alter the aromatase activity (Kraugerud et al., 2011). It is also reported that the exposure to higher level PFOS seemed to inhibit the human 3beta-HSD and 17beta-HSD (Zhao B et al., 2010). Based on the decrease in P4 levels and unchanged T levels, our data indicate that PFOS exposure at low level had minimal effect on the pathway of steroid synthesis from P4 to T. Female P4 also showed the inverse association with maternal PFOS level in the present study. The decreased P4 in this study is consistent with the findings of Barrett et al. 2015, who reported that saliva PFOS levels were inversely associated with serum P4 levels among non-parous women

aged 25 to 35 with natural menstrual cycles (Barrett et al., 2015). There are few studies investigating the impact of PFASs on female progesterone, in spite of the great importance of progesterone for female reproductive health, including the pregnancy. More studies regarding the association between PFASs exposure and female's steroid hormones are needed to assess the importance of hormone levels in cord blood for reproductive health in later life.

We found that the male infants' levels of inhibin B were inversely associated with maternal PFOS levels in both our linear regression, and inverse tendency in quartile analyses. However, previous epidemiological studies among adults have reported no significant association between levels of PFASs and inhibin B (Joensen et al., 2009; Joensen et al., 2013; Vested et al., 2013). These results appear to suggest that the fetus' inhibin B secretion may be sensitive to PFAS exposure. One possible mechanism for PFOS-mediated decreases in inhibin B levels is the vacuolization of Sertoli cells in the seminiferous tubules and the blood-testis barrier disassembly in animals and in vitro tests (Qiu et al., 2013). In another study, higher PFOS exposure reduced the expression of inhibin subunit genes, such as *Inhba* and *Inhbb*, in male adult mice (Wan et al., 2011). These processes may explain how PFOS might inhibit the secretion of inhibin B by Sertoli cells. In contrast, we found that PFOA levels were

positively associated with the male infants' inhibin B levels in both our linear regression analysis and quartile analysis. Thus, given that there is no definitive report regarding the effects of PFOA on Sertoli cells or inhibin B levels, further studies are needed to determine if there is a mechanism for PFOA-induced dysfunction. Inhibin B is produced by Sertoli cells, and mainly down-regulates the FSH synthesis. Inhibin B is reported to be associated with male reproductive health, such as testis volume and cryptorchidism among infants (Main et al., 2006; Suomi et al., 2006). Therefore, the changes of inhibin B level by PFOS and PFOA in this study indicate the disrupted reproductive function in near future.

We observed that maternal PFOS levels were inversely associated with PRL levels in female infants. Because there are no reports regarding the associations between PFASs and PRL levels, the mechanism for PFOS or PFOA-mediated decreases in PRL remains unknown.

The sex difference in the results is predictable due to the different hormone levels and hormone function between males and females. Although the reason for sex differences is not understood, one explanation for the sex difference is the faster elimination time of PFASs in female compared to male in animal studies (Lau et al., 2007; Zhang T et al., 2015).

In the multiple linear regression analysis, potential confounders were selected from Table 1 with $p < 0.50$ and previous literatures. We conducted the backward procedure to exclude some confounders, and we did not find significant difference in the results in Table 4 and 5. In addition, hormone levels are controlled very strictly in the circulation by hypothalamic-pituitary-gonadal axis, hypothalamic-pituitary-adrenal axis. To maintain the hormone balances, the correlations between each hormone level are observed (Ex. T and E2; spearman's $\rho = 0.580$). Moreover, hormones interact with each other, not only reproductive hormones but also other hormones. Although it is almost impossible to consider all correlations in the analyses, our results can demonstrate the disturbed balances of hormone levels by targeting different hormones in the negative-feedback- system and synthesis pathway.

A major strength of the present study is its prospective birth cohort design, which allowed us to estimate the effects of prenatal PFOS and PFOA exposure on fetal reproductive functions using prenatal and perinatal blood samples. However, there are several limitations in this study. Firstly, we included limited number of participants from original cohort because cord blood samples were only obtained from infants who were delivered via vaginal birth. We compared the maternal and infant characteristics of participants in this study with the original cohort population ($n=514$), as well as

participants with PFOS and PFOA data before delivery (n=323) (Supplemental Table 1). The infants in our analyses had a higher gestational age and a heavier birth weight. The median levels of PFOS and PFOA in maternal serum did not significantly differ between the analyzed participants (PFOS: 5.2 ng/mL, PFOA: 1.4 ng/mL) and compared groups, participants who had PFOS and PFOA data before their pregnancy (PFOS: 5.7 ng/mL, PFOA: 1.4 ng/mL) and the original cohort (PFOS: 5.2 ng/mL, PFOA: 1.3 ng/mL). However, it is possible that healthier children were included in our analyses, which may have led us to underestimate the effects of PFOS and PFOA exposure. Secondly, among the 257 mother-infant pairs with available maternal and cord blood samples, we excluded 68 women whose blood samples were obtained after delivery from our final analysis, considering the accurate reflection of prenatal exposure to PFOS and PFOA exposure. However, that might decrease the statistical power of this study. Furthermore, since we believe that the post-delivery samples also reflected those women's PFAS levels during their pregnancy, as PFOS and PFOA have long half-lives in human serum (5.4 years and 3.8 years, respectively), we performed the analyses including those 68 mothers (Supplemental Table 2 to 5). The results did not significantly affect our findings. Thirdly, we could not include all possible confounders in regression analyses due to the lack of some information. We measured the levels of

reproductive hormones from cord blood in this study. It is important to note that hormone levels dramatically change from the end of gestation to after birth (Kuijper et al., 2013) and the levels of reproductive hormones measured in cord blood may be affected by various factors like diurnal cyclicity, gestational week, duration of labor, placenta weight, and the presence of pre-eclampsia (Hollier et al., 2014; Keelan et al., 2012). Although we could use gestational age as a confounding factor and we also checked the seasonal variation of hormone levels, and no significant difference was found, our results in regression analyses might not reflect affected hormone levels accurately. It is also noted that the results of multiple comparison should be carefully considered because there is a 5% chance of incorrectly rejecting the null hypothesis (Hubbard, 2011). However, we did not use a method to counteract the multiple comparison error because it would increase the probability of false negative results.

In addition, simultaneous exposure to other chemical compounds that may affect the levels of reproductive hormones should be considered. Our previous study reported that maternal concentrations of PFOS and PFOA showed weak but significantly correlated with polychlorinated biphenyls (PCBs), and mercury (Hg), due to the similar exposure sources (Miyashita et al., 2015). As additional analyses, we also added the levels of polychlorinated dibenzo-pdioxins and -dibenzofurans

(PCDDs/PCDFs) and polychlorinated biphenyls (PCBs) into final adjusted model in Table 4 and 5 as confounders, and the results were almost same as those in Table 4 and Table 5 in the directions and significance (data not shown). Therefore, we believe that our results indicated that PFOS and PFOA independently influenced the reproductive hormone levels in the different mechanisms from how PCDDs/PCDFs and PCBs did.

5. Conclusion

In summary, this study found that prenatal PFOS exposure significantly increased the male infant's level of E2 and decreased the male infants' levels of T/E2, P4 and inhibin B, and that PFOA associated their increased inhibin B levels. Similarly, PFOS was negatively associated with the female infants' P4 and PRL levels. These results suggest that in utero exposure to PFOS and PFOA, even at relatively low levels, has adverse effects on fetuses' synthesis of steroid hormones, their Leydig cell function, and their Sertoli cell function. However, our group has previously reported an increasing trend for PFAS levels with longer carbon chains (e.g., perfluorononanoic acid and perfluorodecanoic acid) instead of decreasing trend for PFOA and PFOA, in the plasma of pregnant women between 2003 and 2011 (Okada et al. 2013). Therefore, future studies are needed to verify the effects of PFASs with longer carbon chains on

447 fetal reproductive functions. These studies should also include long-term follow-up
448 regarding the fetuses' reproductive development, in order to elucidate the outcomes of
449 altered hormone levels at the fetal stage.
450

References

- Apelberg BJ, Witter FR, Herbstman JB, Calafat AM, Halden RU, Needham LL, Goldman LR (2007) Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in relation to weight and size at birth. *Environmental health perspectives* 115:1670-1676. doi:10.1289/ehp.10334
- Araki A, Mitsui T, Miyashita C, Nakajima T, Naito H, Ito S, Sasaki S, Cho K, Ikeno T, Nonomura K, Kishi R (2014) Association between maternal exposure to di(2-ethylhexyl) phthalate and reproductive hormone levels in fetal blood: the Hokkaido study on environment and children's health. *PloS one* 9:e109039. doi:10.1371/journal.pone.0109039
- Barrett ES, Chen C, Thurston SW, Haug LS, Sabaredzovic A, Fjeldheim FN, Frydenberg H, Lipson SF, Ellison PT, Thune I (2015) Perfluoroalkyl substances and ovarian hormone concentrations in naturally cycling women. *Fertility and sterility* 103:1261-1270 e1263. doi:10.1016/j.fertnstert.2015.02.001
- Bay K, Virtanen HE, Hartung S, Ivell R, Main KM, Skakkebaek NE, Andersson AM, Toppari J (2007) Insulin-like factor 3 levels in cord blood and serum from children: effects of age, postnatal hypothalamic-pituitary-gonadal axis activation, and cryptorchidism. *The Journal of clinical endocrinology and metabolism* 92:4020-4027.

470 doi:10.1210/jc.2007-0974

471 Butenhoff JL, Ehresman DJ, Chang SC, Parker GA, Stump DG (2009) Gestational and
472 lactational exposure to potassium perfluorooctanesulfonate (K+PFOS) in rats:
473 developmental neurotoxicity. *Reproductive toxicology* 27:319-330.
474 doi:10.1016/j.reprotox.2008.12.010

475 Butenhoff JL, Olsen GW, Pfahles-Hutchens A (2006) The applicability of biomonitoring
476 data for perfluorooctanesulfonate to the environmental public health continuum.
477 *Environmental health perspectives* 114:1776-1782.

478 Calafat AM, Kuklennyik Z, Reidy JA, Caudill SP, Tully JS, Needham LL (2007) Serum
479 concentrations of 11 polyfluoroalkyl compounds in the u.s. population: data from the
480 national health and nutrition examination survey (NHANES). *Environmental science &*
481 *technology* 41:2237-2242.

482 Chen MH, Ha EH, Liao HF, Jeng SF, Su YN, Wen TW, Lien GW, Chen CY, Hsieh WS,
483 Chen PC (2013) Perfluorinated compound levels in cord blood and neurodevelopment
484 at 2 years of age. *Epidemiology (Cambridge, Mass)* 24:800-808.
485 doi:10.1097/EDE.0b013e3182a6dd46

486 Collaer ML, Hines M (1995) Human behavioral sex differences: a role for gonadal
487 hormones during early development? *Psychological bulletin* 118:55-107.

488 Crain DA, Janssen SJ, Edwards TM, Heindel J, Ho SM, Hunt P, Iguchi T, Juul A,
 489 McLachlan JA, Schwartz J, Skakkebaek N, Soto AM, Swan S, Walker C, Woodruff TK,
 490 Woodruff TJ, Giudice LC, Guillette LJ, Jr. (2008) Female reproductive disorders: the
 491 roles of endocrine-disrupting compounds and developmental timing. *Fertility and*
 492 *sterility* 90:911-940. doi:10.1016/j.fertnstert.2008.08.067
 493 Fromme H, Tittlemier SA, Volkel W, Wilhelm M, Twardella D (2009) Perfluorinated
 494 compounds--exposure assessment for the general population in Western countries.
 495 *International journal of hygiene and environmental health* 212:239-270.
 496 doi:10.1016/j.ijheh.2008.04.007
 497 Goudarzi H, Nakajima S, Ikeno T, Sasaki S, Kobayashi S, Miyashita C, Ito S, Araki A,
 498 Nakazawa H, Kishi R (2016) Prenatal exposure to perfluorinated chemicals and
 499 neurodevelopment in early infancy: The Hokkaido Study. *The Science of the total*
 500 *environment* 541:1002-1010. doi:10.1016/j.scitotenv.2015.10.017
 501 Harada K, Koizumi A, Saito N, Inoue K, Yoshinaga T, Date C, Fujii S, Hachiya N,
 502 Hirosawa I, Koda S, Kusaka Y, Murata K, Omae K, Shimbo S, Takenaka K, Takeshita T,
 503 Todoriki H, Wada Y, Watanabe T, Ikeda M (2007) Historical and geographical aspects of
 504 the increasing perfluorooctanoate and perfluorooctane sulfonate contamination in
 505 human serum in Japan. *Chemosphere* 66:293-301.

doi:10.1016/j.chemosphere.2006.05.010

Harada KH, Yang HR, Moon CS, Hung NN, Hitomi T, Inoue K, Niisoe T, Watanabe T, Kamiyama S, Takenaka K, Kim MY, Watanabe K, Takasuga T, Koizumi A (2010) Levels of perfluorooctane sulfonate and perfluorooctanoic acid in female serum samples from Japan in 2008, Korea in 1994-2008 and Vietnam in 2007-2008. *Chemosphere* 79:314-319. doi:10.1016/j.chemosphere.2010.01.027

Hollier LP, Keelan JA, Hickey M, Maybery MT, Whitehouse AJ (2014) Measurement of androgen and estrogen concentrations in cord blood: accuracy, biological interpretation, and applications to understanding human behavioral development. *Frontiers in endocrinology* 5:64. doi:10.3389/fendo.2014.00064

Hubbard R (2011) The widespread misinterpretation of p-values as error probabilities. *Journal of Applied Statistics* 38:2617-2626. doi:10.1080/02664763.2011.567245

Inoue K, Okada F, Ito R, Kato S, Sasaki S, Nakajima S, Uno A, Saijo Y, Sata F, Yoshimura Y, Kishi R, Nakazawa H (2004) Perfluorooctane Sulfonate (PFOS) and Related Perfluorinated Compounds in Human Maternal and Cord Blood Samples: Assessment of PFOS Exposure in a Susceptible Population during Pregnancy. *Environmental health perspectives* 112:1204-1207. doi:10.1289/ehp.6864

Joensen UN, Bossi R, Leffers H, Jensen AA, Skakkebaek NE, Jorgensen N (2009) Do

524 perfluoroalkyl compounds impair human semen quality? *Environmental health*
 525 *perspectives* 117:923-927. doi:10.1289/ehp.0800517
 526 Joensen UN, Veyrand B, Antignac JP, Blomberg Jensen M, Petersen JH, Marchand P,
 527 Skakkebaek NE, Andersson AM, Le Bizec B, Jorgensen N (2013) PFOS
 528 (perfluorooctanesulfonate) in serum is negatively associated with testosterone levels,
 529 but not with semen quality, in healthy men. *Human reproduction* 28:599-608.
 530 doi:10.1093/humrep/des425
 531 Juul A, Almstrup K, Andersson AM, Jensen TK, Jorgensen N, Main KM, Rajpert-De
 532 Meyts E, Toppari J, Skakkebaek NE (2014) Possible fetal determinants of male
 533 infertility. *Nature reviews Endocrinology* 10:553-562. doi:10.1038/nrendo.2014.97
 534 Keelan JA, Mattes E, Tan H, Dinan A, Newnham JP, Whitehouse AJ, Jacoby P, Hickey
 535 M (2012) Androgen concentrations in umbilical cord blood and their association with
 536 maternal, fetal and obstetric factors. *PloS one* 7:e42827.
 537 doi:10.1371/journal.pone.0042827
 538 Kishi R, Kobayashi S, Ikeno T, Araki A, Miyashita C, Itoh S, Sasaki S, Okada E,
 539 Kobayashi S, Kashino I, Itoh K, Nakajima S, Members of the Hokkaido Study on E,
 540 Children's H (2013) Ten years of progress in the Hokkaido birth cohort study on
 541 environment and children's health: cohort profile--updated 2013. *Environmental health*

542 and preventive medicine 18:429-450. doi:10.1007/s12199-013-0357-3
 543 Kishi R, Nakajima T, Goudarzi H, Kobayashi S, Sasaki S, Okada E, Miyashita C, Itoh S,
 544 Araki A, Ikeno T, Iwasaki Y, Nakazawa H (2015) The Association of Prenatal Exposure
 545 to Perfluorinated Chemicals with Maternal Essential and Long-Chain Polyunsaturated
 546 Fatty Acids during Pregnancy and the Birth Weight of their Offspring: The Hokkaido
 547 Study. Environmental health perspectives. doi:10.1289/ehp.1408834
 548 Kishi R, Sasaki S, Yoshioka E, Yuasa M, Sata F, Saijo Y, Kurahashi N, Tamaki J, Endo T,
 549 Sengoku K, Nonomura K, Minakami H, Hokkaido Study on E, Children's H (2011)
 550 Cohort profile: the Hokkaido study on environment and children's health in Japan.
 551 International journal of epidemiology 40:611-618. doi:10.1093/ije/dyq071
 552 Knox SS, Jackson T, Javins B, Frisbee SJ, Shankar A, Ducatman AM (2011)
 553 Implications of early menopause in women exposed to perfluorocarbons. The Journal of
 554 clinical endocrinology and metabolism 96:1747-1753. doi:10.1210/jc.2010-2401
 555 Kraugerud M, Zimmer KE, Ropstad E, Verhaegen S (2011) Perfluorinated compounds
 556 differentially affect steroidogenesis and viability in the human adrenocortical carcinoma
 557 (H295R) in vitro cell assay. Toxicology letters 205:62-68.
 558 doi:10.1016/j.toxlet.2011.05.230
 559 Kuijper EA, Ket JC, Caanen MR, Lambalk CB (2013) Reproductive hormone

560 concentrations in pregnancy and neonates: a systematic review. Reproductive
 561 biomedicine online 27:33-63. doi:10.1016/j.rbmo.2013.03.009
 562 Lagiou P, Samoli E, Okulicz W, Xu B, Lagiou A, Lipworth L, Georgila C, Vatten L,
 563 Adami HO, Trichopoulos D, Hsieh CC (2011) Maternal and cord blood hormone levels
 564 in the United States and China and the intrauterine origin of breast cancer. Annals of
 565 oncology : official journal of the European Society for Medical Oncology / ESMO
 566 22:1102-1108. doi:10.1093/annonc/mdq565
 567 Lassance L, Haghiac M, Minium J, Catalano P, Hauguel-de Mouzon S (2015)
 568 Obesity-induced down-regulation of the mitochondrial translocator protein (TSPO)
 569 impairs placental steroid production. The Journal of clinical endocrinology and
 570 metabolism 100:E11-18. doi:10.1210/jc.2014-2792
 571 Lau C, Anitole K, Hodes C, Lai D, Pfahles-Hutchens A, Seed J (2007) Perfluoroalkyl
 572 acids: a review of monitoring and toxicological findings. Toxicological sciences : an
 573 official journal of the Society of Toxicology 99:366-394. doi:10.1093/toxsci/kfm128
 574 Lopez-Espinosa MJ, Mondal D, Armstrong BG, Eskenazi B, Fletcher T (2016)
 575 Perfluoroalkyl Substances, Sex Hormones, and Insulin-like Growth Factor-1 at 6-9
 576 Years of Age: A Cross-Sectional Analysis within the C8 Health Project. Environmental
 577 health perspectives. doi:10.1289/ehp.1509869

578 Main KM, Toppari J, Suomi AM, Kaleva M, Chellakooty M, Schmidt IM, Virtanen HE,
 579 Boisen KA, Kai CM, Damgaard IN, Skakkebaek NE (2006) Larger testes and higher
 580 inhibin B levels in Finnish than in Danish newborn boys. The Journal of clinical
 581 endocrinology and metabolism 91:2732-2737. doi:10.1210/jc.2005-2443
 582 Maisonet M, Calafat AM, Marcus M, Jaakkola JJ, Lashen H (2015) Prenatal Exposure
 583 to Perfluoroalkyl Acids and Serum Testosterone Concentrations at 15 Years of Age in
 584 Female ALSPAC Study Participants. Environmental health perspectives.
 585 doi:10.1289/ehp.1408847
 586 Mastorakos G, Ilias I (2003) Maternal and fetal hypothalamic-pituitary-adrenal axes
 587 during pregnancy and postpartum. Annals of the New York Academy of Sciences
 588 997:136-149.
 589 Matsuzaki M, Haruna M, Ota E, Murayama R, Yamaguchi T, Shioji I, Sasaki S,
 590 Yamaguchi T, Murashima S (2014) Effects of lifestyle factors on urinary oxidative
 591 stress and serum antioxidant markers in pregnant Japanese women: A cohort study.
 592 Bioscience trends 8:176-184.
 593 Midasch O, Drexler H, Hart N, Beckmann MW, Angerer J (2007) Transplacental
 594 exposure of neonates to perfluorooctanesulfonate and perfluorooctanoate: a pilot study.
 595 International archives of occupational and environmental health 80:643-648.

doi:10.1007/s00420-006-0165-9

Miyashita C, Sasaki S, Saijo Y, Okada E, Kobayashi S, Baba T, Kajiwara J, Todaka T, Iwasaki Y, Nakazawa H, Hachiya N, Yasutake A, Murata K, Kishi R (2015) Demographic, behavioral, dietary, and socioeconomic characteristics related to persistent organic pollutants and mercury levels in pregnant women in Japan. *Chemosphere* 133:13-21. doi:10.1016/j.chemosphere.2015.02.062

Nakata A SK, Iwasaki Y, Ito R, Kishi R, Nakazawa H. (2009) Determination of Perfluorinated Compounds in Human Milk and Evaluation of Their Transition from Maternal Plasma. *Bunseki Kagaku* 58:653-659.

Okada E, Kashino I, Matsuura H, Sasaki S, Miyashita C, Yamamoto J, Ikeno T, Ito YM, Matsumura T, Tamakoshi A, Kishi R (2013) Temporal trends of perfluoroalkyl acids in plasma samples of pregnant women in Hokkaido, Japan, 2003-2011. *Environment international* 60:89-96. doi:10.1016/j.envint.2013.07.013

Olsen GW, Burris JM, Ehresman DJ, Froehlich JW, Seacat AM, Butenhoff JL, Zobel LR (2007) Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired fluorochemical production workers. *Environmental health perspectives* 115:1298-1305. doi:10.1289/ehp.10009

614 Procianoy RS, de Oliveira-Filho EA (1996) Aldosterone cord levels in preterm newborn
615 infants. *Acta paediatrica* (Oslo, Norway : 1992) 85:611-613.

616 Qiu L, Zhang X, Zhang X, Zhang Y, Gu J, Chen M, Zhang Z, Wang X, Wang SL (2013)
617 Sertoli cell is a potential target for perfluorooctane sulfonate-induced reproductive
618 dysfunction in male mice. *Toxicological sciences : an official journal of the Society of*
619 *Toxicology* 135:229-240. doi:10.1093/toxsci/kft129

620 Raymer JH, Michael LC, Studabaker WB, Olsen GW, Sloan CS, Wilcosky T, Walmer
621 DK (2012) Concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate
622 (PFOA) and their associations with human semen quality measurements. *Reproductive*
623 *toxicology* 33:419-427. doi:10.1016/j.reprotox.2011.05.024

624 Simpson ER, Mahendroo MS, Means GD, Kilgore MW, Hinshelwood MM,
625 Graham-Lorence S, Amarneh B, Ito Y, Fisher CR, Michael MD, et al. (1994) Aromatase
626 cytochrome P450, the enzyme responsible for estrogen biosynthesis. *Endocrine reviews*
627 15:342-355. doi:10.1210/edrv-15-3-342

628 Suomi AM, Main KM, Kaleva M, Schmidt IM, Chellakooty M, Virtanen HE, Boisen
629 KA, Damgaard IN, Kai CM, Skakkebaek NE, Toppari J (2006) Hormonal changes in
630 3-month-old cryptorchid boys. *The Journal of clinical endocrinology and metabolism*
631 91:953-958. doi:10.1210/jc.2004-2318

632 Szydłarska D, Grzesiuk W, Trybuch A, Kondracka A, Kowalik I, Bar-Andziak E (2012)
 633 Insulin-like factor 3 -- a new hormone related to polycystic ovary syndrome?
 634 Endokrynologia Polska 63:356-361.

635 Tsai MS, Lin CY, Lin CC, Chen MH, Hsu SH, Chien KL, Sung FC, Chen PC, Su TC
 636 (2015) Association between perfluoroalkyl substances and reproductive hormones in
 637 adolescents and young adults. International journal of hygiene and environmental health
 638 218:437-443. doi:10.1016/j.ijheh.2015.03.008

639 Vested A, Ramlau-Hansen CH, Olsen SF, Bonde JP, Kristensen SL, Halldorsson TI,
 640 Becher G, Haug LS, Ernst EH, Toft G (2013) Associations of in utero exposure to
 641 perfluorinated alkyl acids with human semen quality and reproductive hormones in
 642 adult men. Environmental health perspectives 121:453-458, 458e451-455.
 643 doi:10.1289/ehp.1205118

644 Wan HT, Zhao YG, Wong MH, Lee KF, Yeung WS, Giesy JP, Wong CK (2011)
 645 Testicular signaling is the potential target of perfluorooctanesulfonate-mediated
 646 subfertility in male mice. Biology of reproduction 84:1016-1023.
 647 doi:10.1095/biolreprod.110.089219

648 Warembourg C, Debost-Legrand A, Bonvallot N, Massart C, Garlantezec R, Monfort C,
 649 Gaudreau E, Chevrier C, Cordier S (2016) Exposure of pregnant women to persistent

650 organic pollutants and cord sex hormone levels. Human reproduction 31:190-198.
 651 doi:10.1093/humrep/dev260
 652 Washino N, Saijo Y, Sasaki S, Kato S, Ban S, Konishi K, Ito R, Nakata A, Iwasaki Y,
 653 Saito K, Nakazawa H, Kishi R (2009) Correlations between prenatal exposure to
 654 perfluorinated chemicals and reduced fetal growth. Environmental health perspectives
 655 117:660-667. doi:10.1289/ehp.11681
 656 Zhang DY, Xu XL, Shen XY, Ruan Q, Hu WL (2015) Analysis of apoptosis induced by
 657 perfluorooctane sulfonates (PFOS) in mouse Leydig cells in vitro. Toxicology
 658 mechanisms and methods 25:21-25. doi:10.3109/15376516.2014.971140
 659 Zhang Q, Bai Q, Yuan Y, Liu P, Qiao J (2010) Assessment of seminal estradiol and
 660 testosterone levels as predictors of human spermatogenesis. Journal of andrology
 661 31:215-220. doi:10.2164/jandrol.109.007609
 662 Zhang T, Sun H, Qin X, Gan Z, Kannan K (2015) PFOS and PFOA in paired urine and
 663 blood from general adults and pregnant women: assessment of urinary elimination.
 664 Environmental science and pollution research international 22:5572-5579.
 665 doi:10.1007/s11356-014-3725-7
 666 Zhao B, Hu GX, Chu Y, Jin X, Gong S, Akingbemi BT, Zhang Z, Zirkin BR, Ge RS
 667 (2010) Inhibition of human and rat 3beta-hydroxysteroid dehydrogenase and

668 17beta-hydroxysteroid dehydrogenase 3 activities by perfluoroalkylated substances.
669 Chemico-biological interactions 188:38-43. doi:10.1016/j.cbi.2010.07.001
670 Zhao B, Li L, Liu J, Li H, Zhang C, Han P, Zhang Y, Yuan X, Ge RS, Chu Y (2014)
671 Exposure to perfluorooctane sulfonate in utero reduces testosterone production in rat
672 fetal Leydig cells. PloS one 9:e78888. doi:10.1371/journal.pone.0078888
673 Zhao Y, Tan YS, Haslam SZ, Yang C (2010) Perfluorooctanoic acid effects on steroid
674 hormone and growth factor levels mediate stimulation of peripubertal mammary gland
675 development in C57BL/6 mice. Toxicological sciences : an official journal of the
676 Society of Toxicology. pp 214-224
677

Figure Legends

Figure 1. Least square means of each reproductive hormone in cord blood among male infants, according to maternal serum PFOS or PFOA concentration quartiles.

(A) male E2 according to PFOS, (B) male T/E2 according to PFOS, (C) male P4 according to PFOS, (D) male inhibin B according to PFOS, (E) male INSL3 according to PFOS, and (F) male inhibin B according to PFOA. PFOS and PFOA levels were divided into 4 categories. PFOS: $1.5 \leq \text{Quartile (Q)} 1 \leq 3.9$, $3.9 < \text{Q}2 \leq 5.2$, $5.2 < \text{Q}3 \leq 7.1$, $7.1 < \text{Q}4 \leq 16.2$, PFOA : $0.25 \leq \text{Quartile (Q)} 1 \leq 0.9$, $0.9 < \text{Q}2 \leq 1.4$, $1.4 < \text{Q}3 \leq 2.2$, $2.2 < \text{Q}4 \leq 5.3$. Results are based on multiple linear regression models that were adjusted for maternal factors (age, parity, body mass index before pregnancy, annual income, smoking during pregnancy, caffeine consumption during pregnancy, and gestational weeks of blood sampling for PFOS and PFOA measurements) and infant factors (gestational age at birth). * $p < 0.05$, ** $p < 0.01$ compared to the first quartile, as calculated using the Hsu-Dunnett method. LSMs are indicated in rhombus and the black dots depict the upper and lower 95% CI. PFOS: perfluorooctane sulfonate, PFOA: perfluorooctanoate.

Figure 2. Least square means of each reproductive hormone in cord blood among female infants, according to maternal serum PFOS concentration quartiles. (A) female P4 according to PFOS and (B) female PRL according to PFOS. PFOS levels were divided into 4 categories. PFOS: $1.5 \leq \text{Quartile (Q)} 1 \leq 3.9$, $3.9 < Q2 \leq 5.2$, $5.2 < Q3 \leq 7.1$, $7.1 < Q4 \leq 16.2$. Results are based on multiple linear regression models that were adjusted for maternal factors (age, parity, body mass index before pregnancy, annual income, smoking during pregnancy, caffeine consumption during pregnancy, and gestational weeks of blood sampling for PFOS and PFOA measurements) and infant factors (gestational age at birth). * $p < 0.05$, ** $p < 0.01$ compared to the first quartile, as calculated using the Hsu-Dunnett method. LSMs are indicated in rhombus and the black dots depict the upper and lower 95% CI. PFOS: perfluorooctane sulfonate.

1 **Table 1. Maternal perfluorooctane sulfonate and perfluorooctanoate concentrations according to maternal and infant**
2 **characteristics. (n=189)**

Characteristics		N (%)	Mean ± SD	PFOS (ng/mL)		PFOA (ng/mL)	
				Median (IQR)	p-value	Median (IQR)	p-value
Maternal characteristics							
Age at delivery (years)		189	29.7 ± 4.8	ρ = −0.060	0.483	ρ = −0.049	0.500
Pre-pregnancy BMI (kg/m ²)		189	21.2 ± 3.1	ρ = −0.040	0.588	ρ = −0.044	0.547
Parity	Primiparous	103 (54.5)		5.70 (4.20–8.00)	0.001	1.70 (1.20–2.40)	<0.001
	Multiparous	86 (45.5)		4.75 (3.10–6.28)		1.00 (0.70–1.53)	
Annual household income (million yen per year)	<5	132 (69.8)		5.20 (3.63–7.00)	0.802	1.40 (0.90–2.18)	0.277
	≥5	55 (30.2)		5.40 (4.20–7.50)		1.50 (0.90–2.20)	
Educational level (years)	≤12	87 (46.0)		5.20 (4.00–6.80)	0.773	1.30 (0.90–2.00)	0.806
	≥13	102 (54.0)		5.35 (3.65–7.68)		1.50 (1.00–2.30)	
Fish consumption in a week (g/week)		189	47.2 ± 31.3	ρ = −0.051	0.483	ρ = −0.081	0.268
Smoking during pregnancy	Nonsmoker	148 (78.3)		5.30 (4.00–7.45)	0.027	1.40 (0.90–2.20)	0.065
	Smoker	41 (21.7)		4.70 (2.75–6.75)		1.20 (0.75–1.75)	
Alcohol consumption during	Nondrinker	129 (68.3)		5.20 (3.95–7.40)	0.300	1.40 (0.90–2.10)	0.594

pregnancy	Drinker	60 (31.7)		5.25 (3.63–7.08)		1.40 (0.90–2.28)	
Caffeine intake (mg/day)		189	142.3 ± 125.3	$\rho = -0.092$	0.208	$\rho = -0.187$	0.010
Gestational week of blood sampling for PFOS and PFOA		189	33.3 ± 5.3	$\rho = -0.296$	<0.001	$\rho = -0.194$	0.008
Infant characteristics							
Sex	Male	83 (43.9)		5.40 (4.10–7.50)	0.056	1.60 (1.00–2.40)	0.055
	Female	106 (56.1)		5.15 (3.45–7.00)		1.35 (0.80–2.00)	
Birth weight (g)	Male	83 (43.9)	3168.4 ± 298.7	$\rho = -0.007$	0.948	$\rho = -0.146$	0.187
	Female	106 (56.1)	3099.8 ± 350.0	$\rho = -0.211$	0.030	$\rho = -0.162$	0.097
Gestational age (weeks)		189	39.5 ± 1.0	$\rho = 0.004$	0.952	$\rho = 0.058$	0.431

- 1 Statistical analyses were performed using Spearman's correlation test (ρ), or the Mann-Whitney U test and Kruskal-Wallis test, as
- 2 appropriate. SD: standard deviation, PFOS: perfluorooctane sulfonate, PFOA: perfluorooctanoate, IQR: interquartile range, BMI: body
- 3 mass index.

Table 2. Distribution of reproductive hormone levels in cord blood

	Male (N = 83)				Female (N = 106)				p-value
	n	Median	IQR	>LOD (%)	n	Median	IQR	>LOD (%)	
Estradiol (ng/mL)	83	5.37	(3.64–7.80)	100	106	5.01	(3.48–6.67)	100	0.259
Testosterone (pg/mL)	83	93.6	(70.4–122.3)	100	106	71.0	(51.8–105.9)	100	<0.001
T/E2	83	17.2	(12.0–22.6)		106	15.3	(11.7–21.3)		0.273
Progesterone (ng/mL)	83	235.9	(184.8–304.9)	100	106	213.2	(170.8–269.9)	100	0.101
LH (mIU/mL)	80	<LOD	(<LOD–0.83)	32.5	104	<LOD	(<LOD–<LOD)	0.9	<0.001
FSH (mIU/mL)	80	<LOD	(<LOD–0.65)	43.4	103	<LOD	(<LOD–<LOD)	0.0	<0.001
SHBG (nmol/L)	83	16.3	(13.5–19.0)	100	106	15.6	(13.2–18.5)	100	0.382
T/SHBG	83	5.73	(3.85–7.97)		106	4.38	(3.31–6.39)		0.801
PRL (ng/mL)	80	84.7	(67.1–114.8)	100	105	86.9	(56.8–122.5)	99.1	0.693
Inhibin B (pg/mL)	83	43.2	(32.6–56.8)	100	106	<LOD	(<LOD–11.7)	23.6	<0.001
INSL3 (ng/mL)	80	0.28	(0.24–0.32)	100	20	0.20	(0.17–0.25)	100	<0.001

P-values were calculated using the Mann-Whitney U test. IQR: interquartile range, LOD: limit of detection, LH: luteinizing hormone, FSH: follicle stimulating hormone, SHBG: sex hormone-binding globulin, PRL: prolactin, INSL3: insulin-like 3.

Table 3. Correlations between maternal perfluorooctane sulfonate and perfluorooctanoate levels and infant reproductive hormone levels

	Male (N = 83)				Female (N = 106)			
	PFOS		PFOA		PFOS		PFOA	
	Spearman's ρ	p-value	Spearman's ρ	p-value	Spearman's ρ	p-value	Spearman's ρ	p-value
Estradiol (ng/mL)	0.253	0.021	0.085	0.444	0.084	0.391	−0.033	0.739
Testosterone (pg/mL)	−0.006	0.957	−0.094	0.396	0.024	0.805	−0.006	0.950
T/E2	−0.330	0.002	−0.206	0.062	0.005	0.955	−0.042	0.668
Progesterone (ng/mL)	−0.051	0.649	0.260	0.018	−0.147	0.133	0.175	0.072
LH (mIU/mL)	−0.081	0.475	0.027	0.815	n.d.		n.d.	
FSH (mIU/mL)	0.003	0.982	−0.122	0.280	n.d.		n.d.	
SHBG (nmol/L)	−0.038	0.733	0.127	0.253	−0.125	0.201	−0.062	0.527
T/SHBG	0.019	0.865	−0.162	0.143	0.030	0.767	0.005	0.957
PRL (ng/mL)	−0.113	0.318	0.151	0.182	−0.350	<0.001	−0.095	0.333
Inhibin B (pg/mL)	−0.447	<0.001	0.230	0.037	n.d.		n.d.	
INSL3 (ng/mL)	−0.082	0.465	0.146	0.196	n.d.		n.d.	

PFOS: perfluorooctane sulfonate, PFOA: perfluorooctanoate, n.d.: not determined, LH: luteinizing hormone, FSH: follicle stimulating hormone, SHBG: sex hormone-binding globulin, PRL: prolactin, INSL3: insulin-like 3.

Table 4. Linear regression models of maternal perfluorooctane sulfonate and perfluorooctanoate levels and reproductive hormone levels among male infants (n=83)

	Male															
	PFOS								PFOA							
	Crude				Adjusted				Crude				Adjusted			
	B	(95% CI)	p-value	B	(95% CI)	p-value		B	(95% CI)	p-value	B	(95% CI)	p-value			
Estradiol (ng/mL)	0.330	0.027	0.632	0.033	0.372	0.057	0.687	0.021	0.026	−0.223	0.275	0.836	−0.134	−0.436	0.168	0.378
Testosterone (pg/mL)	−0.096	−0.412	0.220	0.548	−0.027	−0.367	0.312	0.872	−0.206	−0.455	0.042	0.103	−0.160	−0.474	0.154	0.313
T/E2	−0.425	−0.679	0.171	0.001	−0.399	−0.643	−0.156	0.002	−0.232	−0.443	−0.022	0.031	−0.041	−0.224	0.143	0.662
Progesterone (ng/mL)	−0.365	−0.675	−0.056	0.021	−0.344	−0.678	−0.010	0.043	0.171	−0.081	0.424	0.181	0.258	−0.056	0.571	0.105
LH (mIU/mL)	−0.132	−0.518	0.253	0.497	−0.243	−0.643	0.158	0.231	−0.085	−0.229	0.399	0.592	0.071	−0.301	0.443	0.704
FSH (mIU/mL)	0.036	−0.247	0.319	0.802	−0.027	−0.321	0.267	0.854	−0.121	−0.349	0.108	0.297	−0.141	−0.410	0.128	0.300
SHBG (nmol/L)	−0.044	−0.160	0.072	0.455	−0.051	−0.167	0.063	0.374	0.052	−0.041	0.144	0.270	0.009	−0.0998	0.116	0.870
T/SHBG	−0.052	−0.377	0.273	0.751	0.024	−0.318	0.366	0.890	−0.258	−0.511	−0.005	0.046	−0.169	−0.484	0.147	0.290
PRL (ng/mL)	−0.102	−0.297	0.094	0.303	−0.132	−0.341	0.077	0.212	0.099	−0.059	0.258	0.217	0.043	−0.152	0.237	0.664
Inhibin B (pg/mL)	−0.423	−0.598	−0.249	<0.001	−0.439	−0.620	−0.257	<0.001	0.182	0.029	0.335	0.021	0.197	0.009	0.384	0.040
INSL3 (ng/mL)	−0.159	−0.317	−0.002	0.047	−0.139	−0.303	−0.025	0.095	0.054	−0.077	0.184	0.418	0.121	−0.030	0.273	0.114

Adjusted for maternal factors (age, parity, body mass index before pregnancy, annual income, smoking during pregnancy, caffeine consumption during pregnancy, and gestational weeks of blood sampling for PFOS/PFOA measurement) and infant factors (gestational age at birth). PFOS: perfluorooctane sulfonate, PFOA: perfluorooctanoate, LH: luteinizing hormone, FSH: follicle stimulating hormone, SHBG: sex hormone-binding globulin, PRL: prolactin, INSL3: insulin-like 3.

Table 5. Linear regression models of maternal perfluorooctane sulfonate and perfluorooctanoate levels and reproductive hormone levels among female infants (n=106)

	Female															
	PFOS								PFOA							
	Crude				Adjusted				Crude				Adjusted			
	B	(95% CI)	p-value		B	(95% CI)	p-value		B	(95% CI)	p-value	B	(95% CI)	p-value		
Estradiol (ng/mL)	0.136	−0.086	0.357	0.227	0.081	−0.148	0.312	0.481	0.033	−0.107	0.173	0.638	−0.040	−0.194	0.114	0.606
Testosterone (pg/mL)	0.044	−0.257	0.344	0.775	0.069	−0.261	0.400	0.679	−0.044	−0.251	0.164	0.678	−0.031	−0.265	0.203	0.796
T/E2	−0.092	−0.349	0.164	0.478	−0.013	−0.284	0.259	0.926	−0.003	−0.181	0.174	0.970	0.069	−0.123	0.261	0.476
Progesterone (ng/mL)	−0.432	−0.766	−0.098	0.012	−0.552	−0.894	−0.210	0.002	0.106	−0.131	0.344	0.377	0.039	−0.216	0.293	0.764
SHBG (nmol/L)	−0.165	−0.386	0.057	0.143	−0.180	−0.417	0.057	0.135	−0.103	−0.256	0.050	0.185	−0.117	−0.286	0.051	0.169
T/SHBG	0.208	−0.171	0.588	0.279	0.249	−0.158	0.657	0.227	0.060	−0.204	0.323	0.656	0.087	−0.203	0.377	0.554
PRL (ng/mL)	−0.426	−0.677	−0.174	0.001	−0.491	−0.764	−0.218	0.001	−0.112	−0.294	0.071	0.229	−0.157	−0.361	0.047	0.131

Adjusted for maternal factors (age, parity, body mass index before pregnancy, annual income, smoking during pregnancy, caffeine consumption during pregnancy, and gestational weeks of blood sampling for PFOS/PFOA measurement) and infant factors (gestational age at birth). PFOS: perfluorooctane sulfonate, PFOA: perfluorooctanoate, SHBG: sex hormone-binding globulin, PRL: prolactin

Figure(s)




Figure 1.




Figure 2.