



HOKKAIDO UNIVERSITY

Title	Driving and photo-regulation of myosin-actin motors at molecular and macroscopic levels by photo-responsive high energy molecules
Author(s)	Menezes, Halley M.; Islam, Md. Jahirul; Takahashi, Masayuki et al.
Citation	Organic & biomolecular chemistry, 15(42), 8894-8903 https://doi.org/10.1039/c7ob01293d
Issue Date	2017-11-14
Doc URL	https://hdl.handle.net/2115/71967
Type	journal article
File Information	Revised_SI.pdf, Supporting Information



Supporting Information

Driving and photo-regulation of myosin-actin motor at molecular and macroscopic level by photo-responsive high energy molecules.

Halley M. Menezes,^{1,2} Md. Jahirul Islam,^{1,2} Masayuki Takahashi³ and Nobuyuki Tamaoki^{1,2*}

¹Research Institute for Electronic Science, Hokkaido University, Kita 20, Nishi 10, Kita-Ku, Sapporo, Hokkaido, 001-0020, Japan.

²Graduate School of Life Science, Hokkaido University, Kita 10, Nishi 8, Kita-ku, Sapporo, Hokkaido, 060-0810, Japan.

³Faculty of Science, Hokkaido University, Kita 13, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810, Japan.


E-mail: tamaoki@es.hokudai.ac.jp

Table of contents

1. Synthetic schemes.....	S2
2. Photoisomerization of AzoTPs.....	S4
3. Isomerization ratio of <i>cis</i> and <i>trans</i> before and after irradiation.....	S5
4. Muscle fibre shortening as a function of AzoTP concentration.....	S6
5. Purity of the AzoTPs by HPLC analysis.....	S7
6. Mass spectra.....	S8
7. Supplementary movie information.....	S14
8. NMR spectral data (¹ H, ¹³ C and ³¹ P NMR).....	S15

1. Synthetic schemes of AzoTP derivatives.

Scheme of AzoethoxyTP (4b)




a) Ethylene glycol, Pd(OAc)₂, PhI(OAc)₂

b) di-*tert*-butyl *N,N*-diisopropylphosphoramidite, 1*H*-tetrazole, dry THF, Ar atmosphere, rt, 6h; then, mCPBA, 0°C, 1h; then rt, 40 min;

c) Trifluoroacetic acid, dry CH₂Cl₂, Ar atmosphere, rt, 6 h; then eluting through DEAE Sephadex A-25 anion exchanger, TEAB;

d) Tributylamine, carbonyldiimidazole, pyrophosphate, dry DMF, Ar atmosphere, rt, overnight.

Scheme of DimethylAzoTP (4c)



(a) mCPBA, EtOAc, 0°C, 3h.

(b) 1,2-phenylenediamine, AcOH, EtOAc, N₂- atmosphere, 50 °C, 72h.

(c) Acetoxyactyl chloride, Triethylamine, DCM, RT, 3h.


(d) K_2CO_3 , MeOH, RT, Overnight.

(e) Di-*tert*-butyl *N,N*-Diisopropylphosphoramidite, 1H-tetrazol, dry THF, Ar-atmosphere, RT, 6h, then mCPBA, 0°C, 1h.

(f) TFA, dry DCM, Ar-atmosphere, RT, 6h.

(g) Tributylamine, carbonyldiimidazole, tributylammonium pyrophosphate, dry DMF, Ar-atmosphere, RT, overnight.

Scheme of AzoethylTP (4)



(a) ArNO, AcOH, Toluene, Ar-atmosphere, 60 °C, 72h.

(b) Di-*tert*-butyl *N,N*-Diisopropylphosphoramidite, 1H-tetrazol, dry THF, Ar-atmosphere, RT, 6h, then mCPBA, 0°C, 1h.

(c) TFA, dry DCM, Ar-atmosphere, RT, 6h.

(d) Tributylamine, carbonyldiimidazole, tributylammonium pyrophosphate, dry DMF, Ar atmosphere, RT, overnight.

2. Photoisomerization of AzoTPs.





Fig. S1. UV-Vis absorption spectra of AzoTP molecules **a)** DimethylAzoTP (**4c**), **b)** AzoethylTP (**4d**) in BRB-80 buffer solution at 25 °C; before irradiation (black line), UV PSS (red line), Vis PSS (green line) and **c)** AzoethoxyTP (**4b**) in water at 25 °C, before irradiation (black line), UV PSS (green line), Vis PSS (red line). Insets show the absorbance changes **a)** at 341 nm, **b)** at 325 nm and **c)** at 324 nm after the alternate irradiations by 365 and 436 nm light for 5 cycles.


3. Isomerization ratio of *cis* and *trans* at UV and Visible PSS

The photo conversion ratio from *trans* to *cis* and *cis* to *trans* isomers of the AzoTPs corresponding to irradiation with 365 nm and 436 nm light was measured with ^1H NMR (400 MHz, D_2O).

a)



b)



c)




Fig. S2. ^1H NMR spectra (400 MHz, D_2O) showing the *trans* and *cis* isomer ratios of **a**) AzoethoxyTP (**4b**), **b**) DimethylAzoTP (**4c**) and **c**) AzoethylTP (**4d**) before irradiation, at UV PSS and Visible PSS.

4. Muscle fibre shortening as a function of AzoTP concentration





Fig. S3. AzoTP (**4a**) concentration dependent shortening of muscle fiber. Lines are best fit through the data trend.

5. Purity of the AzoTPs by HPLC analysis.


a)

mV




b)

mV



c)

mV



d)

mV




Fig. S4. Reverse phase HPLC chromatograms of **a)** AzoTP (**4a**), **b)** AzoethoxyTP (**4b**), **c)** DimethylAzoTP (**4c**), **d)** AzoethylTP (**4d**) showing > 95% purity. Conditions of the RP-HPLC analysis; Column: CN-80Ts, 4.6 × 250 mm (TOSOH). Eluent: Eluent - CH₃CN/NaPi buffer (pH 6.5). Solvent gradient: 15 – 45% of CH₃CN in NaPi buffer. Monitoring wavelength: $\lambda = 327$ nm. Flow rate: 0.5 mL/min at room temperature (23 °C). Injection volume: 20 μ L.

6. Mass spectra of AzoTP molecules.




Fig. S5. ESI mass spectrum of AzoethoxyTP (**4b**), the peaks at $m/z = 504.03$, 503.02 , 479.02 and 481.04 assigned to the $[M-H]^-$ ions of monosodium tetraprotonated, monosodium triprotonated, diprotonated and tetraprotonated AzoethoxyTP species respectively. The peaks at $m/z = 424.05$, 423.05 and 401.07 assigned to $[M-H]^-$ ions of monosodium triprotonated, monosodium diprotonated and triprotonated AzoethoxyTP species respectively.




Fig. S6. ESI mass spectrum of DimethylAzoTP (**4c**), the peaks at $m/z = 668.11$, 544.15 and 522.16 can be assigned to the $[M-H]^-$ ions of disodium-mono triethyl ammonium, monosodium and tetra-protonated DimethylAzoTP species respectively. The peaks at $m/z = 464.16$ and 442.17 can be assigned to the $[M-H]^-$ ions of monosodium and tri-protonated DimethylAzoDP species respectively.




Fig. S7. ESI mass spectrum of AzoethylTP (**4d**), the peaks at $m/z = 487.04$, 465.06 and 385.08 can be assigned to the $[M-H]^-$ ions of monosodium, tetra-protonated AzoethylTP and tri-protonated AzoethylDP species respectively.




Fig. S8. High-resolution mass spectrum of **1b** calculated for $C_{14}H_{16}N_2O_2Na$ $[M + Na]^+$: 265.09475




Fig. S9. High-resolution mass spectrum of **2b** calculated for $C_{22}H_{31}N_2O_5PNa$ $[M + Na]^+$: 457.18628.




Fig. S10. High-resolution mass spectrum of **3b** calculated for $C_{14}H_{14}N_2O_5P$ $[M - H]^-$: 321.06458.

170414_AzoethoxyTP_pn5 #23-25 RT: 0.29-0.31 AV: 2 NL: 3.41E6
T: FTMS (1,1) + p ESI Full ms [150.00-2000.00]




Fig. S11. High-resolution mass spectrum of AzoethoxyTP (**4b**) as tetrasodium salt calculated for $C_{14}H_{14}N_2O_{11}Na_4P_3$ $[M + H]^+$: 570.93957

170417_Dimethyl_1_pn #24-27 RT: 0.30-0.32 AV: 2 NL: 2.88E7
T: FTMS (1,1) + p ESI Full ms [100.00-2000.00]




Fig. S12. High-resolution mass spectrum of **5** calculated for $C_{14}H_{16}N_3$ $[M + H]^+$: 226.13387.

170418_Dimethyl_2_pn3 #17-19 RT: 0.24-0.26 AV: 2 NL: 7.23E7
T: FTMS (1,1) + p ESI Full ms [150.00-2000.00]




Fig. S13. High-resolution mass spectrum of **6** calculated for $C_{18}H_{19}N_3O_3Na$ $[M + Na]^+$: 348.13186.




Fig. S14. High-resolution mass spectrum of **1c** calculated for $C_{16}H_{17}N_3O_2Na [M + Na]^+$: 306.12130.




Fig. S15. High-resolution mass spectrum of **2c** calculated for $C_{24}H_{34}N_3O_5P [M + Na]^+$: 498.21283.




Fig. S16. High-resolution mass spectrum of **3c** calculated for $C_{16}H_{17}N_3O_5P [M - C_6H_{17}N(TEA)]^-$: 362.09113.




Fig. S17. High-resolution mass spectrum of DimethylAzoTP (**4c**) as tetrasodium salt calculated for $C_{16}H_{16}N_3O_{11}Na_4P_3 [M + H]^+$: 611.96612.

170420_Azoethyl_1_pn2#24-27 RT: 0.30-0.32 AV: 2 NL: 1.19E8
T: FTMS (1,1) + p ESI Full ms [100.00-2000.00]




Fig. S18. High-resolution mass spectrum of **1d** calculated for $C_{14}H_{14}N_2ONa$ $[M + Na]^+$: 249.09983.




Fig. S19. High-resolution mass spectrum of **2d** calculated for $C_{22}H_{31}N_2O_4PNa$ $[M + Na]^+$: 441.19137.




Fig. S20. High-resolution mass spectrum of **3d** $C_{14}H_{14}N_2O_4P$ $[M - C_6H_{16}N(TEA)]^-$: 305.06967.

170415_AzoethylTP_pn2#23-25 RT: 0.29-0.31 AV: 2 NL: 2.14E6
T: FTMS (1,1) + p ESI Full ms [150.00-2000.00]




Fig. S21. High-resolution mass spectrum of AzoethylTP (**4d**) as tetrasodium salt calculated for $C_{14}H_{14}N_2O_{10}Na_4P_3$ $[M + H]^+$: 554.94466.


7. Supplementary Movies

Mv01: Movie demonstrates the in-situ photoregulation of F-actin motility by irradiating alternatingly with 365nm(3s) and 510nm (5s). The change in the velocity of fluorescently labelled actin filaments corresponding to the light irradiation could be seen. After UV irradiation the filaments move slowly and after shining Visible light the speed increases drastically. The concentration of **4a** is 40 μ M.

Mv02: This movie demonstrates the effect of *trans* and *cis* isomer of **4a** on the glycerinated muscle fibre. *cis-4a* was added to one fibre and *trans-4a* added to the other. Fibre shortens after addition of *trans-4a* solution while the fibre infused with *cis-4a* doesn't show any change in the length. *cis-4a* was pregenerated before addition on muscle fibre. Concentration of **4a** is 3 mM (total).


Mv03: This movie demonstrates that the photo-induced shortening of muscle fibre is due to the photo-generated *trans* isomer and not due to the heat of irradiation. Buffer solution without any **4a** is added to one fibre and to the fibre *cis-4a* is added. Both the fibres don't shorten even in 20s after addition of buffer and *cis-4a* solution respectively, unlike as seen in the **Mv01** where the fibre shortens significantly in 20s duration after addition of *trans-4a*. So this again reconfirms that *cis-4a* doesn't induce muscle fibre shortening. Both the fibres are irradiated with 510nm light for 10s. Muscle fibre infused with *cis-4a* shortenes about 38% of its initial length while the buffer solution infused fibre doesn't show any change in the length. Concentration of **4a** is 3 mM (total).



Mv04: This movie demonstrates the photoinduced shortening of muscle fibre infused in inactive *cis-4a* solution. To both the muscle fibres *cis-4a* solution was added and one fibre was irradiated with 510nm light for 10s. The irradiated fibre shortens about 31% of its length before irradiation whereas the non-irradiated muscle fibre exhibits no change in length as shown in the snapshots of the video below.





8. NMR spectral data

^1H NMR and ^{13}C NMR spectra of **1b**





¹H NMR and ¹³C NMR spectra of **2b**



^1H NMR and ^{13}C NMR spectra of **3b**





^1H NMR and ^{13}C NMR spectra of **4b**





^1H NMR and ^{13}C NMR spectra of **1d**



¹H NMR and ¹³C NMR spectra of **2d**



¹H NMR and ¹³C NMR spectra of **3d**




^1H NMR and ^{13}C NMR spectra of **4d**

^1H EC AzoTP BI
single_pulse

7.96
7.95
7.94
7.93
7.84
7.83
7.82
7.60
7.59
7.56
7.54
7.53
7.52
7.44
7.42
7.41

4.27
4.25
4.23
4.21
3.50
3.48
3.46




^1H EC AzoTP BI
single_pulse

7.96
7.95
7.94
7.93
7.63
7.62
7.60
7.59
7.56
7.54
7.53
7.52
7.44
7.42
7.41


^1H EC AzoTP BI
single_pulse

3.50
3.48
3.46





^{13}C EC AzoTP


153.3111
151.5076
138.1453
132.6905
132.6031
130.5110
128.7470
123.6646
116.6135
68.1464
68.0853
49.6515
49.3247
49.168
48.988
48.809
48.629
48.3438
32.8882
32.8200




¹H NMR and ¹³C NMR spectra of 5




¹H NMR and ¹³C NMR spectra of 6




^1H NMR and ^{13}C NMR spectra of **1c**



¹H NMR and ¹³C NMR spectra of **2c**




DM sub AzoMP protected ¹³C




¹H NMR and ¹³C NMR spectra of **3c**

Dimethyl Azo monophosphate triethyl salt re
single_pulse






DM sub AzoMP triethyl salt 13C




¹H NMR and ¹³C NMR spectra of **3d**

¹H DM sub AzoTP BI
single_pulse




¹³C DM sub AzoTP



31P NMR spectra of 4c, 4d, 4b.


31P DM sub AzoTP
single pulse decoupled gated NOE




— 0.00 Phosphoric acid

10.84
10.96
12.42
12.54

22.95
23.06
23.17



31P EC AzoTP
single pulse decoupled gated NOE



— 0.00 Phosphoric acid

10.71
10.83
10.89
11.01
11.06
11.19

23.16
23.28
23.40

