



HOKKAIDO UNIVERSITY

Title	The data of change in macrophage gene expression which induced by perilipin 1 overexpression
Author(s)	Yamamoto, Kohei; Miyoshi, Hideaki; Cho, Kyu Yong et al.
Citation	Data in Brief, 19, 179-182 https://doi.org/10.1016/j.dib.2018.05.027
Issue Date	2018-08
Doc URL	https://hdl.handle.net/2115/72098
Rights(URL)	https://creativecommons.org/licenses/by/4.0/
Type	journal article
File Information	1-s2.0-S2352340918305304-main.pdf





Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

The data of change in macrophage gene expression which induced by perilipin 1 overexpression



Kohei Yamamoto^a, Hideaki Miyoshi^{a,*}, Kyu Yong Cho^a,
Akinobu Nakamura^a, Andrew S. Greenberg^b, Tatsuya Atsumi^a

^a Department of Rheumatology, Endocrinology and Nephrology, Faculty of Medicine, Graduate School of Medicine, Hokkaido University, Kita 15 Nishi 7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan

^b Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, 711 Washington Street, Boston, MA 02111, USA

ARTICLE INFO

Article history:

Received 22 January 2018

Received in revised form

17 April 2018

Accepted 7 May 2018

Available online 15 May 2018

Keywords:

perilipin1

PLIN1

Macrophage

Atherosclerosis

ABSTRACT

The data presented here are related to the research article entitled “Overexpression of Perilipin1 protects against atheroma progression in apolipoprotein E knockout mice” [1]. This paper describes data that were obtained from perilipin 1 (PLIN1) transgenic mice (*Plin1Tg*) regarding atherosclerosis. The main aim of collecting the data was to clarify the role of PLIN1 in the pathophysiology of atherosclerosis. The data were collected from C57BL/6J mice, apolipoprotein E knockout mice (*ApoeKO*) and *Plin1Tg/ApoeKO*. The atherosclerotic lesion areas of aorta were $3.3 \pm 1.2\%$ in C57BL/6J mice, $14.2 \pm 3.2\%$ in *ApoeKO*, and $5.6 \pm 1.9\%$ in *Plin1Tg/ApoeKO*. Body weight, gonadal adipose mass and plasma triglyceride concentrations were comparable among the three groups [1]. Furthermore, PLIN1 overexpression did not affect the gene expressions related to cholesterol influx and efflux in macrophage.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <https://doi.org/10.1016/j.atherosclerosis.2018.01.019>

Abbreviations: PLIN1, perilipin 1; *Plin1Tg*, PLIN1 transgenic mice; *ApoeKO*, apolipoprotein E knockout mice; RT-PCR, reverse transcription polymerase chain reaction

* Corresponding author. Fax: +81 11 706 7710.

E-mail address: hmiyoshi@med.hokudai.ac.jp (H. Miyoshi).

<https://doi.org/10.1016/j.dib.2018.05.027>

2352-3409/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Medicine
More specific subject area	Atherosclerosis
Type of data	Table and text file
How data was acquired	Real-Time PCR (Applied Biosystems 7500 Fast Real-Time PCR System, Thermo Fisher)
Data format	Analyzed
Experimental factors	Data obtained from C57BL/6J mice, apolipoprotein E knockout mice and Plin1 transgenic mice
Experimental features	The effect of PLIN1 overexpression on atherosclerosis
Data source location	Hokkaido University, Sapporo, Japan
Data accessibility	The data are available with this article

Value of the data

- Overexpression of PLIN1 in macrophages protected against atheroma progression.
- No major risk factors were altered in PLIN1 transgenic mice fed normal diet.
- Overexpression of PLIN1 did not affect the gene expressions related to cholesterol influx and efflux in macrophage

1. Data

Thioglycollate-elicited peritoneal macrophages were isolated from C57BL/6J mice or perilipin 1 (PLIN1) transgenic mice (*Plin1Tg*). Total RNA was prepared and analyzed by reverse transcription polymerase chain reaction (RT-PCR). Expression of the PLIN1 transgene and endogenous PLIN1 and PLIN2 genes in the *Plin1Tg* macrophages was confirmed. Body weight were 26.6 ± 3.1 g in C57BL/6J mice, 29.0 ± 4.5 g in apolipoprotein E knockout mice (*ApoeKO*), and 27.5 ± 3.9 g in *Plin1Tg/ApoeKO*. Gonadal fat mass were 356 ± 78 mg in C57BL/6J mice, 332 ± 124 mg in *ApoeKO*, and 424 ± 190 mg in *Plin1Tg/ApoeKO*. Plasma total cholesterol were 72 ± 11 mg/dl in C57BL/6J mice, 395 ± 80 mg/dl in *ApoeKO*, and 471 ± 138 mg/dl in *Plin1Tg/ApoeKO*. Plasma tumor necrosis factor-alpha were undetectable in all groups. Plasma interleukin-6 were 22 ± 14 pg/ml in C57BL/6J mice, 64 ± 37 pg/ml in *ApoeKO*, and 28 ± 31 pg/ml in *Plin1Tg/ApoeKO* [1].

The size of the atherosclerotic lesions were examined in the aortic sinus area and in the whole aorta using an en face method with Oil Red O staining. The lesions were quantified as a percentage of total aorta area. The atherosclerotic lesion areas of aorta were $3.3 \pm 1.2\%$ in C57BL/6J mice, $14.2 \pm 3.2\%$ in *ApoeKO*, and $5.6 \pm 1.9\%$ in *Plin1Tg/ApoeKO*. [1].

Peritoneal thioglycollate-elicited macrophages were induced by acute inflammation and might be different in character from macrophages in plaques. We therefore used cultured human macrophages derived from monocytes to assess the expression of genes involved in cholesterol uptake and efflux. Although the CD36 expression level in PLIN1 overexpressed macrophages was 1.2 times higher than the control value, PLIN1 overexpression did not affect gene expression levels of SR-A, ABCA1 or ABCG1 (Fig. 1).




Fig. 1. Relative mRNA Expression (cultured human macrophage).

Table 1

Primers used for detection of transgene expression.

	Forward primer	Reverse primer
<i>hPLIN1</i>	TCTCTCGATACCCGTGCAG	AGGGCTGCTACCTCACTGAA
<i>mPLIN1</i>	TGAAGGGTGTACGGATAACG	ATGTCTCGGAATTCGCTCTC
<i>PLIN2</i>	GATTGAATTCGCCAGGAAGA	TGGCATGTAGTCTGGAGCTG
<i>GAPDH</i>	AACCTTGGCATTGTGGAAGG	ACACATTGGGGTAGGAACA

2. Experimental design, materials, and methods

2.1. Animal experiments

Plin1Tg were generated using the aP2 promotor on a C57BL/6J background as previously described. The *PLIN1* expression level in white adipose tissue in *Plin1Tg* was shown to be two times higher than that in control mice [2]. *ApoeKO* were purchased from the Jackson Laboratory (Bar Harbor, United States). C57BL/6J mice were purchased from Charles River Japan (Yokohama, Japan). *Plin1Tg* and *ApoeKO* were crossed to obtain *Plin1Tg/ApoeKO* mice. The mice were housed at the Graduate School of Medicine's Institute for Animal Experimentation at Hokkaido University in accordance with the institutional guidelines of Hokkaido University Graduate School of Medicine. All mice were housed at room temperature, maintained on a 12 h light/dark cycle, and given free access to water.

All mice received a normal chow diet (MF from Oriental Yeast, Tokyo, Japan) for 20 weeks. Body weight and gonadal fat mass were measured. Blood was collected from inferior vena cava and plasma was obtained by centrifugation for enzymatic determination (Wako, Tokyo, Japan; R&D Systems, Minneapolis, United States) of lipid concentrations and proinflammatory cytokine levels. Aortic sinuses and whole aortas were collected for quantification of atheroma lesions. Data are expressed as means $\pm$ SD.

2.2. RT-PCR analysis

Thioglycollate-elicited macrophages were isolated from C57BL/6J mice and *Plin1Tg* by washing the peritoneal cavity with 3 ml of phosphate-buffered saline one day after the mice were intraperitoneally injected with 50 μ l of 4% thioglycollate in phosphate-buffered saline. Individual cell suspensions were washed with red blood cell lysis buffer (eBioscience, San Diego, United States). Total RNA was isolated from the isolated macrophages using an RNeasy Mini kit (QIAGEN, Venlo, Netherlands) according to the manufacturer's recommendations, and was used as the starting material for cDNA preparation. RT-PCR was performed using ReverTra-Plus (Toyobo, Osaka, Japan) in accordance with the manufacturer's protocols. Primer sequences are shown in Table 1.

Table 2
Primers used for Real-Time PCR.

	Forward primer	Reverse primer
SR-A	CCAGGGACATGGAATGCAA	CCAGTGGGACCTCGATCTCC
CD36	CTGTCATTGGTGCTGTCCTG	CTCAGCGTCCTGGGTACAT
ABCA1	ACCTGCTGCCCTACAGTGAT	ATGGAGACCGAAGTGGTGAG
ABCG1	TGCAATCTTGTCATATTGA	CCAGCCGACTGTTCTGATCA

2.3. Real-Time PCR analysis

Human monocytes were isolated from healthy control. Monocytes were incubated in RPMI 1640 (Thermo Fisher) with 10% pooled human serum at 37 degree 6 days, and derived to macrophage. Total RNA was isolated from the cultured macrophages using an RNeasy Mini kit (QIAGEN, Venlo, Netherlands) according to the manufacturer's recommendations. Real-Time PCR was performed using Applied Biosystems 7500 Fast Real-Time PCR System (Thermo Fisher) in accordance with the manufacturer's protocols. Data are expressed as means $\pm$ SD. Primer sequences are shown in Table 2.

Acknowledgments

We thank N Fujimori and M Watanabe for technical assistance. HM thanks Dr. James W. Perfield II (University of Missouri, MO, USA) for his continued support and mentorship. We thank Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

Financial support

Not applicable.

Transparency document. Supporting information

Transparency data associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2018.05.027>.

References

[1] K. Yamamoto, et al., Overexpression of Perilipin1 protects against atheroma progression in apolipoprotein E knockout mice. *Atherosclerosis*, 2018. "in press".
[2] H. Miyoshi, et al., Perilipin overexpression in mice protects against diet-induced obesity, *J. Lipid Res.* 51 (5) (2010) 975–982.