



HOKKAIDO UNIVERSITY

Title	Comment on "Elevation of Serum Carcinoembryonic Antigen Concentration Caused by Everolimus-Induced Lung Injury : A Case Report"
Author(s)	Nomoto, Hiroshi; Miyoshi, Hideaki; Sekizaki, Tomonori et al.
Citation	Annals of Thoracic and Cardiovascular Surgery, 24(3), 165-166 https://doi.org/10.5761/atcs.lte.18-00018
Issue Date	2018
Doc URL	https://hdl.handle.net/2115/72099
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	24_lte.18-00018.pdf



**Letter to
the Editor**

Comment on “Elevation of Serum Carcinoembryonic Antigen Concentration Caused by Everolimus-Induced Lung Injury: A Case Report”

Hiroshi Nomoto, MD, PhD,¹ Hideaki Miyoshi, MD, PhD,¹ Tomonori Sekizaki, MD,²
and Tatsuya Atsumi, MD, PhD¹

Keywords: everolimus, carcinoembryonic antigen, adverse events

We read with great interest the report of elevated serum carcinoembryonic antigen (CEA) following everolimus administration,¹⁾ in which it was reported that everolimus treatment for renal cell carcinoma with lung metastasis caused pulmonary ground-glass opacities accompanied by elevated serum CEA. Furthermore, interrupting everolimus resulted in improvements of both lung shadows and elevated CEA levels. As there have been no reports of correlations between everolimus and elevated CEA to date, the authors concluded that the elevated CEA could be caused by everolimus-induced lung injury.

CEA is one of the most widely used tumor markers, and serum CEA levels increase in gastrointestinal cancers, as well as other types of carcinomas. However, serum CEA also rises in non-neoplastic conditions, such as inflammatory diseases of the glandular epithelium, biliary obstruction, and metabolic disorders including hyperglycemia.²⁾ Among these metabolic disorders, hypothyroidism is a frequent endocrinological disorder associated with elevated

levels of certain tumor markers. It has been reported that a significantly high proportion of patients with Hashimoto's disease are CEA positive in the hypothyroid condition, and that hormone replacement therapy reduced CEA levels in these patients.³⁾ Indeed, we recently experienced two non-neoplastic patients with severe hypothyroidism accompanied by drastically elevated CEA levels, which decreased with levothyroxine replacement therapy alone (**Table 1**). Although the precise mechanism for such deranged CEA elevation is not yet clear, it has been assumed that hypothyroidism affects CEA metabolism or hepatic clearance because CEA is metabolized and excreted by the liver.^{3,4)} It is also well known that certain multi-kinase inhibitors, including sorafenib, which the authors administered just prior to everolimus, could induce thyroid dysfunction, especially in the Japanese population.^{5,6)} Sorafenib can cause destructive thyroiditis followed by severe hypothyroidism, and even after discontinuing sorafenib, permanent or transient hypothyroidism can occur. From this point of view, we wondered whether hypothyroidism might have occurred after switching from sorafenib to everolimus, contributing, in part, to the elevated CEA.

However, as the authors mentioned, everolimus can cause some adverse metabolic events, including hyperglycemia and dyslipidemia, both of which are known to trigger elevated CEA levels.^{2,7)} Oxidative stress and incrementally increased insulin-like growth factor (IGF)-1 levels due to hyperinsulinemia are assumed to be causes of CEA elevation.^{8,9)} Based on these ideas, we would like to understand the clinical courses and changes in thyroid function and metabolic parameters before and after everolimus administration.

¹Department of Rheumatology, Endocrinology and Nephrology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Sapporo, Japan

²Department of Diabetes and Lipid Metabolism Internal Medicine, Hokkaido Medical Center, Sapporo, Japan

Received: January 18, 2018; Accepted: February 20, 2018

Corresponding author: Hideaki Miyoshi, MD, PhD. Department of Rheumatology, Endocrinology and Nephrology, Faculty of Medicine and Graduate School of Medicine, Hokkaido University, Kita 15, Nishi 7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan
Email: hmiyoshi@med.hokudai.ac.jp

©2018 The Editorial Committee of *Annals of Thoracic and Cardiovascular Surgery*. This work is licensed under a Creative Commons Attribution-NonCommercial-NonDerivatives International License.

Table 1 Relationship between thyroid function and serum carcinoembryonic antigen levels before and after levothyroxine treatment

Variable	Case 1		Case 2	
	Before	After	Before	After
Thyroid stimulating hormone (μ U/mL)	146	2.68	170	1.39
Free T4 (ng/dL)	0.172	1.45	0.087	1.27
Carcinoembryonic antigen (ng/mL)	21.3	4.2	26.9	11.4

Molecularly targeted therapeutic agents are useful tools to treat unresectable advanced carcinomas; however, it is important to appropriately manage adverse endocrinological/metabolic effects. Additionally, we would like to advocate the importance of remembering that some metabolic and endocrinological disorders can increase the levels of certain tumor marker levels, such as unexplainably high serum CEA levels.

Disclosure Statement

H.M. has received honoraria for lectures from Astellas Pharma Inc., AstraZeneca, Dainippon Sumitomo Pharma Co., Ltd., Eli Lilly, Kissei, Kowa Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., MSD, Novartis Pharma, Novo Nordisk Pharma, Takeda Pharmaceutical Co., Ltd., Ono Pharmaceutical Co., Ltd., and Sanofi; and received basic research funding from Astellas Pharma Inc. T.A. has received honoraria for lectures from Mitsubishi Tanabe Pharma Co., Chugai Pharmaceutical Co., Ltd., Astellas Pharma Inc., Takeda Pharmaceutical Co., Ltd., Pfizer Inc., AbbVie Inc., Eisai Co., Ltd., Daiichi Sankyo Co., Ltd., Bristol-Myers Squibb Co., UCB Japan Co., Ltd., and Eli Lilly Japan K.K.; and received basic research funding from Astellas Pharma Inc., Takeda Pharmaceutical Co., Ltd., Mitsubishi Tanabe Pharma Co., Chugai Pharmaceutical Co., Ltd., Daiichi Sankyo, and Otsuka Pharmaceutical Co., Ltd., Pfizer Inc., and Alexion Inc. H.N. and T.S. have no potential conflicts of interest to declare in relation to the publication of this article.

References

- 1) Komatsu H, Izumi N, Tsukioka T, et al. Elevation of serum carcinoembryonic antigen concentration caused by everolimus-induced lung injury: a case report. *Ann Thorac Cardiovasc Surg* [Epub ahead of print] doi: 10.5761/atcs.cr.17-00092.
- 2) Shang X, Song C, Du X, et al. The serum levels of tumor marker CA19-9, CEA, CA72-4, and NSE in type 2 diabetes without malignancy and the relations to the metabolic control. *Saudi Med J* 2017; **38**: 204-8.
- 3) Amino N, Kuro R, Yabu Y, et al. Elevated levels of circulating carcinoembryonic antigen in hypothyroidism. *J Clin Endocrinol Metab* 1981; **52**: 457-62.
- 4) Shuster J, Silverman M, Gold P. Metabolism of human carcinoembryonic antigen in xenogeneic animals. *Cancer Res* 1973; **33**: 65-8.
- 5) van Doorn L, Eskens FA, Visser TJ, et al. Sorafenib induced thyroiditis in two patients with hepatocellular carcinoma. *Thyroid* 2011; **21**: 197-202.
- 6) Miyake H, Kurahashi T, Yamanaka K, et al. Abnormalities of thyroid function in Japanese patients with metastatic renal cell carcinoma treated with sorafenib: a prospective evaluation. *Urol Oncol* 2010; **28**: 515-9.
- 7) Kim KN, Joo NS, Je SY, et al. Carcinoembryonic antigen level can be overestimated in metabolic syndrome. *J Korean Med Sci* 2011; **26**: 759-64.
- 8) Zayed AA, Beano AM, Amer FN, et al. Serum levels of carcinoembryonic antigen in patients with type 2 diabetes. *Endocr Pract* 2016; **22**: 1310-8.
- 9) Ahmed RL, Schmitz KH, Anderson KE, et al. The metabolic syndrome and risk of incident colorectal cancer. *Cancer* 2006; **107**: 28-36.