



HOKKAIDO UNIVERSITY

Title	Inter- and intra-species differences in muscarinic acetylcholine receptor expression in the neural pathways for learned vocalization in songbirds [an abstract of dissertation and a summary of dissertation review]
Author(s)	ASOGWA, Chinweike Norman
Degree Grantor	北海道大学
Degree Name	博士(生命科学)
Dissertation Number	甲第13386号
Issue Date	2018-12-25
Doc URL	https://hdl.handle.net/2115/72357
Rights(URL)	https://creativecommons.org/licenses/by-nc-sa/4.0/
Type	doctoral thesis
File Information	Chinweike_Norman_ASOGWA_abstract.pdf, 論文内容の要旨



Abstract of Doctoral Dissertation

Degree requested Doctor of Life Science Applicant's name: Chinweike Norman ASOGWA

Title of Doctoral Dissertation

Inter- and intra-species differences in muscarinic acetylcholine receptor expression in the neural pathways for learned vocalization in songbirds

(鳴禽類歌神経回路において種内個体差・種間差発現を示すムスカリン性アセチルコリン受容体の研究)

Acetylcholine receptors (AChRs) abound in the central nervous system. Muscarinic AChRs (mAChRs), a functional subclass of AChRs, mediate neuronal responses via intracellular signal transduction. They also play roles in sensorimotor co-ordination and motor skill learning, such as song learning by enhancing cortical plasticity. Songbirds, such as the zebra finch, acquire their songs, a complex sequence of vocal patterns, through sensorimotor coordination during a critical learning period. However, the functions of AChRs in the neural circuits for vocal learning during song learning remain largely unexplored. I investigated the precise expression patterns of chrms in the major brain subdivisions, during the critical period of vocal learning, among songbird species, in early-deafened zebra finches. Further, I examined the effects of overexpressing chrn2 on crystallized adult song in the zebra finch. I found that chrn1 is not expressed in the songbirds whereas there is unique expression of mAChRs subclass (chrn2–5) in the song systems for vocal learning and production in the zebra finch. The expression of excitatory subunits (chrn3 and chrn5) was downregulated in the song nuclei compared with the surrounding regions. In contrast, inhibitory mAChRs (chrn2 and chrn4) were upregulated in the premotor song nucleus HVC relative to the surrounding caudal nidopallium. Chr4 showed developmentally different expression level in HVC during the critical period of song learning. Compared with chrn4, individual differences in chrn2 expression level in HVC emerged during the early stage of the critical period. These individual differences in chrn2 expression persisted despite testosterone administration or auditory deprivation, which altered the timing of song sing stabilization in the zebra finch. Instead, the variability in chrn2 expression level in HVC correlated with parental genetics. In addition, chrn2 expression level in HVC showed species differences and individual variability among songbird species. In early-deafened zebra finches, chrn2 expression level correlated positively and negatively with the motif and repetition indices of song, respectively. In zebra finches at the subsong stage, chrn2 expression level correlated with the coefficient of variation in syllable entropy variance of songs. In adult zebra finches, over-expressing CHRM2 protein had no clear effects on crystallized learned song. These results suggest that mAChRs may contribute to the development of individual and species differences in learned song patterns by modulating the excitability of HVC neurons.