



# HOKKAIDO UNIVERSITY

|                  |                                                                                                                                                                                                          |
|------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Title            | Dietary intakes of fat soluble vitamins as predictors of mortality from heart failure in a large prospective cohort study                                                                                |
| Author(s)        | Eshak, Ehab S.; Iso, Hiroyasu; Yamagishi, Kazumasa et al.                                                                                                                                                |
| Citation         | Nutrition, 47, 50-55<br><a href="https://doi.org/10.1016/j.nut.2017.09.009">https://doi.org/10.1016/j.nut.2017.09.009</a>                                                                                |
| Issue Date       | 2018-03                                                                                                                                                                                                  |
| Doc URL          | <a href="https://hdl.handle.net/2115/72728">https://hdl.handle.net/2115/72728</a>                                                                                                                        |
| Rights           | © 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license<br><a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a> |
| Rights(URL)      | <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>                                                                                      |
| Type             | journal article                                                                                                                                                                                          |
| File Information | Nutrition47_50.pdf                                                                                                                                                                                       |



## **Dietary intakes of fat soluble vitamins as predictors for mortality from heart failure in a large prospective cohort study**

**Running head:** Fat soluble vitamins and heart failure death.

Ehab S. Eshak<sup>1,2\*</sup>, Ph.D; Hiroyasu Iso<sup>1</sup>, Ph.D; Kazumasa Yamagishi<sup>3</sup>, Ph.D; Renzhe Cui<sup>1</sup>, Ph.D; Akiko Tamakoshi<sup>4</sup>, Ph.D.

<sup>1</sup> Public Health, Department of Social Medicine, Osaka University, Graduate School of Medicine, Osaka, Japan

<sup>2</sup> Department of Public Health and Preventive Medicine, Faculty of Medicine, Minia University, Minia, Egypt

<sup>3</sup> Department of Public Health Medicine, Faculty of Medicine, University of Tsukuba, Tsukuba, Japan

<sup>4</sup> Department of Public Health, Hokkaido University Graduate School of Medicine, Hokkaido, Japan.

### **Authors' contributions**

Hiroyasu Iso and Akiko Tamakoshi made a design of this research; Eshak S. Ehab conducted the analyses; Eshak S. Ehab wrote the manuscript; Hiroyasu Iso, Kazumasa Yamagishi, Renzhe Cui and Akiko Tamakoshi made critical revision for the manuscript. All authors read and approved the final manuscript.

**Word Count:** 4,303

**Number of tables:** 2

**Supplementary material:** online supporting materials: 1 Figure, 1 Table.

**Correspondence to** Ehab S Eshak, Public Health, Department of Social Medicine, Osaka University, Graduate School of Medicine, Suita Shi, Osaka 565-0871, Japan.

Phone: +81-6-6879-3911    Fax: +81-6-6879-3919    E-mail: [ehab@pbhel.med.osaka-u.ac.jp](mailto:ehab@pbhel.med.osaka-u.ac.jp)

## **Abstract**

**Objective:** A few reports have investigated the association of dietary vitamin intakes with risk of heart failure in Asia. Therefore, we examined the relation of dietary intakes of fat-soluble vitamins A, K E, and D with mortality from heart failure in Japanese population.

**Research Methods and Procedures:** A total of 23,099 men and 35,597 women aged 40-79 years participated in the Japan Collaborative Cohort Study and completed a food frequency questionnaire, from which dietary intakes of vitamin A, K, E and D were calculated. Cox proportional hazard model was used to estimate the sex-specific risk of heart failure mortality according to increasing quintiles of fat soluble vitamin intakes.

**Results:** During median 19.3-year follow-up, there were 567 deaths from heart failure (395 men and 307 women). Dietary vitamin A intake showed no association with heart failure mortality in both genders; contrary, the reduced risk was observed in women but not in men with dietary intakes of vitamin K, E and D. The multivariable HRs (95% CI) in the highest versus the lowest intake quintiles among women were 0.63 (0.45, 0.87; P for trend=0.006) for vitamin K, 0.55 (0.36, 0.78; P for trend=0.006) for vitamin E and 0.66 (0.48, 0.93; P for trend= 0.01) for vitamin D. The association for each vitamin was slightly attenuated but remained statistically significant after mutual adjustment for other vitamins.

**Conclusions:** High dietary intakes of fat soluble vitamins K, E and D were associated with reduced risk of heart failure mortality in Japanese women but not men.

## **Highlights**

\* Intakes of fish, natto, fruits and vegetables associated with low heart failure risk.

\*These foods are major sources of vitamin D, K, A and E in Japanese diet.

\*Intakes of vitamins K, E and D were associated with reduced death from heart failure. \*The reduced risk of heart failure mortality was evident in Japanese women but not men.

**Keywords:** dietary vitamin A, dietary vitamin K, dietary vitamin E, dietary vitamin D, heart failure mortality, Japanese.

## Introduction

Heart failure is complex syndrome stands on activation of numerous biological mechanisms that follow or coincide with myocardial injury and leads to inadequate systematic perfusion.

(1) In developed countries with high proportion of aging society, like in Japan, heart failure resembles a notable public health burden. (2) Intakes of seafood and fish, (3,4) fermented soybeans (5) and fruits and vegetables (5,6) were associated with lower risk of heart failure.

In Japanese diet, seafood and fish are main dietary contributors of vitamin D; (7) natto (fermented soybeans) is a major source to vitamin K, (8) while vitamins A and E are mainly provided by fruit and vegetables. (9) It remains unclear whether the observed reduced risks with higher intakes of these foods are attributed to the effect of their content of fat soluble vitamins (A, K, E and D).

Dietary antioxidant capacity was inversely associated with risk of cardiovascular diseases including heart failure. (10) However, the association of dietary intakes of vitamin A and E with cardiovascular mortality is questionable.(11-13) There is also some conflicting observations for the impact of vitamin K intake on cardiovascular health. (14-17) In addition there is growing evidence on the protective effects of vitamin D against cardiovascular disease (18,19) including heart failure. (18,20)

Taking together these controvert findings, the associations between the dietary intakes of the fat soluble vitamins and risk of heart failure still need investigation, especially in Asian population. Therefore, we thought to examine the association between dietary intakes of vitamins A, K, E and D with risk of mortality from heart failure among men and women of the Japan Collaborative Cohort study (JACC), a large prospective study.

## Participants and Methods

### *Study Population and baseline covariates*

With a total of 110,585 middle aged (40-79 years) Japanese men (n= 46,395) and women (n=64,190) from 45 communities across Japan, the Ministry of Education, Sports and Science started in 1988 to 1990 to sponsor the JACC Study. Detailed description for the JACC Study was published previously. (21)

Baseline lifestyle and participants' characteristics including demographic data; medical histories of chronic diseases, diabetes mellitus and hypertension; habits of ethanol drinking, smoking, exercise, diet and others were inquired via a self-administered questionnaire that included a 40 food-item food frequency questionnaire (FFQ). If participants did not give answers to  $\geq 4$  items of FFQ or reported histories of either cancer or CVD they were excluded. We also excluded from analysis participants with missing information about fat soluble vitamin intake which left a total of 58,646 eligible individuals for the study (23,099 males and 35,547 females) (**Supplemental Figure I**).

### *Diet by FFQ*

The usual consumption frequency for each food item over the past 12 months, without specifying portion size was inquired through choosing from five frequency responses: rarely, 1-2 times/m, 1-2 times/wk, 3-4 times/wk, and almost every day. (21) These frequencies were transformed to a weekly consumption scores 0, 0.38, 1.5, 3.5, and 7.0/wk, respectively.

Dietary intakes of fat soluble vitamins (A, K, E and D) were calculated by using the Japan Food Tables, Fifth Edition to determine the amount of fat soluble vitamins in each food.

These contents were multiplied by the participant's frequency score for each food item in the FFQ, followed by summing the contents. A validation study among 85 individuals using four 3-d weighed dietary records (DR) over a 1-year period as a reference standard determined the portion size for each food and validated the FFQ intakes. The Spearman rank correlation

coefficients for vitamin A, K, E and D intakes between the FFQ and four 3-days DR were 0.37, 0.40, 0.38 and 0.39. The mean  $\pm$  SD intakes in mg/day from the FFQ were  $967 \pm 446$  for vitamin A,  $180 \pm 64$  for vitamin K,  $4.9 \pm 1.5$  for vitamin E and  $7.8 \pm 3.7$  for vitamin D; whereas the respective intakes from the DR were  $1089 \pm 543$ ,  $279 \pm 104$ ,  $9.1 \pm 1.9$  and  $10.6 \pm 4.5$  mg/day.(22)

### ***Mortality surveillance***

As part of the mortality surveillance in each resident area, investigators conducted a systematic review of death certificates which were centralised at the Ministry of Health and Welfare. The codes of International Classification for Diseases 10th Revision were used to specify the underlying causes of deaths (our primary outcome was heart failure, code 150). This death certificate ascertainment was applied to all deaths within our cohort, except for deaths occurred out of the original resident areas, which were treated as censored cases.

### ***Statistical analysis***

Calorie-adjustment by the residual method (23) was done to the dietary intakes of our exposure variables which were then modeled as categorical (5 quintiles) groups in the main analysis. The significance of the differences in proportion and mean of participants' characteristics and known cardiovascular risk factors across calorie-adjusted quintiles of fat soluble vitamins intakes were tested by  $\chi^2$ -test and the analysis of covariance. Person-years of follow-up were calculated as the period from submission of the questionnaire to either departure of a participant from his/her original residential area, death or termination of follow-up at the end of 2009, whichever came first. Follow-up was terminated in in two areas in 2008, in four study areas in 2003 and in another four areas in 1999.

By means of Cox proportional model, sex-specific age- and multivariable-adjusted hazard ratios (HRs) their 95% confidence intervals (95% CIs) for risk of mortality from heart failure across increasing quintiles of each vitamin. The hypothesized confounders were

history of diabetes and hypertension (yes or no); use of multivitamin supplementation (yes or no); hours of walking (almost never, 0.5, 0.6-0.9, and  $\geq 1$  hour/day); hours of sports (almost never, 1-2, 3-4 and  $\geq 5$  hours/week); smoking category (never, ex-smoker, current smoker of 1-19 and  $\geq 20$  cigarettes/day); quintiles of body mass index (BMI), years of education ( $\leq 15$ , 16-18 and  $>18$  years); perceived mental stress (low, medium and high) and ethanol intake (never, ex-drinker, current drinker of 0.1-22.9, 23.0-45.9, 46.0-68.9, and  $\geq 69.0$  g ethanol/day); calorie-adjusted quintiles of calcium, sodium, potassium, saturated fatty acids, n-3 fatty acids, total dietary fiber and vitamin C intakes and total calorie intakes. The impact of mutual adjustment for each fat soluble vitamin intake was tested in an additional model. We conducted tests for trends across quintiles of vitamins intake by assigning median values for each quintile and testing the significance of this variable. We further analyzed the data after exclusion of persons who died within first 3, 5 and 10 years of follow-up to examine a potential effect of as-yet-undiagnosed diseases at baseline. Two-tailed statistical tests with  $P < 0.05$  regarded as statistically significant were applied by the use of SAS statistical package (version 9.4, SAS).

## Results

As shown in **Table 1** participants in the highest quintile of fat soluble vitamins intake compared with those in the lowest quintile were older, more educated, less likely to smoke and consumed more sodium, potassium, calcium, saturated fat, n-3 fatty acids and dietary fibre; but consumed less alcohol. Moreover, men in the highest intake quintiles of vitamins A, K and D and women in the highest intake quintiles of vitamin A and K were more likely to be hypertensive and diabetics.

During 965, 970 person-years of follow-up, there were 567 deaths (395 men and 307 women) from heart failure. Dietary intake of vitamin A was not associated with heart failure

mortality in either gender. Higher intakes of dietary vitamins K, E and D were associated with lower mortality risk of heart failure among women but not men. The multivariable HRs (95% CI) of heart failure mortality in the highest versus lowest intake quintiles among women were 0.63 (0.45, 0.87; P for trend=0.006) for vitamin K, 0.55 (0.36, 0.78; P for trend=0.006) for vitamin E and 0.66 (0.48, 0.93; P for trend= 0.01) for vitamin D were. The respective HRs (95% CIs) among men were 1.02 (0.67, 1.56; P for trend= 0.61), 1.10 (0.73, 1.86; P for trend= 0.81), and 1.04 (0.72, 1.51; P for trend= 0.26) (**Table 2, model 2**). No material changes were observed in the above reported associations after mutual adjustment for the fat soluble vitamins (**Table 2, model 3**) or after excluding heart failure mortalities that occurred with the first 3, 5 and 10 years of follow-up (**Supplemental Table 1**).

## Discussion

Following 58,646 Japanese men and women for a median 19.3 years in a large prospective cohort study revealed that higher dietary intakes of vitamin K, E and D were associated with reduced risk of heart failure mortality in women only. Dietary intake of vitamin A was not related to heart failure mortality in either gender. These associations remained statistically significant even after controlling for known cardiovascular risk factors, mutual adjustment of fat soluble vitamins and excluding early events within 3-10 years of the follow-up.

The major dietary sources of fat soluble vitamins in Japanese diet are seafood/fish, soybeans and fruit and vegetables;(7-9) dietary recommendations to prevent heart failure suggest increase intakes of these foods.(3-6) Higher intakes of fat soluble vitamins, especially vitamin D may help improving the cardiac function, hypertrophy and contractility.(24,25) Lower intakes of fat soluble vitamins were associated with some evoking factors of heart failure such as diabetes,(26) hypertension,(27) atherosclerosis(28) and coronary heart disease.(29) Moreover, some studies suggested beneficial roles of high dietary intakes of

antioxidants 10 and vitamin D status,(18-20) vitamin E supplementation(30) regarding the risk of heart failure. However, other studies showed no benefits of vitamins E and D supplementation (31-33) or even insignificant increased risk of heart failure with higher long-term supplementation of vitamin E.(34)

The observed gender difference regarding the association between dietary fat soluble vitamins intake and heart failure in our study was also evident in previous Japanese studies that investigated not only the association of food sources of vitamins with cardiovascular incidence and mortality,(35,36) but also in studies investigated the direct associations of water and fat soluble vitamins with cardiovascular mortality.(12, 37) Dietary vitamin C intakes were inversely associated with mortality from cardiovascular diseases in Japanese women but not men; meanwhile, although the associations did not reach a level of significance, the risk of cardiovascular mortality was lower in women than men with higher dietary intakes of vitamins A and E.(12) Another Japanese study showed inverse associations of higher dietary intakes of folate and vitamin B6 with mortality from stroke, coronary heart disease and total cardiovascular diseases in women but not in men; whereas the reduced risk of heart failure was observed in men but not in women.(37) The exact reasons for the gender difference are not clear. The effect of dietary fat soluble vitamins may be suppressed by other factors or was not enough to counteract overwhelming heart failure risk factors in men such as smoking and drinking habits and history of hypertension, which were more prevalent in our men than women as shown in **Table 1**.

To the best of our knowledge this is the first study to show an association between dietary fat soluble vitamins with risk of heart failure in Japanese population. Strengths of our study include its large sample size from a community-based cohort, its prospective design, the use of a validated FFQ and the consistent endpoint determination. Also, the exclusion of participants with known comorbidities before follow-up has reduced bias stem from dietary

modifications due to known chronic diseases. Moreover, the associations did not substantially change when we excluded early heart failure mortalities within 3-10 years from the baseline survey; which indicates few effect of pre-existing unknown illness or reverse causality.

Limitations of our research include the one-time measurement of dietary fat soluble vitamins; consumption is likely to change during the long follow-up period. Second, we adjusted for the use of multivitamin supplementation, yet we have no data about amounts or types of the used supplementary vitamins. We believe that this did not affect our results because in 1980s-90s the use of fat soluble vitamins supplementation was unpopular among Japanese, only 3.3 % of our participants reported the daily use of multivitamin supplementation, and sensitivity analysis by excluding those subjects who used multivitamin supplementation did not change the results materially (data not shown in tables). Another limitation was that only 54% of potential study participants responded to the FFQ; those respondents were younger and more educated than non-respondents; however, we adjusted these variables in our examination. Last, the FFQ underestimated intakes of fat soluble vitamins by at least a quarter, based on a validation study for our cohort; but, this systematic underestimation is supposed to be non-differential. Last, the use of death certificate to ascertain heart failure deaths in the early years of the study, before 1994, was questionable because most deaths of unknown origin like cardiac arrest or arrhythmic death, which are classified as ischemic heart disease deaths in the U.S.(38) were registered as “unspecified heart failure” (I50.9 for ICD-10) in Japan and accounted for 27% to 50% of diagnosed heart failure.(39) Accordingly, heart failure deaths in this study was contaminated with cardiac arrest, and that may have affected the association between dietary intakes of fat soluble vitamins and heart failure; however, sensitivity analyses by excluding heart failure mortalities occurred within 10 years of the follow-up showed similar associations.

## **Conclusions**

In this large community based prospective cohort study, higher dietary intakes of fat soluble vitamins (K, E and D) were associated with reduced risk of mortality from heart failure among Japanese women but not men.

**Funding:** This study was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture of Japan (61010076, 62010074, 63010074, 1010068, 2151065, 3151064, 4151063, 5151069, 6279102, 11181101, 17015022, 18014011, 20014026, and 20390156); and Comprehensive Research on Cardiovascular and Lifestyle Related Diseases (H26-Junkankitou [Seisaku]-Ippan-001).

**Conflict of interest:** none.

## References

- 1-Pacher R. Pathogenesis of heart failure. *Semin Nephrol.* 2001;21:273-7.
- 2-Konishi M, Ishida J, Springer J, et al. Heart failure epidemiology and novel treatments in Japan: facts and numbers. *ESC Heart Fail.* 2016;3:145-151.
- 3-Yamagishi K, Iso H, Date C, et al. Fish, omega-3 polyunsaturated fatty acids, and mortality from cardiovascular diseases in a nationwide community-based cohort of Japanese men and women the JACC (Japan Collaborative Cohort Study for Evaluation of Cancer Risk) Study. *J Am Coll Cardiol.* 2008;52:988-96.
- 4-Djoussé L, Akinkuolie AO, Wu JH, Ding EL, et al. Fish consumption, omega-3 fatty acids and risk of heart failure: a meta-analysis. *Clin Nutr.* 2012;31:846-53.
- 5-Nagura J, Iso H, Watanabe Y, et al. Fruit, vegetable and bean intake and mortality from cardiovascular disease among Japanese men and women: the JACC Study. *Br J Nutr.* 2009;102:285-92.
- 6-Rautiainen S, Levitan EB, Mittleman MA, et al. Fruit and vegetable intake and rate of heart failure: a population-based prospective cohort of women. *Eur J Heart Fail.* 2015;17:20-6.
- 7-Nakamura K, Nashimoto M, Okuda Y, et al. Fish as a major source of vitamin D in the Japanese diet. *Nutrition* 2002, 18, 415–416.
- 8-Kaneki M, Hodges SJ, Hosoi T, et al. Japanese fermented soybean food as the major determinant of the large geographic difference in circulating levels of vitamin K2: possible implications for hip-fracture risk. *Nutrition.* 2001;17:315-21.
- 9-Kato Y, Ikehara S, Maruyama K, et al. Trends in dietary intakes of vitamins A, C and E among Japanese men and women from 1974 to 2001. *Public Health Nutr.* 2009;12:1343–1350.

- 10-Rautiainen S, Levitan EB, Mittleman MA, et al. Total antioxidant capacity of diet and risk of heart failure: a population-based prospective cohort of women. *Am J Med*. 2013;126:494-500.
- 11-Asplund K. Antioxidant vitamins in the prevention of cardiovascular disease: a systematic review. *J Intern Med*. 2002;251:372–392.
- 12-Kubota Y, Iso H, Date C, et al. Dietary intakes of antioxidant vitamins and mortality from cardiovascular disease: the Japan Collaborative Cohort Study (JACC) study. *Stroke*. 2011;42:1665-72.
- 13-Paganini-Hill A, Kawas CH, Corrada MM. Antioxidant vitamin intake and mortality: the Leisure World Cohort Study. *Am J Epidemiol*. 2015;181:120-6.
- 14-Geleijnse JM, Vermeer C, Grobbee DE, et al. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr*. 2004;134:3100-5.
- 15-A Erkkilä AT, Booth SL, Hu FB, et al. Phylloquinone intake as a marker for coronary heart disease risk but not stroke in women. *Eur J Clin Nutr*. 2005;59:196-204.
- 16-Gast GC, de Roos NM, Sluijs I, et al. A high menaquinone intake reduces the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis*. 2009;19:504-10.
- 17-Dam V, Dalmeijer GW, Vermeer C, et al. Association between vitamin K and the metabolic syndrome: A 10-year follow-up study in adults. *J Clin Endocrinol Metab*. 2015;100:2472-9.
- 18-Barnett AG, de Looper M, Fraser JF. The seasonality in heart failure deaths and total cardiovascular deaths. *Aust N Z J Public Health*. 2008;32:408-13.
- 19-Wang L, Song Y, Manson JE, et al. Circulating 25-hydroxy-vitamin D and risk of cardiovascular disease: a meta-analysis of prospective studies. *Circ Cardiovasc Qual Outcomes* 2012;5:819-829.

- 20-Agarwal M, Phan A, Willix R Jr, et al. Is vitamin D deficiency associated with heart failure? A review of current evidence. *J Cardiovasc Pharmacol Ther.* 2011;16:354-63.
- 21-Tamakoshi A, Ozasa K, Fujino, et al. Cohort profile of the Japan Collaborative Cohort Study at final follow-up. *J Epidemiol.* 2013;23:227-232.
- 22-Date C, Fukui M, Yamamoto A, et al. Reproducibility and validity of a self-administered food frequency questionnaire used in the JACC Study. *J Epidemiol.* 2005;15:S9-23.
- 23-Willet WC, Stampfer MJ. Total energy intake: implication for epidemiological analysis. *Am J Epidemiol.* 1986;124:17-27.
- 24-Achinger SG, Ayus JC. The role of vitamin D in left ventricular hypertrophy and cardiac function. *Kidney Int Suppl.* 2005;(95):S37-42.
- 25-Berezin AE. Vitamin D deficiency and heart failure: possible mechanisms and predictive value. *J Dis Markers.* 2014;1: 1014.
- 26-Martine C. The role of vitamins in the prevention and treatment of type 2 diabetes and its complications. *Journal of diabetes Nursing.* 2013;17:376-381.
- 27-Llopis-González A, Rubio-López N, Pineda-Alonso M, et al. Hypertension and the fat-soluble vitamins A, D and E. *Int J Environ Res Public Health.* 2015;12:2793-809.
- 28-Ozkanlar S, Akcay F. Antioxidant vitamins in atherosclerosis--animal experiments and clinical studies. *Adv Clin Exp Med.* 2012;21:115-23.
- 29-Moats C, Rimm EB. Vitamin intake and risk of coronary disease: observation versus intervention. *Curr Atheroscler Rep.* 2007;9:508-14.
- 30-Bauersachs J, Fleming I, Fraccarollo D, et al. Prevention of endothelial dysfunction in heart failure by vitamin E: attenuation of vascular superoxide anion formation and increase in soluble guanylyl cyclase expression. *Cardiovasc Res.* 2001;51:344-50.
- 31-Chae CU, Albert CM, Moorthy MV, et al. Vitamin E supplementation and the risk of heart failure in women. *Circ Heart Fail.* 2012;5:176-82.

- 32-Keith ME, Jeejeebhoy KN, Langer A, et al. A controlled clinical trial of vitamin E supplementation in patients with congestive heart failure. *Am J Clin Nutr.* 2001;73:219-24.
- 33-Joris PJ, Mensink RP. Effects of supplementation with the fat-soluble vitamins E and D on fasting flow-mediated vasodilation in adults: A meta-analysis of randomized controlled trials. *Nutrients.* 2015 ;7:1728-43.
- 34-Lonn E, Bosch J, Yusuf S, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *Journal of American Medical Associations* 2005; 293: 1338-47.
- 35-Kokubo Y, Iso H, Ishihara J, et al. Association of dietary intake of soy, beans, and isoflavones with risk of cerebral and myocardial infarctions in Japanese populations: the Japan Public Health Center-Based (JPHC) study cohort I. *Circulation.* 2007;116: 2553 – 2562.
- 36-Nakamura K, Nagata C, Oba S, et al. Fruit and vegetable intake and mortality from cardiovascular disease are inversely associated in Japanese women but not in men. *J Nutr.* 2008;138:1129 – 1134.
- 37-Cui R, Iso H, Date C, et al. Dietary folate and vitamin b6 and B12 intake in relation to mortality from cardiovascular diseases: Japan collaborative cohort study. *Stroke.* 2010;41:1285-9.
- 38-Saito I, Folsom AR, Aono H, et al. Comparison of fatal coronary heart disease occurrence based on population surveys in Japan and the USA. *Int J Epidemiol.* 2000;29:837-844
- 39-Saito I. Review of death certificate diagnosis of coronary heart disease and heart failure in Japan. *Nippon Koshu Eisei Zasshi (Jpn J Public Health).* 2004;51: 909-916.

**Table 1.** Participants’ characteristics according to energy-adjusted quintiles of dietary intake of fat soluble vitamins

|                                    | Vitamin A               |             |             | Vitamin K   |             |             | Vitamin E   |             |             | Vitamin D   |             |             |
|------------------------------------|-------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
|                                    | Q1                      | Q3          | Q5          | Q1          | Q3          | Q5          | Q1          | Q3          | Q5          | Q1          | Q3          | Q5          |
| <b>Men, <i>n</i></b>               | 4619                    | 4620        | 4620        | 4619        | 4620        | 4620        | 4619        | 4620        | 4620        | 4619        | 4620        | 4620        |
| Age, year                          | 53.9 ± 9.2 <sup>a</sup> | 55.8 ± 10.1 | 57.9 ± 9.8  | 53.5 ± 9.2  | 56.1 ± 10.0 | 58.7 ± 9.9  | 53.5 ± 9.2  | 54.7 ± 9.7  | 58.8 ± 9.9  | 54.3 ± 9.6  | 56.0 ± 10.2 | 57.9 ± 9.4  |
| History of hypertension, %         | 18                      | 19          | 20          | 17          | 19          | 21          | 18          | 19          | 19          | 17          | 19          | 20          |
| History of diabetes, %             | 4                       | 7           | 7           | 5           | 6           | 7           | 5           | 6           | 7           | 5           | 7           | 7           |
| Body mass index, kg/m <sup>2</sup> | 22.8 ± 2.8              | 22.8 ± 2.7  | 22.5 ± 2.7  | 22.8 ± 2.8  | 22.7 ± 2.8  | 22.6 ± 2.7  | 22.8 ± 2.8  | 22.7 ± 2.8  | 22.7 ± 2.7  | 22.7 ± 2.8  | 22.6 ± 2.7  | 22.7 ± 2.8  |
| Current smoker, %                  | 59                      | 54          | 51          | 60          | 54          | 48          | 62          | 53          | 47          | 57          | 53          | 53          |
| Ethanol intake, gm/day             | 40.1 ± 24.6             | 32.7 ± 21.3 | 33.2 ± 21.5 | 39.6 ± 24.9 | 32.3 ± 21.6 | 30.4 ± 19.8 | 41.2 ± 25.2 | 32.5 ± 20.9 | 28.2 ± 19.2 | 37.1 ± 24.1 | 32.1 ± 21.3 | 33.4 ± 21.2 |
| Sports ≥5 hour/week, %             | 18                      | 21          | 20          | 19          | 20          | 21          | 19          | 20          | 20          | 19          | 21          | 20          |
| Walking ≥5 hour/week, %            | 51                      | 48          | 52          | 49          | 49          | 53          | 49          | 49          | 53          | 51          | 48          | 51          |
| >18 years education,%              | 14                      | 17          | 17          | 14          | 17          | 18          | 14          | 18          | 19          | 15          | 17          | 18          |
| High mental stress,%               | 25                      | 25          | 22          | 26          | 25          | 22          | 26          | 25          | 22          | 24          | 25          | 24          |
| Vitamin supplementation, %         | 3                       | 3           | 4           | 3           | 4           | 4           | 3           | 4           | 4           | 3           | 4           | 4           |
| Sodium intake, mg/day              | 1763 ± 765              | 2229 ± 690  | 2478 ± 775  | 1515 ± 611  | 2286 ± 644  | 2746 ± 657  | 1433 ± 572  | 2231 ± 569  | 2893 ± 645  | 1741 ± 751  | 2191 ± 663  | 2674 ± 691  |
| Calcium intake, mg/day             | 383 ± 132               | 510 ± 130   | 581 ± 162   | 369 ± 129   | 505 ± 127   | 626 ± 137   | 352 ± 119   | 504 ± 116   | 463 ± 130   | 419 ± 147   | 507 ± 143   | 578 ± 152   |
| Potassium intake, mg/day           | 1794 ± 448              | 2354 ± 450  | 2721 ± 626  | 1710 ± 412  | 2328 ± 401  | 2955 ± 475  | 1653 ± 372  | 2313 ± 337  | 3025 ± 443  | 1910 ± 524  | 2332 ± 500  | 2748 ± 574  |
| Saturated fat intake, mg/day       | 7.5 ± 2.7               | 9.7 ± 2.7   | 11.2 ± 3.4  | 7.7 ± 2.9   | 9.7 ± 2.9   | 11.3 ± 3.3  | 7.0 ± 2.5   | 9.6 ± 2.5   | 12.3 ± 3.0  | 7.8 ± 3.0   | 9.7 ± 2.9   | 11.2 ± 3.2  |
| N-3 fatty acids intake, mg/day     | 1.3 ± 0.5               | 1.7 ± 0.5   | 2.0 ± 0.6   | 1.1 ± 0.4   | 1.7 ± 0.5   | 2.1 ± 0.6   | 1.0 ± 0.3   | 1.6 ± 0.3   | 2.4 ± 0.5   | 1.0 ± 0.4   | 1.6 ± 0.3   | 2.4 ± 0.4   |
| Dietary fiber intake, g/day        | 8.1 ± 2.2               | 9.5 ± 2.2   | 11.9 ± 3.1  | 7.1 ± 1.7   | 10.4 ± 1.7  | 13.6 ± 2.2  | 7.1 ± 1.6   | 10.2 ± 1.6  | 13.7 ± 2.2  | 8.7 ± 2.5   | 10.4 ± 2.6  | 11.8 ± 3.0  |
| Total energy intake, kcal/day      | 1942 ± 486              | 1666 ± 459  | 1779 ± 505  | 1812 ± 495  | 1674 ± 478  | 1783 ± 447  | 1824 ± 498  | 1687 ± 490  | 1803 ± 492  | 1860 ± 464  | 1640 ± 497  | 1809 ± 462  |

|                                    |                         |             |             |             |             |            |             |             |             |             |             |             |
|------------------------------------|-------------------------|-------------|-------------|-------------|-------------|------------|-------------|-------------|-------------|-------------|-------------|-------------|
|                                    |                         |             |             |             |             |            |             |             |             |             |             |             |
|                                    |                         |             |             |             |             |            |             |             |             |             |             |             |
|                                    |                         |             |             |             |             |            |             |             |             |             |             |             |
| Vitamin A intake, g/day            | 461 ± 142               | 920 ± 52    | 2241 ± 1194 | 731 ± 656   | 1071 ± 712  | 1530 ± 974 | 695 ± 574   | 4067 ± 697  | 1554 ± 1029 | 826 ± 645   | 1102 ± 734  | 1347 ± 1018 |
| Vitamin K intake, g/day            | 119 ± 45                | 185 ± 54    | 230 ± 73    | 93 ± 22     | 175 ± 11    | 286 ± 27   | 105 ± 36    | 179 ± 42    | 299 ± 48    | 138 ± 60    | 185 ± 61    | 225 ± 68    |
| Vitamin E intake, g/day            | 3.6 ± 1.1               | 5.0 ± 1.1   | 6.1 ± 1.6   | 3.2 ± 0.9   | 5.0 ± 0.9   | 6.8 ± 1.2  | 3.0 ± 0.6   | 4.9 ± 0.2   | 7.2 ± 0.9   | 3.9 ± 1.2   | 5.0 ± 1.1   | 6.3 ± 1.4   |
| Vitamin D intake, g/day            | 6.1 ± 3.2               | 7.8 ± 3.1   | 9.1 ± 3.6   | 5.7 ± 3.0   | 7.9 ± 3.0   | 9.9 ± 3.5  | 4.9 ± 2.4   | 7.7 ± 2.7   | 10.9 ± 3.2  | 3.5 ± 1.1   | 7.3 ± 0.5   | 13.0 ± 1.8  |
|                                    |                         |             |             |             |             |            |             |             |             |             |             |             |
| <b>Women, <i>n</i></b>             | 7019                    | 7019        | 7019        | 7019        | 7019        | 7019       | 7019        | 7019        | 7019        | 7019        | 7019        | 7019        |
| Age, year                          | 55.5 ± 9.8 <sup>a</sup> | 56.5 ± 10.1 | 56.7 ± 9.6  | 54.7 ± 9.8  | 56.0 ± 10.0 | 58.4 ± 9.5 | 55.7 ± 9.9  | 56.0 ± 10.0 | 57.7 ± 9.4  | 56.1 ± 10.1 | 55.8 ± 10.0 | 57.0 ± 9.4  |
| History of hypertension, %         | 20                      | 22          | 18          | 19          | 20          | 23         | 20          | 21          | 20          | 20          | 20          | 20          |
| History of diabetes, %             | 3                       | 4           | 5           | 3           | 3           | 4          | 3           | 4           | 4           | 3           | 3           | 4           |
| Body mass index, kg/m <sup>2</sup> | 23.1 ± 3.2              | 23.0 ± 3.1  | 22.7 ± 3.0  | 23.0 ± 3.2  | 22.9 ± 3.1  | 23.0 ± 3.1 | 23.0 ± 3.2  | 22.9 ± 3.0  | 22.9 ± 3.0  | 22.9 ± 3.2  | 22.9 ± 3.1  | 23.1 ± 3.1  |
| Current smoker, %                  | 6                       | 4           | 4           | 6           | 4           | 4          | 7           | 5           | 3           | 6           | 5           | 4           |
| Ethanol intake, gm/day             | 13.8 ± 19.3             | 9.2 ± 10.8  | 9.1 ± 11.7  | 13.8 ± 19.8 | 8.7 ± 9.7   | 8.6 ± 10.5 | 14.9 ± 20.5 | 8.8 ± 9.6   | 7.8 ± 8.9   | 12.5 ± 18.7 | 9.1 ± 11.3  | 9.8 ± 11.6  |
| Sports ≥5 hour/week, %             | 19                      | 21          | 22          | 20          | 21          | 21         | 19          | 22          | 21          | 20          | 22          | 20          |
| Walking ≥5 hour/week, %            | 53                      | 50          | 53          | 50          | 50          | 55         | 52          | 51          | 53          | 54          | 50          | 53          |
| >18 years education,%              | 7                       | 10          | 12          | 8           | 10          | 11         | 8           | 10          | 11          | 9           | 10          | 10          |
| High mental stress,%               | 22                      | 22          | 20          | 23          | 21          | 19         | 21          | 21          | 19          | 21          | 22          | 20          |
| Vitamin supplementation, %         | 3                       | 3           | 4           | 3           | 3           | 3          | 3           | 3           | 3           | 3           | 3           | 3           |
| Sodium intake, mg/day              | 1760 ± 729              | 2126 ± 623  | 2202 ± 656  | 1529 ± 589  | 2147 ± 595  | 2506 ± 577 | 1434 ± 565  | 2099 ± 519  | 2612 ± 281  | 1641 ± 688  | 2204 ± 569  | 2493 ± 600  |
| Calcium intake, mg/day             | 397 ± 134               | 530 ± 125   | 587 ± 143   | 400 ± 136   | 521 ± 123   | 618 ± 126  | 378 ± 129   | 524 ± 115   | 627 ± 123   | 444 ± 152   | 519 ± 136   | 575 ± 139   |
| Potassium intake, mg/day           | 1874 ± 435              | 1461 ± 43   | 2730 ± 508  | 1854 ± 430  | 2414 ± 370  | 2928 ± 397 | 1784 ± 392  | 2423 ± 316  | 2961 ± 383  | 2037 ± 524  | 2403 ± 444  | 2741 ± 484  |
| Saturated fat intake, mg/day       | 7.7 ± 2.9               | 10.0 ± 2.7  | 11.3 ± 3.0  | 8.3 ± 2.0   | 10.0 ± 2.8  | 11.0 ± 3.1 | 7.4 ± 2.7   | 10.0 ± 2.5  | 12.0 ± 2.9  | 8.3 ± 3.2   | 10.0 ± 2.58 | 10.9 ± 2.9  |
| N-3 fatty acids intake, mg/day     | 1.3 ± 0.5               | 1.7 ± 0.5   | 1.8 ± 0.5   | 1.2 ± 0.5   | 1.7 ± 0.4   | 2.0 ± 0.5  | 1.0 ± 0.4   | 1.6 ± 0.3   | 2.3 ± 0.4   | 1.1 ± 0.4   | 1.6 ± 0.3   | 2.3 ± 0.4   |
| Total dietary fiber intake/day     | 8.6 ± 2.2               | 10.9 ± 2.3  | 11.7 ± 2.5  | 7.8 ± 1.8   | 10.7 ± 1.6  | 13.5 ± 1.9 | 7.8 ± 1.7   | 10.7 ± 1.6  | 13.3 ± 2.1  | 9.3 ± 2.5   | 10.6 ± 2.3  | 11.9 ± 2.5  |

|                               |            |            |             |            |            |            |            |           |            |            |            |            |
|-------------------------------|------------|------------|-------------|------------|------------|------------|------------|-----------|------------|------------|------------|------------|
| Total energy intake, kcal/day | 1603 ± 399 | 1379 ± 331 | 1448 ± 373  | 1479 ± 404 | 1391 ± 350 | 1475 ± 336 | 1503 ± 422 | 1380 ± 33 | 1514 ± 352 | 1523 ± 381 | 1358 ± 358 | 1505 ± 337 |
| Vitamin A intake, g/day       | 494 ± 158  | 942 ± 51   | 2227 ± 1237 | 775 ± 726  | 1093 ± 742 | 1480 ± 951 | 747 ± 674  | 111 ± 743 | 1489 ± 979 | 903 ± 772  | 1127 ± 751 | 1265 ± 879 |
| Vitamin K intake, g/day       | 130 ± 45   | 201 ± 56   | 232 ± 63    | 105 ± 25   | 186 ± 11   | 286 ± 22   | 120 ± 40   | 191 ± 43  | 165 ± 44   | 180 ± 57   | 193 ± 58   | 226 ± 62   |
| Vitamin E intake, g/day       | 3.9 ± 1.1  | 5.4 ± 1.1  | 6.1 ± 1.3   | 3.7 ± 1.0  | 5.3 ± 0.8  | 6.8 ± 1.0  | 3.4 ± 0.7  | 5.2 ± 0.2 | 7.1 ± 0.7  | 4.1 ± 1.2  | 5.3 ± 1.0  | 6.4 ± 1.2  |
| Vitamin D intake, g/day       | 6.4 ± 3.3  | 8.2 ± 3.0  | 8.9 ± 3.3   | 6.1 ± 3.0  | 8.0 ± 2.9  | 9.7 ± 3.3  | 5.3 ± 2.5  | 8.0 ± 2.6 | 10.8 ± 3.1 | 3.8 ± 1.2  | 7.5 ± 0.5  | 12.9 ± 1.7 |

<sup>a</sup> Means ± standard deviation, all such variables.

| Table 2. Sex-specific associations between dietary intakes of fat soluble vitamins and risk of heart failure mortality |        |                  |                  |                  |                  |                      |        |                  |                  |                  |                  |                      |
|------------------------------------------------------------------------------------------------------------------------|--------|------------------|------------------|------------------|------------------|----------------------|--------|------------------|------------------|------------------|------------------|----------------------|
|                                                                                                                        | Men    |                  |                  |                  |                  |                      | Women  |                  |                  |                  |                  |                      |
|                                                                                                                        | 1(low) | 2                | 3                | 4                | 5(high)          | P-trend <sup>a</sup> | 1(low) | 2                | 3                | 4                | 5(high)          | P-trend <sup>a</sup> |
| <b>Vitamin A</b>                                                                                                       |        |                  |                  |                  |                  |                      |        |                  |                  |                  |                  |                      |
| Person years, n                                                                                                        | 77213  | 75429            | 74935            | 73202            | 71439            |                      | 120743 | 120743           | 120743           | 120743           | 120743           | 120743               |
| Cases, n                                                                                                               | 39     | 34               | 49               | 67               | 51               |                      | 66     | 76               | 70               | 53               | 62               |                      |
| Model 1 <sup>b</sup>                                                                                                   | 1.00   | 0.75 (0.47-1.18) | 0.96 (0.63-1.46) | 1.13 (0.76-1.68) | 0.86 (0.57-1.31) | 0.13                 | 1.00   | 1.00 (0.72-1.38) | 0.87 (0.62-1.21) | 0.68 (0.47-0.98) | 0.91 (0.64-1.29) | 0.47                 |
| Model 2 <sup>c</sup>                                                                                                   | 1.00   | 0.79 (0.50-1.25) | 1.03 (0.67-1.58) | 1.27 (0.84-1.91) | 0.92 (0.60-1.41) | 0.14                 | 1.00   | 1.00 (0.72-1.39) | 0.88 (0.62-1.23) | 0.66 (0.46-0.95) | 0.92 (0.65-1.31) | 0.51                 |
| Model 3 <sup>d</sup>                                                                                                   | 1.00   | 0.83 (0.51-1.36) | 1.14 (0.70-1.85) | 1.40 (0.85-2.32) | 0.96 (0.60-1.62) | 0.16                 | 1.00   | 1.11 (0.77-1.60) | 1.05 (0.71-1.55) | 0.83 (0.54-1.29) | 1.23 (0.80-1.89) | 0.33                 |
| <b>Vitamin K</b>                                                                                                       |        |                  |                  |                  |                  |                      |        |                  |                  |                  |                  |                      |
| Person years, n                                                                                                        | 75354  | 74890            | 75232            | 73719            | 73028            |                      | 117502 | 117502           | 117502           | 117502           | 117502           |                      |
| Cases, n                                                                                                               | 35     | 42               | 53               | 46               | 64               |                      | 70     | 60               | 67               | 55               | 75               |                      |
| Model 1 <sup>b</sup>                                                                                                   | 1.00   | 0.98 (0.62-1.53) | 0.99 (0.64-1.52) | 0.82 (0.53-1.28) | 0.94 (0.62-1.42) | 0.40                 | 1.00   | 0.70 (0.50-0.99) | 0.72 (0.52-1.01) | 0.57 (0.40-0.80) | 0.62 (0.45-0.86) | 0.005                |
| Model 2 <sup>c</sup>                                                                                                   | 1.00   | 0.99 (0.63-1.56) | 1.06 (0.69-1.63) | 0.91 (0.58-1.42) | 1.02 (0.67-1.56) | 0.61                 | 1.00   | 0.70 (0.50-1.00) | 0.73 (0.52-1.02) | 0.57 (0.40-0.81) | 0.63 (0.45-0.87) | 0.006                |
| Model 3 <sup>d</sup>                                                                                                   | 1.00   | 1.04 (0.62-1.74) | 1.11 (0.67-2.11) | 1.06 (0.58-1.94) | 1.14 (0.59-2.23) | 0.90                 | 1.00   | 0.68 (0.46-1.00) | 0.68 (0.44-1.04) | 0.54 (0.33-0.86) | 0.63 (0.38-0.94) | 0.02                 |
| <b>Vitamin E</b>                                                                                                       |        |                  |                  |                  |                  |                      |        |                  |                  |                  |                  |                      |
| Person years, n                                                                                                        | 74774  | 74358            | 75262            | 74429            | 73380            |                      | 115792 | 115792           | 115792           | 115792           | 115792           |                      |
| Cases, n                                                                                                               | 32     | 47               | 50               | 43               | 68               |                      | 76     | 54               | 62               | 76               | 59               |                      |
| Model 1 <sup>b</sup>                                                                                                   | 1.00   | 1.19 (0.76-1.87) | 1.07 (0.68-1.67) | 0.80 (0.51-1.27) | 1.07 (0.70-1.63) | 0.38                 | 1.00   | 0.69 (0.48-0.97) | 0.70 (0.50-0.97) | 0.82 (0.60-1.03) | 0.55 (0.39-0.77) | 0.005                |
| Model 2 <sup>c</sup>                                                                                                   | 1.00   | 1.28 (0.81-2.01) | 1.16 (0.74-1.83) | 0.92 (0.58-1.47) | 1.10 (0.73-1.86) | 0.81                 | 1.00   | 0.67 (0.47-0.95) | 0.70 (0.50-0.98) | 0.80 (0.58-1.01) | 0.55 (0.36-0.78) | 0.006                |
| Model 3 <sup>d</sup>                                                                                                   | 1.00   | 1.35 (0.74-2.45) | 1.11 (0.63-2.34) | 0.87 (0.49-1.68) | 0.99 (0.64-1.62) | 0.20                 | 1.00   | 0.65 (0.41-1.02) | 0.65 (0.39-1.04) | 0.76 (0.47-1.04) | 0.57 (0.29-0.98) | 0.04                 |
| <b>Vitamin D</b>                                                                                                       |        |                  |                  |                  |                  |                      |        |                  |                  |                  |                  |                      |
| Person years, n                                                                                                        | 75176  | 73968            | 73847            | 74875            | 74356            |                      | 116251 | 116251           | 116251           | 116251           | 116251           |                      |

| Cases, n             | 47   | 36               | 39               | 46               | 72               |      | 78   | 67               | 65               | 54               | 63               |      |
|----------------------|------|------------------|------------------|------------------|------------------|------|------|------------------|------------------|------------------|------------------|------|
| Model 1 <sup>b</sup> | 1.00 | 0.65 (0.42-1.01) | 0.63 (0.41-0.96) | 0.75 (0.50-1.12) | 1.00 (0.69-1.45) | 0.29 | 1.00 | 0.82 (0.59-1.14) | 0.82 (0.59-1.15) | 0.64 (0.45-0.90) | 0.68 (0.49-0.95) | 0.01 |
| Model 2 <sup>c</sup> | 1.00 | 0.68 (0.44-1.05) | 0.65 (0.43-1.00) | 0.77 (0.51-1.16) | 1.04 (0.72-1.51) | 0.26 | 1.00 | 0.77 (0.55-1.07) | 0.79 (0.57-1.11) | 0.60 (0.42-0.85) | 0.66 (0.48-0.93) | 0.01 |
| Model 3 <sup>d</sup> | 1.00 | 0.67 (0.42-1.08) | 0.62 (0.36-1.06) | 0.67 (0.37-1.22) | 0.85 (0.44-1.65) | 0.92 | 1.00 | 0.78 (0.55-1.09) | 0.83 (0.49-1.17) | 0.63 (0.33-0.92) | 0.69 (0.39-0.99) | 0.05 |

<sup>a</sup> Median values of vitamins intakes in each quintile were used to test for a linear trend across quintiles.

<sup>b</sup> Model 1 Age-adjusted hazard ratio (95% CI) by Cox proportional model.

<sup>c</sup> Model 2 Adjusted further for histories of hypertension and diabetes, smoking status, body mass index, hours of walking, hours of sports, educational status, perceived mental stress and alcohol intake, multivitamin supplementation, quintiles of energy-adjusted sodium, calcium, potassium, saturated fatty acids, n-3 fatty acids, dietary fiber and vitamin C, and total energy intakes.

<sup>d</sup> Model 3 Adjusted further mutually for the other 3 fat soluble vitamins.