



HOKKAIDO UNIVERSITY

Title	Intra-ooplasmic injection of a multiple number of sperm to induce androgenesis and polyploidy in the dojo loach <i>Misgurnus anguillicaudatus</i> (Teleostei: Cobitidae)
Author(s)	Yasui, George Shigueki; Saito, Taiju; Zhao, Yan et al.
Citation	Zygote, 26(5), 408-416 https://doi.org/10.1017/S0967199418000448
Issue Date	2018-10
Doc URL	https://hdl.handle.net/2115/73420
Rights	This article has been published in a revised form in Zygote https://doi.org/10.1017/S0967199418000448 . This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © copyright holder.
Type	journal article
File Information	Zygote_26(5)_408.pdf



Intra-ooplasmic injection of a multiple number of sperm to induce androgenesis and polyploidy in the dojo loach *Misgurnus anguillicaudatus* (Teleostei: Cobitidae)

Running title: Androgenesis resulting from polyspermy

George Shigueki Yasui^{1,2}, *Taiju Saito*^{3,4}, *Yan Zhao*^{1,5}, *Takafumi Fujimoto*^{1,*}, *Etsuro Yamaha*³, and *Katsutoshi Arai*¹

¹Faculty and Graduate School of Fisheries Sciences, Hokkaido University, Hakodate, Hokkaido, Japan.

²Laboratory of Fish Biotechnology, National Center for Research and Conservation of Continental Fish, Chico Mendes Institute of Biodiversity Conservation, Rodovia Pref. Euberto Nemésio Pereira de Godoy, Pirassununga, Sao Paulo State, Brazil (present address).

³Nanae Fresh-Water Laboratory, Field Science Center for Northern Biosphere, Sakura, Nanae, Kameda, Hokkaido, Japan.

⁴South Ehime Fisheries Research Center, Ehime University, Uchidomari, Ainan, Ehime, Japan (present address).

⁵College of Fisheries and Life Science, Shanghai Ocean University, Shanghai, China (present address).

***Corresponding author**

Takafumi Fujimoto

Faculty and Graduate School of Fisheries Sciences, Hokkaido University,
Hakodate, Hokkaido, Japan.

Email: fujimoto@fish.hokudai.ac.jp.

Tel/Fax: +81-138-40-5536

Summary

Polyspermy was initiated by microinjecting a multiple number of sperm into the activated and dechorionated eggs of dojo loach *Misgurnus anguillicaudatus* (Teleostei: Cobitidae). A 10-nL sperm suspension from an albino (recessive trait) male (10^5 , 10^6 , 10^7 , or 10^8 sperm mL^{-1}) was microinjected into eggs from a wild-type female. Although the rates of embryos developing into the blastula stage in the injection group at the highest sperm concentration were similar to that of the control group, the hatching rates of the injection group were much lower. A large proportion of embryos that developed from the injected eggs were haploid and mosaics containing haploid cells. Most of the haploid and mosaic embryos inherited only paternally derived alleles in the microsatellite markers (i.e., androgenesis was initiated by injecting multiple sperm). In contrast, some haploid embryos contained both paternal and maternal alleles despite haploidy, suggesting that they were mosaics consisting of cells with either paternal or maternal inheritance. The injected eggs displayed diploid, hypotriploid, and triploid cells, all of which included both maternally and paternally derived alleles. One albino tetraploid with only paternal alleles was also observed from the injected eggs. These results suggest that part of the sperm microinjected into the ooplasm should form a male pronucleus(-i), which could develop by androgenesis or could fuse with the female pronucleus(-i). Thus, microinjecting multiple sperm is considered a potential technique by which to induce androgenesis and polyploidy.

Keywords: Fertilization, Fish, ICSI, Pronucleus, Spermatozoa

Introduction

Artificial fertilization by intracytoplasmic sperm injection (ICSI) is a technique initiated by microinjecting a sperm into the cytoplasm of an egg. ICSI has been applied not only in assisted reproduction technology to resolve problems related to human male infertility, but also as a potential tool to analyze basic biological issues; however, some technical constraints to this technique are specific to teleosts, and thus only a small number of trials on these fish have been reported (Poleo *et al.*, 2001, 2005; Otani *et al.*, 2009). It is difficult to transfer the protocol established in mammals to teleosts because the fish egg envelope (chorion) is generally too thick, hard, and tough for microinjection. In addition, a mammalian egg can accept sperm at any site on the egg, but a teleost sperm can enter the teleost egg only through the micropyle, an entrance located at the animal pole of the egg; therefore, all previous ICSI experiments on teleosts adopted the protocol of sperm microinjection through the micropyle with precise control of the injection point. The efficacies of ICSI in teleosts, or the successful rates of fertilization in these studies, were 1.6% in zebrafish (Poleo *et al.*, 2001), 8.5% in tilapia (Poleo *et al.*, 2005), and 13.4% in medaka (Otani *et al.*, 2009), which were lower than those recorded in mammals (e.g., 50% in mice) (Yanagimachi, 2005). Although injecting the sperm into teleosts through the micropyle at the animal pole of the egg is a prerequisite for successful ICSI, the differences in microscope magnifications between eggs and sperm apparently reduce microinjection maneuverability. Under the microscope, a single sperm is very small and the egg is comparatively large; therefore, it is impossible to observe both the egg and sperm simultaneously under the same microscopic view. Even with these challenges, ICSI remains an attractive approach to promote basic and applied studies in various fishes and environments, from model and endangered species to aquaculture. If sperm, including immature spermatids and non- or poorly motile sperm, can be appropriately cryopreserved for an extended period of time, commercially important and endangered genotypes could be reconstituted using ICSI (Wakayama & Yanagimachi, 1998; Mazur *et al.*, 2008); however, the major constraint to applying ICSI to teleosts is the technical difficulty of microinjecting a sperm through the micropyle and into a very thin layer of ooplasm at the animal pole of the egg.

When routine ICSI in teleosts is difficult, an alternative method is to inject several sperm into the egg. As far as we know, no previous attempts have been made to fertilize a teleost egg by microinjecting more than a single sperm and observing the subsequent production of viable progeny.

Thus, the purpose of this study was to precisely evaluate the biological effects of injecting multiple sperm into the teleost egg ooplasm. To this end, we induced embryogenesis by microinjecting multiple sperm into the ooplasm of the dojo loach (*Misgurnus anguillicaudatus* [Teleostei: Cobitidae]) egg, and studied the developmental ability, ploidy status, phenotypic expression (albino, recessive trait), and microsatellite DNA genotypes of the resultant progeny.

Materials and Methods

Ethics

This study was performed in accordance with the Guide for the Care and Use of Laboratory Animals in Hokkaido University, Japan. All animal experiments were approved by the Animal Studies Ethical Committee of Hokkaido University (approval number 19-2).

Induced maturation and fertilization procedures

Fertilization procedures were similar to those of our previous works (Fujimoto *et al.*, 2006). Adult diploid males with an albino phenotype (recessive trait) and wild-type diploid females (dominant trait) were used as the parents. Spermiation and ovulation were artificially induced by a single dose of 100 IU human chorionic gonadotropin for males and 500 IU for females (Aska Pharmaceuticals, Tokyo, Japan). After 12 h at 27°C, the gametes were sampled by stripping. Sperm was collected into capillary hematocrit tubes and the sperm content was homogenized in a 1.5-mL microtube containing 200 µL sperm medium (128.4 mM NaCl, 2.7 mM KCl, 1.4 mM CaCl₂, 2.4 mM NaHCO₃, pH 8.2; Kurokura *et al.*, 1984). The sperm samples were maintained at 4°C for storage. The motility of fresh sperm was always >80%.

Groups of 100–300 eggs were stripped onto plastic 90-mm-diameter Petri dishes covered with polyvinylidene chloride film (Saran wrap, Asahi Kasei, Tokyo, Japan). The eggs were fertilized using the routine dry method. Sperm (50 µL) was added to the egg mass and then activated by a 40-fold dilution with tap water dechlorinated by aging. After 10 min, the eggs were transferred to another Petri dish without the polyvinylidene chloride film and the water content was increased to 50 mL. The eggs were then incubated at 20°C. Any dead eggs, which were identified by their white appearance or delayed developmental stages, were counted and eliminated at 6-h intervals.

Sperm motility in embryo and sperm media

In this experiment, we confirmed whether the loach sperm was immotile in both the 200 mM KCl injection medium and the embryo medium comprising 1.6% egg albumen, 100 IU streptomycin, and

100 IU penicillin in Ringer's solution (7.5 g/L NaCl, 0.2 g/L KCl, and 0.2 g/L CaCl₂). Both media were diluted by distilled water to provide concentrations ranging from zero (i.e., distilled water) to 100% (undiluted). Sperm from three wild-type diploid males was also diluted with sperm media, as described above. To evaluate sperm motility, we used a glass slide coated with 0.01% bovine serum albumin to prevent the attachment of sperm cells to the glass surface. A small aliquot of sperm was placed on the glass slide and diluted 20-fold by sperm medium (Kurokura *et al.*, 1984). Sperm motility was measured as in previous studies (Yasui *et al.*, 2009). The percentages of all motile cells (total motility), all cells with straightforward movement along a linear track (progressive motility), and all non-motile cells were evaluated.

Optimization of injection procedure

The eggs were microinjected according to the protocol described in our previous studies (Saito *et al.*, 2006). Aliquots of sperm were diluted using a solution containing 200 mM KCl and 0.5% rhodamine. The dilutions provided sperm concentrations at 10⁵, 10⁶, 10⁷, and 10⁸ sperm mL⁻¹. Each sperm suspension was placed in a borosilicate capillary micropipette, with an open inner diameter of 10 µm. The micropipette was then loaded into the Eppendorf Celltram Vario microinjector with the Narishige M-150 micro-manipulator set (Japan) and the injection procedure was assessed using the Leica M165 FC stereoscopic microscope (Germany).

The dechorionated unfertilized eggs were prepared similar to the procedure described by Tanaka *et al.* (2009). Groups of 100–120 eggs were stripped onto a Petri dish, followed by the addition of 50 mL dechlorinated tap water. After 2 min, the perivitelline space was observed and the water was replaced with dechoriation medium (0.12% trypsin and 0.4% urea in Ringer's solution, modified from Yamaha *et al.*, 1986). After dechoriation (~5 min), the eggs were transferred to an agar-plated Petri dish containing embryo medium.

Approximately 10 nL sperm suspension was injected into the upper central area of the animal pole of each dechorionated egg. Considering the injection volume and sperm dilutions, the number of injected sperm was estimated to be 1, 10, 100, and 1,000 cells per injection.

We induced six types of control groups. The first control (Intact control: Co) was induced by fertilization between eggs and sperm before the injection trials. The second control (Intact control: Cf) was induced by fertilization between eggs and sperm after the injection trials. In these two controls, the fertilized eggs with their chorions intact were cultured in dechlorinated tap water. The third control (Dechorionated control) was induced by fertilization between eggs and sperm before

the injection trials, after which the chorion of the fertilized eggs was removed using the dechoriation medium, and the eggs were cultured in the embryo medium. The fourth control (Unfertilized control) consisted of eggs activated by only water. The fifth control (Uninjected control) was that treated with the sperm suspension at 10^8 sperm mL^{-1} poured onto the animal pole of the dechorionated eggs. The sixth control (Injected control) was that treated with a solution containing 200 mM KCl and 0.5% rhodamine microinjected into the dechorionated eggs.

Observation of embryonic development

To examine the developmental ability of eggs microinjected with the sperm solution, the eggs were observed at the following stages: 2-cell, blastula, 12-somite, and hatching of the control according to the criteria proposed by Fujimoto *et al.* (2006). Some eggs (approximately 30 embryos) were fixed at the blastula stage with 2% paraformaldehyde and 2% glutaraldehyde in phosphate buffer for later observation. Normal and abnormal larvae were counted at the hatching stage based on their external appearance. The presence or absence of melanophore pigments was examined on the surface of the larvae.

Ploidy determination and microsatellite DNA genotyping

DNA-content flow cytometry was used to measure the ploidy status of each embryo using the Ploidy Analyzer (Partec). Samples for flow cytometry were prepared according to the method of Fujimoto *et al.* (2007). DNA samples were extracted and purified from the parent fish and their progeny (20 individuals from the injection group at the highest sperm concentration and 10 individuals from the control) and then amplified by polymerase chain reaction using the primer sets of *Mac24*, *Mac345*, and *Mac449* according to Morishima *et al.* (2008). Genotyping was conducted using the I3130xdl automated sequencer (Applied Biosystems [ABI]) with GeneMapper v 3.7 (ABI).

Statistical analyses

Data are shown as the mean $\pm$ standard deviation. All experiments were conducted in triplicate using different egg sources. The obtained data were checked for normality using the Lilliefors test and compared using analysis of variance, followed by Tukey's multiple range test ($P > .05$).

Results

Loach sperm at 80–100% concentration were not motile in either the embryo or injection medium (Fig. 1). The embryo medium at 70% and the injection medium at 40% triggered motility, which was fully activated at lower concentrations (Fig. 1). Percentage of sperm showing progressive motility

was $84.0 \pm 2.6\%$, when the sperm was activated by contact with distilled water (Fig. 1). Thus, the quality of the sperm used here was good.

Co, Cf, and Dechorionated controls showed similar trends in all developmental stages, and hatching rates ranged between 68.6% and 76.5% (Table 1). Among the groups that contained eggs injected with sperm, the fertilization success expressed as a percentage of embryos at the two-cell stage increased with an increasing amount of sperm cells injected into the ooplasm. When only one sperm was injected into an egg, the fertilization rate was $7.2 \pm 5.9\%$, with no hatching. Injecting 10 sperm provided similar results, with a fertilization rate of $7.8 \pm 6.8\%$ and only two hatched embryos ($0.5 \pm 0.6\%$)—one non-pigmented abnormal albino larva and one pigmented normal wild-type larva. Approximately 100 sperm injected per egg increased both the fertilization rate ($24.9 \pm 9.9\%$) and hatching ($1.4 \pm 3.2\%$), but all were abnormal larvae (six non-pigmented albinos and one pigmented wild-type larva). The highest fertilization rate ($87.4 \pm 5.1\%$) was obtained at the highest sperm concentration (1,000 sperm per egg), with a hatching rate of $5.9 \pm 5.7\%$ (28 larvae) comprising five pigmented normal wild-type, three pigmented abnormal wild-type, and 20 non-pigmented abnormal albino larvae.

The eggs in the Unfertilized control did not show any cleavage. Some eggs began to exhibit cleavage-like segmentation in the Injected control, but did not develop further. In the Uninjected control eggs, some developed and viable larvae hatched.

The ploidy status of the progeny sampled from all the treatments before hatching is provided in Table 2. The progeny used for ploidy analysis comprised normal embryos with wild-type pigmentation and abnormal embryos with/without pigmentation. In the controls (i.e., Co, Cf, and Dechorionated), normal embryos were measured, and the majority (98–99%) were diploid, with only a small number of triploid progeny. In groups containing eggs injected with sperm, the progeny died before the beginning of pigmentation, but their ploidy status was estimated. The uninjected control gave six haploid, five mosaic, and two triploid progeny, in which normally pigmented wild-type progeny with haploid–triploid mosaicism ($n = 1$) and triploidy ($n = 2$) were observed. Injecting 1 sperm and 10 sperm gave two abnormal haploids (1 fish in each group). Injecting 100 sperm gave haploids ($n = 8$), a haploid–diploid mosaic ($n = 1$), and a hyperdiploid ($n = 1$), but all were abnormal. When approximately 1,000 sperm were injected, a predominant appearance of haploids was observed ($n = 44$), together with diploids ($n = 2$), triploids ($n = 7$), and various types of mosaicism ($n = 7$). In this group, the normal progeny were diploid ($n = 1$), triploid ($n = 6$), and haploid–triploid

mosaics (n = 1).

Most injected eggs exhibited abnormal development that was expressed especially during the blastula stage. Among the embryos developed from the injected eggs, unusual development was detected by observing blastomeres of different sizes as well as those containing anuclear cells (Fig. 2).

Figure 3 shows the external gross appearance, presence or absence of melanophore pigments (visible marker of paternal genome), and ploidy status based on flow cytometric analyses of progeny developed from fertilized eggs in the control groups and those injected with sperm. Control progeny with normal appearance and wild-type body pigmentation showed diploidy (Table 3, Figs. 3A1–3). Among the progeny of the injected groups (Table 3), haploid (Figs. 3G1–3), diploid (Figs. 3E1–3), triploid (Figs. 3B1–3), tetraploid (Figs. 3F1–3), hypotriploid (Figs. 3D1–3), and haploid–hypodiploid–hyperdiploid mosaic (Figs. 3C1–3) progeny were observed. Abnormal diploid (Fig. 3E2) and normal triploid progeny (Fig. 3B2) gave apparent expressions of melanophore pigments on their body surface, but other normal haploid–hypodiploid–hyperdiploid mosaic (Fig. 3C2) and abnormal hypotriploid (Fig. 3D2) progeny exhibited few pigments within a limited area on their body surfaces. Abnormal haploid (Figs. 3G1–3, H1, H2, H4), abnormal haploid–triploid mosaic (Figs. 3H1, H3, H5), and normal-looking tetraploid (Figs. 3F1–3) progeny without melanophore expression were also observed in the injected group.

Microsatellite DNA genotyping was conducted in the above-mentioned progeny from the injected group of approximately 1,000 sperm injected per egg (nos. 1–6 in Table 3), and the haploid, haploid–diploid, haploid–triploid, and haploid–pentaploid progeny of the same injected group showed extremely abnormal appearance and were sampled before hatching (nos. 7–20 in Table 3). All control group (normal wild-type) progeny had heterozygous genotypes consisting of alleles from both the female and male parents in the three independent loci that we examined. On the contrary, in the injected group, 12 (nos. 5, 7, 8, 10–14, and 17–20 in Table 3) of 20 progeny demonstrated only paternally derived alleles, and thus were androgenotes with an all-male inheritance (Table 3). Among these androgenetically developed progeny, only the tetraploid progeny (no. 5 in Table 3) exhibited a normal appearance (Figs. 3F1–3); the haploid, haploid–diploid, haploid–triploid, and haploid–pentaploid mosaics demonstrated inviable abnormalities (Figs. 3G1–3, H1–5, Table 3). From the Injected group, three haploid (nos. 6, 9, and 15 in Table 3) and one haploid–diploid mosaic (no. 16 in Table 3) had both maternally and paternally derived alleles and did not develop androgenetically

(Table 3). Among these non-androgenotes, one haploid (no. 6 in Table 3) demonstrated the albino phenotype, but also had microsatellite alleles from both parents. Four fully or partially pigmented progeny with triploidy (Figs. 3B1–3, no. 1 in Table 3), haploid–hypodiploid–hyperdiploidy (Figs. 3C1–3, no. 2 in Table 3), hypotriploidy (Figs. 3D1–3, no. 3 in Table 3), and diploidy (Figs. 3E1–3, no. 4 in Table 3) were presumably initiated by fertilization with injected sperm because they showed both maternally and paternally derived alleles at three loci (Table 3). Among these progeny, triploid (Figs. 3B1–3, no. 1 in Table 3) and haploid–hypodiploid–hyperdiploid mosaic progeny (Figs. 3C1–3, no. 2 in Table 3) displayed a normal appearance (Table 3).

Discussion

We confirmed that sperm were not motile in either the embryo or the injection medium. We injected a sperm suspension into the ooplasm of each dojo loach egg that was activated in advance with ambient water, followed by dechoriation. As with zebrafish (Paleo *et al.*, 2001), the dojo loach eggs are activated by contact with water, as evidenced in Tanaka *et al.* (2009). Embryos developed in the groups that contained eggs injected with sperm. This suggests that fertilization or embryogenesis should be the result of injected or non-injected non-motile cells reaching the oocytes. Increasing the number of injected sperm increased the fertilization success, as indicated by the number of cleaved eggs. Presumably, a larger number of sperm increased the probability of forming the male pronucleus. Because the breakdown of the sperm plasma membrane (Morozumi *et al.*, 2006) and nuclear envelope (Yamashita *et al.*, 1990) is a prerequisite for the decondensation of the sperm nucleus and the subsequent formation of the male pronucleus for successful fertilization, part of the injected sperm are likely to transform into male pronuclei and then proceed with the process of fertilization and subsequent embryogenesis.

Small but significant numbers of progeny developed and sometimes hatched from the eggs injected with sperm. Most resultant embryos were abnormal and inviable, but a few progeny exhibited normal to near-normal appearance. The highest eupolyploid was a tetraploid and one mosaic that included pentaploid and hexaploid cell populations, although injecting 1,000 sperm per egg was conducted in this study. Haploidy was the most frequent ploidy in eggs injected with sperm. These results indicate that activated eggs still retained the ability to continue with the fertilization reaction; however, it is unclear why only one sperm or several sperm contributed to embryogenesis,

even after injecting hundreds and thousands of them into the ooplasm.

Microsatellite genotyping revealed that there are two types of haploids. The first was generated by androgenesis resulting from inheriting only paternally derived alleles. Androgenotes were haploid–diploid, haploid–triploid, and haploid–pentaploid mosaics and tetraploids. These results suggest that a single sperm or multiple sperm injected into the ooplasm became male pronuclei and contributed to embryogenesis without incorporating the egg nucleus. The second type of haploid exhibited both maternally and paternally derived alleles. These haploids are considered mosaics comprising the following two types of haploid cell populations: androgenetically developed haploid cells from the male nucleus and gynogenetically developed haploid cells from the female nucleus. This suggests that each haploid cell population that independently formed from the sperm or egg pronucleus developed into an embryo without syngamy.

Injecting multiple numbers of sperm into an egg of the dojo loach gave rise to progeny with various ploidy statuses, a large proportion of which were androgenotes developed from the male pronucleus(-i) transformed from the injected sperm. Another small proportion of progeny developed from the fusion of both female and male pronuclei or probable mosaics comprising maternally derived cells from the female pronucleus(-i) and paternally derived cells from the male pronucleus(-i). No pure gynogenetic progeny were observed. These results suggest that injecting multiple numbers of sperm into the egg ooplasm should have potential as a new biological technique to induce artificial androgenesis.

Acknowledgments

This study was supported in part by KAKENHI Grants-in-Aid for Young Scientist (B) (grant number 18780138) to T.F. and for Challenging Exploratory Research (grant number 25660161) to K.A. from the Japan Society for the Promotion of Science (JSPS).

References

- Detlaf, T. A. (1961). Rate of propagation of the fertilization impulse and dynamics of completion of the second maturation division in sturgeon eggs. *Doklady Akademii Nauk SSSR* **140**, 967–9 (In Russian).
- Fujimoto, T., Kataoka, T., Sakao, S., Saito, T., Yamaha, E. & Arai, K. (2006). Developmental stages and germ cell lineage of the loach (*Misgurnus anguillicaudatus*). *Zool. Sci.* **23**, 977–89.
- Fujimoto, T., Sakao, S., Yamaha, E. & Arai, K. (2007). Evaluation of different doses of UV irradiation to loach eggs for genetic inactivation of the maternal genome. *J. Exp. Zool. Part A*

304 **307**, 449–62.

305 Kurokura, H., Hirano, R., Tomita, M. & Iwahashi, M. (1984). Cryopreservation of carp sperm.
306 *Aquaculture* **37**, 267–73.

307 Mazur, P., Leibo, S. P., Seidel, Jr. G. E. (2008). Cryopreservation of the germplasm of animals used
308 in biological and medical research: importance, impact, status, and future directions. *Biol.*
309 *Reprod.* **78**, 2–12.

310 Morishima, K., Nakayama, I. & Arai, K. (2008). Genetic linkage map of the loach *Misgurnus*
311 *anguillicaudatus* (Teleostei: Cobitidae). *Genetica* **132**, 227–41.

312 Morozumi, K., Shikano, T., Miyazaki, S. & Yanagimachi, R. (2006). Simultaneous removal of sperm
313 plasma membrane and acrosome before intracytoplasmic sperm injection improves oocyte
314 activation/embryonic development. *Proc. Natl. Acad. Sci. USA* **103**, 17661–6.

315 Otani, S., Iwai, T., Nakahata, S., Sakai, C. & Yamashita, M. (2009). Artificial fertilization by
316 intracytoplasmic sperm injection in a teleost fish, the medaka (*Oryzias latipes*). *Biol. Reprod.*
317 **80**, 175–83.

318 Poleo, G. A., Denniston, R. S., Reggio, B. C., Godke, R. A. & Tiersch, T. R. (2001). Fertilization of
319 eggs of zebrafish, *Danio rerio*, by intracytoplasmic sperm injection. *Biol. Reprod.* **65**, 961–6.

320 Poleo, G. A., Godke, R. A. & Tiersch, T. R. (2005). Intracytoplasmic sperm injection using
321 cryopreserved, fixed, and freeze-dried sperm in eggs of Nile tilapia. *Mar. Biotechnol.* **7**, 104–
322 11.

323 Saito, T., Fujimoto, T., Maegawa, S., Inoue, K., Arai, K. & Yamaha, E. (2006). Visualization of
324 primordial germ cells in vivo using GFPnos1 3'UTR mRNA. *Int. J. Dev. Biol.* **50**, 691–9.

325 Tanaka, D., Takahashi, A. & Ueno, K. (2009). Morphometric characteristics and reproductive
326 capacity of nuclear transplants derived from embryonic cells of loach, *Misgurnus*
327 *anguillicaudatus*, *J. Exp. Zool.* **311A**, 11–9.

328 Wakayama, T. & Yanagimachi, R. (1998). Development of normal mice from oocytes injected with
329 freeze-dried spermatozoa. *Nature Biotechnol.* **16**, 639–41.

330 Yamaha, E., Usui, K., Onozato, H. & Hamada K. (1986). A method for dechoriation in goldfish
331 *Carassius auratus*. *Bull. Japan. Soc. Sci. Fish.* **52**, 1929–34.

332 Yamashita, M., Onozato, H., Nakanishi, T. & Nagahama, Y. (1990). Breakdown of the sperm nuclear
333 envelope is a prerequisite for male pronucleus formation: direct evidence from the gynogenetic
334 crucian carp *Carassius auratus langsdorfii*. *Dev. Biol.* **137**, 155–60.

335 Yanagimachi, R. (2005). Intracytoplasmic injection of spermatozoa and spermatogenic cells: Its
336 biology and applications in humans and animals. *Reprod. Biomed. Online* **10**, 247–88.


337 Yasui, G. S., Arias-Rodriguez, L., Fujimoto, T. & Arai, K. (2009). A sperm cryopreservation protocol
338 the loach *Misgurnus anguillicaudatus* and its applicability for other related species. *Anim.*
339 *Reprod. Sci.* **116**, 335–45.







Figures Legends:


Figure 1. Motility parameters of loach sperm after mixing with embryo medium (EM) or injection medium (IM) at increasing concentrations ranging from 0 (distilled water, DW) to 100% (undiluted medium). Solid, open, and gray rectangles indicate immotility, non-progressive motility, and progressive motility, respectively.

Figure 2. Blastula embryos from the control (A) and injected groups (B). Top (A1, B1): embryos under a stereomicroscope using normal light. Middle (A2, B2): embryos under epi-illumination and a dark background. Bottom (A3, B3): boxed area with dashed lines from A2 and B2. Note the abnormal embryos with irregular cell proliferation, reduced or divided blastoderms (solid black arrowheads in B1), acellular development (solid white arrowheads in B3), and asynchronous cell development showing blastomeres of different sizes (open white arrowheads).

Figure 3. External appearance (left, 1), its magnified image in a boxed area (center, 2), and flow cytometric histogram (right, 3) showing the ploidy status of progeny 4 d after fertilization using wild-type female (dominant phenotype) and albino male (recessive phenotype) (A), and progeny developed from injecting multiple numbers of sperm (B–H). Arrows indicate pigmentation. In each flow cytometric histogram, the gray peak denotes somatic cells of goldfish as a standard reference. Progeny shown in B–G correspond to Injected group, nos. 1–6 in Table 3.







1 **Table 1. Egg number; proportion of survived embryos at two-cell, blastula, gastrula, somite, and hatched stages; and proportion of normal**
2 **(wild-type/albino) and abnormal (wild-type/albino) progeny in six types of control groups and sperm-injected (1, 10, 100, 1,000 sperm per egg) groups.**

Treatments	Egg number	No cleavage (%)	Two-cell (%)	Blastula (%)	Gastrula (%)	Somite (%)	Hatch (%)	Normal (%)	Abnormal (%)
								wild-type (%) / albino (%)	wild-type (%) / albino (%)
Intact control, Co	544	11.9 ± 8.7	88.1 ± 8.7	77.3 ± 17.8	77.1 ± 18.9	76.6 ± 18.3	76.5 ± 18.1	77% / 0%	1.5% / 0%
Intact control, Cf	652	21.4 ± 15.1	78.6 ± 15.1	72.1 ± 16.1	69.9 ± 14.6	69.0 ± 14.1	68.6 ± 13.9	71% / 0%	0.7% / 0%
Dechorionated control	506	17.3 ± 4.7	82.9 ± 4.7	75.6 ± 12.6	73.8 ± 11.2	73.2 ± 11.0	72.3 ± 10.9	72% / 0%	2.7% / 0%
Unfertilized control	344	100.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	-	-
Uninjected control	329	52.5 ± 19.1	47.5 ± 19.0	33.4 ± 2.7	19.0 ± 11.2	10.9 ± 4.0	2.3 ± 2.8	1.2% / 0%	0% / 1.8%
Injected control	317	89.2 ± 12.4	10.8 ± 12.4	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	-	-
1 sperm egg ⁻¹	389	92.8 ± 5.9	7.2 ± 5.9	0.3 ± 0.6	0.3 ± 0.6	0.3 ± 0.6	0.0 ± 0.0	-	-
10 sperm egg ⁻¹	414	92.2 ± 6.8	7.8 ± 6.8	3.9 ± 4.6	1.8 ± 3.1	1.0 ± 1.5	0.5 ± 0.6	0.2% / 0%	0% / 0.2%
100 sperm egg ⁻¹	409	75.1 ± 9.9	24.9 ± 9.9	20.1 ± 2.9	11.1 ± 4.1	3.8 ± 3.6	1.4 ± 3.2	0% / 0%	0.2% / 1.4%
1,000 sperm egg ⁻¹	431	12.5 ± 5.1	87.4 ± 5.1	81.5 ± 4.1	50.1 ± 6.6	17.9 ± 6.9	5.9 ± 5.7	1.1% / 0%	0.7% / 4.6%

3 The first control (Intact control: Co) was induced by fertilization between eggs and sperm before the injection trials. The second control (Intact control: Cf)
4 was induced by fertilization between eggs and sperm after the injection trials. In these two controls, the fertilized eggs were cultured with their chorion. The
5 third control (Dechorionated control) was induced by fertilization between eggs and sperm before the injection trials, after which the chorion of the fertilized
6 eggs was removed using the dechoriation medium and then cultured in the culture medium. The fourth control (Unfertilized control) comprised eggs
7 activated by only water. The fifth-control (Uninjected control) comprised eggs fertilized by sperm suspension at 10⁸ sperm mL⁻¹ that was poured onto the
8 animal pole of the dechorionated eggs. The sixth control (Injected control) comprised dechorionated eggs microinjected with a solution containing 200 mM
9 KCl and 0.5% rhodamine.
10

11 **Table 2. Ploidy status of progeny in control groups and sperm-injected (1, 10, 100, 1,000 sperm per egg) groups.**

Treatments	Ploidy status of progeny										
	1n	1n-hyper1n*	1n-2n*	1n-2n- hyper2n*	1n-3n*	1n-5n*	2n	Hyper 2n	3n	5n-6n	Total
Intact control	0	0	0	0	0	0	78(78)	0	2(2)	0	80
Dechorionated control	0	0	0	0	0	0	88(88)	0	1(1)	0	89
Uninjected control	6	2	0	1	2(1)	0	0	0	2(2)	0	13
1 sperm egg ⁻¹	1	0	0	0	0	0	0	0	0	0	1
10 sperm egg ⁻¹	1	0	0	0	0	0	0	0	0	0	1
100 sperm egg ⁻¹	8	0	1	0	0	0	0	1	0	0	10
1,000 sperm egg ⁻¹	44	1	1	0	3(1)	1	2(1)	0	7(6)	1	60

12 Number in parenthesis denotes the number of normal and pigmented wild-type progeny.

13 *Dash indicates mosaicism of cell populations with different ploidy status.

14

15

16

Table 3. Ploidy status, color phenotype, external appearance, and microsatellite DNA genotypes of progeny from control (normal fertilization) and microinjection of approximately 1,000 sperm from albino male into the ooplasm of an egg from wild-type female.

Group/ Larvae no.	Ploidy	Color phenotype*	External appearance	<i>Mac 24</i> **	<i>Mac 345</i> **	<i>Mac 449</i> **
Control 1	2n	wild-type	normal	a / c	a / c	a / c
2	2n	wild-type	normal	a / c	a / c	a / c
3	2n	wild-type	normal	a / c	b / c	b / c
4	2n	wild-type	normal	a / c	a / c	a / c
5	2n	wild-type	normal	a / c	a / c	a / c
6	2n	wild-type	normal	a / c	a / c	b / c
7	2n	wild-type	normal	a / c	b / c	b / c
8	2n	wild-type	normal	a / c	a / c	a / c
9	2n	wild-type	normal	a / c	a / c	a / c
10	2n	wild-type	normal	a / c	a / c	a / c
Injected 1 ^a	3n	wild-type	normal	a / c	b / c	a / c
2 ^a	1n-1.9n-2.1n	albino/+	normal	a / c	a / c	b / c
3 ^a	2.8n	albino/+	abnormal	a / c	b / c	b / c
4 ^a	2n	wild-type	abnormal	a / c	a / c	b / c
5 ^a	4n	albino	normal	c	c	c
6 ^a	1n	albino	abnormal	a / c	b / c	a / c
7 ^b	1n	n.d.	abnormal	c	c	c
8 ^b	1n	n.d.	abnormal	c	c	c
9 ^b	1n	n.d.	abnormal	a / c	a / c	a / c
10 ^b	1n-3n	n.d.	abnormal	c	c	c
11 ^b	1n-2n	n.d.	abnormal	c	c	c
12 ^b	1n	n.d.	abnormal	c	c	c
13 ^b	1n-2n	n.d.	abnormal	c	c	c
14 ^b	1n-2n	n.d.	abnormal	c	c	c
15 ^b	1n	n.d.	abnormal	a / c	a / c	a / c
16 ^b	1n-2n	n.d.	abnormal	a / c	a / c	b / c
17 ^b	1n	n.d.	abnormal	c	c	c
18 ^b	1n-5n	n.d.	abnormal	c	c	c
19 ^b	1n	n.d.	abnormal	c	c	c
20 ^b	1n	n.d.	abnormal	c	c	c

* wild-type: normal pigmentation, albino: no pigmentation, +: partially pigmented, n.d.: not determined

** Female genotypes (a/b or a/a): 120/120 at *Mac 24* (linkage group 16), 514/546 at *Mac 345* (linkage group 6), 228/240 at *Mac 449* (linkage group 14). Male genotypes (c/c): 112/112 at *Mac 24*, 506/506 at *Mac 345*, 236/236 at *Mac449*.

^aProgeny 1–6 correspond to individuals shown in Figs. 3B–G.

^bProgeny 7–20 have the same abnormal morphology patterns shown in Fig. 3H.