



HOKKAIDO UNIVERSITY

Title	Detection of novel gammaherpesviruses from fruit bats in Indonesia
Author(s)	Wada, Yuji; Sasaki, Michihito; Setiyono, Agus et al.
Citation	Journal of medical microbiology, 67(3), 415-422 https://doi.org/10.1099/jmm.0.000689
Issue Date	2018-03
Doc URL	https://hdl.handle.net/2115/74394
Type	journal article
File Information	manuscript (revise).pdf



**Detection of novel gammaherpesviruses from fruit bats in
Indonesia**

Yuji Wada¹, Michihito Sasaki¹, Agus Setiyono², Ekowati Handharyani², Ibenu
Rahmadani³, Siswatiana Taha⁴, Sri Adiani⁵, Munira Latief⁶, Zainal Abidin
Kholilullah⁷, Mawar Subangkit², Shintaro Kobayashi^{1,#a}, Ichiro Nakamura⁸,
Takashi Kimura^{1,#b}, Yasuko Orba¹ and Hirofumi Sawa^{1,9,10}

¹ Division of Molecular Pathobiology, Research Center for Zoonosis Control,
Hokkaido University, Sapporo, Hokkaido, Japan

² Laboratory of Veterinary Pathology, Faculty of Veterinary Medicine, Bogor
Agricultural University, Bogor, Indonesia

³ Veterinary Investigation and Diagnostic Center, Bukittinggi, Indonesia

⁴ Faculty of Agriculture, Gorontalo State University, Gorontalo, Indonesia

⁵ Faculty of Animal Husbandry, Sam Ratulangi University, Manado, Indonesia

⁶ Office of Animal Husbandry and Fisheries, Soppeng, Indonesia

⁷ Veterinary Medicine Study Program, Faculty of Medicine, Hasanudin University,
Makassar, Indonesia

⁸ Unit of International Cooperation, Research Center for Zoonosis Control,
Hokkaido University, Sapporo, Hokkaido Japan

⁹ Global Institution for Collaborative Research and Education (GI-CoRE),
Hokkaido University, Sapporo, Hokkaido, Japan

¹⁰ Global Virus Network, Baltimore, MD 21201, USA

24

25 Current addresses:

26 #^a Laboratory of Public Health, Department of Preventive Veterinary Medicine,
27 Graduate School of Veterinary Medicine, Hokkaido University, Sapporo, Hokkaido,
28 Japan.

29 #^b Laboratory of Comparative Pathology, Graduate School of Veterinary Medicine,
30 Hokkaido University, Sapporo, Hokkaido, Japan

31

32 Corresponding author:

33 Hirofumi Sawa (h-sawa@czc.hokudai.ac.jp)

34

35 Key words:

36 Gammaherpesvirus, Fruit bat, PCR screening, Long-distance PCR, Phylogenetic
37 analysis

38

39 Word count for abstract: 124

40 Word count for main text: 2150

41

42 Database depository:

43 The DDBJ accession numbers for the MgBHV and MgGHV *DPOL*, *gB* and
44 *DPOL-gB* sequences are LC268906-LC268990

Abstract

Bats are an important natural reservoir of zoonotic viral pathogens. We previously isolated an alphaherpesvirus in fruit bats in Indonesia, and here establish the presence of viruses belonging to other taxa of the family *Herpesviridae*. We screened the same fruit bat population with pan-herpesvirus PCR and discovered 68 sequences of novel gammaherpesvirus, designated 'megabat gammaherpesvirus' (MgGHV). A phylogenetic analysis of approximately 3.4 kbp of continuous MgGHV sequences encompassing the glycoprotein B gene and DNA polymerase gene revealed that the MgGHV sequences are distinct from those of other reported gammaherpesviruses. Further analysis suggested the existence of coinfections of herpesviruses in Indonesian fruit bats. Our findings extend our understanding of the infectious cycles of herpesviruses in bats in Indonesia and the phylogenetic diversity of the gammaherpesviruses.

Text

Outbreaks of emerging and re-emerging zoonoses have compromised public health and damaged the global economy in recent years. Bats are known to be carriers of lyssaviruses, henipaviruses, severe acute respiratory syndrome-like coronaviruses, and filoviruses, so they are regarded as an important natural reservoirs of zoonotic viral pathogens [1-5].

Herpesviruses are enveloped, double-stranded DNA viruses belonging to the family *Herpesviridae*. These viruses include three genera: *Alphaherpesvirinae*, *Betaherpesvirinae*, and *Gammapherpesvirinae*. *Gammapherpesvirinae* contains viruses that are oncogenic to humans, including Human gammaherpesvirus 4 (also known as Epstein-Barr virus) and Human gammaherpesvirus 8 (also known as Kaposi's sarcoma-associated herpesvirus). Therefore, the viruses of the genus *Gammapherpesvirinae* are important in human health.

Many gammaherpesvirus sequences have recently been detected in a broad range of bat species worldwide [6-9]. *Myotis* gammaherpesvirus 8 was isolated from a tumor cell line derived from *Myotis velifer incautus* (mouse-eared bat), and this virus has been shown to cause cytopathic effects in Vero cells and can replicate in some human cell lines [10, 11].

In a previous study, we reported a high prevalence of fruit bat alphaherpesvirus 1 (FBAHV1) in fruit bats in Indonesia [12]. Here, to extend our understanding of the herpesviruses carried by fruit bats, we used PCR to screen for a broad spectrum of herpesviruses, and followed by phylogenetically analyzed the sequences detected.

In this study, 183 bat tissue samples were analyzed: 69 from *Pteropus vampyrus* (large flying fox), 61 from unassigned *Pteropus* bats (*Pteropus* sp.), 17 from *Dobsonia moluccensis* (Moluccan bare-backed fruit bat), and 36 from *Acerodon celebensis* (Sulawesi fruit bat). The samples were collected in the Lima Puluh Kota, Magelang, Panjalu, Paguyaman, Popayato, Sidrap, Soppeng, and Surabaya regions of Indonesia in 2010-2014, as shown in Fig. 1 and Table 1. Several healthy bats were captured from flocks at each location and euthanized with an overdose of ketamine and xylazine, with the permission of the Directorate General of Livestock and Animal Health Services, Ministry of Agriculture, Republic of Indonesia. All procedures were approved by the Animal Care and Use Committee of the Veterinary Teaching Hospital, Bogor Agricultural University (permit number 05-2010 RSHP-IPB). The samples were stored at -80°C and have been used previously to screen for FBAHV1 and other viruses [12-15]. The bat species were identified according to their morphological characteristics, and a nucleotide sequence analysis of their mitochondrial 16S rRNA and cytochrome *b* genes, as previously described [12, 15]. The mitochondrial nucleotide sequences of *Pteropus* sp. showed greatest identity (96% 16S rRNA and 95% cytochrome *b* genes) to the corresponding sequences of *P. hypomelanus* (small flying fox; accession numbers AF069537 and AB062472). Therefore, we concluded that these bats were closely related to *P. hypomelanus*, but should be classified as a different species.

Total DNA was extracted from spleen tissue samples with DNAzol (Molecular Research Center) or the QIAamp DNA Mini Kit (Qiagen). We screened for

herpesviruses with semi-nested PCR using TaKaRa Ex Taq Hot Start Version (Takara Bio) and degenerate primers targeting the DNA polymerase (*DPOL*) gene of the herpesviruses [16]. This primer set has been used to detect herpesviruses in a broad range of wild animals [17-19]. The PCR products, of approximately 250 bp, were purified from the gel after electrophoresis and sequenced directly with the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems). The sequences were analyzed with BLASTN, and 71 of the 183 tested samples were positive for herpesvirus sequences. The BLASTN results suggested that the 67 of these 71 sequences (172 bp in length) originated from gammaherpesviruses and that the remaining four sequences (181 bp in length) originated from betaherpesviruses, as summarized in Table S1 (available in the online Supplementary Material). Therefore, the origins of these herpesvirus sequences were designated 'megabat gammaherpesvirus' (MgGHV) and 'megabat betaherpesvirus' (MgBHV), respectively. The *DPOL* sequences determined were deposited in GenBank under accession numbers LC268906-LC268972. The sequences of MgGHV were detected in the bat samples collected at every location that we sampled, suggesting that MgGHVs are geographically distributed throughout Indonesia, as summarized in Figure 1 and Table 1.

Because the majority of the sequences belonged to MgGHV, distributed geographically throughout Indonesia, we focused our investigation on MgGHV. A multiple-sequence alignment by CLUSTAL W and a subsequent Maximum likelihood phylogenetic analysis of the partial MgGHV *DPOL* sequences were performed by the MEGA 7 program [20]. The MgGHVs were phylogenetically

divided into seven groups (MgGHV Group-1 to -7), as shown in Fig. 2(a). The MgGHV sequences clustered according to the bat species in which they were detected, except MgGHV Group-1, which contained sequences detected in *Pteropus* sp. and *Acerodon celebensis*. All of the MgGHV groups were phylogenetically related to previously described gammaherpesviruses detected in *Pteropus* species. MgGHV Group-1, -2 and -3 formed a cluster with *Pteropus giganteus* gammaherpesvirus isolate-3 (PgHV-3); MgGHV Group-4 and -5 formed a cluster with PgHV-2; and MgGHV Group-6 was closely related to PgHV-1. These PgHVs were detected in *P. giganteus* in Bangladesh [21]. MgGHV Group-7 contained a single sequence that was identical to that of Pteropine herpesvirus 2, which was detected in *Pteropus vampyrus* kept in a captive breeding facility in the United States [22].

To investigate the MgGHV phylogeny in more detail, we additionally screened for glycoprotein B (*gB*) gene, which is located upstream from the *DPOL* gene, and subsequent long-distance PCR (LD-PCR). Both of *DPOL* and *gB* are commonly used for the phylogenetic analysis of the gammaherpesviruses [23, 24]. The target sequence of the LD-PCR was approximately 3.4 kbp in length and flanked by the partial *gB* and *DPOL* sequences obtained from the screening [17, 25].

First, we used nested PCR to screen for the gammaherpesvirus *gB* gene using Platinum Taq DNA Polymerase (Invitrogen) and degenerate primers, as previously described [26]. With this procedure, the 417-456 bp partial *gB* sequences were obtained from 16 of the 67 spleen DNA samples for gammaherpesvirus *DPOL*, and a BLASTN analysis showed that all of them were

derived from gammaherpesviruses, as summarized in Table 1. The *gB* sequences determined were deposited in the GenBank under accession numbers LC268973-LC268980 and LC268982-LC268986. A phylogenetic analysis of these *gB* sequences revealed that the MgGHVs formed a discrete cluster, separate from known gammaherpesviruses clades, as shown in Fig. 2(b). Based on the *gB* phylogeny, the MgGHVs were tentatively assigned to six groups, MgGHV Group-A to -F. MgGHV Group-B, -C, -E and -F were congruent with the bat species in which they were detected, whereas MgGHV Group-A and -D contained sequences detected in both *Pteropus vampyrus* and *Pteropus* sp., as summarized in Table S1. Remarkably, the phylogenetic relationships of each MgGHV group were not consistent between the phylogenetic trees constructed with the *DPOL* and *gB* sequences, as shown in Fig. 2 (a, b). For instance, *DPOL* from several bat samples was assigned to MgGHV Group-1, whereas the *gB* sequences detected in the same samples were assigned to several different groups (MgGHV Group-A, B, C and F), as summarized in Table S2 (available in the online Supplementary Material). These observations suggest the presence of coinfections with different MgGHVs in individual fruit bats. Mühldorfer *et al.* have previously reported coinfections of BatGHV-3, -4, and -5 in *Nyctalus noctula* (noctule), but this is the first report of coinfections of gammaherpesviruses in *Pteropus vampyrus*, *Pteropus* sp. and *Dobsonia moluccensis* [7]. A genomic recombination event has previously been reported in the human gammaherpesviruses [27]. Therefore, it is possible that different gammaherpesviruses might interact each other and play an important role in their survival or evolution in bats. Therefore, coinfection warrants

attention in studies of the prevalence or evolutionary history of gammaherpesviruses in bats in future epidemiological researches.

LD-PCR was then performed on the samples doubly positive for samples *DPOL* and *gB*. Of the 16 samples tested, four were positive for the 3,351-3,544 bp LD-PCR target sequence between *DPOL* and *gB*. The origins of these *DPOL-gB* sequences were designated MgGHV_IFB11-41 (LC268987), MgGHV_IFB12-05 (LC268988), MgGHV_IFB12-16 (LC268989), and MgGHV_IFB13-11 (LC268990). A phylogenetic analysis based on the *DPOL-gB* sequences grouped the viruses into two clusters, as shown in Fig. 2 (c). MgGHV_IFB11-41 and MgGHV_IFB13-11 were detected in *Pteropus* sp., and were phylogenetically located relatively close to Murid gammaherpesvirus 4. In an analysis of nucleotide identity with the GENETYX software v.10 (Genetyx Corporation), MgGHV_IFB11-41 and MgGHV_IFB13-11 showed highest identity (61%-62%) to *Diceros bicornis* gammaherpesvirus 1. MgGHV_IFB12-05 and MgGHV_IFB12-16 were detected in *Dobsonia moluccensis*, and formed a cluster with *Hexaprotodon liberiensis* gammaherpesvirus 1 and Human gammaherpesvirus 4. MgGHV_IFB12-05 and MgGHV_IFB12-16 shared highest identity (66%) with *Hexaprotodon liberiensis* gammaherpesvirus 1. All four MgGHVs identified here were segregated from currently known gammaherpesviruses. To date, the nucleotide sequences corresponding to the *DPOL-gB* region are available for six bat gammaherpesviruses: *Eptesicus serotinus* rhadinovirus 1, *Nyctalus noctula* rhadinovirus 2, *Myotis* gammaherpesvirus 8, *Myotis ricketti* herpesvirus 1, *Myotis ricketti* herpesvirus 2, and *Pipistrellus nathusii* rhadinovirus 1 [10, 28, 29].

However, the four MgGHVs identified here are clearly distinct from the other reported gammaherpesviruses detected in bats.

In the study, we noted the existence of two different *gB* sequences in the DNA sample positive for MgGHV_IFB11-41. The first *gB* sequence was designated MgGHV_IFB11-41_a (LC268980), and was detected with nested PCR screening for *gB*. However, the amplicon from the subsequent LD-PCR, using a gene-specific primer targeting MgGHV_IFB11-41_a, generated a different *gB* sequence from MgGHV_IFB11-41_a. The existence of these two sequences was reproducibly confirmed with PCR. In addition to the comparative phylogenetic analysis of the partial *DPOL* and *gB* sequences, this finding also suggests the presence of a coinfection with two different MgGHVs in a single fruit bat.

To investigate possible coinfections with MgGHV and MgBHV, we performed another nested PCR to screen for *gB* in the same samples in which the MgBHV sequences was detected. A gammaherpesviral *gB* sequence was detected in one (MgGHV_IFB11-45; LC268981) of the four samples. We also detected a coinfection with MgGHV and FBAHV when we amplified a partial *DPOL* sequence (MgGHV_IFB11-32: LC268932) from a fruit bat sample from which FBAHV was isolated in our previous study [12]. These observations suggest that MgGHVs can establish coinfections with other viruses of the family *Herpesviridae*.

In the phylogenetic analysis, partial incongruence was observed between the *DPOL* and *gB* trees. We also failed to recover the long continuous *DPOL-gB* region of MgGHV with LD-PCR in 12 of the 16 bats that were positive for both the gammaherpesvirus *DPOL* and *gB* genes. This suggests that the *DPOL* and *gB*

sequences obtained from the individual bats were derived from different MgGHV origins. Coinfections with several beta- or gammaherpesviruses have also been reported in other animals [7, 30]. Considering the broad reactivity of the degenerate primers used in the present study, the coexistence of different herpesviruses might be an obstacle to the effective use of the LD-PCR approach [16].

In this study, we detected 67 partial *DPOL* sequences from MgGHVs, and all of them were phylogenetically related to previously reported gammaherpesviruses detected in *Pteropus* bats. The sequences of MgGHV Group-3 shared 96%-100% identity with those of PgHV-3, and those of MgGHV Group-4 shared 99% identity with those of PgHV-2, even though the sequences were detected in different bat species. This suggests that these groups of gammaherpesviruses are transmitted independently of the host species. This hypothesis is also supported by another finding in the phylogenetic analysis of MgGHV Group-1. In this group, identical partial MgGHV *DPOL* sequences were detected in different bat species, *Pteropus* sp. and *Acerodon celebensis* as summarized in Table S1. It has also been reported that spillover events of alpha- and gammaherpesviruses beyond the species in the past [26, 31, 32]. It is possible that these groups of gammaherpesviruses are frequently transmitted between species and are maintained in a broad range of bat species inhabiting the Asian region.

The MgGHV Group-7 nucleotide sequence was identical to that of Pteropine herpesvirus 2, and both sequences were detected in *Pteropus vampyrus*, as shown in Fig. 2(a) and Table S1. Detailed information about the bat in which the

243 Pteropine herpesvirus 2 sequence was detected is not available, so it is unclear
244 whether this bat was imported from an Asian country or was born at the captive
245 breeding facility in the United States [22]. Escalera-Zamudio, *et al.* suggested that
246 the existence of endogenous gammaherpesviruses in the bat genome was
247 unlikely [31]. Therefore, we infer that *Pteropus vampyrus* is susceptible to
248 infection with this group of gammaherpesviruses.

249 In summary, using PCR screening, we detected genomic fragments of four
250 novel betaherpesviruses and 67 novel gammaherpesviruses, designated
251 MgBHVs and MgGHVs, respectively, in fruit bats inhabiting Indonesia. Our results
252 demonstrate the geographic distribution and genetic diversity of the MgGHVs in
253 these bat populations. The LD-PCR approach allowed us to determine a relatively
254 long region of the MgGHV genomic sequence and to infer a deep phylogeny for
255 the identified MgGHVs. Our results also suggest that coinfections with MgGHVs
256 and FBAHV and with MgBHV and different MgGHVs occur, and confirm the
257 interspecies transmission of MgGHV Group-1, -3 and -4. This study contribute to
258 our current knowledge of the herpesviruses in the bats of Indonesia, and the
259 phylogenetic diversity of the gammaherpesviruses.

260 **Funding information**

261 This research was supported in part by a Grant-in-Aid for Scientific Research
262 on Innovative Areas from the Ministry of Education, Culture, Science, Sports and
263 Technology (MEXT) of Japan (16H06429, 16H06431 and 16K21723); and grants
264 from MEXT/JSPS KAKENHI (16H05805).

265 **Acknowledgements**

266 The authors would like to thank Ms. Satoko Sasaki for her excellent technical
267 support.

268 **Conflicts of interest**

269 The authors declare that there are no conflicts of interest.

Ethical statement

All experiments involving animals in Indonesia were performed in accordance with the ethical guidelines of the Animal Care and Use Committee of the Animal Teaching Hospital, Bogor Agricultural University, which are based on the Guide for the Care and Use of Laboratory Animals (7th and 8th editions) by the National Research Council of the National Academies, and the Guideline on the Care and Use of Animals for Scientific Purposes by National Advisory Committee for Laboratory Animal Research. The protocol was approved by the Animal Care and Use Committee of the Veterinary Teaching Hospital, Bogor Agricultural University (permit number 05-2010 RSHP-IPB).

All of the bat samples were collected with the permission of the Directorate General of Livestock and Animal Health Services, Ministry of Agriculture, Republic of Indonesia. The samples were stored at -80°C and have been used previously to screen for other viruses.

References

1. **Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T.** Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev* 2006;19(3):531-545.
2. **Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A et al.** Fruit bats as reservoirs of Ebola virus. *Nature* 2005;438(7068):575-576.
3. **Li W, Shi Z, Yu M, Ren W, Smith C et al.** Bats are natural reservoirs of SARS-like coronaviruses. *Science* 2005;310(5748):676-679.
4. **Field H, Young P, Yob JM, Mills J, Hall L et al.** The natural history of Hendra and Nipah viruses. *Microbes Infect* 2001;3(4):307-314.
5. **Johnson N, Vos A, Freuling C, Tordo N, Fooks AR et al.** Human rabies due to lyssavirus infection of bat origin. *Vet Microbiol* 2010;142(3-4):151-159.
6. **Zheng XY, Qiu M, Chen SW, Xiao JP, Ma LZ et al.** High prevalence and diversity of viruses of the subfamily Gammaherpesvirinae, family Herpesviridae, in fecal specimens from bats of different species in southern China. *Arch Virol* 2016;161(1):135-140.
7. **Mühldorfer K, Speck S, Kurth A, Lesnik R, Freuling C et al.** Diseases and causes of death in European bats: dynamics in disease susceptibility and infection rates. *PLoS One* 2011;6(12):e29773.
8. **Sano K, Okazaki S, Taniguchi S, Masangkay JS, Puentespina R et al.** Detection of a novel herpesvirus from bats in the Philippines. *Virus Genes* 2015;51(1):136-139.
9. **Jánoska M, Vidovszky M, Molnár V, Liptovszky M, Harrach B et al.**

308 Novel adenoviruses and herpesviruses detected in bats. *Vet J*
309 2011;189(1):118-121.

310 10. **Shabman RS, Shrivastava S, Tsibane T, Attie O, Jayaprakash A et al.**
311 Isolation and Characterization of a Novel Gammaherpesvirus from a Microbat Cell
312 Line. *mSphere* 2016;1(1).

313 11. **Host KM, Damania B.** Discovery of a Novel Bat Gammaherpesvirus.
314 *mSphere* 2016;1(1).

315 12. **Sasaki M, Setiyono A, Handharyani E, Kobayashi S, Rahmadani I et**
316 **al.** Isolation and characterization of a novel alphaherpesvirus in fruit bats. *J Virol*
317 2014;88(17):9819-9829.

318 13. **Anindita PD, Sasaki M, Setiyono A, Handharyani E, Orba Y et al.**
319 Detection of coronavirus genomes in Moluccan naked-backed fruit bats in
320 Indonesia. *Arch Virol* 2015;160(4):1113-1118.

321 14. **Kobayashi S, Sasaki M, Nakao R, Setiyono A, Handharyani E et al.**
322 Detection of novel polyomaviruses in fruit bats in Indonesia. *Arch Virol*
323 2015;160(4):1075-1082.

324 15. **Sasaki M, Setiyono A, Handharyani E, Rahmadani I, Taha S et al.**
325 Molecular detection of a novel paramyxovirus in fruit bats from Indonesia. *Viol J*
326 2012;9:240.

327 16. **Rose TM, Strand KB, Schultz ER, Schaefer G, Rankin GW et al.**
328 Identification of two homologs of the Kaposi's sarcoma-associated herpesvirus
329 (human herpesvirus 8) in retroperitoneal fibromatosis of different macaque
330 species. *J Virol* 1997;71(5):4138-4144.

- 331 17. **Troyer RM, Beatty JA, Stutzman-Rodriguez KR, Carver S, Lozano CC**
332 **et al.** Novel gammaherpesviruses in North American domestic cats, bobcats, and
333 pumas: identification, prevalence, and risk factors. *J Virol* 2014;88(8):3914-3924.
- 334 18. **Lacoste V, Mauciere P, Dubreuil G, Lewis J, Georges-Courbot MC et**
335 **al.** Simian homologues of human gamma-2 and betaherpesviruses in mandrill and
336 drill monkeys. *J Virol* 2000;74(24):11993-11999.
- 337 19. **Delang L, Segura Guerrero N, Tas A, Quérat G, Pastorino B et al.**
338 Mutations in the chikungunya virus non-structural proteins cause resistance to
339 favipiravir (T-705), a broad-spectrum antiviral. *J Antimicrob Chemother*
340 2014;69(10):2770-2784.
- 341 20. **Kumar S, Stecher G, Tamura K.** MEGA7: Molecular Evolutionary
342 Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol*
343 2016;33(7):1870-1874.
- 344 21. **Anthony SJ, Epstein JH, Murray KA, Navarrete-Macias I,**
345 **Zambrana-Torrel CM et al.** A strategy to estimate unknown viral diversity in
346 mammals. *MBio* 2013;4(5):e00598-00513.
- 347 22. **Paige Brock A, Cortés-Hinojosa G, Plummer CE, Conway JA, Roff SR**
348 **et al.** A novel gammaherpesvirus in a large flying fox (*Pteropus vampyrus*) with
349 blepharitis. *J Vet Diagn Invest* 2013;25(3):433-437.
- 350 23. **VanDevanter DR, Warrener P, Bennett L, Schultz ER, Coulter S et al.**
351 Detection and analysis of diverse herpesviral species by consensus primer PCR.
352 *J Clin Microbiol* 1996;34(7):1666-1671.
- 353 24. **Auerbach MR, Czajak SC, Johnson WE, Desrosiers RC, Alexander L.**

354 Species specificity of macaque rhadinovirus glycoprotein B sequences. *J Virol*
355 2000;74(1):584-590.

356 25. **Ehlers B, Küchler J, Yasmum N, Dural G, Voigt S et al.** Identification of
357 novel rodent herpesviruses, including the first gammaherpesvirus of *Mus*
358 *musculus*. *J Virol* 2007;81(15):8091-8100.

359 26. **Ehlers B, Dural G, Yasmum N, Lembo T, de Thoisy B et al.** Novel
360 mammalian herpesviruses and lineages within the Gammaherpesvirinae:
361 cospeciation and interspecies transfer. *J Virol* 2008;82(7):3509-3516.

362 27. **Poole LJ, Zong JC, Ciufo DM, Alcendor DJ, Cannon JS et al.**
363 Comparison of genetic variability at multiple loci across the genomes of the major
364 subtypes of Kaposi's sarcoma-associated herpesvirus reveals evidence for
365 recombination and for two distinct types of open reading frame K15 alleles at the
366 right-hand end. *J Virol* 1999;73(8):6646-6660.

367 28. **Wibbelt G, Kurth A, Yasmum N, Bannert M, Nagel S et al.** Discovery of
368 herpesviruses in bats. *J Gen Virol* 2007;88(Pt 10):2651-2655.

369 29. **Wu Z, Ren X, Yang L, Hu Y, Yang J et al.** Virome analysis for
370 identification of novel mammalian viruses in bat species from Chinese provinces.
371 *J Virol* 2012;86(20):10999-11012.

372 30. **Prepens S, Kreuzer KA, Leendertz F, Nitsche A, Ehlers B.** Discovery of
373 herpesviruses in multi-infected primates using locked nucleic acids (LNA) and a
374 bigenic PCR approach. *Virology* 2007;4:84.

375 31. **Escalera-Zamudio M, Rojas-Anaya E, Kolokotronis SO, Taboada B,**
376 **Loza-Rubio E et al.** Bats, Primates, and the Evolutionary Origins and

377 Diversification of Mammalian Gammaherpesviruses. *MBio* 2016;7(6).
378 32. **Wertheim JO, Smith MD, Smith DM, Scheffler K, Kosakovsky Pond**
379 **SL.** Evolutionary origins of human herpes simplex viruses 1 and 2. *Mol Biol Evol*
380 2014;31(9):2356-2364.
381

Table 1. Results of PCR screening for gammaherpesviruses

	Sampling location	DPOL gene	gB gene
<i>Pteropus vampyrus</i>	Lima Puluh Kota	11/20	3/11
	Magelang	2/20	1/2
	Panjalu	10/26	1/10
	Surabaya	2/3	0/2
Subtotal		25/69	5/25
<i>Pteropus</i> sp.	Paguyaman	20/35	9/20
	Popayato	3/4	0/3
	Sidrap	6/15	0/6
	Soppeng	5/7	1/5
Subtotal		34/61	10/34
<i>Dobsonia moluccensis</i>	Paguyaman	6/17	2/6
<i>Acerodon celebensis</i>	Paguyaman	2/36	0/2
Total		67/183	17/67

382

Fig. 1. Geographic information on the numbers and species of captured bats.

In total, 183 bats were captured in the Lima Puluh Kota, Magelang, Panjalu, Paguyaman, Popayato, Sidrap, Soppeng and Surabaya regions of Indonesia between 2010 and 2014. The bats were classified as four species: *Pteropus vampyrus*, *Pteropus* sp., *Dobsonia moluccensis*, and *Acerodon celebensis*.

Fig. 2. Phylogenetic analysis of partial *DPOL*, *gB* and *gB-DPOL* nucleotide sequences of MgGHV.

Phylogenetic trees of partial gammaherpesvirus (a) *DPOL*, (b) *gB* and (c) *gB-DPOL* nucleotide sequences were constructed by Maximum likelihood phylogenetic analyses in the MEGA7 program. The 67 partial MgGHV *DPOL* sequences (172 bp in length) were used for the analysis and clustered into seven groups (MgGHV Group-1 to -7) when the entire tree was visualized. When the 18 partial MgGHV *gB* sequences (417-456 bp in length) were analyzed, they clustered into six groups (MgGHV Group-A to -F). Four MgGHV *gB-DPOL* sequences (3,351-3,544 bp in length) were analyzed. All the MgGHV sequences are highlighted in gray. The bootstrap values obtained after 1,000 replicates are indicated at the major tree roots. The scale bar represents a distance of 0.1 bp of substitutions per site. The sequences detected in bats are indicated with asterisks.