



HOKKAIDO UNIVERSITY

Title	Human serum N-glycans as highly sensitive cancer biomarkers : Potential benefits and the risks [an abstract of entire text]
Author(s)	Gebrehiwot, Abrha Gebreselema
Description	この博士論文全文の閲覧方法については、以下のサイトをご参照ください。【担当:理学部図書室】 https://www.lib.hokudai.ac.jp/dissertations/copy-guides/
Degree Grantor	北海道大学
Degree Name	博士(生命科学)
Dissertation Number	甲第13604号
Issue Date	2019-03-25
Doc URL	https://hdl.handle.net/2115/74475
Type	doctoral thesis
File Information	Abrha_Gebreselema_Gebrehiwot_summary.pdf



Summary of Doctoral Dissertation

Degree requested Doctor of Life Science Applicant's name Abrha Gebreselema Gebrehawot

Title of Doctoral Dissertation

Human serum *N*-glycans as highly sensitive cancer biomarkers: Potential benefits
and the risks
(高感度がんマーカーとしてのヒト血清*N*-グリカン：その潜在的な恩恵と危険性)

Apart from the cancer genome, deciphering posttranslational protein glycosylation alterations has been of critical importance as a promising target for discovering novel diagnostic and therapeutic agents. In the present study, the author performed large-scale glycomics research of human whole serum and purified IgG in Ethiopian breast cancer patients and matched controls using a glycoblotting-assisted MALDI-TOF/MS-based quantitative analysis, aiming at identification of new biomarkers for early detection of breast cancer. With an overall quantitative up-regulation in all the breast cancer stages compared to NC, highest abundance of core-fucosylated, highly branched, and sialylated *N*-glycans with strong distinguishing power (AUC = 0.8-1) was clearly found in the earlier stage (stage I and II) patients. IgG-focused *N*-glycomic profiling further revealed increased fucosylation and agalactosylation of IgG glycoforms that could specifically discriminate stage-II patients from controls. Increased fucosylation and agalactosylation of IgG reduce its affinity with the Fc receptors and thus its effector function, a suggested mechanism allowing the tumor cells to escape from the immune system. By addressing serum and IgG *N*-glycosylation signatures of native black Africans for the first time, this study revealed novel candidate glyco-biomarkers markedly associated with early stage BC.

Using MALDI-TOF/MS- and HPLC-based quantitative approaches, the author extended his research by performing inter-ethnic serum *N*-glycome and sialic acid variations among US origin control, Japanese, Indian, and Ethiopian healthy volunteers in association with the identified glycan biomarkers, aiming to investigate the influence of confounding factors like ethnic variation (which is often overlooked in biomarker research). Informative findings of ethnic-associated variations in the serum *N*-glycans and free sialic acid were found with consistent highest abundance primarily demonstrated in the Ethiopian group compared to the other ethnicities.

Surprisingly, some of the *N*-glycans greatly elevated in the Ethiopian subjects have been identified as sensitive serum biomarkers of various cancers, implicating the critical impact of ethnic differences in human serum *N*-glycome variation, the ignorance of which may provide unclear and imprecise conclusion during diagnosis by using glycan-related disease biomarkers.