



HOKKAIDO UNIVERSITY

Title	Identification of group A rotaviruses from Zambian fruit bats provides evidence for long-distance dispersal events in Africa
Author(s)	Sasaki, Michihito; Kajihara, Masahiro; Changula, Katendi et al.
Citation	Infection, genetics and evolution, 63, 104-109 https://doi.org/10.1016/j.meegid.2018.05.016
Issue Date	2018-09
Doc URL	https://hdl.handle.net/2115/75328
Rights	© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	R1_RVA Manuscript final.pdf



Identification of group A rotaviruses from Zambian fruit bats provides evidence for long-distance dispersal events in Africa

Michihito Sasaki^{a*}, Masahiro Kajihara^b, Katendi Changula^c, Akina Mori-Kajihara^b, Hirohito Ogawa^{d,1}, Bernard M. Hang'ombe^c, Aaron S. Mweene^{e,f,g}, Martin Simuunza^e, Reiko Yoshida^b, Michael Carr^{h,i}, Yasuko Orba^a, Ayato Takada^{b,h} and Hirofumi Sawa^{a,g,h*}

^aDivision of Molecular Pathobiology, Research Center for Zoonosis Control, Hokkaido University, Sapporo 001-0020, Japan

^bDivision of Global Epidemiology, Research Center for Zoonosis Control, Hokkaido University, Sapporo 001-0020, Japan

^cDepartment of Paraclinical Studies, School of Veterinary and Medicine, University of Zambia, PO Box 32379, Lusaka, Zambia

^dHokudai Center for Zoonosis Control in Zambia, School of Veterinary Medicine, University of Zambia, PO Box 32379, Lusaka, Zambia

^eDepartment of Disease Control, School of Veterinary and Medicine, University of Zambia, PO Box 32379, Lusaka, Zambia

^fAfrica Center of Excellence for Infectious Diseases of Humans and Animals, University of Zambia, PO Box 32379, Lusaka, Zambia

^gGlobal Virus Network, Baltimore, Maryland 21201, USA

^hGlobal Institution for Collaborative Research and Education (GI-CoRE), Hokkaido University, Sapporo 001-0020, Japan

ⁱNational Virus Reference Laboratory, School of Medicine, University College Dublin, Dublin 4, Ireland

25 **Footnote**

26 Present address: ¹Department of Virology, Okayama University Graduate School of
27 Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan

28

29 *Corresponding authors:

30 Michihito Sasaki, Tel: +81-11-706-9513, E-mail: m-sasaki@czc.hokudai.ac.jp

31 Hirofumi Sawa, Tel: +81-11-706-5185, E-mail: h-sawa@czc.hokudai.ac.jp

32

33 Running title: Long-distance dispersal of rotaviruses in fruit bats

34

Abstract

Group A rotavirus (RVA) is a major cause of diarrhea in children worldwide. Although RVA infects many animals, little is known about RVA in bats. The present study investigated the genetic diversity of RVA in Zambian bats. We identified RVA from two straw-colored fruit bats (*Eidolon helvum*) and an Egyptian fruit bat (*Rousettus aegyptiacus*), and analyzed the genome sequences of these strains. Genome segments of the RVA strains from Zambian *E. helvum* showed 97%–99% nucleotide sequence identity with those of other RVA strains from *E. helvum* in Cameroon, which is 2,800 km from the sampling locations. These findings suggest that migratory straw-colored fruit bat species, distributed across sub-Saharan Africa, have the potential to disseminate RVA across long distances. By contrast, the RVA strain from Zambian *R. aegyptiacus* carried highly divergent *NSP2* and *NSP4* genes, leading us to propose novel genotypes N21 and E27, respectively. Notably, this RVA strain also shared the same genotype for *VP6* and *NSP3* with the RVA strains from Zambian *E. helvum*, suggesting interspecies transmission and genetic reassortment may have occurred between these two bat species in the past. Our study has important implications for RVA dispersal in bat populations, and expands our knowledge of the ecology, diversity and evolutionary relationships of RVA.

Keywords

Rotavirus; African fruit bats; Long-distance dispersal; Interspecies transmission; Phylogenetic analysis; Novel RVA genotypes

1. Introduction

Rotavirus is a major causative agent of gastroenteritis in children under five, with more than 120,000 cases of diarrheal death annually, worldwide (Clark et al., 2017). Among nine species of rotavirus (groups A to I), group A rotavirus (RVA) is the major species and the most well studied to date. RVA has a genome of 11 segments of double-stranded RNA, which encode the viral structural proteins (VP1–4, VP6 and VP7) and the non-structural proteins (NSP1–6). The current nomenclature system of RVA defines the genotype as: Gx-P[x]-Ix-Rx-Cx-Mx-Ax-Nx-Tx-Ex-Hx for the VP7-, VP4-, VP6-, VP1-, VP2-, VP3-, NSP1-, NSP2-, NSP3-, NSP4- and NSP5/6-encoding genes, respectively (Matthijssens et al., 2011a; Matthijssens et al., 2008). Based on the genome sequence, all RVA isolates are classified into genotypes in accordance with the recommendations of the Rotavirus Classification Working Group (RCWG) to ensure uniformity (Matthijssens et al., 2008). This classification system has been widely adopted and has greatly facilitated the analysis of RVA sequence data, which has uncovered high genetic diversity and proposed new genotypes (Esona et al., 2018; He et al., 2017; Ianiro et al., 2017; Li et al., 2016; Rojas et al., 2016; Rojas et al., 2017; Yinda et al., 2016).

Bats harbor numerous pathogens and act as reservoir hosts of high-consequence zoonotic viruses (Hayman, 2016; Olival et al., 2017). A limited number of studies have reported on RVA from frugivorous bats: *Eidolon helvum* in Kenya and Cameroon (Esona et al., 2010; Yinda et al., 2016), *Rousettus aegyptiacus* in Kenya (Waruhiu et al., 2017) and *Rousettus leschenaultii* in China (He et al., 2017), and insectivorous bats: *Molossus molossus* in Brazil (Asano et al., 2016), *Rhinolophus simulator* in Zambia (Sasaki et al., 2016), *Taphozous mauritanus* in Kenya (Waruhiu et al., 2017), and *Rhinolophus hipposideros*, *Aselliscus stoliczkanus*, *Scotophilus kuhlii*, *Hipposideros pomona* and

81 *Taphozous melanopogon* in China (He et al., 2017; He et al., 2013; Xia et al., 2014).
82 Genetic characterization of bat RVAs has led to discoveries of new RVA genotypes. In
83 addition, these studies revealed that bat RVAs not only carry unique genotypes exclusively
84 observed in bats, but also share some genome segments with RVAs derived from humans
85 and other mammals, indicative of interspecies transmission and the zoonotic potential of
86 bat-borne RVA (Esona et al., 2010; He et al., 2017; Sasaki et al., 2016). Although the
87 sporadic detection of RVA from bats worldwide has demonstrated that RVA infection can
88 occur in some bat species, thus far, the genotypic tropism(s) and transmission cycle of RVA
89 in bat populations are poorly understood.

90 Previously, we reported RVA strain LUS12-14 from the insectivorous horseshoe bat
91 species, *R. simulator*, in Zambia (Sasaki et al., 2016). In the present study, we screened
92 insectivorous and frugivorous bat species in Zambia to investigate the prevalence of RVA
93 infection and also to determine host species susceptible to RVA infection. Three RVA
94 strains were newly identified from the fruit bats, *E. helvum* and *R. aegyptiacus*. Our
95 findings have important implications for RVA dissemination across long distances in
96 African fruit bats and provide evidence of interspecies transmission and genetic
97 reassortment events among African bat RVAs.

2. Materials and Methods

2.1. Sample collection and ethics statement

From 2014 to 2015, 60 frugivorous and 40 insectivorous bat species were captured at five different locations in Zambia, with permission from the Department of National Parks and Wildlife (formerly the Zambia Wildlife Authority), Ministry of Tourism and Arts (Act No. 12 of 1998). Spleen, liver, kidney and colon tissues were collected through dissection. Bats were speciated based on morphology and sequencing of ribosomal RNA and cytochrome b loci, as previously described (Sasaki et al., 2012). Sample information is summarized in Table 1.

2.2 Nested RT-PCR screening for RVA

Total RNA was extracted from bat colon tissue using the QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. For nested RT-PCR screening, cDNA was synthesized using random hexamers and SuperScript IV Reverse Transcriptase (Invitrogen; Thermo Fisher Scientific, Waltham, MA, USA), and subjected to nested PCR amplification employing the Tks Gflex DNA polymerase (Takara Bio, Kusatsu, Japan) and oligonucleotide primers targeting RVA VP7 as follows: RotexoF (5'- MDCGGWTAGMYBYBT TTWAATG -3') and RotexoR (5'- CCCATNGMDATCCAYTTTRTT -3') for the 1st round PCR, and RotinF (5'- TAGCYBYBT TTTRATGTATGGKAT -3') and RotinR (5'- TCCATNGGRTTRCAHARCC -3') for the 2nd round PCR (Li et al., 2016). The thermocycling conditions were: 1 cycle of 94°C for 2 min followed by 35 cycles of 98°C for 10 s, 46°C (1st PCR) or 50°C (2nd PCR) for 15 s and 68°C for 30 s. Amplicons were purified with the MonoFas DNA Purification Kit I (GL Sciences, Tokyo, Japan) and sequenced using BigDye Terminator v3.1 Cycle

Sequencing Kit (Applied Biosystems; Thermo Fisher Scientific).

2.3 Amplification and sequencing of RVA genome segments

Each genome segment was separately amplified by a nested RT-PCR strategy. After denaturation at 95°C for 5 min, RNA samples were reverse transcribed with SuperScript IV Reverse Transcriptase and specific primer sets targeting the 5' and 3' ends of each of the 11 RVA genome segments, referred to as *exoF* or *exoR*, as described previously (Li et al., 2016). The 1st round PCR was performed with Tks Gflex DNA polymerase and the gene-specific primer pairs that were used in the reverse transcription step. The 2nd round PCR was performed with Tks Gflex DNA polymerase and the inner primer set, referred to as *inF* or *inR* as described previously (Li et al., 2016). The PCR amplicons were sequenced as described above.

2.4 Assignment of RVA genotypes

Genotypes of the identified segments were determined using the online tool RotaC (<http://rotac.regatools.be>) or following the judgment of RCWG (<https://rega.kuleuven.be/cev/viralmetagénomics/virus-classification/rcwg>) (Maes et al., 2009).

2.5 Phylogenetic analysis

Maximum likelihood phylogenetic trees with 500 bootstrap replicates were inferred from multiple nucleotide sequence alignments of full-length genes of RVA reference strains and bat RVAs using MEGA7 software (Kumar et al., 2016). For the maximum likelihood analyses, the GTR+G+I model for *VP1* and *VP6*, the GTR+G model for *VP7*, *NSP2* and

147 *NSP3*, and the TN93+G model for *NSP4* were employed based on the “Find best
148 DNA/protein model” in the MEGA7 software.

149

150 2.6 Nucleotide sequence accession numbers

151 The determined RVA genome sequences were deposited in the DDBJ/EMBL/GenBank
152 database under accession no. LC277159–LC277170.

153

3. Results

3.1 Detection of RVA VP7 genome segments in Zambian fruit bats

During 2014–2015, three frugivorous bat species (*E. helvum*, *Epomophorus crypturus*, *R. aegyptiacus*) and three insectivorous bat species (*Hipposideros gigas*, *Nycteris* sp., *Miniopterus schreibersii*) were captured in Zambia (Table 1). No bats showed signs of serious infection, including diarrhea. RNA was extracted from 100 bat colon samples and subjected to nested RT-PCR screening targeting the conserved VP7 gene of RVA. The screening identified three VP7 positive samples from Zambian fruit bats: strain ZFB14-52 from an adult male *E. helvum*, ZFB14-135 from an adult female *E. helvum* and ZFB14-126 from an adult female *R. aegyptiacus*. To determine the genotype, we attempted to amplify the near-complete sequence of the VP7 gene and recovered it from ZFB14-52 and ZFB14-135, but not ZFB14-126.

3.2 Detection of RVA genome segments from VP7-positive bats

To further characterize the RVA strains detected in Zambian fruit bats, we sought to identify the remaining 10 genome segments of RVA in the VP7-positive specimens. The genome segments were amplified by nested RT-PCR. All RT-PCR products were sequenced directly and multiple peaks were not observed in the sequencing electropherogram, suggesting each amplicon originated from a single RVA strain. We determined the sequences of VP6 and NSP3 from strain ZFB14-52, VP6 and NSP2-4 from strain ZFB14-126, and VP1, VP6 and NSP3 from strain ZFB14-135 (Table 2). Despite multiple attempts by RT-PCR, the sequences of the other RVA genome segments remain to be elucidated.

3.3 Sequence comparison and phylogenetic analysis of *VP7*, *VP1*, *VP6* and *NSP3*

Genotype identification was performed employing the RotaC online tool, which indicated that *VP7* of ZFB14-52 and ZFB14-135 could be assigned to the G31 genotype (Table 2). The sequence of these *VP7* genome segments showed 98% nucleotide identity to RVA strain BatLi08, belonging to the G31 genotype, which was discovered previously from *E. helvum* in the South West region of Cameroon (Yinda et al., 2016). Phylogenetic analysis of *VP7* showed that ZFB14-52 and ZFB14-135 clustered with BatLi08 and were distantly related to BatLi09, BatLi10 and BatLy17 belonging to the G30 genotype (Figure 1), which were also identified from *E. helvum* in Cameroon (Yinda et al., 2016).

The *VP1* of ZFB14-135 showed 94% nucleotide identity with that of BatLi08 belonging to the R15 genotype. The *VP6* of ZFB14-52, ZFB14-126 and ZFB14-135 showed 97%, 90% and 99% nucleotide identities with that of BatLi08 belonging to the I22 genotype, respectively. The *NSP3* of ZFB14-52, ZFB14-126 and ZFB14-135 showed 98%, 90% and 99% nucleotide identities with that of BatLi08 belonging to the T17 genotype, respectively. Phylogenetic analyses of these genome segments revealed that ZFB14-52, ZFB14-126 and ZFB14-135 formed a discrete cluster with Cameroonian bat RVAs (BatLi08, BatLi09, BatLi10 and BatLy17) and were clearly separable from other previously described bat RVAs (Figure 1). Collectively, these results indicated that Zambian fruit bat RVAs harbor the same genotypes of *VP1*, *VP6*, *VP7* and *NSP3* as Cameroonian fruit bat RVAs and exhibit high nucleotide sequence identities with these genome segments.

3.4 Identification of novel *NSP2* and *NSP4* genotypes

The *NSP2* and *NSP4* genotypes of ZFB14-126 could not be determined by RotaC due to their nucleotide sequence divergence. BLAST search analyses indicated that both *NSP2*

and NSP4 of ZFB14-126 showed <80% nucleotide sequence identity with all available RVA sequence data deposited in the DDBJ/EMBL/GenBank public databases. Therefore, these sequences were submitted to RCWG and were approved as new genotypes: N21 for NSP2 and E27 for NSP4 (RCWG, 2018). Phylogenetic analyses revealed that NSP2 of ZFB14-126 was distantly related to other RVAs and segregated in a different clade from the Cameroonian bat RVA N15 genotype (Figure 2). Furthermore, NSP4 of ZFB14-126 was highly divergent from all other RVAs and represented a distinct lineage of NSP4 (Figure 2). These findings indicate that RVA strain ZFB14-126 possessed discordant NSP2 and NSP4 gene segments when compared with other genome segments.

4. Discussion

E. helvum is distributed across sub-Saharan Africa and previous studies revealed that the mean migratory distance of *E. helvum* was 860 km with a range from 270–3,000 km (Ossa et al., 2012; Richter and Cumming, 2008). Prior genetic studies revealed a panmictic population of *E. helvum* across continental Africa, suggesting that this bat species travels and interbreeds over long distances (Peel et al., 2013). In Zambia, over one million *E. helvum* roost from October to December (Peel et al., 2017). Previous reports have suggested that migration of *E. helvum* facilitates the introduction of viruses into the bat population, such as filoviruses, henipaviruses, lyssaviruses and coronaviruses (Drexler et al., 2012; Leopardi et al., 2016; Ogawa et al., 2015; Peel et al., 2013).

In this study, we identified bat RVA strains ZFB14-52 and ZFB14-135 from *E. helvum* in Zambia, which belong to genotypes G31 for VP7, R15 for VP1, I22 for VP6, and T17 for NSP3. These genotypes were initially identified from Cameroonian *E. helvum* by another research group who proposed that novel RVA genotype constellations exist in *E. helvum*, such as has been determined in humans and domesticated animals (Matthijnssens et al., 2011b; Matthijnssens and Van Ranst, 2012; Yinda et al., 2016). Our results support this view that certain RVA genotype constellations exist in this bat species. Interestingly, these Zambian bat RVA strains (ZFB14-52 and ZFB14-135) carried VP7, VP6 and NSP3 genome segments that shared 97%–99% nucleotide sequence identity with those of BatLi08 from *E. helvum* in Limbe, Cameroon, at least 2,800 km apart from our sampling locations. Notably, it has been reported that RVA strain BatLy03 from Cameroonian *E. helvum* shared the same genotypes for VP2, VP6, VP7, NSP2, NSP3 and NSP5 as strain KE4852 from Kenyan *E. helvum* (Yinda et al., 2016). These findings suggest that the migration of *E. helvum* may have the potential to spread RVA across long distances and impact on the viral ecology.

Recent genetic analyses of bat RVAs have discovered new genotypes of this virus (Asano et al., 2016; Esona et al., 2010; He et al., 2017; Yinda et al., 2016). In this study, we identified the previously unrecognized genotypes N21 for *NSP2* and E27 for *NSP4* in RVA strain ZFB14-126 from *R. aegyptiacus* in Zambia. Both N21 and E27 were distinguished from other mammalian RVAs by long branch lengths in their phylogenies (Figure 2). These results indicate that previously unrecognized genotypes are harbored by bats with unique evolutionary histories. In addition, ZFB14-126 shared the same I22 and T17 genotypes with ZFB14-52 and ZFB14-135 from *E. helvum* (Table 2), suggesting interspecies transmission and genetic reassortment may have occurred between these two bat species in the past. However, we could not formally exclude the possibility of mixed infection with different RVA strains in this individual bat.

Unfortunately, we failed to recover all RVA gene segments of these RVA strains and their complete genotype constellations remain to be elucidated. Although several universal primer sets targeting the 5' and 3' regions of each genome segment were employed to amplify RVA genomes and determine the genotypes (Fujii et al., 2012; Gentsch et al., 1992; Gouvea et al., 1990; Li et al., 2016), there are significant nucleotide mismatches between these primers and recently described bat RVA genomes (Yinda et al., 2016). A high-throughput sequencing approach may help to identify divergent bat RVA genomes and determine the genotype constellations (He et al., 2017; Yinda et al., 2016).

Previous studies reported that bat RVAs carry genome segments closely related to other mammalian RVAs, including human RVAs (Asano et al., 2016; He et al., 2017; Sasaki et al., 2016). In this study, all detected genome segments were divergent to those of other mammalian RVAs and are tentatively considered to be bat-specific. However, Esona *et al.* recently identified RVA strain KE4852 from *E. helvum*, which carried *VP4* and *NSP4* genes

with shorter genetic distances to other mammalian RVAs providing evidence of interspecies transmission between *E. helvum* and other mammal host species (Esona et al., 2010). Therefore, further studies with increasing numbers of specimens are required to evaluate the public health risk of RVA harbored by Zambian bats. and to delineate further the genetic diversity and evolutionary history of these viruses.

Acknowledgments

We thank Sakae Kashiwara, Emiko Nakagawa, Edgar Simulundu, Ryo Nakao, Wakako Furuyama, Chiho Kaneko, Joseph Ndebe, Penjaninge Kapila, Ladslav Moonga, John Yabe and the Department of National Parks and Wildlife of the Ministry of Tourism and Arts (formerly ZAWA) for technical assistance in Zambia. We also thank Dr. Jelle Matthijnsens and RCWG for help with RVA genotyping, and Kate Fox from Edanz Group for editing a draft of this manuscript.

Funding

This work was supported by the Japan Initiative for Global Research Network of Infectious Diseases (J-GRID) from Japan Agency for Medical Research and Development (AMED) (JP18fm0108008); AMED/Japan International Cooperation Agency (JICA) within the framework of the Science and Technology Research Partnership for Sustainable Development (SATREPS); Grants-in-Aid for Scientific Research on Innovative Areas from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan (16H06429, 16H06431, 16K21723); and Japan Society for the Promotion of Science (JSPS) KAKENHI (16H05805).

Declaration of interest

The authors declare that they have no conflicts of interest.

References

- Asano, K.M., Gregori, F., Hora, A.S., Scheffer, K.C., Fahl, W.O., Iamamoto, K., Mori, E., Silva, F.D., Taniwaki, S.A., Brandão, P.E., 2016. Group A rotavirus in Brazilian bats: description of novel T15 and H15 genotypes. *Arch Virol* 161(11), 3225-3230.
- Clark, A., Black, R., Tate, J., Roose, A., Kotloff, K., Lam, D., Blackwelder, W., Parashar, U., Lanata, C., Kang, G., Troeger, C., Platts-Mills, J., Mokdad, A., Sanderson, C., Lamberti, L., Levine, M., Santosham, M., Steele, D., Network, G.R.S., 2017. Estimating global, regional and national rotavirus deaths in children aged <5 years: Current approaches, new analyses and proposed improvements. *PLoS One* 12(9), e0183392.
- Drexler, J.F., Corman, V.M., Müller, M.A., Maganga, G.D., Vallo, P., Binger, T., Gloza-Rausch, F., Cottontail, V.M., Rasche, A., Yordanov, S., Seebens, A., Knörnschild, M., Oppong, S., Adu Sarkodie, Y., Pongombo, C., Lukashev, A.N., Schmidt-Chanasit, J., Stöcker, A., Carneiro, A.J., Erbar, S., Maisner, A., Fronhoffs, F., Buettner, R., Kalko, E.K., Kruppa, T., Franke, C.R., Kallies, R., Yandoko, E.R., Herrler, G., Reusken, C., Hassanin, A., Krüger, D.H., Matthee, S., Ulrich, R.G., Leroy, E.M., Drosten, C., 2012. Bats host major mammalian paramyxoviruses. *Nat Commun* 3, 796.
- Esona, M.D., Mijatovic-Rustempasic, S., Conrardy, C., Tong, S., Kuzmin, I.V., Agwanda, B., Breiman, R.F., Banyai, K., Niezgoda, M., Rupprecht, C.E., Gentsch, J.R., Bowen, M.D., 2010. Reassortant group A rotavirus from straw-colored fruit bat (*Eidolon helvum*). *Emerg Infect Dis* 16(12), 1844-1852.
- Esona, M.D., Roy, S., Rungsisuriyachai, K., Gautam, R., Hermelijn, S., Rey-Benito, G.,

Bowen, M.D., 2018. Molecular characterization of a human G20P[28] rotavirus a strain with multiple genes related to bat rotaviruses. *Infect Genet Evol* 57, 166-170.

Fujii, Y., Shimoike, T., Takagi, H., Murakami, K., Todaka-Takai, R., Park, Y., Katayama, K., 2012. Amplification of all 11 RNA segments of group A rotaviruses based on reverse transcription polymerase chain reaction. *Microbiol Immunol* 56(9), 630-638.

Gentsch, J.R., Glass, R.I., Woods, P., Gouvea, V., Gorziglia, M., Flores, J., Das, B.K., Bhan, M.K., 1992. Identification of group A rotavirus gene 4 types by polymerase chain reaction. *J Clin Microbiol* 30(6), 1365-1373.

Gouvea, V., Glass, R.I., Woods, P., Taniguchi, K., Clark, H.F., Forrester, B., Fang, Z.Y., 1990. Polymerase chain reaction amplification and typing of rotavirus nucleic acid from stool specimens. *J Clin Microbiol* 28(2), 276-282.

Hayman, D.T., 2016. Bats as Viral Reservoirs. *Annu Rev Virol* 3(1), 77-99.

He, B., Huang, X., Zhang, F., Tan, W., Matthijnsens, J., Qin, S., Xu, L., Zhao, Z., Yang, L., Wang, Q., Hu, T., Bao, X., Wu, J., Tu, C., 2017. Group A Rotaviruses in Chinese Bats: Genetic Composition, Serology, and Evidence for Bat-to-Human Transmission and Reassortment. *J Virol* 91(12).

He, B., Yang, F., Yang, W., Zhang, Y., Feng, Y., Zhou, J., Xie, J., Bao, X., Guo, H., Li, Y., Xia, L., Li, N., Matthijnsens, J., Zhang, H., Tu, C., 2013. Characterization of a novel G3P[3] rotavirus isolated from a lesser horseshoe bat: a distant relative of feline/canine rotaviruses. *J Virol* 87(22), 12357-12366.

Ianiro, G., Di Bartolo, I., De Sabato, L., Pampiglione, G., Ruggeri, F.M., Ostanello, F., 2017. Detection of uncommon G3P[3] rotavirus A (RVA) strain in rat possessing a human RVA-like VP6 and a novel NSP2 genotype. *Infect Genet Evol* 53, 206-211.

335 Kumar, S., Stecher, G., Tamura, K., 2016. MEGA7: Molecular Evolutionary Genetics
336 Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol*.

337 Leopardi, S., Oluwayelu, D., Meseko, C., Marciano, S., Tassoni, L., Bakarey, S., Monne, I.,
338 Cattoli, G., De Benedictis, P., 2016. The close genetic relationship of lineage D
339 Betacoronavirus from Nigerian and Kenyan straw-colored fruit bats (*Eidolon*
340 *helvum*) is consistent with the existence of a single epidemiological unit across
341 sub-Saharan Africa. *Virus Genes* 52(4), 573-577.

342 Li, K., Lin, X.D., Huang, K.Y., Zhang, B., Shi, M., Guo, W.P., Wang, M.R., Wang, W.,
343 Xing, J.G., Li, M.H., Hong, W.S., Holmes, E.C., Zhang, Y.Z., 2016. Identification of
344 novel and diverse rotaviruses in rodents and insectivores, and evidence of
345 cross-species transmission into humans. *Virology* 494, 168-177.

346 Maes, P., Matthijnsens, J., Rahman, M., Van Ranst, M., 2009. RotaC: a web-based tool for
347 the complete genome classification of group A rotaviruses. *BMC Microbiol* 9, 238.

348 Matthijnsens, J., Ciarlet, M., McDonald, S.M., Attoui, H., Bányai, K., Brister, J.R., Buesa,
349 J., Esona, M.D., Estes, M.K., Gentsch, J.R., Iturriza-Gómara, M., Johne, R.,
350 Kirkwood, C.D., Martella, V., Mertens, P.P., Nakagomi, O., Parreño, V., Rahman,
351 M., Ruggeri, F.M., Saif, L.J., Santos, N., Steyer, A., Taniguchi, K., Patton, J.T.,
352 Desselberger, U., Van Ranst, M., 2011a. Uniformity of rotavirus strain nomenclature
353 proposed by the Rotavirus Classification Working Group (RCWG). *Arch Virol*
354 156(8), 1397-1413.

355 Matthijnsens, J., Ciarlet, M., Rahman, M., Attoui, H., Bányai, K., Estes, M.K., Gentsch,
356 J.R., Iturriza-Gómara, M., Kirkwood, C.D., Martella, V., Mertens, P.P., Nakagomi,
357 O., Patton, J.T., Ruggeri, F.M., Saif, L.J., Santos, N., Steyer, A., Taniguchi, K.,
358 Desselberger, U., Van Ranst, M., 2008. Recommendations for the classification of

359 group A rotaviruses using all 11 genomic RNA segments. Arch Virol 153(8),
 360 1621-1629.

361 Matthijnssens, J., De Grazia, S., Piessens, J., Heylen, E., Zeller, M., Giammanco, G.M.,
 362 Bányai, K., Buonavoglia, C., Ciarlet, M., Martella, V., Van Ranst, M., 2011b.
 363 Multiple reassortment and interspecies transmission events contribute to the
 364 diversity of feline, canine and feline/canine-like human group A rotavirus strains.
 365 Infect Genet Evol 11(6), 1396-1406.

366 Matthijnssens, J., Van Ranst, M., 2012. Genotype constellation and evolution of group A
 367 rotaviruses infecting humans. Curr Opin Virol 2(4), 426-433.

368 Ogawa, H., Miyamoto, H., Nakayama, E., Yoshida, R., Nakamura, I., Sawa, H., Ishii, A.,
 369 Thomas, Y., Nakagawa, E., Matsuno, K., Kajihara, M., Maruyama, J., Nao, N.,
 370 Muramatsu, M., Kuroda, M., Simulundu, E., Changula, K., Hang'ombe, B.,
 371 Namangala, B., Nambota, A., Katampi, J., Igarashi, M., Ito, K., Feldmann, H.,
 372 Sugimoto, C., Moonga, L., Mweene, A., Takada, A., 2015. Seroepidemiological
 373 Prevalence of Multiple Species of Filoviruses in Fruit Bats (*Eidolon helvum*)
 374 Migrating in Africa. J Infect Dis 212 Suppl 2, S101-108.

375 Olival, K.J., Hosseini, P.R., Zambrana-Torrel, C., Ross, N., Bogich, T.L., Daszak, P., 2017.
 376 Host and viral traits predict zoonotic spillover from mammals. Nature 546(7660),
 377 646-650.

378 Ossa, G., Kramer-Schadt, S., Peel, A.J., Scharf, A.K., Voigt, C.C., 2012. The movement
 379 ecology of the straw-colored fruit bat, *Eidolon helvum*, in sub-Saharan Africa
 380 assessed by stable isotope ratios. PLoS One 7(9), e45729.

381 Peel, A.J., Sargan, D.R., Baker, K.S., Hayman, D.T.S., Barr, J.A., Crameri, G., Suu-Ire, R.,
 382 Broder, C.C., Lembo, T., Wang, L.F., Fooks, A.R., Rossiter, S.J., Wood, J.L.N.,

383 Cunningham, A.A., 2013. Continent-wide panmixia of an African fruit bat
384 facilitates transmission of potentially zoonotic viruses. *Nat Commun* 4, 2770.

385 Peel, A.J., Wood, J.L., Baker, K.S., Breed, A.C., Carvalho, A.D., Fernández-Loras, A.,
386 Gabrieli, H.S., Gembu, G.-C., Kakengi, V.A., Kaliba, P.M., 2017. How Does
387 Africa's Most Hunted Bat Vary Across the Continent? Population Traits of the
388 Straw-Coloured Fruit Bat (*Eidolon helvum*) and Its Interactions with Humans. *Acta*
389 *Chiropterologica* 19(1), 77-92.

390 Rotavirus Classification Working Group (RCWG), 2018. List of accepted genotypes.
391 <https://rega.kuleuven.be/cev/viralmetagénomics/virus-classification/rcwg>.

392 Richter, H., Cumming, G., 2008. First application of satellite telemetry to track African
393 straw - coloured fruit bat migration. *Journal of Zoology* 275(2), 172-176.

394 Rojas, M., Gonçalves, J.L., Dias, H.G., Manchego, A., Pezo, D., Santos, N., 2016.
395 Whole-genome characterization of a Peruvian alpaca rotavirus isolate expressing a
396 novel VP4 genotype. *Vet Microbiol* 196, 27-35.

397 Rojas, M.A., Gonçalves, J.L.S., Dias, H.G., Manchego, A., Santos, N., 2017. Identification
398 of two novel Rotavirus A genotypes, G35 and P[50], from Peruvian alpaca faeces.
399 *Infect Genet Evol* 55, 71-74.

400 Sasaki, M., Orba, Y., Sasaki, S., Gonzalez, G., Ishii, A., Hang'ombe, B.M., Mweene, A.S.,
401 Ito, K., Sawa, H., 2016. Multi-reassortant G3P[3] group A rotavirus in a horseshoe
402 bat in Zambia. *J Gen Virol* 97(10), 2488-2493.

403 Sasaki, M., Setiyono, A., Handharyani, E., Rahmadani, I., Taha, S., Adiani, S., Subangkit,
404 M., Sawa, H., Nakamura, I., Kimura, T., 2012. Molecular detection of a novel
405 paramyxovirus in fruit bats from Indonesia. *Virol J* 9, 240.

406 Waruhiu, C., Ommeh, S., Obanda, V., Agwanda, B., Gakuya, F., Ge, X.Y., Yang, X.L., Wu,

L.J., Zohaib, A., Hu, B., Shi, Z.L., 2017. Molecular detection of viruses in Kenyan bats and discovery of novel astroviruses, caliciviruses and rotaviruses. *Virol Sin* 32(2), 101-114.

Xia, L., Fan, Q., He, B., Xu, L., Zhang, F., Hu, T., Wang, Y., Li, N., Qiu, W., Zheng, Y., Matthijnssens, J., Tu, C., 2014. The complete genome sequence of a G3P[10] Chinese bat rotavirus suggests multiple bat rotavirus inter-host species transmission events. *Infect Genet Evol* 28, 1-4.

Yinda, C.K., Zeller, M., Conceição-Neto, N., Maes, P., Deboutte, W., Beller, L., Heylen, E., Ghogomu, S.M., Van Ranst, M., Matthijnssens, J., 2016. Novel highly divergent reassortant bat rotaviruses in Cameroon, without evidence of zoonosis. *Sci Rep* 6, 34209.

Figure Legends

Fig. 1 Phylogenetic analysis based on the nucleotide sequences of the *VP7*, *VP6*, *VP1* and *NSP3* genes

The group A rotavirus (RVA) strains ZFB14-52, -126 and -135 identified in this study are highlighted in red. Other reference bat RVAs included in the analysis are colored in blue. Genotypes are shown to the left of each taxon. The bootstrap values above 70 after 500 replicates are shown at tree nodes. The scale bars represent the numbers of nucleotide substitutions per site.

Fig. 2 Phylogenetic analysis based on the nucleotide sequences of the *NSP2* and *NSP4* genes

The group A rotavirus (RVA) strain ZFB14-126 identified in this study is highlighted in red. Other reference bat RVAs included in the analysis are colored in blue. Genotypes are shown to the left of each taxon. The bootstrap values above 70 after 500 replicates are shown at tree nodes. The scale bars represent the numbers of nucleotide substitutions per site.

437 **Table 1. Sample information and RT-PCR screening results for rotavirus**

Bat species	Location	RT-PCR positive/total
Fruit bats		
<i>Eidolon helvum</i>	Ndola	1/10
<i>Eidolon helvum</i>	Kasanka national park	1/10
<i>Epomophorus crypturus</i>	Monze	0/20
<i>Rousettus aegyptiacus</i>	Lusaka	1/20
Insectivorous bats		
<i>Hipposideros gigas</i>	Lusaka	0/10
<i>Miniopterus schreibersii</i>	Lusaka	0/10
<i>Nycteris</i> sp.	Livingstone	0/20

438

439 **Table 2. Genotype constellations of African bat-borne RVA strains.**

Strain name	Host	Location	Genotype										
			VP7	VP4	VP6	VP1	VP2	VP3	NSP1	NSP2	NSP3	NSP4	NSP5
RVA/Bat-wt/ZMB/ZFB1	<i>Eidolon</i>												
4-52/2014/G31P[x]	<i>helvum</i>	Zambia	G31	P[x]	I22	Rx	Cx	Mx	Ax	Nx	T17	Ex	Hx
RVA/Bat-wt/ZMB/ZFB1	<i>Eidolon</i>												
4-135/2014/G31P[x]	<i>helvum</i>	Zambia	G31	P[x]	I22	R15	Cx	Mx	Ax	Nx	T17	Ex	Hx
RVA/Bat-wt/ZMB/ZFB1	<i>Rousettus</i>												
4-126/2014/GxP[x]	<i>aegyptiacus</i>	Zambia	Gx	P[x]	I22	Rx	Cx	Mx	Ax	N21	T17	E27	Hx
RVA/Bat-wt/KEN/KE485	<i>Eidolon</i>												
2/07/2007/G25P[6]	<i>helvum</i>	Kenya	G25	P[6]	I15	Rx	C8	Mx	Ax	N8	T11	E2	H10
RVA/Bat-wt/CMR/BatLi0	<i>Eidolon</i>												
8/2014/G31P[42]	<i>helvum</i>	Cameroon	G31	P[42]	I22	R15	C15	M14	A25	N15	T17	E22	H17
RVA/Bat-wt/CMR/BatLi0	<i>Eidolon</i>												
9/2014/G30P[42]	<i>helvum</i>	Cameroon	G30	P[42]	I22	R15	C15	M14	A25	N15	T17	E22	H17
RVA/Bat-wt/CMR/BatLi1	<i>Eidolon</i>												
0/2014/G30P[42]	<i>helvum</i>	Cameroon	G30	P[42]	I22	R15	C15	M14	A25	N15	T17	E22	H17
RVA/Bat-wt/CMR/BatLy1	<i>Eidolon</i>												
7/2014/G30P[47]	<i>helvum</i>	Cameroon	G30	P[47]	I22	R15	C15	M14	A25	N15	T17	E22	H17
RVA/Bat-wt/ZMB/LUS12	<i>Rhinolophus</i>												
-14/2012/G3P[3]	<i>simulator</i>	Zambia	G3	P[3]	I3	R2	C2	M3	A9	N2	T3	E2	H3

440 *Strains and genotypes reported in the present study are shown in bold.

Figure 1




Figure 2

