



HOKKAIDO UNIVERSITY

Title	Accumulation of the myosin-II-spectrin complex plays a positive role in apical extrusion of Src-transformed epithelial cells
Author(s)	Takagi, Mikio; Ikegawa, Masaya; Shimada, Takashi et al.
Citation	Genes to cells, 23(11), 974-981 https://doi.org/10.1111/gtc.12643
Issue Date	2018-11
Doc URL	https://hdl.handle.net/2115/75973
Rights	This is the peer reviewed version of the following article: Takagi M, Ikegawa M, Shimada T, et al., Accumulation of the myosin-II-spectrin complex plays a positive role in apical extrusion of Src-transformed epithelial cells, Genes Cells, 2018;23:974-981, which has been published in final form at https://doi.org/10.1111/gtc.12643 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	journal article
File Information	GenesCells23_974.pdf



1 **Accumulation of the myosin-II-spectrin complex plays a**
2 **positive role in apical extrusion of Src-transformed**
3 **epithelial cells**

4

5 Mikio Takagi¹ | Masaya Ikegawa² | Takashi Shimada³ | Susumu Ishikawa¹ | Mihoko
6 Kajita¹ | Takeshi Maruyama¹ | Tomoko Kamasaki¹ | Yasuyuki Fujita¹

7

8 ¹Division of Molecular Oncology, Institute for Genetic Medicine, Graduate School of
9 Chemical Sciences and Engineering, Hokkaido University, Sapporo, Japan

10 ²Genomics, Proteomics and Biomedical Functions, Department of Life and Medical
11 Systems, Faculty of Life and Medical Sciences, Doshisha University, Kyoto, Japan

12 ³SHIMADZU Bioscience Research Partnership, Innovation Center, Shimadzu
13 Scientific Instruments, Bothell, USA

14

15 **Correspondence**

16 Yasuyuki Fujita, Division of Molecular Oncology, Institute for Genetic Medicine,
17 Graduate School of Chemical Sciences and Engineering, Hokkaido University,
18 Sapporo, Japan.

19 Email: yasu@igm.hokudai.ac.jp

20

1 **ABSTRACT**

2 **At the initial stage of carcinogenesis, transformation occurs in single cells within**
3 **the epithelium. Recent studies have revealed that the newly emerging**
4 **transformed cells are often apically eliminated from epithelial tissues. However,**
5 **the underlying molecular mechanisms of this cancer preventive phenomenon still**
6 **remain elusive. In this study, we first demonstrate that myosin-II accumulates in**
7 **Src-transformed cells when they are surrounded by normal epithelial cells.**
8 **Knockdown of the heavy chains of myosin-II substantially diminishes apical**
9 **extrusion of Src cells, suggesting that accumulated myosin-II positively regulates**
10 **the apical elimination of transformed cells. Furthermore, we have identified β -**
11 **spectrin as a myosin-II-binding protein under the co-culture of normal and Src-**
12 **transformed epithelial cells. β -Spectrin is also accumulated in Src cells that are**
13 **surrounded by normal cells, and the β -spectrin accumulation is regulated by**
14 **myosin-II. Moreover, knockdown of β -spectrin significantly suppresses apical**
15 **extrusion of Src cells. Collectively, these results indicate that accumulation of the**
16 **myosin-II-spectrin complex plays a positive role in apical extrusion of Src-**
17 **transformed epithelial cells. Further elucidation of the molecular mechanisms of**
18 **apical extrusion would lead to establishment of a novel type of cancer preventive**
19 **medicine.**

20

1 **1 | INTRODUCTION**

2 In epithelial tissues, abnormal or functionally suboptimal cells are often eliminated
 3 from epithelia. This homeostatic mechanism is, at least partly, mediated by cell
 4 competition. Cell competition is a process by which cells with different properties
 5 compete with each other for survival and space (Amoyel and Bach, 2014; Baker,
 6 2017; Claveria and Torres, 2016; Di Gregorio et al., 2016; Johnston, 2009; Maruyama
 7 and Fujita, 2017; Merino et al., 2016; Morata and Ballesteros-Arias, 2015; Ohsawa et
 8 al., 2018; Vincent et al., 2013; Wagstaff et al., 2013). For example, when oncogenic
 9 Ras- or Src-transformed cells are surrounded by normal epithelial cells, transformed
 10 cells are apically extruded from the epithelial layer. The apical eradication of
 11 transformed cells has been observed in cell culture systems (Hogan et al., 2009; Kajita
 12 et al., 2010; Leung and Brugge, 2012; Wu et al., 2014) as well as in *Drosophila*
 13 (Vidal et al., 2006), zebrafish (Kajita et al., 2010), and mouse (Kon et al., 2017;
 14 Sasaki et al., 2018) *in vivo*. In this process, normal epithelial cells recognize the
 15 presence of the neighboring transformed cells and actively eliminate them from
 16 epithelia, implying that normal epithelia have anti-tumor activity that does not involve
 17 immune cells, which is termed EDAC (epithelial defense against cancer) (Kajita et al.,
 18 2014). However, the underlying molecular mechanisms still remain largely obscure.

19 Non-muscle myosin-II binds to actin and generates mechanical forces, thereby
 20 regulating various cellular processes such as cell migration, cytokinesis, and cell
 21 shape changes (Heissler and Manstein, 2013; Pecci et al., 2018; Vicente-Manzanares
 22 et al., 2009). Myosin-II consists of a homodimer of heavy chains, two regulatory light
 23 chains, and two essential light chains, and activity of myosin-II is regulated by
 24 phosphorylation of regulatory light chains. Mammalian cells express three isoforms of
 25 non-muscle myosin-II, NM IIA, IIB, and IIC, which differ in their heavy chains;

1 among them NM IIA is most widely distributed in tissues. To investigate the
2 competitive interaction between normal and transformed cells, we have established
3 Madin-Darby canine kidney (MDCK) epithelial cells stably expressing oncogenic
4 cSrcY527F or RasV12 in a tetracycline-inducible manner (Hogan et al., 2009; Ohoka
5 et al., 2015). In Src- or Ras-transformed cells that are surrounded by normal cells,
6 phosphorylation of myosin light chain (MLC) of myosin-II is elevated (Hogan et al.,
7 2009; Kajita et al., 2010). The increased MLC phosphorylation is not observed when
8 transformed cells alone are cultured, indicating that activation of myosin-II is induced
9 by the presence of the neighboring normal cells in a non-cell-autonomous manner.
10 Addition of the myosin-II inhibitor blebbistatin or expression of the dominant-
11 negative form of MLC (MLC-AA) in transformed cells suppresses apical extrusion
12 (Hogan et al., 2009; Kajita et al., 2010), indicating that activity of myosin-II plays a
13 crucial role in the elimination of transformed cells.

14 In this study, we demonstrate that not only activity of myosin-II but also its
15 **accumulation** is involved in apical extrusion of transformed cells.

16

2 | RESULTS

2.1 | Accumulation of myosin-IIA plays a positive role in apical extrusion of Src-transformed cells

To further understand the role of myosin-II in apical extrusion of Src-transformed cells, we examined the localization of heavy chains of myosin-II (NMHC-IIA: non-muscle myosin heavy chain IIA) by immunofluorescence. When Src-transformed cells were surrounded by normal epithelial cells, NMHC-IIA was profoundly accumulated at the apical side of Src cells (Figure 1a, b). When Src-transformed cells alone were present, the accumulation of NMHC-IIA was not observed, and the immunofluorescence level of NMHC-IIA was comparable to that in normal cells (Figure 1a, b), suggesting that the presence of the neighboring normal cells induces accumulation of NMHC-IIA in Src cells in a non-cell-autonomous fashion. These results imply that not only activity of myosin-II (Hogan et al., 2009; Kajita et al., 2010) but also **accumulation** of myosin-II is regulated during apical extrusion. Accumulation of NMHC-IIA also occurred in RasV12-transformed cells that were surrounded by normal cells (Figure S1a, b), but less prominently than that in Src-transformed cells; therefore, in this study we focused on the functional role of myosin-II in apical extrusion of Src-transformed cells.

Previous studies have shown that F-actin is not accumulated in Src- or RasV12-transformed cells that are surrounded by normal cells (Hogan et al., 2009; Kajita et al., 2010). Accordingly, addition of cytochalasin D, the inhibitor of actin polymerization, did not significantly affect the accumulation of NMHC-IIA in Src cells (Figure S3a, b). In contrast, the myosin-II inhibitor blebbistatin significantly suppressed the accumulation of NMHC-IIA (Figure S3e, f), **suggesting the presence of certain molecular machinery that recruits active myosin-II to the apical side of Src cells.**

1 Collectively, these results suggest that NMHC-IIA is accumulated in Src-transformed
 2 cells surrounded by normal cells in an F-actin-independent, but myosin-activity-
 3 dependent fashion.

4 To examine the functional role of the myosin-II accumulation in apical extrusion of
 5 Src-transformed cells, we established cSrcY527F-transformed cells stably expressing
 6 NMHC-IIA-shRNA (Figure 1c). When NMHC-IIA-knockdown Src cells were
 7 surrounded by normal cells, apical extrusion was substantially diminished (Figure 1d),
 8 demonstrating that myosin-II plays a crucial role in apical elimination of Src-
 9 transformed cells.

10

11 **2.2 | Accumulation of the myosin-II-spectrin complex plays a positive role in** 12 **apical extrusion of Src-transformed epithelial cells**

13 Previous studies have reported that myosin IIA interacts with the spectrin-actin
 14 membrane skeleton (Puszkin et al., 1978; Smith et al., 2018). Spectrins are large
 15 skeletal molecules that exist mainly as heterotetramers consisting of α -spectrin and β -
 16 spectrin. They form a filamentous network on the cytoplasmic face of the membrane,
 17 thereby providing a scaffold for various proteins (Machnicka et al., 2014; Machnicka
 18 et al., 2012). By immunoprecipitation, we showed that β -spectrin interacted with
 19 NMHC-IIA under the co-culture condition (Figure 2a). Next, subcellular localization
 20 of β -spectrin was examined by immunofluorescence. We found that β -spectrin was
 21 accumulated at the apical side of Src-transformed cells that were surrounded by
 22 normal cells (Figure 2b, c). In contrast, accumulation of β -spectrin was not observed
 23 when Src cells alone were cultured (Figure 2b, c). The non-cell-autonomous
 24 accumulation of β -spectrin was diminished by blebbistatin (Figure S3e, g), but not by

1 cytochalasin D (Figure S3c, d). Thus, the mode of accumulation of β -spectrin was
2 akin to that of myosin-II.

3 To examine the functional role of spectrin in apical extrusion, we established
4 cSrcY527F-transformed cells stably expressing β -spectrin-shRNA (Figure S2a). **The**
5 **amount of co-immunoprecipitated β -spectrin was profoundly diminished when β -**
6 **spectrin was knocked down in Src-transformed cells (Figure S2b), suggesting that the**
7 **interaction between NMHC-IIA and β -spectrin predominantly occurs in Src-**
8 **transformed cells under the mix culture condition.** We found that knockdown of β -
9 spectrin did not significantly affect accumulation of myosin-II (Figure 3a, b). In
10 contrast, knockdown of NMHC-IIA substantially diminished accumulation of β -
11 spectrin (Figure 3c, d), indicating that myosin-II regulates localization of β -spectrin in
12 Src cells surrounded by normal cells. Finally, we investigated the effect of
13 knockdown of β -spectrin on the behavior of Src-transformed cells. When β -spectrin-
14 knockdown Src cells were surrounded by normal cells, apical extrusion was
15 significantly suppressed (Figure 3e). Collectively, these results suggest that
16 accumulation of the myosin-II-spectrin complex plays a positive role in apical
17 extrusion of Src-transformed epithelial cells.

18

1 **3 | DISCUSSION**

2 In this study, we demonstrate that myosin-II and β -spectrin accumulate at the apical
3 region of transformed cells surrounded by normal cells, which positively regulates
4 apical extrusion. As shown in Figure 2a, the higher amount of β -spectrin binds to
5 myosin-II in a single culture of Src-transformed cells than that of normal cells, but is
6 further elevated under the co-culture condition, suggesting that the interaction
7 between normal and transformed cells somehow promotes the complex formation
8 between β -spectrin and myosin-II. Both myosin-II and spectrin have actin-binding
9 property. However, we have not observed accumulation of F-actin at the apical region
10 of Src-transformed cells that are surrounded by normal cells (Kajita et al., 2010).
11 Consistently, the actin-polymerization inhibitor cytochalasin D does not significantly
12 affect the accumulation of myosin-II or spectrin, implying that the interaction between
13 myosin-II and spectrin is regulated by an F-actin-independent mechanism.

14 Previous studies have shown that the paxillin-plectin-EPLIN complex is
15 accumulated in RasV12- or Src-transformed cells when they are surrounded by
16 normal epithelial cells, which promotes apical elimination of the transformed cells
17 (Kadeer et al., 2017; Kasai et al., 2018; Ohoka et al., 2015). But, knockdown of
18 EPLIN does not significantly influence accumulation of NMHC-IIA, and knockdown
19 of NMHC-IIA does not affect accumulation of EPLIN (Figure S4a-e). Thus, it is
20 likely that the myosin-II-spectrin complex regulates apical extrusion of Src-
21 transformed cells independently of the paxillin-plectin-EPLIN complex. It has been
22 revealed that other than the accumulation of the paxillin-plectin-EPLIN complex, a
23 variety of non-cell-autonomous changes can occur in transformed cells during apical
24 extrusion such as induction of the Warburg effect-like metabolic changes and elevated

1 endocytosis (Kon et al., 2017; Saitoh et al., 2017). In future studies, the upstream and
2 downstream regulators of the myosin-II-spectrin complex need to be identified.

3 Apical extrusion of transformed cells has been also observed in various epithelial
4 tissues in mouse *in vivo* systems (Kon et al., 2017; Sasaki et al., 2018). In addition,
5 environmental factors such as obesity and inflammation profoundly influence the
6 frequency of elimination of transformed cells (Sasaki et al., 2018). It should be
7 determined whether and how the myosin-II-spectrin complex is involved in these
8 processes *in vivo*. Further investigation of apical extrusion of transformed cells would
9 lead to a novel approach for cancer prevention.

10

11

4 | EXPERIMENTAL PROCEDURES

4.1 | Antibodies and Materials

Rabbit anti-NMHC-IIA (LS-B2802) antibody was purchased from LifeSpan BioSciences, Inc. Rabbit anti-NMHC-IIA (M 8064) antibody was from Sigma-Aldrich. The former was used for immunofluorescence, and the latter was for western blotting and immunoprecipitation. Mouse anti-EPLIN (sc-136399) and mouse anti- β -spectrin (sc-136074) antibodies were from Santa Cruz Biotechnology. Mouse anti-actin (Clone C4) antibody was from Millipore. Alexa-Fluor-568- and -647-conjugated secondary antibodies were from ThermoFisher Scientific. Hoechst 33342 (Life Technologies) was used at a dilution of 1:5,000. For immunofluorescence, the primary antibodies described above were diluted in phosphate-buffered saline (PBS) containing 1% BSA at 1:100. All secondary antibodies were used at 1:200. Alexa-Fluor-647-conjugated phalloidin (Life Technologies) was used at 1.0 U ml^{-1} . For western blotting, all primary antibodies were used at 1:1,000, and secondary antibodies were at 1:1,000. The following inhibitors were used where indicated: cytochalasin D (Sigma-Aldrich, 100 nM) and blebbistatin (Millipore, 30 μM). DMSO (Sigma-Aldrich) was added as a control.

4.2 | Cell Culture

MDCK, MDCK-pTR GFP-cSrcY527F, and MDCK-pTR GFP-RasV12 cells were cultured as previously described (Hogan et al., 2009; Ohoka et al., 2015). MDCK-pTR GFP-cSrcY527F cells stably expressing EPLIN-shRNA were established as previously described (Ohoka et al., 2015). MDCK-pTR GFP-cSrcY527F cells stably expressing NMHC-IIA-shRNA or β -spectrin-shRNA were established as follows:

1 Double-stranded DNA fragments coding NMHC-IIA-shRNA sequences (NMHC-IIA
 2 -shRNA1: 5'- GATCCCCGCTGCTAAGAAGCTGGTATTTCAAGAGA
 3 ATACCAGCTTCTTAGCAGCTTTTTC -3' and 5'-
 4 TCGAGAAAAAGCTGCTAAGAAGCTGGTATTCTCTTGAAATACCAGCTTCT
 5 TAGCAGCGGG -3' or NMHC-IIA-shRNA2: 5'- GATCCCC
 6 GCAACTGACAGCCATGAAATTCAAGAGATTTTCATGGCTGTCAGTTGCTTTT
 7 TC -3' and 5'-
 8 TCGAGAAAAAGCAACTGACAGCCATGAAATCTCTTGAAATTTTCATGGCTGT
 9 CAGTTGCGGG -3') or β -spectrin-shRNA sequences (β -spectrin-shRNA1: 5'-
 10 GATCCCCGCACAGGTTTGAGAGCCTTTTCAAGAGAAAGGCTCTCAAACCT
 11 GTGCTTTTTTC -3' and 5'-
 12 TCGAGAAAAAGCACAGGTTTGAGAGCCTTTCTCTTGAAAAGGCTCTCAAA
 13 CCTGTGCGGG -3' or β -spectrin-shRNA2: 5'-
 14 GATCCCCGCACACTACATTTGAGCATTTCAGAGAATGCTCAAATGTAGT
 15 GTGCTTTTTTC -3' and 5'-
 16 TCGAGAAAAAGCACACTACATTTGAGCATTCTCTTGAAATGCTCAAATGT
 17 AGTGTGCGGG -3') were inserted into the *Bgl*II and *Xho*I site of pSUPER.neo + gfp
 18 (Oligoengine). MDCK-pTR GFP-cSrcY527F cells were transfected with
 19 pSUPER.neo + gfp NMHC-IIA-shRNA or β -spectrin-shRNA using Lipofectamine
 20 2000 (Invitrogen), followed by antibiotic selection in the medium containing
 21 5 $\mu\text{g ml}^{-1}$ blasticidin (InvivoGen), 400 $\mu\text{g ml}^{-1}$ zeocin (InvivoGen), and
 22 800 $\mu\text{g ml}^{-1}$ G418 (Life Technologies). To induce the expression of GFP-cSrcY527F,
 23 the tetracycline-inducible MDCK-pTR GFP-cSrcY527F cells were treated with
 24 2 $\mu\text{g ml}^{-1}$ tetracycline (Sigma-Aldrich). For inhibitor treatment, the indicated

inhibitors were added together with tetracycline, and cells were further cultured for 16 h or 24 h. For analyses of immunofluorescence or apical extrusion, cells were seeded onto Type-I collagen-mounted coverslips. Type-I collagen (Cellmatrix Type I-A) was obtained from Nitta Geratin and was neutralized on ice to a final concentration of 2 mg ml^{-1} according to the manufacturer's instructions.

4.3 | Immunofluorescence

MDCK-pTR GFP-cSrcY527F or MDCK-pTR GFP-RasV12 cells were mixed with MDCK cells at a ratio of 1:50 and cultured on the collagen matrix as previously described (Hogan et al., 2009). For immunofluorescence analyses, the mixture of cells was incubated for 8–12 h until they formed a monolayer, followed by tetracycline treatment for 16 h. Cells were fixed with 4% paraformaldehyde in PBS and permeabilized with 0.5% Triton X-100 in PBS, followed by blocking with 1% BSA in PBS. Primary or secondary antibodies were incubated for 2 h or 1 h, respectively at ambient temperature. Immunofluorescence images were acquired by the Olympus FV1200 system and the Olympus FV10-ASW software. For quantification of immunofluorescence intensity, more than 30 transformed cells were analyzed for each experiment using the MetaMorph software (Molecular Devices). For analyses of apical extrusions, the samples were prepared as described above, except that cells were treated with tetracycline for 24 h. More than 30 cells were analyzed for each experiment, and apically extruded cells were quantified.

4.4 | Immunoprecipitation and western blotting

For immunoprecipitation, 0.9×10^7 MDCK cells and 0.9×10^7 MDCK-pTR GFP-cSrcY527F cells for mix culture or 1.8×10^7 MDCK or MDCK-pTR GFP-cSrcY527F

1 cells for single culture were seeded in 14.5-cm dishes (one or two dishes for each
2 experimental condition) (Greiner-Bio-One) and cultured for 9 h until a monolayer was
3 formed. Cells were then incubated with tetracycline to induce c-SrcY527F expression.
4 After 16 h culture in the presence of tetracycline, cells were washed with ice-cold
5 PBS containing 1 mM Na_3VO_4 and lysed for 30 min in Triton X-100 lysis buffer
6 (20 mM Tris-HCl [pH 7.5], 150 mM NaCl, and 1% Triton X-100) containing the
7 following inhibitors: 1 mM Na_3VO_4 , 10 mM NaF, 5 $\mu\text{g ml}^{-1}$ leupeptin, 1 mM
8 phenylmethylsulfonyl fluoride, and 7.2 trypsin inhibitor units of aprotinin. After
9 centrifugation of the cell lysates at 21,500 g for 10 min, the supernatant was first pre-
10 cleared with Protein G sepharose (Sigma-Aldrich) at 4°C for 30 min. The pre-cleared
11 cell lysates were then incubated with Dynabeads protein G (Life Technologies)
12 conjugated to control rabbit IgG (5 or 10 μg) at 4°C for 30 min and finally subjected
13 to immunoprecipitation with Dynabeads Protein G conjugated to rabbit anti-NMHC-
14 IIA antibody (5 or 10 μg) at 4°C for 1 h. Immunoprecipitated proteins were subjected
15 to SDS-PAGE, followed by western blotting with anti- β -spectrin antibody. Western
16 blotting was performed as described previously (Hogan et al., 2004). Primary
17 antibodies were used at 1:1,000. Western blotting data were analyzed using
18 ImageQuant™ LAS4010 (GE healthcare). To examine the knockdown efficiency of
19 NMHC-IIA, β -spectrin, or EPLIN, MDCK-pTR GFP-cSrcY527F cells stably
20 expressing NMHC-IIA-shRNA, β -spectrin-shRNA, or EPLIN-shRNA were seeded
21 onto 6-well dishes (Corning). After 12-24 h, the incubated cells were lysed with
22 Triton X-100 lysis buffer containing protease inhibitors (5 $\mu\text{g ml}^{-1}$ leupeptin, 1 mM
23 phenylmethylsulfonyl fluoride, and 7.2 trypsin inhibitor units of aprotinin) and
24 directly boiled with SDS-PAGE sample buffer.

1

2 **4.5 | Data Analyses**3 Two-tailed Student's t -tests were used to determine P -values for statistical analyses.

4

5 **4.6 | Data Availability**6 The datasets generated and/or analyzed during the current study are available from the
7 corresponding author on reasonable request.

8

1

2 **REFERENCES**

3

4 Amoyel, M., and Bach, E.A. (2014). Cell competition: how to eliminate your
5 neighbours. *Development* 141, 988-1000.

6 Baker, N.E. (2017). Mechanisms of cell competition emerging from *Drosophila*
7 studies. *Curr Opin Cell Biol* 48, 40-46.

8 Claveria, C., and Torres, M. (2016). Cell Competition: Mechanisms and Physiological
9 Roles. *Annual review of cell and developmental biology* 32, 411-439.

10 Di Gregorio, A., Bowling, S., and Rodriguez, T.A. (2016). Cell Competition and Its
11 Role in the Regulation of Cell Fitness from Development to Cancer. *Dev Cell*
12 38, 621-634.

13 Heissler, S.M., and Manstein, D.J. (2013). Nonmuscle myosin-2: mix and match. *Cell*
14 *Mol Life Sci* 70, 1-21.

15 Hogan, C., Dupre-Crochet, S., Norman, M., Kajita, M., Zimmermann, C., Pelling,
16 A.E., Piddini, E., Baena-Lopez, L.A., Vincent, J.P., Itoh, Y., *et al.* (2009).
17 Characterization of the interface between normal and transformed epithelial
18 cells. *Nat Cell Biol* 11, 460-467.

19 Hogan, C., Serpente, N., Cogram, P., Hosking, C.R., Bialucha, C.U., Feller, S.M.,
20 Braga, V.M., Birchmeier, W., and Fujita, Y. (2004). Rap1 regulates the
21 formation of E-cadherin-based cell-cell contacts. *Mol Cell Biol* 24, 6690-6700.

22 Johnston, L.A. (2009). Competitive interactions between cells: death, growth, and
23 geography. *Science* 324, 1679-1682.

24 Kadeer, A., Maruyama, T., Kajita, M., Morita, T., Sasaki, A., Ohoka, A., Ishikawa, S.,
25 Ikegawa, M., Shimada, T., and Fujita, Y. (2017). Plectin is a novel regulator for
26 apical extrusion of RasV12-transformed cells. *Scientific reports* 7, 44328.

27 Kajita, M., Hogan, C., Harris, A.R., Dupre-Crochet, S., Itasaki, N., Kawakami, K.,
28 Charras, G., Tada, M., and Fujita, Y. (2010). Interaction with surrounding
29 normal epithelial cells influences signalling pathways and behaviour of Src-
30 transformed cells. *J Cell Sci* 123, 171-180.

31 Kajita, M., Sugimura, K., Ohoka, A., Burden, J., Suganuma, H., Ikegawa, M.,
32 Shimada, T., Kitamura, T., Shindoh, M., Ishikawa, S., *et al.* (2014). Filamin acts
33 as a key regulator in epithelial defence against transformed cells. *Nature*
34 *communications* 5, 4428.

35 Kasai, N., Kadeer, A., Kajita, M., Saitoh, S., Ishikawa, S., Maruyama, T., and Fujita,
36 Y. (2018). The paxillin-plectin-EPLIN complex promotes apical elimination of
37 RasV12-transformed cells by modulating HDAC6-regulated tubulin acetylation.
38 *Scientific reports* 8, 2097.

39 Kon, S., Ishibashi, K., Katoh, H., Kitamoto, S., Shirai, T., Tanaka, S., Kajita, M.,
40 Ishikawa, S., Yamauchi, H., Yako, Y., *et al.* (2017). Cell competition with
41 normal epithelial cells promotes apical extrusion of transformed cells through
42 metabolic changes. *Nat Cell Biol* 19, 530-541.

43 Leung, C.T., and Brugge, J.S. (2012). Outgrowth of single oncogene-expressing cells
44 from suppressive epithelial environments. *Nature* 482, 410-413.

45 Machnicka, B., Czogalla, A., Hryniewicz-Jankowska, A., Boguslawska, D.M.,
46 Grochowalska, R., Heger, E., and Sikorski, A.F. (2014). Spectrins: a structural
47 platform for stabilization and activation of membrane channels, receptors and
48 transporters. *Biochimica et biophysica acta* 1838, 620-634.

- 1 Machnicka, B., Grochowalska, R., Boguslawska, D.M., Sikorski, A.F., and Lecomte,
2 M.C. (2012). Spectrin-based skeleton as an actor in cell signaling. *Cell Mol Life*
3 *Sci* 69, 191-201.
- 4 Maruyama, T., and Fujita, Y. (2017). Cell competition in mammals - novel
5 homeostatic machinery for embryonic development and cancer prevention. *Curr*
6 *Opin Cell Biol* 48, 106-112.
- 7 Merino, M.M., Levayer, R., and Moreno, E. (2016). Survival of the Fittest: Essential
8 Roles of Cell Competition in Development, Aging, and Cancer. *Trends Cell*
9 *Biol* 26, 776-788.
- 10 Morata, G., and Ballesteros-Arias, L. (2015). Cell competition, apoptosis and tumour
11 development. *Int J Dev Biol* 59, 79-86.
- 12 Ohoka, A., Kajita, M., Ikenouchi, J., Yako, Y., Kitamoto, S., Kon, S., Ikegawa, M.,
13 Shimada, T., Ishikawa, S., and Fujita, Y. (2015). EPLIN is a crucial regulator
14 for extrusion of RasV12-transformed cells. *J Cell Sci* 128, 781-789.
- 15 Ohsawa, S., Vaughen, J., and Igaki, T. (2018). Cell Extrusion: A Stress-Responsive
16 Force for Good or Evil in Epithelial Homeostasis. *Dev Cell* 44, 532.
- 17 Pecci, A., Ma, X., Savoia, A., and Adelstein, R.S. (2018). MYH9: Structure, functions
18 and role of non-muscle myosin IIA in human disease. *Gene*.
- 19 Puszkun, S., Maimon, J., and Puszkun, E. (1978). Erythrocyte actin and spectrin.
20 Interactions with muscle contractile and regulatory proteins. *Biochimica et*
21 *biophysica acta* 513, 205-220.
- 22 Saitoh, S., Maruyama, T., Yako, Y., Kajita, M., Fujioka, Y., Ohba, Y., Kasai, N.,
23 Sugama, N., Kon, S., Ishikawa, S., *et al.* (2017). Rab5-regulated endocytosis
24 plays a crucial role in apical extrusion of transformed cells. *Proc Natl Acad Sci*
25 *U S A* 114, E2327-E2336.
- 26 Sasaki, A., Nagatake, T., Egami, R., Gu, G., Takigawa, I., Ikeda, W., Nakatani, T.,
27 Kunisawa, J., and Fujita, Y. (2018). Obesity Suppresses Cell-Competition-
28 Mediated Apical Elimination of RasV12-Transformed Cells from Epithelial
29 Tissues. *Cell Rep* 23, 974-982.
- 30 Smith, A.S., Nowak, R.B., Zhou, S., Giannetto, M., Gokhin, D.S., Papoin, J., Ghiran,
31 I.C., Blanc, L., Wan, J., and Fowler, V.M. (2018). Myosin IIA interacts with the
32 spectrin-actin membrane skeleton to control red blood cell membrane curvature
33 and deformability. *Proc Natl Acad Sci U S A* 115, E4377-E4385.
- 34 Vicente-Manzanares, M., Ma, X., Adelstein, R.S., and Horwitz, A.R. (2009). Non-
35 muscle myosin II takes centre stage in cell adhesion and migration. *Nat Rev*
36 *Mol Cell Biol* 10, 778-790.
- 37 Vidal, M., Larson, D.E., and Cagan, R.L. (2006). Csk-deficient boundary cells are
38 eliminated from normal *Drosophila* epithelia by exclusion, migration, and
39 apoptosis. *Dev Cell* 10, 33-44.
- 40 Vincent, J.P., Fletcher, A.G., and Baena-Lopez, L.A. (2013). Mechanisms and
41 mechanics of cell competition in epithelia. *Nat Rev Mol Cell Biol* 14, 581-591.
- 42 Wagstaff, L., Kolahgar, G., and Piddini, E. (2013). Competitive cell interactions in
43 cancer: a cellular tug of war. *Trends in Cell Biology* 23, 160-167.
- 44 Wu, S.K., Gomez, G.A., Michael, M., Verma, S., Cox, H.L., Lefevre, J.G., Parton,
45 R.G., Hamilton, N.A., Neufeld, Z., and Yap, A.S. (2014). Cortical F-actin
46 stabilization generates apical-lateral patterns of junctional contractility that
47 integrate cells into epithelia. *Nat Cell Biol* 16, 167-178.

1 **ACKNOWLEDGEMENTS**

2 This study was supported by Japan Society for the Promotion of Science (JSPS)
3 Grant-in-Aid for Scientific Research on Innovative Areas 26114001, Grant-in-Aid for
4 Scientific Research (A) 18H03994, Strategic Japanese-Swiss Science and Technology
5 Program, AMED under Grant Number JP17ck0106361 and JP18cm0106234, SAN-
6 ESU GIKEN CO. LTD, the Naito Foundation, and the Takeda Science Foundation.

7

8

1

2 **FIGURE LEGENDS**

3 **FIGURE 1** Accumulation of myosin-II positively regulates apical extrusion of Src-
 4 transformed cells that are surrounded by normal epithelial cells. (a)
 5 Immunofluorescence images of MDCK-pTR GFP-cSrcY527F cells mixed with
 6 normal MDCK cells or cultured alone. Cells were incubated with tetracycline for 16 h
 7 and stained with anti-NMHC-IIA antibody (red) and Hoechst (blue). Scale bar, 10 μ m.
 8 (b) Quantification of the immunofluorescence intensity of NMHC-IIA. Data are mean
 9 $\pm$ SD from three independent experiments. * P <0.05 (Student t -test). More than 30
 10 cells were analyzed in each experimental condition. (c) Establishment of MDCK-pTR
 11 GFP-cSrcY527F cells stably expressing NMHC-IIA-shRNA1 or -shRNA2. Cell
 12 lysates of each NMHC-IIA-knockdown cell line were analyzed by western blotting
 13 with the indicated antibodies. (d) Effect of knockdown of NMHC-IIA on apical
 14 extrusion. MDCK-pTR GFP-cSrcY527F, MDCK-pTR GFP-cSrcY527F NMHC-IIA-
 15 shRNA1 or -shRNA2 cells were co-cultured with normal MDCK cells, and at 24 h
 16 after tetracycline addition, apical extrusion was analyzed. Data are mean $\pm$ SD from
 17 three independent experiments. * P <0.05; ** P <0.005 (Student t -test). More than 30
 18 cells were analyzed in each experimental condition.

19

20 **FIGURE 2** β -Spectrin binds to myosin-II and accumulates in Src-transformed cells
 21 surrounded by normal cells. (a) Co-immunoprecipitation of β -spectrin with anti-
 22 NMHC-IIA antibody. Cells were cultured under three different conditions: i) normal
 23 MDCK cells alone, ii) 1:1 mix culture of normal MDCK and MDCK-pTR GFP-
 24 cSrcY527F cells, and iii) MDCK-pTR GFP-cSrcY527F cells alone. At 16 h after
 25 induction of cSrcY527F expression, the cell lysates were subjected to

1 immunoprecipitation with control IgG or anti-NMHC-IIA antibody, followed by
 2 western blotting with anti- β -spectrin antibody. **Note that the doublet bands of β -**
 3 **spectrin were not observed in a reproducible manner (e.g. Figure S2b), and the**
 4 **significance and identity of these bands are not known at present.** (b)
 5 Immunofluorescence images of MDCK-pTR GFP-cSrcY527F cells mixed with
 6 normal MDCK cells or cultured alone. Cells were incubated with tetracycline for 16 h
 7 and stained with anti- β -spectrin antibody (red) and Hoechst (blue). Scale bar, 10 μ m.
 8 (c) Quantification of the immunofluorescence intensity of β -spectrin. Data are mean $\pm$
 9 SD from three independent experiments. * P <0.05 (Student t -test). More than 30 cells
 10 were analyzed in each experimental condition.

11

12 **FIGURE 3** β -Spectrin also plays a positive role in apical extrusion of Src-
 13 transformed cells. (a and c) Immunofluorescence images of MDCK-pTR GFP-
 14 cSrcY527F or MDCK-pTR GFP-cSrcY527F β -spectrin-shRNA1 (a) or NMHC-IIA-
 15 shRNA1 (c) cells mixed with normal MDCK cells. Cells were incubated with
 16 tetracycline for 16 h and stained with anti- β -spectrin (red) and anti-NMHC-IIA (gray)
 17 antibodies and Hoechst (blue). (b and d) Quantification of the immunofluorescence
 18 intensity of NMHC-IIA (b) or β -spectrin (d). Data are mean $\pm$ SD from three
 19 independent experiments. n.s.: not significant. * P <0.05 (Student t -test). More than 30
 20 cells were analyzed in each experimental condition. (e) Effect of knockdown of β -
 21 spectrin on apical extrusion. MDCK-pTR GFP-cSrcY527F, MDCK-pTR GFP-
 22 cSrcY527F β -spectrin-shRNA1 or -shRNA2 cells were co-cultured with normal cells,
 23 and at 24 h after tetracycline addition, apical extrusion was analyzed. Data are mean $\pm$
 24 SD from three independent experiments. * P <0.05 (Student t -test). More than 30 cells
 25 were analyzed in each experimental condition.






Figure 1 Takagi et al.

a



b



c




Figure 2 Takagi et al.




Figure 3 Takagi et al.