



Title	Analysis for genetic loci controlling protoscolex development in the Echinococcus multilocularis infection using congenic mice
Author(s)	Islam, Md Atiqul; Torigoe, Daisuke; Kameda, Yayoi et al.
Citation	Infection, Genetics and Evolution, 65, 65-71 https://doi.org/10.1016/j.meegid.2018.07.017
Issue Date	2018-11
Doc URL	https://hdl.handle.net/2115/76048
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Infection, Genetics and Evolution65_65_72.pdf



1 **Analysis for genetic loci controlling protoscolex development in the *Echinococcus multilocularis***
2 **infection using congenic mice**

3

4 Md Atiqul Islam^a, Daisuke Torigoe^{a,1}, Yayoi Kameda^{a,2}, Takao Irie^b, Hirokazu Kouguchi^b, Ryo
5 Nakao^c, Md Abdul Masum^d, Osamu Ichii^d, Yasuhiro Kon^d, Hassan T. Tag-EL-Din-Hassan^{a,e,f},
6 Masami Morimatsu^{a,f}, Kinpei Yagi^b, Takashi Agui^{a,f,*}

7

8 ^aLaboratory of Laboratory Animal Science and Medicine, Department of Applied Veterinary Sciences,
9 Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan

10 ^bDepartment of Infectious Diseases, Hokkaido Institute of Public Health, Sapporo 060-0819, Japan

11 ^cLaboratory of Parasitology, Department of Disease Control, ^dLaboratory of Anatomy, Department of
12 Basic Veterinary Sciences, and ^fLaboratory of Laboratory Animal Science and Medicine, Department
13 of Applied Veterinary Sciences, Faculty of Veterinary Medicine, Hokkaido University, Sapporo 060-
14 0818, Japan

15 ^ePoultry Production Department, Mansour University, Mansoura 35516, Egypt.

16

17 ¹Present address: Division of Microbiology and Genetics, Institute of Resource Development and
18 Analysis, Kumamoto University.

19 ²Present address: Division of Bioresources, Research Center for Zoonosis Control, Hokkaido
20 University.

21

22

23

24

25 *Corresponding author at: Laboratory of Laboratory Animal Science and Medicine, Department of
26 Applied Veterinary Sciences, Faculty of Veterinary Medicine, Hokkaido University, Kita 18, Nishi 9,
27 Kita-ku, Sapporo 060-0818, Japan

28 Phone: - +81-11-706-5106. E-mail: agui@vetmed.hokudai.ac.jp

29

30 Abstract

31 The resistance/susceptibility to *Echinococcus multilocularis* infection in mice is genetically controlled.
32 However, genetic factors responsible for these differences remain unknown. Our previous study in
33 genetic linkage analysis has revealed that there is a significant quantitative trait locus (QTL) for the
34 establishment of cyst (*Emcys1*), and a highly significant QTL for the development of protoscolex of *E.*
35 *multilocularis* larvae (*Empsc1*), on mouse chromosomes 6 and 1, respectively. The current study
36 aimed to confirm these QTLs and narrow down the critical genetic region that controls
37 resistance/susceptibility to *E. multilocularis* infection by establishing congenic and subcongenic lines
38 from C57BL/6 (B6) and DBA/2 (D2) mice. For protoscolex development phenotype, two congenic
39 lines, B6.D2-*Empsc1* and D2.B6-*Empsc1* were developed, where responsible QTL, *Empsc1* was
40 introgressed from D2 into B6 background and *vice versa*. For cyst establishment phenotype, two
41 congenic lines, B6.D2-*Emcys1* and D2.B6-*Emcys1* were developed, where responsible QTL, *Emcys1*
42 was introgressed from D2 into B6 background and *vice versa*. Because there was no significant
43 difference in cyst establishment between B6.D2-*Emcys1* and D2.B6-*Emcys1* mice after challenge
44 with *E. multilocularis*, it is suggested that the *Emcys1* does not solely control the cyst establishment in
45 mouse liver. However, infection experiments with B6.D2-*Empsc1* and D2.B6-*Empsc1* mice showed a
46 significant difference in protoscolex development in the cyst. It confirms that the *Empsc1* controls
47 phenotype of the protoscolex development in the cyst. Subsequently, two subcongenic lines, B6.D2-
48 *Empsc1.1* and B6.D2-*Empsc1.2* from B6.D2-*Emcys1* and one subcongenic line, D2.B6-*Empsc1.1*
49 from D2.B6-*Empsc1* were developed to narrow down the critical region responsible for protoscolex
50 development. From the results of infection experiments with *E. multilocularis* in these subcongenic
51 mice, it is concluded that a gene responsible for protoscolex development is located between
52 *D1Mit290* (68.1 cM) and *D1Mit511* (97.3 cM).

53

54 **Key Words:** *Echinococcus multilocularis*; protoscolex; congenic mouse; quantitative trait locus

55

56

58 **1. Introduction:**

59 The aim of the current research is to verify genetic loci responsible for controlling the development
60 of alveolar echinococcosis (AE) by using congenic and subcongenic mice. AE is a zoonotic disease
61 caused by the metacestode stage of tapeworm *Echinococcus multilocularis* characterized by a severe
62 hepatic disorder in many parts of the northern hemisphere (Craig et al., 1996). Infiltrating parasitic
63 growth (hematogenous spread) can cause severe organ damage. Long term treatment is one of the
64 expansive issues and causing health problems (Kern et al., 2003; Reuter et al., 2000). Mainly rodents
65 and occasionally humans become the intermediate hosts being infected by oral ingestion of eggs
66 excreted from contaminated feces of the definitive host carnivores in the environment. The eggs hatch
67 out into oncospheres in the intestine of intermediate hosts. The oncospheres migrate to the liver by
68 penetrating the intestinal wall and develop to metacestode. Mature metacestodes produce huge
69 numbers of protoscoleces in intermediate hosts. (Gottstein et al., 1995). Mice have been widely used
70 as an experimental model to study the host-parasite interplay in the *E. multilocularis* infection.
71 Matsumoto et al (2010) demonstrated that two inbred strains, C57BL/6 (B6) and DBA/2 (D2) mice,
72 differed markedly in their susceptibility to *E. multilocularis* infection. The oral administration of *E.*
73 *multilocularis* eggs to D2 mice established a higher number of cysts in the liver as compared to B6
74 mice. Moreover, a significant number of protoscoleces were observed in the cysts of D2 mice,
75 whereas protoscoleces were completely absent in B6 mice. Additionally, the observation from other
76 studies have supported the results that the susceptibility to *E. multilocularis* infection is genetically
77 controlled both in animals and humans (Nakaya et al., 1997; Hildreth and Granholm, 2003; Vuitton,
78 2003). Several studies have reported that B6 and D2 mice have susceptibility difference against
79 various infectious diseases (David et al., 1995; Adrianus et al., 2009; Marquis et al., 2009; Simon et
80 al., 2009). The genetic factors responsible for difference in the susceptibility to *E. multilocularis*
81 parasite still remains unknown. Considering the marked difference between B6 and D2 mice in
82 susceptibility to *E. multilocularis* infection, quantitative trait locus (QTL) analysis was conducted in
83 backcrossed progenies from B6 and D2 mice (Nakao et al., 2011). QTL mapping is a promising tool

84 for the detection of genetic loci that contribute to the determination of differences in phenotypic
85 variation. QTL analysis has revealed a significant QTL, *Emcys1* for the establishment of *E.*
86 *multilocularis* cysts in the mouse liver, suggesting that it controls the number of cysts in the liver
87 (Nakao et al., 2011). Furthermore, the development of protoscoleces in cysts in the mouse liver were
88 controlled by a distinct highly significant QTL, *Empsc1*, indicating a role of different host factor
89 interplaying with parasites at each developmental stage (Nakao et al., 2011). Lack of information in
90 the interplay between parasite and the intermediate host makes it difficult to focus on certain genes
91 responsible for resistance/susceptibility to *E. multilocularis* infection. It is critically important to
92 address the role of genetic factors to understand the course of infection as well as to get better
93 treatment strategies. To verify that previously identified QTLs are indeed responsible for
94 susceptibility or resistance to *E. multilocularis* infection, making congenic mouse strains is one of the
95 most reliable strategies. Congenic mouse strains are defined as those, in which genetic alteration or
96 mutation is transferred into a standard inbred mouse strain (Markel et al., 1997). Several researchers
97 have published valuable findings using congenic mice that could confirm the responsible genetic
98 regions to address candidate genes in various diseases, including infectious and autoimmune diseases
99 (Shimizu et al., 2007; Marquis et al., 2008; Allen et al., 2006). The purpose of the current study is to
100 verify that previously identified QTLs are indeed responsible for resistance/susceptibility to *E.*
101 *multilocularis* infection by establishing congenic mouse strains from B6 and D2 mice, where the
102 significant QTLs are introgressed from B6 to D2-genetic background and *vice versa* from D2 to B6-
103 genetic background. Furthermore, subcongenic lines were generated to narrow down the critical
104 region, including QTLs responsible for resistance/susceptibility to *E. multilocularis* infection.
105

106 2. Materials and methods

107

108 2.1. Mice

109

110 Specific pathogen-free inbred mice, C57BL/6NSlc (B6) and DBA/2CrSlc (D2) were purchased
111 from Japan SLC (Shizuoka, Japan) to generate congenic and subcongenic lines. For generation of
112 congenic and subcongenic lines, we were adhered to the AAALAC International-accredited program
113 and the Regulation for the Care and Use of Laboratory Animals, Hokkaido University, and animal use
114 protocol was approved by the President of Hokkaido University after the review by the Institutional
115 Animal Care and Use Committee. Parental inbred, congenic, and subcongenic mice were subjected to
116 infection experiments with *E. multilocularis*. Infection experiments were performed in accordance
117 with the regulation of Hokkaido Institute of Public Health and animal use protocol was approved by
118 President of Hokkaido Institute of Public Health after the review by the Ethics Committee of the
119 Institute.

120

121 2.1.1. Congenic lines

122

123 For the *Emcys1*, two congenic lines were generated by the introgression of *Emcys1* in chromosome
124 (Chr) 6 from B6 to D2-genetic background and *vice versa* from D2 to B6-genetic background. The
125 (B6 x D2) F_1 mice were backcrossed to B6 or D2 parental strain [(B6 x D2) F_1 x B6 or (B6 x D2) F_1 x
126 D2]. Congenic lines were generated by the speed congenic method (Markel et al., 1997). Briefly, in
127 case of B6-genetic background, backcrossing was repeated up to 5 generations and homozygous
128 founders were established by brother-sister mating. Similarly, for the D2-genetic background,
129 backcrossing was repeated up to 7 generations and homozygous founders were established by brother-
130 sister mating. For the *Empsc1*, two congenic lines were generated according to the same method as for
131 the *Emcys1*-congenic lines, where *Empsc1* in Chr 1 was introgressed from B6 to D2-genetic
132 background and *vice versa* from D2 to B6-genetic background. For both genetic backgrounds (B6 and
133 D2), backcrossing was repeated up to 6 generations and homozygous founders were established by

brother-sister mating. Introgression of chromosomal regions was confirmed by genotyping of microsatellite markers shown in Figs. 1 and 5. Congenic lines were named according to the international nomenclature guidelines and abbreviated as in parenthesis; B6.D2-(*D6Mit188-D6Mit15*)/NSlcHkv (B6.D2-*Emcys1*), D2.B6-(*D6Mit188-D6Mit15*)/CrSlcHkv (D2.B6-*Emcys1*), B6.D2-(*D1Mit191-D1Mit291*)/NSlcHkv (B6.D2-*Empsc1*), and D2.B6-(*D1Mit191-D1Mit291*)/CrSlcHkv (D2.B6-*Empsc1*).

2.1.2. Subcongenic lines

Two subcongenic lines were developed from B6.D2-*Empsc1*. B6.D2-*Empsc1* was mated with B6 parental strain to generate heterozygous F₁ and then F₁ generations were backcrossed to B6 to establish heterozygous subcongenic lines. After that, each line was intercrossed (brother-sister mating) to get homozygous subcongenic lines. Another subcongenic line was developed from D2.B6-*Empsc1* according to the same method as subcongenic line from B6.D2-*Empsc1*. D2.B6-*Empsc1* was mated with D2 parental strain to generate F₁ heterozygous subcongenic mice and homozygous subcongenic line from D2.B6-*Empsc1* was established by brother-sister mating. Introgression of chromosomal regions was confirmed by genotyping of microsatellite markers as shown in Fig. 1. Subcongenic lines were named according to the international nomenclature guidelines and abbreviated as in parenthesis; B6.D2-(*D1Mit191-D1Mit290*)/NSlcHkv (B6.D2-*Empsc1.1*), B6.D2-(*D1Mit201-D1Mit291*)/NSlcHkv (B6.D2-*Empsc1.2*), and D2.B6-(*D1Mit191-D1Mit14*)/CrSlcHkv (D2.B6-*Empsc1.1*).

2.2 Microsatellite markers and genotyping

A total of 134 microsatellite markers (Supplementary Table 1) were selected from the database in the Mouse Genome Informatics (MGI), The Jackson laboratory, ME, USA (<http://www.informatics.jax.org/>) to generate congenic and subcongenic lines. Twenty-eight

161 microsatellite markers (Figs. 1 and 5) were used to confirm the introgressed regions for cyst
162 establishment and protoscolex development in congenic and subcongenic mice. Briefly, a piece of
163 ear-punched tissue was collected and genomic DNA was extracted by incubating samples at 54 °C for
164 3 h in 0.5 ml of lysis buffer [10 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10 mM
165 ethylenediaminetetraacetic acid (EDTA) and 1% sodium dodecyl sulfate (SDS)] with 5 µl of 10
166 mg/ml proteinase K. After genomic DNA extraction the touchdown PCR was performed with *Taq*
167 DNA polymerase (Ampliqon A/S, Odense M, Denmark) as follows; denaturing at 95 °C for 1 min,
168 followed by 10 cycles of denaturing at 95 °C for 30 sec, primer annealing at 65 °C for 30 sec (-2 °C in
169 2 cycles), and extension at 72 °C for 30 sec, and then, 35 cycles of denaturing at 95 °C for 30 sec,
170 primer annealing at 55 °C for 30 sec, and extension at 72 °C for 30 sec, and final extension at 72 °C
171 for 1 min, using a T100TM thermal cycler (Bio-Rad Laboratories Inc., Hercules, CA, USA). The
172 amplified samples were electrophoresed in 12% polyacrylamide gels in TBE buffer (89 mM Tris, 89
173 mM boric acid, and 2 mM EDTA), and stained with ethidium bromide for the visualization under a
174 UV light. The genotype was identified by the size of the PCR product.

175

176 2.3 Infection experiments and phenotype assessment

177

178 The infection experiments were performed according to Nakao et al., (2011). Briefly, *E.*
179 *multilocularis* eggs were collected from feces of the infected dogs. Experimental infection was
180 conducted in parental inbred strains (B6 and D2), congenic, and subcongenic mice by oral
181 administration of *E. multilocularis* eggs. For the assessment of cyst establishment, mice were infected
182 with 200 eggs, sacrificed, and necropsied at 4 weeks after infection. Mouse liver was cut into small
183 slices with approximately 0.5-mm thickness and the total number of cysts was counted in each liver.
184 For the assessment of protoscolex development in the cyst, mice were infected with 2,000 eggs,
185 sacrificed, and necropsied at 16 weeks after infection, because larger number of eggs and longer period
186 were needed to assess protoscolex development in the cyst. Livers were collected, parasitic cysts were
187 dissected from the liver, and total weight of the cysts was measured. A part of the cysts (1-2 g) was

188 minced, passed through a 125- μ m sieve, and washed repeatedly with saline. The number of mature
189 protoscoleces was counted under a stereoscopic microscope and expressed as number/gram of cysts.
190 Moreover, protoscoleces were counted in histopathological sections of liver to confirm the protoscolex
191 development in cysts.

192

193 *2.4. Histopathology*

194

195 Liver tissues were fixed in 10% formalin in phosphate-buffered saline (pH 7.4). The specimen was
196 dehydrated in consecutive stages with increasing concentrations of alcohol and cleaned by xylene.
197 Finally, liver samples were processed for paraffin embedding and cut into 2- μ m-thick sections. Tissue
198 sections were stained with haematoxylin and eosin for microscopic examinations with an All-in-One
199 Fluorescence Microscope BZ-X710 (Keyence, Osaka, Japan) to confirm protoscolex development in
200 the liver section.

201

202 *2.5. Statistical analysis*

203

204 All data were subjected to non-parametric test, Mann-Whitney *U*-test to compare between two
205 groups and one-way ANOVA with Bonferroni post-hoc analysis for more than two groups. Statistical
206 analyses were performed using a GraphPad Prism 5 software version 5 for windows (GraphPad
207 software Inc., San Diego, CA, USA). $P < 0.05$ was considered to be significant.

208

209 **3. Results**

210

211 *3.1 Confirmation of *Empsc1* using congenic mice*

212

213 In congenic mice, B6.D2-*Empsc1* and D2.B6-*Empsc1*, introgressed chromosomal region (*DIMit7*
214 to *DIMit511*) was confirmed by genotyping microsatellite markers (Fig. 1). This region covered

highly significant region of the QTL peak detected in the previous QTL analysis (Nakao et al., 2011). Other chromosomes except for Chr 1 were confirmed to be recipient-genetic background by genotyping microsatellite markers located in other chromosomes as shown in Supplementary Table 1. To confirm the *Empsc1* for protoscolex development, B6.D2-*Empsc1* and D2.B6-*Empsc1* mice were subjected to examination at 16 weeks after oral administration of 2,000 eggs of *E. multilocularis*. The number of mature protoscoleces in the cysts was calculated in B6.D2-*Empsc1* and D2.B6-*Empsc1* as well as B6 and D2 mice (parental strains). The number of protoscolex in the cyst of B6.D2-*Empsc1* was $28,285 \pm 15,819$, comparable to that of D2 ($28,255 \pm 5,124$), and significantly greater than that in B6 mice, which was completely absent (Fig. 2A). On the other hand, the number of protoscolex in the cyst of D2.B6-*Empsc1* mice was $5,128 \pm 2,902$ and significantly less than that of D2 control mice ($239,709 \pm 65,461$) (Fig. 2B). These results indicate that the *Empsc1* is definitely located between *D1Mit7* and *D1Mit511* and contributes to the difference in protoscolex development between B6 and D2 mice.

3.2 Narrowing the critical genetic region

To narrow down the length of critical region, three subcongenic lines were developed from congenic mice, B6.D2-*Empsc1* and D2.B6-*Empsc1*. Genotyping of microsatellite markers revealed that two lines of B6.D2-subcongenic mice, B6.D2-*Empsc1.1* and B6.D2-*Empsc1.2*, were introgressed D2-derived chromosomal region between *D1Mit7* and *D1Mit399* (38.6 cM and 70.3 cM) and *D1Mit496* and *D1Mit511* (63.1 cM and 97.3 cM), respectively (Fig.1). On the other hand, D2.B6-subcongenic line, D2.B6-*Empsc1.1*, was confirmed to be introgressed B6-derived chromosomal region between *D1Mit7* and *D1Mit290* (38.6 cM and 68.1 cM) in the D2-genetic background (Fig.1).

These three lines of subcongenic mice were challenged with *E. multilocularis* infection and protoscoleces in the cysts were counted under a light microscope. The number of protoscolex in the cyst of B6.D2-*Empsc1.2* and D2.B6-*Empsc1.1* were $22,700 \pm 7,651$ and $55,660 \pm 8,588$, respectively, indicating a significant development of protoscoleces as well as D2 mice ($104,367 \pm 48,349$), whereas

242 B6.D2-*Empsc1.1* and B6 mice did not develop protoscoleces (Fig. 3). The number of B6.D2-
243 *Empsc1.2* and D2.B6-*Empsc1.1* was significantly different, although protoscoleces were significantly
244 developed in these two subcongenic strains. This difference may be due to the difference in genetic
245 background. Thus, some genetic factors contributing to the protoscolex development may be present
246 in the D2 genetic background besides the introgressed regions. Alternatively, another genetic factor
247 affecting the protoscolex development may exist in the D2-derived region existing in B6.D2-
248 *Empsc1.2* but not in D2.B6-*Empsc1.1*, that is a region between *D1Mit496* and *D1Mit290*. The third
249 possibility is retaining some fragmental regions derived from the donor strain in congenic strains that
250 affect protoscolex development. Nonetheless, these results conclude that the *Empsc1* is located in the
251 critical region between *D1Mit290* and *D1Mit511* (68.1 cM and 97.3 cM) and is secured in the
252 introgressed region of B6.D2-*Empsc1.2* (Fig.1).

253

254 3.3 Confirmation of protoscolex in subcongenic mice with histopathological analysis

255

256 Protoscolex development in subcongenic mice was confirmed by histopathological analysis of liver
257 sections. Protoscoleces in the cyst of B6.D2-*Empsc1.2* were developed as well as in D2 mice, whereas
258 B6.D2-*Empsc1.1* mice did not develop any protoscolex as seen in B6 mice (Fig. 4A). The mean
259 number of protoscoleces in the cyst of D2 and B6.D2-*Empsc1.2* was 6.2 ± 1.1 and 7.8 ± 1.9 ,
260 respectively, and statistically different from that of B6 and B6.D2-*Empsc1.1* (Fig. 4B). These results
261 are consistent with results from the actual counting under a light microscope.

262

263 3.4 Congenic lines for cyst establishment

264

265 Congenic lines, B6.D2-*Emcys1* and D2.B6-*Emcys1*, for cyst establishment phenotype were
266 confirmed introgressed region by genotyping microsatellite markers (Fig. 5). Both congenic lines
267 were replaced chromosomal region between *D6Mit188* and *D6Mit15* with donor haplotype, in which
268 *Emcys1* locus could be included based on the QTL analysis data in the previous paper (Nakao et al.,

2011). Other chromosomes except for Chr 6 were confirmed to be recipient-genetic background by genotyping of microsatellite markers locating in other chromosomes as shown in Supplementary Table 1. These congenic mice were then challenged with *E. multilocularis* infection and investigated cyst establishment (Fig. 6). The number of cysts established in liver of B6.D2-*Emcys1* and D2.B6-*Emcys1* congenic mice was 7.5 ± 2.3 and 35.5 ± 5.9 , respectively, and was not different from that of each parental strain, B6 (13.2 ± 3.2) and D2 (41.2 ± 4.7), indicating that the replacement of *Emcys1* locus with the donor genotype did not alter the recipient phenotype. The reason for the ineffectiveness of the *Emcys1* introgression is unknown; however, our result indicates that the number of cysts is not controlled by the *Emcys1* alone.

4. Discussion

A previous investigation showed that susceptibility or resistance to *E. multilocularis* infection was genetically controlled and D2 mice were more susceptible to infection than B6 mice (Matsumoto et al., 2010). Further, another study identified a significant QTL, *Emcys1* in Chr 6 and a highly significant QTL, *Empsc1* in Chr 1 as responsible for cyst establishment and protoscolex development, respectively (Nakao et al., 2011).

In this study, congenic lines, B6.D2-*Empsc1* and D2.B6-*Empsc1*, were established from B6 and D2 parental strains and it was confirmed that the *Empsc1* is responsible for the difference in the protoscolex development. As 834 protein-coding genes are present in the introgressed chromosomal region between *DIMit7* and *DIMit511*, narrowing down the introgressed region could lead to the identification of candidate genes responsible for protoscolex development. Therefore, subcongenic lines, B6.D2-*Empsc1.1* and B6.D2-*Empsc1.2*, were generated from B6.D2-*Empsc1* and D2.B6-*Empsc1.1* line was generated from D2.B6-*Empsc1* (Fig.1). These subcongenic mice were challenged with *E. multilocularis* infection and we found protoscolex development in B6.D2-*Empsc1.2* mice, but not in B6.D2-*Empsc1.1*. On the other hand, D2.B6-*Empsc1.1* mice showed protoscolex development as seen in D2 mice. From these results, we could narrow down a critical region between *DIMit290*

296 and *DIMit511*, in which a gene responsible for protoscolex development must be located. Based on
297 MGI database, there are still 331 protein-coding genes located in the critical region, among which
298 several candidate genes such as those relating to the immune response or inflammation are listed in
299 Table 1. However, at this point it is difficult to address responsible gene(s) until functional
300 verification of gene(s) controlling the protoscolex development is carried out. Yang et al (2006)
301 reported that host genetic factors could contribute to the susceptibility of humans to *E. multilocularis*.
302 Moreover, some genes showed quantitative genetic variation in mice as well as in humans (Korstanje
303 et al. 2004; Hellebrandt et al., 2005). Many researchers have proposed that the interplay between
304 parasites and hosts has been developed by an evolutionarily conserved signaling system, specifically
305 the interaction between parasitic receptors and host-derived molecules (Gelmedin et al., 2008; Brehm,
306 2010). Our results from this study could facilitate the identification of candidate gene(s) involving this
307 signaling pathway for protoscolex development in both mouse and human, leading to the
308 implementation of the most effective control strategy for AE. Other researchers have successfully
309 addressed candidate genes responsible for some genetic phenotypes with the linkage analysis using
310 congenic and subcongenic lines (Pelletier et al., 2016; Kanagaratham et al., 2014) and gene
311 expression profiling in subcongenic mouse lines (Sander et al., 2007; Rennie et al., 2008; Ahn et al.,
312 2010; Stark et al., 2010). Therefore, we have started to produce backcrosses from B6.D2-*Empsc1.2* to
313 identify responsible gene(s) with genetic linkage analysis.

314 In case of cyst establishment phenotype, two congenic lines, B6.D2-*Emcys1* and D2.B6-*Emcys1*,
315 were generated from parental strains B6 and D2, respectively. The results from the current study
316 could not confirm that the genetic region between *D6Mit188* and *D6Mit15* was responsible for the
317 cyst establishment in the mouse liver (Fig. 6). This result suggests that other genetic factors may be
318 involved in the cyst establishment of *E. multilocularis* in the mouse liver.

319 In conclusion, a QTL *Empsc1* for protoscolex development of *E. multilocularis* was confirmed
320 by the generation of congenic lines. Moreover, we could narrow down the critical genetic region to
321 the length of 34.2 Mb between *DIMit290* and *DIMit511* and secured candidate gene(s) in the
322 introgressed region of B6.D2-*Empsc1.2*. These findings will lead to the identification of the candidate

323 gene(s) for protoscolex development of *E. multilocularis*. Identification of candidate gene(s) will
324 enrich the scientific knowledge of this zoonotic disease echinococcosis and would lead to the
325 development of new and effective drugs against echinococcosis.

326

327

328 **References**

329

- 330 Ahn, S.H., Deshmukh, H., Johnson, N., Cowell, L.G., Rude, T.H., Scott, W.K., Nelson, C.L., Zaas,
331 A.K., Marchuk, D.A., Keum, S., Lamlerthton, S., Sharma-Kuinkel, B.K., Sempowski, G.D.,
332 Fowler, V.G., 2010. Two genes on A/J chromosome 18 are associated with susceptibility to
333 *Staphylococcus aureus* infection by combined microarray and QTL analyses. PLoS Pathog. 6,
334 e1001088.
- 335 Adrianus, C.M., Jennifer, B., Anna, H., Jennifer, L., Malak, K., Sarah, R., David, F., Geoffrey, N., Lu,
336 L., Robert, W.W., Richard, J.W., 2009. Host genetic variation affects resistance to infection
337 with a highly pathogenic H5N1 influenza A virus in mice. J. Virol. 83, 10417-10426.
- 338 Allen, W., Cowley, J.R., 2006. The genetic dissection of essential hypertension. Nat. Rev. Genet. 7,
339 829-840.
- 340 Brehm, K., 2010. The role of evolutionarily conserved signaling systems in *Echinococcus*
341 *multilocularis* development and host-parasite interaction. Med. Microbiol. Immunol. 199,
342 247-259.
- 343 Craig, P.S., Rogan, M.T., Allan, J.C., 1996. Detection, screening and community epidemiology of
344 taeniid cestode zoonoses: cystic echinococcosis, alveolar echinococcosis and neurocysticercosis.
345 Adv. Parasitol. 38, 169-250.
- 346 David, G., Brownstein, Lisa, G., 1995. Chromosome mapping of *Rmp-4*, a gonad-dependent gene
347 encoding host resistance to mousepox. J. Virol. 69, 6958-6964.
- 348 Gelmedin, V., Caballero-Gamiz, R., Brehm, K., 2008. Characterization and inhibition of a p38-like
349 mitogen-activated protein kinase (MAPK) from *Echinococcus multilocularis*: antiparasitic
350 activities of p38 MAPK inhibitors. Biochem. Pharmacol. 76, 1068-1081.
- 351 Gottstein, B., Felleisen, R., 1995. Protective immune mechanisms against the metacestode of
352 *Echinococcus multilocularis*. Parasitol. Today 11, 320-326.
- 353 Hillebrandt, S., Wasmuth, H.E., Weiskirchen, R., Hellerbrand, C., Keppeler, H., Werth, A., Schirin-
354 Sokhan, R., Wilkens, G., Geier, A., Lorenzen, J., Köhl, J., Gressner, A.M., Matern, S., Lammert,

355 F., 2005. Complement factor 5 is a quantitative trait gene that modifies liver fibrogenesis in
356 mice and humans. *Nat. Genet.* 37, 835-843.

357 Hildreth, M.B., Granholm, N.H., 2003. Effect of mouse strain variations and cortisone treatment on
358 the establishment and growth of primary *Echinococcus multilocularis* hydatid cysts. *J. Parasitol.*
359 89, 493-495.

360 Kanagaratham, C., Marino, R., Camateros, P., Ren, J., Houle, D., 2014. Mapping of a chromosome 12
361 region associated with airway hyper responsiveness in a recombinant congenic mouse strain and
362 selection of potential candidate genes by expression and sequence variation analyses. *PLoS ONE*
363 9, e104234.

364 Kern, P., Bardonnet, K., Renner, E., Auer, H., Pawlowski, Z., Ammann, R.W., 2003. European
365 echinococcosis registry: human alveolar echinococcosis, Europe, 1982- 2000. *Emerg. Infect. Dis.*
366 9, 343-349.

367 Korstanje, R., Eriksson, P., Samnegård, A., Olsson, P.G., Forsman-Semb, K., Sen, S., Churchill,
368 G.A., Rollins, J., Harris, S., Hamsten, A., Paigen, B., 2004. Locating *Ath8*, a locus for murine
369 atherosclerosis susceptibility and testing several of its candidate genes in mice and humans.
370 *Atherosclerosis* 177, 443-450.

371 Matsumoto, J., Kouguchi, H., Oku, Y., Yagi, K., 2010. Primary alveolar echinococcosis: course of
372 larval development and antibody responses in intermediate host rodents with different genetic
373 backgrounds after oral infection with eggs of *Echinococcus multilocularis*. *Parasitol. Int.* 59,
374 435-444.

375 Markel, P., Shu, P., Ebeling, C., Carlson, G., Nagle, D., Smutko, J., 1997. Theoretical and empirical
376 issues for marker-assisted breeding of congenic mouse strains. *Nat. Genet.* 17, 280-284.

377 Marquis, J.F., Gros, P., 2008. Genetic analysis of resistance to infections in mice: A/J meets
378 C57BL/6J. *Curr. Top. Microbiol. Immunol.* 321, 27-57.

379 Marquis, J.F., Ronald, L., Lynn R., Robert, J. N., Philippe, G., 2009. Genetic and functional
380 characterization of the mouse *Trl3* locus in defense against Tuberculosis. *J. Immunol.* 182,
381 3757-3767.

382 Nakao, R., Kameda, Y., Kouguchi, H., Matsumoto, J., Dang, Z., Simon, A.Y., Torigoe, D., Sasaki, N.,
383 Oku, Y., Sugimoto, C., Agui, T., Yagi, K., 2011. Identification of genetic loci affecting the
384 establishment and development of *Echinococcus multilocularis* larvae in mice. *Int. J. Parasitol.*
385 41, 1121-1128.

386 Nakaya, K., Nakao, M., Ito, A., 1997. *Echinococcus multilocularis*: mouse strain difference in hydatid
387 development. *J. Helminthol.* 71, 53-56.

388 Pelletier, A.N., Guilbault, L., Guimont-Desrochers, F.G., Hillhouse, E.E., Lesage, S., 2016. NK cell
389 proportion and number are influenced by genetic loci on chromosomes 8, 9, and 17. *J. Immunol.*
390 196, 2627-2636.

391 Peters, L.L., Robledo, R.F., Bult, C.J., Churchill, G.A., Paigen, B.J., Svenson, K.L., 2007. The mouse
392 as a model for human biology: a resource guide for complex trait analysis. *Nat. Rev. Genet.* 8, 58-
393 69.

394 Rennie, C., Hulme, H., Fisher, P., Halp, L., Agaba, M., Noyes, H.A., Kemp, S.J., Brass, A., 2008. A
395 systematic, data-driven approach to the combined analysis of microarray and QTL data. *Dev. Biol.*
396 (Basel) 132, 293-299.

397 Reuter, S., Jensen, B., Buttenschoen, K., Kratzer, W., Kern P., 2000. Benzimidazoles in the treatment
398 of alveolar echinococcosis: a comparative study and review of the literature. *J. Antimicrob.*
399 *Chemother.* 59, 451-456.

400 Rogner, U.C., Avner, P., 2003. Congenic mice: cutting tools for complex immune disorders. *Nat. Rev.*
401 *Immunol.* 3, 243-252.

402 Sander, B., Rob, J., Vandebriel, Jeroen, L.A., Pennings, Eric, R., Gremmer, Piet, W. W., Henk, J. K.,
403 Timo, M.B., Peter, D., Frits, R.M., Barbara, H., Tjeerd, G. K., 2007. Comparative gene
404 expression profiling in two congenic mouse strains following *Bordetella pertussis* infection.
405 *BMC Microbiol.* 7, 1186/1471-2180-7-88.

406 Simon, A.Y., Moritoh, K., Torigoe, D., Asano, A., Sasaki, N., Agui, T., 2009. Multigenic control of
407 resistance to Sendai virus infection in mice. *Infect. Genet. Evol.* 9, 1253-1259.

408 Shimizu, V.S., Khan, R., Mostowy, S., Larivie`re, L., Wilkinson, R., Riendeau, N., Behr, M., Malo, D.,
 409 2007. Molecular genetic analysis of two loci (*Ity2* and *Ity3*) involved in the host response to
 410 infection with *Salmonella Typhimurium* using congenic mice and expression profiling. *Genetics*
 411 177, 1125-1139.

412 Spiliotis, M., Konrad, C., Gelmedin, V., Tappe, D., Brückner, S., Mösch, H.U., Brehm, K., 2006.
 413 Characterisation of EmMPK1, an ERK-like MAP kinase from *Echinococcus multilocularis* which
 414 is activated in response to human epidermal growth factor. *Int. J. Parasitol.* 36, 1097-1112.

415 Stark, J.M., Barmada, M.M., Winterberg, A.V., Majumber, N., Gibbons, W.J., Stark, M.A., Sartor,
 416 M.A., Medvedovic, M., Kolls, J., Bein, K., Mailaparambil, B., Krueger, M., Heinzmann, A.,
 417 Leikauf, G.D., Prows, D.R., 2010. Genomewide association analysis of respiratory syncytial
 418 virus infection in mice. *J. Virol.* 84, 2257-2269.

419 Vuitton, D.A., 2003. The ambiguous role of immunity in echinococcosis: protection of the host or of
 420 the parasite? *Acta Trop.* 85, 119-132.

421 Vuitton, D.A., Gottstein, B., 2010. *Echinococcus multilocularis* and its intermediate
 422 host: A model of parasite-host interplay. *J. Biomed. Biotechnol.* 2010, 923193.

423 Yang, Y.R., Ellis, M., Sun, T., Li, Z., Liu, X., Vuitton, D.A., Bartholomot, B., Giraudoux, P., Craig,
 424 P.S., Boufana, B., Wang, Y., Feng, X., Wen, H., Ito, A., McManus, D.P., 2006. Unique family
 425 clustering of human echinococcosis cases in a Chinese community. *Am. J. Trop. Med. Hyg.* 74,
 426 487-494.

427

428 **Legends to figures**

429

430 **Fig. 1.** Schematic diagram of Chr1 in congenic and subcongenic mice for *Empsc1*. Shaded, open, and
431 black columns indicate B6-derived, D2-derived, and unknown chromosomal portions, respectively.

432

433 **Fig. 2.** Protoscolex development in the liver of B6, D2, and congenic mice for *Empsc1*. The number
434 of protoscolex in cystic lesions of the liver was counted and expressed as number of protoscolex per 1
435 gram of cystic lesion. (A) Data represent the mean $\pm$ SEM for B6 (n=4), D2 (n=4), and B6.D2-
436 *Empsc1* (n=5). (B) Data represent the mean $\pm$ SEM for D2 (n=9) and D2.B6-*Empsc1* (n=3). Data are
437 representative of two independent experiments. ** and * indicate $P < 0.01$ and $P < 0.05$, respectively.

438

439 **Fig. 3.** Protoscolex development in the liver of B6, D2, and subcongenic mice for *Empsc1*. The
440 number of protoscolex in cystic lesions of the liver was counted and expressed as number of
441 protoscolex per 1 gram of cystic lesion. Data represent the mean $\pm$ SEM for B6 (n=3), D2 (n=3) and
442 subcongenic mice for *Empsc1*, B6.D2-*Empsc1.1* (n=3), B6.D2-*Empsc1.2* (n=3), and D2.B6-*Empsc1.1*
443 (n=5). * indicates $P < 0.05$.

444

445 **Fig. 4.** (A) Photographs of liver sections in B6, D2, and subcongenic mice for *Empsc1*. Arrows
446 indicate protoscolecces. (B) Protoscolex counts in liver sections in B6 (n=6), D2 (n=6), B6.D2-
447 *Empsc1.1* (n=6), and B6.D2-*Empsc1.2* (n=8). The number of protoscolex was counted in 10 cysts
448 selected randomly in each mouse liver and the mean value was calculated in each mouse. Data
449 represent the mean $\pm$ SEM. ** and * indicate $P < 0.01$ and $P < 0.05$, respectively.

450

451 **Fig. 5.** Schematic diagram of Chr 6 in congenic mice for *Emcys1*; B6.D2- *Emcys1* and D2.B6- *Emcys1*.
452 Shaded, open, and black columns indicate B6-derived, D2-derived, and unknown chromosomal
453 portions, respectively.

454

455 **Fig. 6.** Cyst establishment in the liver of B6 (n=6), D2 (n=6), and congenic mice for *Emcys1*; B6.D2-
456 *Emcys1* (n=5) and D2.B6- *Emcys1* (n=6). The data represent mean $\pm$ SEM. Data are representative of
457 two independent experiments. ** indicates $P < 0.01$.