HOKKAIDO UNIVERSITY

Title	Serodiagnosis of elephant tuberculosis: a useful tool for early identification of infected elephants at the captive-wild interface
Author(s)	Paudel, Sarad; Mikota, Susan K.; Thapa, Jeewan et al.
Citation	European journal of wildlife research, 64(6), 70 https://doi.org/10.1007/s10344-018-1229-3
Issue Date	2018-12
Doc URL	https://hdl.handle.net/2115/76245
Type	journal article
File Information	European journal of wildlife research64(6)_70-.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Title: Serodiagnosis of elephant tuberculosis: A useful tool for early identification of infected elephants at the captive-wild interface

Authors: Sarad Paudel[1, 2]*, Susan K. Mikota[3]*, Jeewan Thapa[4]*, Konstantin P. Lyashchenko[5], Kamal Prasad Gairhe[6], Ishwari Prasad Dhakal[7], Naresh Subedi[8], Bhagwan Maharjan[9], Suraj Subedi[8], Gretchen E. Kaufman[10], Toshio Tsubota[1]

Author's affiliation

[1] Laboratory of Wildlife Biology and Medicine
Graduate School of Veterinary Medicine
Hokkaido University
Kita 18, Nishi 9, Kita-ku,
Sapporo 060-0818, Japan

[2] Current Address
Department of Cell Physiology
Faculty of Medicine and Graduate School of Medicine
Hokkaido University
Kita 15, Nishi 7, Kita-ku
Sapporo, Hokkaido, 060-8638, Japan

[3] Elephant Care International
166 Limo View Lane
Hohenwald TN 38462, USA

[4] Hokkaido University Research Center for Zoonosis Control
Kita 20, Nishi 10, Kita-ku, Sapporo 001-0020, Japan

[5] Chembio Diagnostic Systems, Inc.
3661 Horseblock Road
Medford, NY 11763, USA

[6] Department of National Parks and Wildlife Conservation
Chitwan National Park, Chitwan, Nepal

[7] Agriculture and Forestry University
Rampur, Chitwan, Nepal

[8] National Trust for Nature Conservation
Khumaltar, Lalitpur, Nepal

[9] German Nepal Tuberculosis Project (GENETUP), Kalimati, Kathmandu, Nepal

[10] Veterinary Initiative for Endangered Wildlife
1627 W. Main St., Suite 445, Bozeman, MT USA 59715

*Sarad Paudel, Susan K. Mikota and Jeewan Thapa contributed equally to this work

Corresponding author:

Toshio Tsubota
Laboratory of Wildlife Biology and Medicine
Graduate School of Veterinary Medicine
Hokkaido University
Kita 18, Nishi 9, Kita-ku,
Sapporo 060-0818, Japan
Phone: +81-11-706-5101; Fax: +81-11-706-5569
Email: tsubota@vetmed.hokudai.ac.jp

Abstract Tuberculosis (TB) is an emerging disease in elephants primarily caused by *Mycobacterium tuberculosis* (*M. tb)* and in some occassion by *M. bovis*. We performed culture and three serological tests –the Elephant TB STAT-PAK,® DPP VetTB® Assay, and MAPIA (multi-antigen print immunoassay)–prospectively on samples from eight elephants in Nepal that died of suspected or confirmed tuberculosis (TB) between 2007 and 2013. Among them, all elephants were reactive to DPP VetTB® Assay, five to Elephant TB STAT-PAK,® and two were reactive to MAPIA. Similarly, six elephants were positive on culture on samples collected antemortem or post-mortem. We observed antibody responses months to years before culture confirmation of TB which shows that serological tests can be highly useful for the early diagnosis of TB in elephants. Validated point-of-care serological tests are easily performed in the field and hold promise for improved TB surveillance in other non-domestic species.

Key words Asian elephants, tuberculosis, DPP VetTB Assay, ElephantTB STAT-PAK, serological assay.

Introduction

Tuberculosis (TB) is an emerging disease in elephants caused primarily by *Mycobacterium tuberculosis* (*M. tb).* Numerous cases have occurred among captive elephants, but it is only since 2013 that *M. tb* has been found in wild elephants (Obanda et al. 2013; Perera et al. 2014; Zachariah et al. 2017; Chandranaik et al. 2017). The occurrence of this human pathogen in the wild is cause for concern and further research.

In captive elephants, TB is diagnosed by isolating *M. tb* from respiratory samples collected using a trunk wash procedure (Lyashchenko et al. 2006). This method has been widely used in the U.S. and Europe. In elephant range countries, *M. tb* has been cultured from captive elephants in India (Venugopal and Abraham 2015), Nepal (Paudel et al. 2014), Thailand (Angkawanish et al. 2010), and from an elephant in Australia that was imported from Thailand (Stephens et al. 2013). Although positive culture provides a definitive diagnosis, the method has significant limitations (Lyashchenko et al. 2006; Mikota et al. 2015). In many cases, repeated ante-mortem culture fails to identify infected elephants that are then diagnosed postmortem. Supplementary Table 1 lists publications that rreport the poor recovery of *M. tb* from multiple trunk wash samples from known TB-infected elephants.

Specific serological point-of-care tests offer a practical TB screening method in elephants and other wildlife species. The ElephantTB STAT-PAK® (STAT-PAK) and Dual Path Platform (DPP) VetTB® assays (Chembio Diagnostic Systems, Inc., Medford NY, USA), licensed by the United States Department of Agriculture, have been used for TB screening in several Asian elephant range countries (Abraham et al. 2008; Mar et al. 2012; Ong et al. 2013; Lassausaie et al. 2015). The more accurate DPP replaced the STAT-PAK in 2012.

105 TB was first diagnosed in elephants in Nepal in 2002 and between 2002 and 2015, 13
106 elephants died of suspected or confirmed TB. The Nepal Elephant TB Control and Management
107 Action Plan was initiated in 2011 to mitigate the transmission of TB between free-ranging
108 wildlife such as rhinoceros and wild elephants and captive elephants used for tourism, patrolling
109 of protected areas, and research (Mikota et al. 2016). This Plan was approved by the Department
110 of National Parks and Wildlife Conservation, the Ministry of Forestry and provides guidelines
111 for routine testing, treatment, and reintegration of elephants into active service following
112 treatment when possible.

113 We present the clinical, serological, and culture data from eight cases and show that
114 specific serological tests are a useful tool to manage TB in captive elephants and decrease TB
115 risk to wild populations.

116 **Materials and Methods**

117 **Study subjects and sample collection**

118 Four male and four female captive elephants, aged 31–70 years were included in the
119 study (Supplementary Table 2). Seven dead elephants were owned by the government and one
120 elephant was privately owned. Health care and clinical examination was provided by Elephant
121 TB program veterinarians, supervised by the government's Senior Wildlife Veterinarian. All
122 animals died from suspected TB based on necropsy lesions and none was euthanized.

123 Pre-mortem blood was collected from the auricular vein of seven elephants. Twenty-one
124 trunk wash samples for culture were collected at various times ante-mortem from six elephants
125 according to published guidelines (USAHA 2010) or using a modified procedure (Abraham and

126 Davis 2008). In one case (E8), an excreted respiratory discharge sample was submitted for
127 culture.

128 **Serology testing**

129 The multi-antigen print immuno assay (MAPIA) was originally developed as a tool to
130 identify seroreactive antigens in human TB and has been adapted to identify the
131 immunodominant proteins of *M. tb* or *M. bovis* recognized in elephants and other species
132 (Lyashchenko et al. 2006). The selected antigens were used for the development of the
133 serological tests described below. In this study, the MAPIA was performed at the Chembio
134 laboratory on serum samples collected in 2006 from six elephants (E1, E2, E3, E6, E7 and E8)
135 as previously described (Lyashchenko et al. 2006).

136 The STAT-PAK is a lateral flow screening test that incorporates a cocktail of several *M.*
137 *tb* and/or *M. bovis* antigens impregnated on nitrocellulose membrane housed in a disposable
138 plastic cassette. Sample and buffer solution migrate to a test pad by capillary action. If
139 antibodies are present, they bind to the antigen and the colored latex bead-based signal detection
140 system results in a visible blue line in the test window (Lyashchenko et al. 2006).

141 The DPPVetTB Assay is an immune-chromatographic screening test that detects
142 antibodies to ESAT-6/CFP 10 and MPB83 antigens. The DPP technology uses two
143 nitrocellulose strips allowing independent delivery of the test sample and antibody-detecting
144 agent of colloidal gold particles (Greenwald et al. 2009). The ElephantTB STAT-PAK® and
145 DPP VetTB® assays (Chembio Diagnostic Systems, Inc., Medford, NY) were performed
146 according to the manufacturer's instructions using blood collected ante-mortem or banked serum.
147 In two cases (E5 and E7) the DPP was performed using frozen-thawed post-mortem lung fluid.

148 **Postmortem examination**

149 Post-mortem examinations were performed on seven elephants. Elephant E4 was
150 returned to her owner in India where she died after several months; a postmortem examination
151 was not performed. All staff used personal protective equipment including N-95 masks during
152 necropsy. The necropsy was performed using procedures recommended by Montali (2006).
153 Representative lung lesions were collected in sterile tubes for culture. Tissue samples from
154 elephant E1 were submitted to the Central Veterinary Laboratory in Kathmandu, Nepal for
155 histopathology.

156 **Culture of trunk discharges and lung tissues**

157 Cultures of lung tissue samples were conducted following the guidelines of European
158 Society for Mycobacteriology (Groothius and Yates 1991) and Lowenstein- Jensen (L-J) media
159 was used to culture the trunk wash samples and lung lesions. Decontamination procedure was
160 used according to (USAHA, 2010).

161 **Results**

162 The relationship between serology and culture results of eight elephants to time of death
163 is illustrated in Figure 1. Clinical, diagnostic, postmortem, and culture data of eight TB-suspect
164 elephants in Nepal is presented in Supplementary Table 2. Five of eight elephants showed
165 clinical signs of TB including coughing, trunk discharge, or weight loss.

166 **Serological testing**

167 Two of six elephants tested by MAPIA in 2006 were seropositive (E2 and E3). E3 was
168 seropositive more than three years prior to diagnosis by culture. Elephants E6, E7, and E8 were
169 non-reactive on MAPIA in 2006, but showed positive reactivity on the DPP test and had

evidence of TB at postmortem examination 6-7 years later, suggesting that they were infected and seroconverted after 2006.

Between 2006 and 2012, seven elephants were tested using the STAT-PAK. Elephants E1, E2, E3 and E4 were seropositive, elephant E7 had equivocal results, and elephants E6 and E8 were seronegative.

All eight elephants reacted on the DPP VetTB assay at some point between 2006 and 2013, including two elephants that were tested using postmortem lung fluid. This is an off-label use of the test but shows that TB antibodies are present in other body fluids.

Postmortem

All seven elephants that underwent postmortem examination had gross lung lesions compatible with TB. Histopathology performed on tissues from E1 showed acid-fast bacilli in the lung, scattered granulomatous foci, and a chronic inflammatory response.

Culture

Twenty-one trunk wash samples collected antemortem from six elephants were negative for *M. tb*. Five lung tissue samples and one pre-mortem trunk wash sample from six elephants (E3-E8) were culture positive for *M. tb*. Ante-mortem trunk respiratory discharge from E4 was positive on culture shortly before her death. Elephant E1 was negative on ante- and postmortem cultures despite gross postmortem findings compatible with TB. Elephant E2 also had distinct TB-like lung gross lesions at postmortem examination; however, samples were not submitted for culture.

Discussion

In our study, TB was confirmed at the time of death by culture in six cases. Culture was negative in the case of E1, however, samples may have been compromised during shipment because of high ambient temperatures. Samples for culture were transported to the TB laboratory within 24 h of collection. Culture was also negative for TB in all cases where it was performed at times significantly preceding death and consequently was not a predictor of disease.

By contrast, Elephants E3 and E4 were reactive on the STAT-PAK 44 and 28 months respectively before diagnosis by culture, supporting previous findings that elephants can be seropositive months to years in advance of TB diagnosis by culture (Lyashchenko et al. 2012). We did not detect seroconversion in E6 and E7 until shortly before they died of TB.

We observed a strong correlation between serology and post-mortem confirmation by culture and poor correlation between antemortem trunk wash culture and post-mortem diagnosis. In our study, 21 trunk wash samples collected at various times from six elephants were negative for mycobacteria; only the expectorated discharge from E4 was positive shortly before her death. All the elephants that were culture positive postmortem were reactive on one or more serological tests but antemortem trunk wash samples were never culture positive (Figure 1).

Other studies have reported similar findings (Supplementary Table 1). In Thailand, *M. tb* was isolated from only two of 60 trunk wash samples from three elephants that were culture positive post-mortem (Angkawanish et al. 2010). In Sweden, only 7 of 189 trunk wash sample were positive from 5 elephants confirmed TB-infected at postmortem (Moller et al.2005). Although trunk wash culture is considered the “gold standard” for the diagnosis of TB in elephants, these examples clearly show the deficiencies of culture for surveillance and intervention.

There are 109 to 142 resident wild elephants and several elephant herds that migrate between Nepal and India (Pradhan et al. 2011). Captive and wild elephants interact during natural breeding and intermingle with other wildlife during grazing in protected areas. Captive elephants have close contact with wild rhinos during tourist safaris, patrolling, translocations, and during the annual rhino count which is conducted on elephant-back. This provides ample opportunities for TB to be transmitted to endangered wildlife and the potential for the development of a wild reservoir of *M. tb*.

While *M. bovis* is a known threat to a variety of species in the wild such as Eurasian badgers (*Meles meles*) (Drewe et al. 2010), and cape buffalo (*Syncerus caffer*) where in Kruger National Park TB has spilled over to other mammalian species, including lions (Miller et al. 2012), it is only since 2013 that *M. tb* has been found in any wild species. A wild African elephant that had been under human care was the first reported case although it is not clear whether the mycobacterial species was confirmed (Obanda et al. 2013). Since then, four cases with no history of captivity have been reported in wild Asian elephants in India (Zachariah et al. 2017; Chandranaik et al. 2017) and one additional case in Sri Lanka (Perera et al. 2014).

When successful, culture and isolation of *M. tb* is valuable to confirm infection, identify the mycobacterial species and strain, and test it for drug susceptibility (Mikota 2008). But in developing countries culture method is not practical, nor reliable, as a primary technique for TB screening.

The serological tests have been shown to be early and accurate correlatives of active TB in elephants (Lyashchenko et al. 2006, 2012). Prompt diagnosis of TB at the captive-wild interface facilitates initiation of appropriate management strategies to prevent transmission to herd mates, wild elephants, rhinos, and other susceptible species, including humans. These tests have been instrumental in the success of the Nepal Elephant Tuberculosis Control and

Management Action Plan. Through on-going surveillance and treatment, no deaths related to TB were reported between 2013 and 2018.

The value of these serological tests is not limited to elephants although their use in most of other wildlife species is off-label. The DPP VetTB Assay has been used in rhinos (Duncan et al. 2009; Miller et al. 2015), lions (Miller et al. 2012), warthogs (Miller et al. 2016), badgers (Drewe et al. 2010), and wood bison (Himsworth et al. 2010). In the United States, the DPP VetTB Assay is approved by the USDA Bovine TB Eradication Program for testing several species of captive cervids (Lyashchenko et al. 2013, 2018). This study shows serology as promising "non-culture" based TB diagnosis method in elephants, however, more samples size are required to determine a statistical correlation between serology and culture in elephants.

Acknowledgments We acknowledge Chitra Bahadur Khadka, Purushottam Pandey and Kiran Rijal at Chitwan National Park, Nepal for helping with blood collection and sample collection during the necropsy. We highly acknowledge the Department of National Parks and Wildlife Conservation (DNPWC), Government of Nepal, National Trust of Nature Conservation (NTNC), Nepal and WWF-Nepal for their kind support in this study. We thank Christy Williams, Asian Rhino and Elephant Action Strategy (AREAS), WWF-Nepal for providing the funding support to the elephant TB study in Nepal.

References

Abraham D, Cheeran JV, Sukumar R, Mikota SK, Rao S, Ganguly S, Varma S (2008) Health assessment of captive Asian elephants in India with special reference to tuberculosis.

261 Report to Project Elephant, Ministry of Environment and Forests, Government of India,
262 New Delhi

263 Abraham D, Davis J (2008) Revised trunk wash collection procedure for captive elephants in a
264 range country setting. Gajah 28:53-54

265 Angkawanish T, Wajjwalku W, Sirimalaisuwan A, Mahasawangkul S, Kaewsakhorn T, Boonsri
266 K, Rutten VP (2010) *Mycobacterium tuberculosis* infection of domesticated Asian
267 elephants, Thailand. Emerg Infect Dis16:1949-1951

268 Chandranaik BM, Shivashankar BP, Umashankar KS, Nandini P, Giridhar P,Byregowda SM,
269 Shrinivasa BM (2017). *Mycobacterium tuberculosis* infection in free-roaming wild Asian
270 elephant. Emerg Infect Dis 23:555-557

271 Drewe JA, Tomlinson AJ, Walker NJ, Delahay RJ (2010) Diagnostic accuracy and optimal use
272 of three tests for tuberculosis in live badgers. PLoS ONE 5(6):e11196.
273 doi:10.1371/journal.pone.0011196

274 Duncan AE, Lyashchenko K, Greenwald R, Miller M, Ball R (2009) Application of elephant TB
275 STAT-PAK assay and MAPIA (multi-antigen print immunoassay) for detection of
276 tuberculosis and monitoring of treatment in black rhinoceros (*Diceros bicornis*). J Zoo
277 Wildl Med 40(4):781-785

278 Greenwald R, Lyashchenko O, Esfandiari J, Miller M, Mikota S, Olsen JH, Ball R, Dumonceaux
279 G, Schmitt D, Moller T, Payeur JB, Harris B, Sofranko D, Waters WR, Lyashchenko KP
280 (2009) Highly accurate antibody assays for early and rapid detection of tuberculosis in
281 African and Asian elephants. Clin Vaccine Immunol 16:605-612

282 Groothuis DG, Yates MD (1991) Decontamination, microscopy and isolation. In: Groothius D G,
283 Yates M D, editors. Diagnostic and Public Health Mycobacteriolgy. 2nd ed. London:
284 Bureau of Hygiene and Tropical Diseases, European Society for Mycobacteriology; p. 63

285 Himsworth CG, Elkin BT, Nishi JS, Epp T, Lyashchenko KP, Surujballi O, Turcotte C,
286 Esfandiari J, Greenwald R, Leighton FA (2010) Comparison of test performance and
287 evaluation of novel immunoassays for tuberculosis in a captive herd of wood bison
288 naturally infected with *Mycobacterium bovis*. J Wild Dis 46(1):78-86

289 Lassausaie J, Bret A, Bouapao X, Chanthavong V, Castonguay-Vanier J, Quet F, Mikota SK,
290 Théorêt C, Buisson Y, Bouchard B (2015) Tuberculosis in Laos, who is at risk: the
291 mahouts or their elephants? Epidemiol Infect 143:922-931

292 Lyashchenko KP, Greenwald R, Esfandiari J, Olsen JH, Ball R, Dumonceaux G, Dunker F,
293 Buckley C, Richard M, Murray S, Payeur JB, Andersen P, Pollock JM, Mikota S, Miller
294 M, Sofranko D, Waters WR (2006) Tuberculosis in elephants: antibody responses to
295 defined antigens of *Mycobacterium tuberculosis*, potential for early diagnosis, and
296 monitoring of treatment. Clin Vaccine Immunol 13:722-732

297 Lyashchenko KP, Greenwald R, Esfandiari J, Mikota S, Miller M, Moller T, Volgelnest L,
298 Gairhe K, Robbe-Austerman S, Gai J, Waters WR (2012) Field application of
299 serodiagnostics to identify elephants with tuberculosis prior to case confirmation by
300 culture. Clin Vaccine Immunol 19:1269-1275

301 Lyashchenko KP, Greenwald R, Esfandiari J, O'Brien DJ, Schmitt SM, Palmer MV, Waters WR
302 (2013) Rapid detection of serum antibody by dual-path platform VetTB assay in white-
303 tailed deer infected with *Mycobacterium bovis*. ClinVaccine Immunol 20: 907-911

304 Lyashchenko KP, Gortazar C, Miller MA, Waters WR (2018) Spectrum of antibody profiles in
305 tuberculous elephants, cervids, and cattle. Vet Microbiol 214:89-92

306 Mar KU, Htut W, Zar MN, Soe AT, Mikota SK (2012) Report to Myanma Timber Enterprise on
307 Tuberculosis Testing of MTE Elephants

308 Mikota SK (2008) Tuberculosis in elephants. In: Fowler ME, Miller RE (eds) Zoo and Wild
309 Animal Medicine, Current Therapy, 6th Ed. Saunders/Elsevier, St. Louis, pp. 355-364

310 Mikota SK, Lyashchenko KP, Lowenstine L, Agnew D, Maslow JN (2015). Mycobacterial
311 infections in elephants. In: Mukundan H, Chambers MA, Waters WR Larsen M (eds)
312 Many hosts of mycobacteria, tuberculosis, leprosy, and other mycobacterial diseases of
313 man and animals. CABI Publishing House, Wallingford, U.K. pp 259-276

314 Mikota SK, Kaufman GE, Subedi N, Dhakal IP (2016) *Mycobacterium tuberculosis* in elephants
315 in Asia - taking a one health approach. In: Cork S, Hall DC, Liljebjelke (eds) One health
316 case studies: addressing complex problems in a changing world. 5M Publishing Ltd,
317 Sheffield, UK, pp 54-64

318 Miller M, Joubert J, Mathebula N, de-Klerk-Lorist LM, Lyashchenko KP, Bengis R, van Helden
319 P, Hofmeyr M, Olea-Popelka F, Greenwald R, Esfandiari J, Buss P (2012) J Zoo Wild
320 Med 43(2):317-323

321 Miller MA, Greenwald R, Lyashchenko KP (2015) Potential for serodiagnosis of tuberculosis in
322 black rhinoceros (*Diceros bicornis*). J Zoo Wild Med 46(1):100-104

323 Miller M, Buss P, de-Klerk-Lorist LM, Hofmeyr J, Hausler G, Lyashchenko K, Lane EP, Botha
324 L, Parsons S, van Helden P (2016) Application of rapid serologic tests for detection of

325 *Mycobacterium bovis* infection in free-ranging warthogs (*Phacochoerus africanus*) —
326 implications for antemortem disease screening. J Wild Dis 52(1):180-182

327 Moller T, Roken B, Petersson L, Vitaud C, Lyashchenko K (2005) Preliminary results of a new
328 serological test for detection of TB-infection (*Mycobacterium tuberculosis*) in elephants
329 (*Elaphas maximus* and *Loxodonta africanum*) Swedish case studies. Verh ber Erkg
330 Zootiere 42:173e81

331 Montali RJ (2006) Postmortem diagnostics. In: Fowler M, Mikota S K, editors. Biology,
332 Medicine, and Surgery of elephants. Ames, IA: Blackwell Publishing; Pp. 199- 209

333 Obanda VJ, Poghon M, Yongo M, Mulei I, Ngotho K, Waititu K, Makumi J, Gakuya F,
334 Osmondi P, Soriguer RC, Alasaad S (2013). First reported case of fatal tuberculosis in a
335 wild African elephant with past human-wildlife contact. Epidemiol Infect 141:1476-1480

336 Ong BL, Ngeow YF, Razak MF, Yakubu Y, Zakaria Z, Mutalib AR, Hassan L, Ng HF,
337 Verasahib K (2013) Tuberculosis in captive Asian elephants (*Elephas maximus*) in
338 Peninsular Malaysia. Epidemiol Infect 141:1481-1487

339 Paudel S, Mikota SK, Nakajima C, Gairhe KP, Maharjan B, Thapa J, Poudel A, Shimozuru M,
340 Suzuki Y, Tsubota T (2014) Molecular characterization of *Mycobacterium tuberculosis*
341 isolates from elephants of Nepal. Tuberculosis 94:287-292

342 Perera BVP, Salgadu MA, Gunawardena GSPS, Smith NH, Jinadasa HRN (2014) First
343 confirmed case of fatal tuberculosis in a wild Sri Lankan elephant. Gajah 41:28-31

344 Pradhan NMB, Williams CA, Dhakal M (2011) Current status of Asian elephants in Nepal.
345 Gajah 35:87-92

346 Steinmetz H, Rutten M (2016) TB or Not TB: Diagnosis of tuberculosis in a group of Asian
347 elephants (*Elephas maximus*). Proc American Assoc Zoo Veterinarians, European Assoc
348 Zoo and Wildl Veterinarians, Institute for Zoo and Wild Research, Joint Conference
349 p116

350 Stephens N, Vogelnest L, Lowbridge C, Christensen A, Marks GB, Sintchenko V, McAnulty J
351 (2013). Transmission of *Mycobacterium tuberculosis* from an Asian elephant *(Elephas*
352 *maximus*) to a chimpanzee (*Pan troglodytes*) and humans in an Australian zoo.
353 Epidemiol Infect 141:1488-1497

354 USAHA Proceedings (2010) Guidelines for the control of tuberculosis in elephants. Proceedings
355 of 114th Annual Meeting of the United States Animal Health Association. 114: 578-639
356 Venugopal, K.P. and Abraham, D. (2015) Zoonotic tuberculosis: the case of mahouts and
357 captive elephants in southern India, Poster Presentation, The 46th Union World Congress
358 on Lung Health, International Union Against Tuberculosis and Lung Diseases, 2-6
359 December, Cape Town, South Africa.
360 Vogelnest L, Hulst F, Thompson P, Lyashchenko KP, Herrin KA (2015) Diagnosis and
361 management of tuberculosis (*Mycobacterium tuberculosis*) in an Asian elephant (*Elephas*
362 *maximus*) with a newborn calf. J Zoo Wildl Med 46(1): 77-85
363 Zachariah A, Pandiyan J, Madhavilatha GK, Mundayoor S, Chandramohan B, Sajesh PK,
364 Santhosh S, Mikota SK (2017). *Mycobacterium tuberculosis* in Wild Asian Elephants,
365 Southern India. Emerg Infect Dis 23:504-506

366

367

368

Fig 1 Relationship of serology and culture results to time of death

Footnote: serology shown includes results of one or more tests during the time period indicated. The MAPIA was only run in 2006.

Title: Serodiagnosis of elephant tuberculosis: A useful tool for early identification of infected elephants at the captive-wild interface

Table 1. Reports Demonstrating the Poor Recovery of TB from Multiple Trunk Wash Samples From Known TB-Infected Elephants

Country	Number of elephants with confirmed TB	Number of trunk washes performed	Number of TB-positive cultures	Reference
Sweden	5	189	7	Moller et al. 2005
Thailand	3	60	2	Angkawanish at al. 2010
Australia	1	371 (includes samples collected from 6 elephants total)	1	Vogelnest et al. 2015
Switzerland	3	> 200 (includes samples collected from 10 elephants total)	0 (3 elephants were euthanized and TB confirmed postmortem)	Steinmetz and Rutten 2016

Table 2. Clinical serological and culture data of eight elephants in Nepal

Elephant	Gender	Age (Years)	Clinical Signs	MAPIA Results	STAT-PAK Results	DPP Results	AM culture results	PM culture results	Date of Death	Comments
E1	M	47	weight loss; reflux	NR 2006	R 2006	R 2006 (banked serum)	N 2006 (n=3)	Negative	Aug 2007	Suspect culture samples damaged during transport
E2	M	32	weight loss; coughing	R 2006	R 2006	R 2006 (banked serum)	N 2006 (n=3)	NA	Sept 2007	Samples for culture not submitted
E3	F	65	weight loss	R 2006	R 2006, 2008	R 2009	N 2006 (n=2) N 2008 (n=1)	*M. tb*	Aug 2009	Too old to treat
E4	F	56	coughing, weakness, trunk discharge	NA	R 2008, 2010	R 2010	P 2010 (n=1) *M.tb*	NA	2010; month unknown	*M.tb* isolated from trunk discharge; no PM or PM culture performed
E5	F	60	none	NA	NA	R 2009 (lung fluid)	NA	*M. tb*	Sept 2009	AM blood not available
E6	M	31	none	NR 2006	NR 2006, 2008, 2010, 2011	R 2012	N 2006 (n=3) N 2010 (n=1)	*M. tb*	Sept 2012	Extensive TB lesions; late to sero-convert compared to other cases
E7	F	65 or 70	loss of muscle mass noted at necropsy	NR 2006	NR 2006, 2008, 2010; equivocal 2011	NR 2011; R 2013 (lung fluid)	N 2006 (n=3) *M. terrae* isolated from one sample	*M. tb*	Feb 2013	Extensive TB lesions; late to sero-convert compared to other cases
E8	M	32	weight loss	NR 2006	NR 2006, 2007, 2008	R 2013	N 2006 (n=3) N 2013 (n=1)	*M. tb*	March 2013	Five year serology testing gap

R: reactive ; NR: non-reactive ; N: negative ; P: positive ; NA: not applicable ; AM: Ante-mortem ; PM : Post-mortem


References:

Angkawanish T, Wajjwalku W, Sirimalaisuwan A, Mahasawangkul S, Kaewsakhorn T, Boonsri K, Rutten VP (2010) *Mycobacterium tuberculosis* infection of domesticated Asian elephants, Thailand. Emerg Infect Dis16:1949-1951

Moller T, Roken B, Petersson L, Vitaud C, Lyashchenko K (2005) Preliminary results of a new serological test for detection of TB-infection (*Mycobacterium tuberculosis*) in elephants (*Elaphas maximus* and *Loxodonta africanum*) Swedish case studies. Verh ber Erkg Zootiere 42:173e81

Steinmetz H, Rutten M (2016) TB or Not TB: Diagnosis of tuberculosis in a group of Asian elephants (*Elephas maximus*). Proc American Assoc Zoo Veterinarians, European Assoc Zoo and Wildl Veterinarians, Institute for Zoo and Wild Research, Joint Conference p116

Vogelnest L, Hulst F, Thompson P, Lyashchenko KP, Herrin KA (2015) Diagnosis and management of tuberculosis (*Mycobacterium tuberculosis*) in an Asian elephant (*Elephas maximus*) with a newborn calf. J Zoo Wildl Med 46(1): 77-85


Elephant
2006
2007
2008
2009
2010
2011
2012
2013
E1 culture
serology
StatPak/DPP+
E2 culture
serology
StatPak/MAPIA/DPP+
E3 culture
serology
StatPak/MAPIA/DPP+
E4 culture
serology
StatPak/DPP+
E5 culture
serology
DPP+
E6 culture
serology
StatPak/MAPIA/DPP-
DPP+
E7 culture
serology
StatPak/MAPIA/DPP-
DPP+
E8 culture
serology
StatPak/MAPIA/DPP-
DPP+