



HOKKAIDO UNIVERSITY

Title	Ascomycete <i>Aspergillus oryzae</i> Is an Efficient Expression Host for Production of Basidiomycete Terpenes by Using Genomic DNA Sequences
Author(s)	Nagamine, Shota; Liu, Chengwei; Nishishita, Jumpei et al.
Citation	Applied and Environmental Microbiology, 85(15), e00409-19 https://doi.org/10.1128/AEM.00409-19
Issue Date	2019-08
Doc URL	https://hdl.handle.net/2115/76550
Type	journal article
File Information	Appl. Environ. Microbiol.pdf



Ascomycota *Aspergillus oryzae* is an efficient expression host for production of Basidiomycota terpenes using genomic DNA sequences

Shota Nagamine¹, Chengwei Liu¹, Jumpei Nishishita¹, Takuto Kozaki¹, Kaho Sogahata¹, Yoshiro Sato¹, Atsushi Minami^{1,*}, Taro Ozaki¹, Claudia Schmidt-Dannert², Jun-ichi Maruyama,³ and Hideaki Oikawa^{1,4,*}

1. Department of Chemistry, Faculty of Science, Hokkaido University, Kita-ku Kita 10 Jo Nishi 8 Chome, Sapporo 060-0810, Japan.
2. Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, St. Paul, Minnesota, USA.
3. Department of Biotechnology, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan.
4. Lead Contact

Running title (not to exceed 54 characters) 53 characters

Expression of Basidiomycota gene using genomic DNA sequences

Keywords

Basidiomycota, sesquiterpene synthase, functional analysis, *Aspergillus oryzae*

Corresponding authors

Hideaki Oikawa and Atsushi Minami

E-mail addresses

hoik@sci.hokudai.ac.jp, aminami@sci.hokudai.ac.jp

Abstract

Basidiomycota fungi are an attractive resource for biologically active natural products for use in pharmaceutically relevant compounds. Recently, genome projects of mushroom fungi have provided a great deal of biosynthetic gene cluster information. However, functional analyses of the gene clusters for natural products were largely unexplored because of the difficulty of cDNA preparation and lack of gene manipulation tools for Basidiomycota fungi. To develop a versatile host for Basidiomycota genes, we examined gene expression using genomic DNA sequences in the robust Ascomycota host *Aspergillus oryzae*, which is frequently used for the production of metabolites from filamentous fungi. Exhaustive expression of 30 terpene synthase genes from Basidiomycota *Clitopilus pseudo-pinsitus* and *Stereum hirsutum* showed two splicing patterns, completely spliced cDNAs giving terpenes (15 cases) and mostly spliced cDNAs, indicating that *A. oryzae* correctly spliced most of introns at the predicted positions and lengths. The mostly spliced cDNAs were expressed after PCR-based removal of introns, resulting in the successful production of terpenes (14 cases). During this study, we observed relatively frequent mispredictions in the automated program. Hence, the complimentary use of AO-expression and automated prediction will be a powerful tool for genome mining.

Importance

The recent large influx of genome sequences from Basidiomycota, which are prolific producers of bioactive natural products may provide opportunities to develop novel drug candidates. The development of a reliable expression system is essential for the genome mining of natural products because of the lack of a tractable host for heterologous expression of Basidiomycota genes. For this purpose, we applied the Ascomycota *Aspergillus oryzae* system for the direct expression of fungal natural product biosynthetic genes from genomic DNA. Using this system, 29 sesquiterpene synthase genes and diterpene biosynthetic genes for bioactive pleuromutilin were successfully expressed. Together with the use of computational tools for intron-prediction, this *Aspergillus oryzae* system represents a practical method for the production of Basidiomycota natural products.

1 Introduction

2 Mushroom-forming Basidiomycota fungi are known to be prolific producers of structurally
3 diverse, bioactive natural products (NPs) and have been used since ancient times in traditional
4 medicine (1). This group of fungi is particularly known for their terpenoid NPs, which appear to be
5 the major class of NPs produced by them (1, 2). Well-known examples of pharmaceutically
6 relevant terpenoid-derived compounds include the widely used livestock antibiotic pleuromutilin
7 (3), the anticancer compound illudin S (4), the melleolide antibiotics (5), and the nerve growth
8 factor-synthesis promoting erinacines (6) (Figure 1). The 1000 fungal genomes project performed
9 by the Joint Genome Institute (JGI) (7) has resulted in a large influx of Basidiomycota genome
10 sequences and genomic data from hundreds of Basidiomycota taxa that can now be searched *in*
11 *silico* for NP biosynthetic gene clusters.

12
13 However, despite their incredible potential for NP discovery, Basidiomycota fungi are a largely
14 unexplored territory for drug discovery compared to Ascomycota fungi. While an abundance of
15 genetic tools and techniques for transformation and genetic manipulation are available for yeast and
16 filamentous fungi, such methods are mostly lacking for Basidiomycota fungi (2, 8). In addition,
17 many Basidiomycota strains are difficult to grow under laboratory conditions, requiring the
18 determination of suitable growth conditions and often long fermentation times. Characterization of
19 their biosynthetic genes therefore typically requires heterologous expression in a more genetically
20 tractable host. Unfortunately, Basidiomycota have very intron-rich genomes and genes that contain
21 very small and unpredictable exons, especially in cytochrome P450 monooxygenase genes, that
22 currently necessitate the amplification of genes from cDNA. Even if a gene is expressed, alternative
23 splice variants may be produced that are not functional (2). All pioneering works to access the large
24 diversity of terpenoid NPs made by Basidiomycota have therefore relied on the amplification of
25 functional terpene synthase (TS) genes from cDNA for expression and characterization in *E. coli* or
26 yeasts, limiting the complete functional analysis of all computationally identified TSs in
27 Basidiomycota genomes (9-13). Being able to directly express Basidiomycota biosynthetic genes
28 from genomic DNA in a suitable fungal surrogate host would therefore greatly accelerate the
29 functional characterization of NP pathways.

30
31 We reasoned that a genetically tractable Ascomycota could potentially be used as a heterologous
32 host for the functional expression of Basidiomycota NP biosynthetic genes. Unlike yeasts, which
33 have relatively few genes with introns, a splicing machinery of filamentous fungi such as
34 *Aspergillus* is more similar to that of Basidiomycota (14). Heterologous expression of biosynthetic
35 genes from filamentous fungi directly from genomic DNA has become a powerful approach to
36 elucidate biosynthetic pathways. We have used this strategy with the Ascomycota *Aspergillus*
37 *oryzae* (AO) for the heterologous production of several NPs and their pathway intermediates
38 (15-18) and to access difficult to obtain pathway intermediates for the study of intriguing enzyme
39 reactions (19-21). Recently, we successfully produced in AO the Basidiomycota-derived diterpene
40 pleuromutilin from cDNA amplified biosynthetic genes. Like others, we found the cloning of
41 functional genes from Basidiomycota cDNA to be difficult and tedious. We and others also showed

that the native AO NADPH cytochrome P450 reductase supports the activity of the Basidiomycota P450 monooxygenases pathway (22, 23). Given our success with using AO for the direct expression of fungal NP biosynthetic genes from genomic DNA and the ability of AO to functionally express a Basidiomycota terpenoid NP biosynthetic enzymes, including its P450 monooxygenase, we therefore hypothesized that this Ascomycota would make an excellent platform strain for the characterization and production of Basidiomycota NP pathways, in particular its terpenoid pathways.

Here, we describe a new strategy using AO as an expression host for the heterologous production of Basidiomycota NP biosynthetic genes. We selected two different terpenoid classes (pleuromutilin diterpene and sesquiterpenoids) from two different Basidiomycota and performed a systematic analysis of the predicted and experimentally obtained splicing patterns in AO. Our results indicated that a significant fraction of genes was correctly spliced by AO, while many partially spliced genes could be corrected based on gene model predictions and functionally expressed in AO.

Results and Discussion

Expression of pleuromutilin biosynthetic genes from genomic DNA in Aspergillus oryzae

We previously achieved the heterologous production of the Basidiomycota diterpene pleuromutilin using an *A. oryzae* expression system (22). Therefore, we initially examined the heterologous expression of pleuromutilin biosynthetic genes using genomic DNA sequences in this Ascomycota host. When we tested the gene expression of GGPP synthase *ple4* and terpene synthase *ple3* in *A. oryzae*, the resultant transformant (TF) successfully produced premutilin (**1**) (Figure 2). Sequencing of cDNA sequences recovered from the TF AO-*ple3/4* showed that all introns (*ple3/4*: 3/4) were correctly spliced (Figure 2). We next examined the expression of intron rich P450 genes (*ple1/5/6*: 10/13/11). cDNA sequence analysis showed that all 11 introns of *ple6* were successfully spliced while *ple1/5* each contained one unspliced introns (*ple1*: intron-6 out of 10; *ple5*: intron-4 out of 13), which can be readily removed by PCR-based methods. Based on these promising results, we speculated that gene splicing in Basidiomycota resembles that of Ascomycota fungi (14, 24).

Functional gene expression analysis of STS genes

Given that almost all introns of *ple* genes were successfully spliced in AO, we next performed a comprehensive analysis of Basidiomycota gene expression from genomic DNA in this host. To establish the versatility of Basidiomycota heterologous expression using AO, we chose to test the expression of sesquiterpene synthases (STSs) because they are prolific in Basidiomycota genomes and are small enzymes that convert abundant farnesyl pyrophosphate (FPP) in host cells into various cyclic, volatile products that can be readily identified by GC-MS analysis of cultures (25). In this work, we selected 31 STS genes for characterization from two Basidiomycota fungi, our pleuromutilin producer *C. pseudo-pinsitus* and for comparison, *Stereum hirsutum* which was previously investigated for sesquiterpenoid production (11) (Figure 3, Table S1).

1 **1. STSs from *Clitopilus pseudo-pinsitus***

2 Our Local-Blast search using aristolochene synthase as a query sequence identified 18 STS
3 candidate genes (CpSTS1-18) in *C. pseudo-pinsitus* (Figure 3, Table S1). Open reading frames
4 (ORFs) were manually predicted by comparison with functionally characterized fungal STS genes.
5 Except for CpSTS10, which lacked the conserved motifs (DDXXD; NSE) of STSs, 17 ORFs were
6 selected for expression in AO. To ensure the expression of target genes, we used the recently
7 developed fungal CRISPR-Cas9 system specifically optimized for AO (26). In the presence of the
8 CRISPR/Cas9 plasmid with the protospacer sequence for the SC103 locus, which was found in a
9 strain that produced betaenone in good yield (27), *ligD*-deficient *A. oryzae* NSPID1 (28) was
10 transformed using a donor DNA plasmid carrying a targeted STS gene. Usually, the knock-in rate
11 ranged from 80 to 90%, much higher than in normal random integrations. GC-MS analysis of
12 headspace volatiles from these TFs revealed that nine AO TFs expressed functional STS that
13 cyclized FPP into the cyclic products (Figures 3, S1, and S2). Sesquiterpenes were identified by
14 retention indices (RI) and comparison of MS fragmentation patterns with those reported in
15 databases or the literature (Figure S1 and Table S2).

16
17 To find out why the remaining eight TFs did not produce sesquiterpenoids, cDNA of STS were
18 recovered and sequenced. The results showed that although AO correctly spliced the vast majority
19 of introns that were computationally predicted by Augustus (29), some introns were (partially)
20 skipped, thereby generating multiple splicing patterns leading to incomplete protein (ex. one intron;
21 CpSTS5/7/13/14/16, two introns; CpSTS8/17) (Figures 3 and S3, and Table S3). Following the
22 removal of nonspliced introns in amplified cDNA by PCR-based techniques,
23 CpSTS5/7/8/13/14/16/17 were expressed by *E. coli* TFs and yielded terpenes (Figures 3, S1, and S2,
24 and Table S2). Only CpSTS15 did not give any product although all introns were likely spliced
25 correctly.

26 Sesquiterpenes produced by CpSTSs include 1) aromadendrene type sesquiterpenes featuring
27 the 5-7-3 tricyclic skeleton, such as ledene (**2**) and virifloridol (**3**) (C1,C10-cyclization products)
28 (Figure 3b), 2) structurally unique sesquiterpenes such as 6-protoilludene (**7**), sterpurene (**8**),
29 pentalenene (**9**), and β -caryophyllene (**10**) (C1,C11-cyclization products) (Figure 3c), 3) cadinene
30 type sesquiterpenes possessing the 6-6 bicyclic structure, such as δ -cadinene (**12**) and γ -muurolene,
31 and biosynthetically related ones such as α -cubebene (**13**) (C1,C10-cyclization products) (Figures
32 3e and S2), and 4) linear and monocyclic sesquiterpenes such as α -farnesene (**11**) and β -elemene
33 (Figures 3c and S2). Among them, **7** and cadinenes are frequently isolated from fungi and the
34 corresponding STSs have been functionally characterized (2). By contrast, **8** produced by CpSTS1
35 is known as a fungi specific sesquiterpene and a sterpurene synthase has not been identified.
36 Successful functional characterization of CpSTS1 enabled us to search putative sterpurene
37 synthases in public database and identify homologs in *Psilocybe cyanescens* (72%, PPQ78014.1),
38 *Gymnopilus dilepis* (72%, PPQ64797.1), and *Fibularhizoctonia* sp. CBS 109695 (66%,
39 KZP33092.1) although **8** and the related metabolites have not been isolated from them.

Based on a previous phylogenetic analysis (10), 69 Basidiomycota STSs are divided into clades I-IV (Figure 4). The majority of characterized CpSTS (11/17) in this work groups with fungal STS in clades II and III that are proposed to catalyze the C1,C10- and C1,C11-cyclization of FPP, respectively (Figures 3, 4, and S2). The remaining STS group with clade I (2 STSs), clade IV (2 STSs), and unclassified clade (1 STS). We used this phylogenetic framework as a guide to investigate the relationship of homologous STSs in the context of sequence homology, gene structures, and chemical structures/reaction mechanisms. The species-specific STSs CpSTS8/9/11/12/13/16 (sequence identity: >40%) belonging to clade II have similar gene structures (Figure 3a, Table S3) (30), possibly arising by gene duplication of ancestral STS, a well known evolutionary mechanism in this class of fungi (31). These STSs produced the aromadendrene family members and aristolene (**6**), which are frequently found as plant volatiles (Figures 3b and S4), but only one corresponding STS from *Citrus* (CsSesquiTPS5) has been characterized (32). All of them are most likely biosynthesized via bicyclogermacrene (**IM1**) (30, 32-34). These results suggested a strong relationship of the gene structure and the cyclization mechanism. It should be emphasized that comprehensive analysis of 17 STSs provides direct evidence regarding the gene structure and the enzyme function.

2. STSs from *S. hirsutum*

Previously, 18 STS genes were bioinformatically predicted in the mushroom *S. hirsutum* and five STS genes were successfully amplified from cDNA for functional expression in *E. coli* (11, 35). The remaining STS were not studied, partially because of cDNA availability limitations. We sought this suite of STS from *S. hirsutum* would be another good test system for our direct gene expression strategy. Of the 18 STS genes (ShSTS1-18), we excluded five genes from our study to avoid the expression of ShSTS2/9/14/15/6 that showed >60% homology to ShSTS1/8/13/16 and CpSTS3 (Table S4). GC-MS analysis of culture headspace showed that seven AO TFs expressing (ShSTS-1/4/7/8/11/16/17) produced sesquiterpenes such as **7**, **12**, **15**, β -barbatene, and hirsutene, (Figures 3, S1, S2 and Table S2). As in the case of CpSTSs, sequencing of cDNA showed intron skipping at one (ShSTS5/12/13/18) or two positions (ShSTS3/10) (Figure 3d). Removal of predicted introns in ShSTS3/5/10/12/13/18 gave functional STSs in *E. coli* transformants that synthesized terpenes such as **10**, **11**, **13**, **14**, **16**, and γ -cadinene (Figures 3, S1, S2 and Table S2). The metabolite profiles revealed that clade III ShSTSs as well as those of CpSTSs produce structurally unique sesquiterpene such as **7-10** as a single product while other STSs appear to give multiple biosynthetically related products.

A strong correlation between gene structure and function was also found in *Stereum* TSs; clade II ShSTS8/9/10/11/12 revealed a relatively high sequence identity (>48%) and the gene structures of ShSTS10/11/12 are nearly identical, whereas that of the ShSTS8 gene is likely modified by the insertion of an extra intron (Figures 3d and 4). These STSs generated cadinene-type sesquiterpenes biosynthesized via *Z,E*-germacrenedieryl cation (**IM2**) (Figure 3e). In addition, a conservation of sequence identity (>54%) and gene structure is observed for clade III ShSTSs, ShSTS15/16/17/18 (Figures 3c and 4), which produce **7** as a characteristic and major sesquiterpene made by

mushrooms. The gene structure of ShSTS13 resembles with those of Clade III ShSTSs except that an extra *N*-terminal intron is inserted (Figure 3d). The structural difference between **7** (ShSTS15/16/17/18) and **10** (ShSTS13) might reflect to alter the transition states derived from common intermediate **IM3**. When we searched for putative biosynthetic gene clusters surrounding clade II and III Cp- and Sh-STS paralogs/orthologs, we found quite divergent arrangements of putative biosynthetic genes for modifying terpene scaffold (Figure S5).

Splicing of Basidiomycota genes in AO.

To gain insights into the splicing patterns of Basidiomycota genes in AO, we analyzed the sequences of spliced and nonspliced introns of genes investigated in this study (Table S3). Overall, CpSTS have a slightly smaller mean intron length (56 nt, size range: 47-69 nt) and average number of introns (4 introns) compared to ShSTS with an intron length of 64 nt (size range: 48-122 nt) and 5 introns on average. The 5'- and 3'-splice site consensus sequences (GUxxGU and YAG, respectively) matched those from *C. neoformans* (Basidiomycota fungi) and *Fusarium* sp. (Ascomycete) (14, 24). Analysis of branch sites revealed that the majority introns (82%, 108/132) has a CTNAN motif, while 13% (17/132) contain a TTNAN motif and for the remaining (7/132) no motif could be determined. Branch site motif appears to be the most important predictor for splicing in AO, as 96% of introns with a CTNAN are spliced in AO, while only 65% of introns with a TTNAN or 43% of discernable motif are spliced. Only one spliced intron (CpSTS8-intron1) out of all introns spliced in AO (132 number) was mis-spliced in AO (Figure S3), demonstrating that introns in Basidiomycota genes that are recognized for splicing are faithfully and reliably spliced by AO. Intron skipping tends to occur at one or two positions in STS genes. A similar pattern was observed for pleuromutilin genes investigated in this work (Figure 2).

Estimation of reliability for computational prediction of introns and ORFs

Comparison of the cDNA sequences of the large number of 30 functional STS sequences characterized in this work with computationally predicted gene structures provides an excellent assessment of the reliability of bioinformatics tools such as Augustus (29) for Basidiomycota gene prediction (Tables S3 and S5). In Augustus, the accuracy of gene structure predictions depended on the gene model used; predictions with the mushroom *Coprinus cinereus* model instead of AO (expression host) were much more accurate. Even with the mushroom gene model, Augustus mispredicted 11 out 30 STS gene structures (Figures 3 and S3, Table S5). Misprediction included i) insertion of an extra intron in *CpSTS16*; ii) skipping of an essential intron in *ShSTS5*; iii) prediction of shorter/longer introns in *CpSTS18* and *ShSTS12/16*; iv) misprediction of introns closely located in the 3'-end of *CpSTS4/13*, *ShSTS17* (extra intron), and *CpSTS9* (intron skipping) and the 5'-end of *CpSTS8* and *ShSTS11* (Figure 5). Most errors concerning ORFs and introns located close to the 3'-terminal end of the genes (example iv) can be avoided by manual prediction by considering the amino acid sequences of the known STSs and the gene structures of paralogous STS genes (Figure 3). Therefore, automated prediction may be useful when these measures are taken. Actually, we used Augustus prediction when we needed to remove nonspliced intron(s) in the AO expression. We successfully used this strategy to identify and subsequently remove nonspliced introns by AO in

recovered cDNA for expression of active STS by EC-TFs (F1). Consequently, direct expression of Basidiomycota genes from genomic DNA in AO in conjunction with bioinformatics proved to be an effective and reliable strategy to quickly obtain functional NP biosynthetic genes.

Concluding remarks

In this work we developed an efficient approach for heterologous expression of Basidiomycota NP biosynthetic genes directly from genomic DNA in the genetically tractable and well-studied Ascomycota host *A. oryzae*. We show that direct expression of unspliced genes from genomic DNA from Basidiomycota in AO results in almost correctly spliced introns and that a combination of computational and manual gene structure prediction can be used to identify and correct nonspliced introns in recovered cDNA by a quick PCR step. Out of 30 STS expressed in AO using this method, half of them were functionally expressed and a simple PCR step yielded functional cDNA for the remaining STS. Comprehensive analysis of gene structures indicated a strong correlation between gene structure preservation and evolution of enzyme function. Comparison of intron sequences revealed intron features such as branch sites that are less likely to be spliced by an Ascomycota such as AO. This information will aid in the prediction of Basidiomycota STS activities, provide insights in the diversification of NP cluster in this class of fungi and improve computational gene prediction methods. Further, by knowing which introns are less likely to be processed, genomic sequences can be preemptively, spot corrected prior to introduction into a versatile, heterologous Ascomycota production host – thereby opening the door for the accessing the largely undiscovered NP diversity of Basidiomycota mushrooms.

Materials and Methods

General. All reagents commercially supplied were used as received. Column chromatography was carried out on 60N silica gel (Kanto Chemicals). Optical rotations were recorded on JASCO P-2200 digital polarimeter. ¹H-NMR spectra were recorded on Bruker DRX-500 or Bruker AMX-500 spectrometer (500 MHz for ¹H-NMR). NMR spectra were recorded in CDCl₃ (99.8 atom% enriched, Kanto). ¹H chemical shifts were reported in δ value based on residual chloroform (7.26 ppm) as a reference. Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad), coupling constant (Hz), and integration. GC-MS analyses were conducted with MS-2010 (Shimadzu). Mass spectra were obtained with a Waters ACQUITY QDa (ESI mode).

Oligonucleotides for polymerase chain reactions (PCRs) were purchased from Hokkaido System Science Co., Ltd. PCRs were performed with a BioRad S1000 thermal cycler. All PCR reactions were performed with the KOD-Plus-Neo (TOYOBO). The assembly of DNA fragments was performed by using either In-Fusion Advantage PCR cloning kit (Clontech Laboratories) or HiFi DNA Assembly Master Mix (New England Biolabs).

Strains and culture conditions. *Escherichia coli* HST08 was used for cloning and following standard recombinant DNA techniques. *E. coli* BL21-Gold(DE3) was used for protein expression. A fungal host strain used in this study was *A. oryzae* NSAR1 (36), a quadruple auxotrophic mutant

(*niaD*⁻, *sC*⁻, *ΔargB*, *adeA*⁻), and *A. oryzae* NSPID1 (28), a strain with a highly efficient gene-targeting background (*niaD*⁻, *sC*⁻, *ΔpyrG*, *ΔligD*) for expression. AO-*bet123* (27), a betaenone B highly producing transformant, was used for genome sequencing.

Genome Sequencing and Analysis. Genome sequencing of AO-*bet123* was performed by Hokkaido System Science Co., Ltd. (Hokkaido, Japan) with an Illumina HiSeq 2000 system. Read mapping was performed with the following programs, BWA, Samtools, GATK, and Picard. The results showed a clear gap between AO090103000023 and AO090103000025 on the chromosome 8. This site is tentatively defined as SC103 region in this study.

Construction of a Cas9 plasmid pC9SC103. The Cas9 plasmid, pC9SC103, harbouring guide RNA sequences for SC103 was constructed as follows; The gRNA fragments (G-Fr1: U6P-SC103 and G-Fr2: SC103-U6T) were amplified from ppAsACas9gwA (26) with primer set as shown in Supplemental Table S6. These primary products were used as a template for the fusion PCR to afford guide RNA sequences flanked by U6 promoter/terminator set. The PCR product (U6P-guide RNA sequence for SC103-U6T) was then inserted into *Sma*I-digested ppAsACas9 to construct pC9SC103.

Construction of donor vectors, pDP1031 and pDP1032. The donor vectors, pDP1031 and pDP1032, harbouring homologous arms of SC103 was constructed as follows; Two DNA fragments, SC103-up and SC103-down, were amplified from genomic DNA of *Aspergillus oryzae* NSAR1 with primer sets as shown in Supplemental Table S6. Each PCR product was inserted into *Hind*III/*Eco*RI-digested pUC19 to afford pUC19-SC103. The DNA fragment harbouring primer-terminator sets for overexpression in *A. oryzae* were amplified from either pTAex3 or pUARA2 with primer set as shown in Supplemental Table S6. Each PCR product was inserted into *Bam*HI-digested pUC19-SC103 to afford pDP1031 (single primer/terminator set) and pDP1032 (tandem primer/terminator sets).

Bioinformatics analysis. STS gene annotation, alignment, and gene structure prediction were performed as described previously (11). Briefly, gene predictions were made in Augustus (29) with different fungal gene models using the genomic region 10-15 kb flanking identified STS ORFs. STS gene predictions were manually aligned with functionally characterized fungal STS. All sequence alignments and phylogenetic analysis were performed in MEGA7 (37) using MUSCLE (38) for protein alignments. Phylogenetic analysis of fungal STSs (**Figure 2B**) was done using the Neighbor-joining method using the Poisson correction methods with 500 bootstrap replications (39, 40).

Accession numbers and references protein sequences are as follows:

Basidiomycota STS: *Coprinus cinereus* (Cop1-4, Cop6) [XP 001832573, XP 001836556, XP 01832925, XP 01836356, XP 01832548], *Omphalotus olearius* (Omp1-10), *Fomitopsis pinicola* [FomPi84944], *Stereum hirsutum* [Steh1|159379, 128017, 25180, 64702, 73029], *Armillaria gallica* (ArmGa1) [P0DL13], and *Postia placenta* [PpSTS01 [BBD74517.1], 02 [BBD74518.1], 03

[BBD74519.1], 06 [BBD74520.1], 07 [BBD74521.1], 08 [BBD74522.1], 09 [BBD74523.1], 10 [BBD74524.1], 13 [BBD74525.1], 14 [BBD74526.1], 16 [BBD74527.1], 18 [BBD74528.1], 22 [BBD74529.1], 24 [BBD74530.1], 25 [BBD74531.1], 29 [BBD74532.1].

Ascomycota STS: *Fusarium fujikori* (Ffsc4) [HF563560.1] and Ffsc6 [HF563561.1], *Fusarium gramineareum* (FgCLM1) [GU123140]), *Aspergillus terreus* (atAS) [Q9UR08], *Penicillium roqueforti* (prAS) [W6Q4Q9], and *Botrytis cinerea* (BcBOT2) [AAQ16575.1].

Accession number: The sequences of sesquiterpene synthase genes found in *C. pseudo-pinsitus* ATCC20527 have been deposited in the DNA Data Bank of Japan (DDBJ). The accession numbers are summarized in Table S1.

Construction of *A. oryzae* expression plasmids of *ple* and sesquiterpene synthase genes. Pleuromutin biosynthetic genes (*ple13456*) were amplified from the genomic DNA of *C. pseudo-pinsitus* ATCC20527 (22) with primer set as shown in Supplemental Table S6. Each PCR product was inserted into appropriate restriction site (site 1 and/or site 2) of pDP1031 or pDP1032 to construct the following expression plasmids; pDP1031-*ple5* (*KpnI* site), pDP1032-*ple6* (*NheI* site), pDP1032-*ple3* (*KpnI*)/4 (*SpeI*), and pDP1032-*ple7* (*KpnI*)-*ple1* (*SpeI*).

Sesquiterpene synthase genes were amplified from the genomic DNA of *C. pseudo-pinsitus* ATCC20527 and *S. hirsutum* FP-91666 S1 (11) with primer set as shown in Supplemental Table S6. Each PCR product was then inserted into the appropriate restriction sites of expression vectors to construct the following expression plasmids; pTAex3-*CpSTS1*, pUARA2-*CpSTS2*, pUARA2-*CpSTS3*, pTAex3-*CpSTS4*, pDP1031-*CpSTS5*, pDP1031-*CpSTS6*, pDP1031-*CpSTS7*, pDP1031-*CpSTS8*, pDP1031-*CpSTS9*, pDP1031-*CpSTS10*, pDP1031-*CpSTS11*, pDP1031-*CpSTS12*, pUSA2-*CpSTS13*, pDP1031-*CpSTS14*, pDP1031-*CpSTS15*, pUSA2-*CpSTS16*, pDP1031-*CpSTS17*, pDP1031-*CpSTS18*, pDP1032-*ShSTS1/18*, pDP1031-*ShSTS3*, pDP1031-*ShSTS4*, pDP1031-*ShSTS5*, pDP1031-*ShSTS7*, pDP1031-*ShSTS8*, pDP1031-*ShSTS10*, pDP1032-*ShSTS11/16*, pDP1031-*ShSTS12*, pDP1031-*ShSTS13*, and pDP1031-*ShSTS17*.

Transformation of *Aspergillus oryzae* (genome-editing method). Most transformants used in this study was constructed by genome-editing method; a spore suspension of *A. oryzae* NSPID1 (1.0 x 10⁸ cells) were inoculated into CD (0.3 % of NaNO₃, 0.2 % of KCl, 0.1 % of K₂HPO₄, 0.05 % of MgSO₄·7H₂O, 2 % of dextrin, 0.002 % of FeSO₄·7H₂O, 0.15 % of methionine, 0.488 % of uracil, 0.2 % of uridine, 100 mL, pH 5.5) medium supplemented with appropriate nutrients. After 3 days incubation at 30 °C (200 rpm), mycelia was collected by filtration and washed with water. Protoplasting was performed using Yatalase (Takara; 5.0 mg mL⁻¹) in Solution 1 (0.8 mM of NaCl, 10 mM of NaH₂PO₄, pH 6.0) at 30 °C for 2 h. Protoplasts were centrifuged at 2,000 rpm (Beckman JLA10.500) for 5 min and washed with 0.8 M of NaCl solution. Then, protoplasts were adjusted to 2.0 x 10⁸ cells/mL by adding Solution 2 (0.8 M of NaCl, 10 mM of CaCl₂, 10 mM of Tris-HCl, pH 8.0) and Solution 3 (40 % (w/v) of PEG4000, 50 mM of CaCl₂, 50 mM of Tris-HCl, pH 8.0) in 4/1 volume ratio. To the protoplast solution (200 μ L) was added a cas9 plasmid (2 μ g) and a donor plasmid (5 μ g). The aliquot was incubated on ice for 20 min and then Solution 3 (1 mL) added to

the aliquot. After 20 min incubation at room temperature, Solution 2 (10 mL) added to the mixtures and the mixture was centrifuged at 2,000 rpm (Beckman JLA10.500) for 5 min. After decantation, the residue was diluted with Solution 2 (500 μ L) and the mixture (100 μ L) was poured onto the CD agar plate (1.5 %) supplemented with 4.7% of NaCl and then overlaid with the soft-top CD agar (0.6 %) containing 21.8 % of sorbitol. The plates were incubated at 30 °C for 3-7 days.

Analysis of the metabolites from AO-transformants harboring a STS gene. Mycelia of each transformant was inoculated into a MPY medium (1 mL) containing 20 mM of uridine and 0.2% of uracil in 10 mL test tube. Each culture was incubated at 30 °C for 4 days. The volatile organic compounds were extracted by a SPME fiber (50/30 μ m DVB/CAR/PDMS, Stableflex, 24Ga, Manual Holder), which was conditioned by inserting it into the GC injector to prevent contamination, during 30 min at room temperature. After extraction, the fiber was pulled into the needle sheath and the SPME device was removed from the vial and then inserted into the injection port of a GC-MS QP2010 apparatus (Shimadzu, Kyoto, Japan) with a HP-5 capillary column (0.32 mm $\times$ 30 m, 0.25 μ m film thickness; J&W Scientific, Folsom, CA). Each sample was injected onto the column at 60 °C in the splitless mode. The column temperature was increased by 4 °C min⁻¹ to 180 °C. The flow rate of the helium carrier gas was 0.66 mLmin⁻¹ (**method A**).

Isolation of sesquiterpenes.

Sterpurene; Mycelia of AO-*CpSTS1* were inoculated into a solid medium containing polished rice (100 g) and adenine (10 mg) in 500 mL Erlenmeyer flasks. Each culture was incubated at 30 °C for 12 days. After extraction with ethyl acetate, the extract was concentrated in vacuo to afford crude extracts. The crude extracts were purified with silica gel column chromatography (hexane) to give sterpurene (20.0 mg from 1.3 kg of rice medium from AO-*CpSTS1*). The NMR data are in good agreement with the reported data (Figure S6, 41).

9-Alloaromadendrene; Mycelia of AO-*CpSTS11* was inoculated into a MPY medium (30 mL) containing 20 mM of uridine and 0.2 % of uracil in 200 mL Erlenmeyer flasks. Each culture was incubated at 30 °C for 5 days. The crude extracts were purified with silica gel column chromatography (hexane) to give 9-alloaromadendrene (1.0 mg from 150 mL of MPY medium containing uridine and uracil). The NMR data are in good agreement with the reported data (Figure S6, 42).

cDNA preparation. Each transformant was grown on MPY medium containing 20 mM of uridine and 0.2 % of uracil for 3-5 days at 30 °C. Total RNA was extracted from each dried mycelia using TRIzol® Reagent (Invitrogen) according to the manufacturer's instructions and then treated with DNase I (Life Technologies) for reverse transcription. Complementary DNA (cDNA) was synthesized with PrimeScript™ II 1st strand cDNA synthesis kit (Takara) using the oligo dT primer according to the manufacturer's instructions. The cDNA was used as a template of the PCR reactions for direct sequencing and subcloning of each STS gene into the pColdI vector.

Estimation of sequences of functionally active STSs. cDNA sequences derived from AO transformant were aligned with the predicted sequence by Augustus for identifying the nonspliced

introns in cDNA sequences. Alignments were computed by using ClustalW. The resultant transcripts of putative intronless STSs were then manually examined for the presence of and proper alignment of the conserved motifs characteristic for sesquiterpene synthases.

Construction of *E. coli* expression plasmids. Intronless DNA clone was prepared from PCR-based removal of the nonspliced introns (Figures 3 and 4) as follows; two DNA fragments (upstream- and downstream regions of the nonspliced intron) were separately amplified from cDNA with primer set as shown in Supplemental Table S6. Each PCR product was inserted into *Nde*I-, *Eco*RI-, or *Kpn*I-digested pColdI to construct the following expression plasmids; pColdI-*CpSTS5*, pColdI-*CpSTS7*, pColdI-*CpSTS8*, pColdI-*CpSTS13*, pColdI-*CpSTS14*, pColdI-*CpSTS16*, pColdI-*CpSTS17*, pColdI-*ShSTS3*, pColdI-*ShSTS5*, pColdI-*ShSTS10*, pColdI-*ShSTS12*, pColdI-*ShSTS13*, and pColdI-*ShSTS18*.

Analysis of the metabolites from EC-transformants. The constructed plasmids were separately introduced into *E. coli* BL21-Gold(DE3) for overexpression. The transformant was grown at 37 °C at an OD₆₀₀ of ~0.6 in 500 mL flask. After cooling at 4 °C, isopropyl β-D-thiogalactopyranoside (0.1 mM) was added to the culture. After incubation at 16 °C for 17 h, the volatile organic compounds were extracted by a SPME fiber (50/30um DVB/CAR/PDMS. Stableflex, 24Ga, Manual Holder) during 30 min at room temperature. The extracts were then analyzed by a GC-MS apparatus according to **method A**.

Supplemental Material

Supplemental material for this article may be found at xxx.

Acknowledgments

This work was financially supported by Grants-in-Aid for Japan Society for the Promotion of Science Grants JP15H01835 to H.O., JP16H03277 and JP16H06446 to A.M., and JP18K14342 to L.C.

References

1. Lorenzen K, Anke T. 1998. Basidiomycetes as a source for new bioactive natural products. *Curr Org Chem* 2;329-364.
2. Quin MB, Flynn CM, Schmidt-Dannert C. 2014. Traversing the fungal terpenome. *Nat Prod Rep* 31;1449-1473.
3. Kavanagh F, Hervey A, Robbins WJ. 1951. Antibiotic substances from Basidiomycetes: VIII. *Pleurotus multilus* (Fr.) Sacc. and *Pleurotus passeckerianus* Pilat. *Proc Natl Acad Sci USA* 37;570-574.
4. McMorris TC, Anchel M. 1963. Structures of Basidiomycete metabolites illudin S and illudin M. *J Am Chem Soc* 85;831-832.

- 1 5. Midland SL, Izac RR, Wing RM., Zaki AI, Munnecke DE, Sims JJ. 1982. Melleolide, a new
2 antibiotic from *Armillaria mellea*. Tetrahedron Lett 23;2515-2518.
- 3 6. Kawagishi H, Shimada A, Hosokawa S, Mori H, Sakamoto H, Ishiguro Y, Sakemi S, Bordner J,
4 Kojima N, Furukawa S. 1996. Erinacines E, F, and G, stimulators of nerve growth factor
5 (NGF)-synthesis, from the mycelia of *Hericium erinaceum*. Tetrahedron Lett 37;7399-7402.
- 6 7. Martin F, Cullen D, Hibbett D, Pisabarro A, Spatafora JW, Baker SE, Grigoriev IV. 2011.
7 Sequencing the fungal tree of life. New Phytol 190;818-821.
- 8 8. Stadler M, Hoffmeister D. 2015. Fungal natural products-the mushroom perspective. Front
9 Microbiol 6;127.
- 10 9. Agger S, Lopez-Gallego F, Schmidt-Dannert C. 2009. Diversity of sesquiterpene synthases in
11 the basidiomycete *Coprinus cinereus*. Mol Microbiol 72;1181-1195.
- 12 10. Wawrzyn GT, Quin MB, Choudhary S, Lopez-Gallego F, Schmidt-Dannert C. 2012. Draft
13 genome of *Omphalotus olearius* provides a predictive framework for sesquiterpenoid natural
14 product biosynthesis in Basidiomycota. Chem Biol 19;772-783.
- 15 11. Quin MB, Flynn CM, Wawrzyn GT, Choudhary S, Schmidt-Dannert C. 2013. Mushroom
16 hunting by using bioinformatics: Application of a predictive framework facilitates the selective
17 identification of sesquiterpene synthases in Basidiomycota. ChemBioChem 14;2480-2491.
- 18 12. Ichinose H, Kitaoka T. 2018. Insight into metabolic diversity of the brown-rot basidiomycete
19 *Postia placenta* responsible for sesquiterpene biosynthesis: semi-comprehensive screening of
20 cytochrome P450 monooxygenase involved in protoilludene metabolism. Microb Biotechnol
21 11;952-965.
- 22 13. Burkhardt I, Kreuzenbeck N, Beemelmans C, Dickschat JS. 2019. Mechanistic
23 characterization of three sesquiterpene synthases from the termite-associated fungus
24 Termitomyces. Org Biomol Chem 17;3348-3355.
- 25 14. Kupfer DM, Drabenstot SD, Buchanan KL, Lai HS, Zhu H, Dyer DW, Roe BA, Murphy JW.
26 2004. Introns and splicing elements of five diverse fungi. Eukaryot Cell 3;1088-1100.
- 27 15. Fujii R, Minami A, Tsukagoshi T, Sato N, Sahara T, Ohgiya S, Gomi K, Oikawa H. 2011. Total
28 biosynthesis of diterpene aphidicolin, a specific inhibitor of DNA polymerase alpha:
29 Heterologous expression of four biosynthetic genes in *Aspergillus oryzae*. Biosci Biotechnol
30 Biochem 75;1813-1817.
- 31 16. Tagami K, Liu CW, Minami A, Noike M, Isaka T, Fueki S, Shichijo Y, Toshima H, Gomi K,
32 Dairi T, Oikawa H. 2013. Reconstitution of biosynthetic machinery for indole-diterpene
33 paxilline in *Aspergillus oryzae*. J Am Chem Soc 135;1260-1263.
- 34 17. Ye Y, Minami A, Mandi A, Liu C, Taniguchi T, Kuzuyama T, Monde K, Gomi K, Oikawa H.
35 2015. Genome mining for sesterterpenes using bifunctional terpene synthases reveals a unified
36 intermediate of di/sesterterpenes. J Am Chem Soc 137;11846-11853.

- 1 18. Heneghan MN, Yakasai AA, Halo LM, Song Z, Bailey AM, Simpson TJ, Cox RJ, Lazarus CM.
2 2010. First heterologous reconstruction of a complete functional fungal biosynthetic multigene
3 cluster. *ChemBioChem* 11;1508-1512.
- 4 19. Liu C, Tagami K, Minami A, Matsumoto T, Frisvad JC, Suzuki H, Ishikawa J, Gomi K, Oikawa
5 H. 2015. Reconstitution of biosynthetic machinery for the synthesis of the highly elaborated
6 indole diterpene penitrem. *Angew Chem Int Ed* 54;5748-5752.
- 7 20. Ye Y, Minami A, Igarashi Y, Izumikawa M, Umemura M, Nagano N, Machida M, Kawahara T,
8 Shin-ya K, Gomi K, Oikawa H. 2016. Unveiling the biosynthetic pathway of the ribosomally
9 synthesized and post-translationally modified peptide ustiloxin B in filamentous fungi. *Angew*
10 *Chem Int Ed* 55;8072-8075.
- 11 21. Matsuda Y, Iwabuchi T, Wakimoto T, Awakawa T, Abe I. 2015. Uncovering the unusual D-ring
12 construction in terretonin biosynthesis by collaboration of a multifunctional cytochrome P450
13 and a unique isomerase. *J Am Chem Soc* 137;3393-3401.
- 14 22. Yamane M, Minami A, Liu C, Ozaki T, Takeuchi I, Tsukagoshi T, Tokiwano T, Gomi K, Oikawa
15 H. 2017. Biosynthetic machinery of diterpene pleuromutilin isolated from Basidiomycete fungi.
16 *ChemBioChem* 18;2317-2322.
- 17 23. Bailey AM, Alberti F, Kilaru S, Collins CM, de Mattos-Shipley K, Hartley AJ, Hayes P, Griffin
18 A, Lazarus CM, Cox RJ, Willis CL, O'Dwyer K, Spence DW, Foster GD. 2016. Identification
19 and manipulation of the pleuromutilin gene cluster from *Clitopilus passeckerianus* for increased
20 rapid antibiotic production. *Sci Rep* 6;25202.
- 21 24. Phasha MM, Wingfield BD, Coetzee MPA, Santana QC, Fourie G, Steenkamp ET. 2017.
22 Architecture and distribution of introns in core genes of four *Fusarium* species. *G3-Genes*
23 *Genom Genet* 7;3809-3820.
- 24 25. Dickschat JS. 2014. Capturing volatile natural products by mass spectrometry. *Nat Prod Rep*
25 31;838-861.
- 26 26. Katayama T, Tanaka Y, Okabe T, Nakamura H, Fujii W, Kitamoto K, Maruyama J. 2016.
27 Development of a genome editing technique using the CRISPR/Cas9 system in the industrial
28 filamentous fungus *Aspergillus oryzae*. *Biotechnol Lett* 38;637-642.1
- 29 27. Ugai T, Minami A, Fujii R, Tanaka M, Oguri H, Gomi K, Oikawa H. 2015. Heterologous
30 expression of highly reducing polyketide synthase involved in betaenone biosynthesis. *Chem*
31 *Commun* 51;1878-1881.
- 32 28. Maruyama J, Kitamoto K. 2008. Multiple gene disruptions by marker recycling with highly
33 efficient gene-targeting background (Delta ligD) in *Aspergillus oryzae*. *Biotechnol Lett*
34 30;1811-1817.
- 35 29. Hoff K J, Stanke M. 2013. WebAUGUSTUS-a web service for training AUGUSTUS and
36 predicting genes in eukaryotes. *Nucleic Acids Res* 41;W123-W128.
- 37 30. Miller DJ, Allemann RK. 2012. Sesquiterpene synthases: passive catalysts or active players?
38 *Nat Prod Rep* 29;60-71.

- 1 31. Syed K, Shale K, Pagadala NS, Tuszynski J. 2014. Systematic identification and evolutionary
2 analysis of catalytically versatile cytochrome P450 monooxygenase families enriched in model
3 Basidiomycete fungi. Plos One 9;e86683.
- 4 32. Alquezar B, Rodriguez A, de la Pena M, Pena L. 2017. Genomic analysis of terpene synthase
5 family and functional characterization of seven sesquiterpene synthases from *Citrus sinensis*.
6 Front Plant Sci 8;1481.
- 7 33. Chou WKW, Fanizza I, Uchiyama T, Komatsu M, Ikeda H, Cane DE. 2010. Genome mining in
8 *Streptomyces avermitilis*: Cloning and characterization of SAV_76, the synthase for a new
9 sesquiterpene, avermitilol. J Am Chem Soc 132;8850-8851.
- 10 34. Dickschat JS, Martens T, Brinkhoff T, Simon M, Schulz S. 2005. Volatiles released by a
11 *Streptomyces* species isolated from the North Sea. Chem Biodivers 2;837-865.
- 12 35. Flynn CM, Schmidt-Dannert C. 2018. Sesquiterpene synthase-3-hydroxy-3-methylglutaryl
13 coenzyme A synthase fusion protein responsible for hirsutene biosynthesis in *Stereum hirsutum*.
14 Appl Environ Microb 84;e00036-18.
- 15 36. Jin FJ, Maruyama J, Juvvadi PR, Arioka M, Kitamoto K. 2004. Development of a novel
16 quadruple auxotrophic host transformation system by argB gene disruption using adeA gene and
17 exploiting adenine auxotrophy in *Aspergillus oryzae*. FEMS Microbiol Lett 239;79-85.
- 18 37. Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis
19 version 7.0 for bigger datasets. Mol Biol Evol 33;1870-1874.
- 20 38. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
21 throughput. Nucleic Acids Res 32;1792-1797.
- 22 39. Jones DT, Taylor WR, Thornton JM. 1992. The rapid generation of mutation data matrices from
23 protein sequences. Comput Appl Biosci 8;275-282.
- 24 40. Felsenstein J. 1985. Confidence limits on phylogenies: An approach using the bootstrap.
25 Evolution 39;783-791.
- 26 41. Ayer WA, Saeedi-Ghomi MH. 1981. 1-sterpurene-3,12,14-triol and 1-sterpurene, metabolites of
27 silver-leaf disease fungus *Stereum purpureum*. Can J Chem 59;2536-2538.
- 28 42. Friedel HD, Matusch R. 1987. Neue aromadendran-derivate aus tolu-balsam. Helv Chim Acta
29 70;1753-1759.
- 30




Figure 1. Chemical structures of biologically active terpenoids produced by Basidiomycota fungi.







Figure 2. (A) Biosynthetic pathway of pleuromutilin. The numbers in parentheses are the nonspliced introns in the cDNA recovered from AO-TF and the total introns of the corresponding gene. (B) GC-MS profiles of the metabolites from (a) AO-*ple34* and (b) synthetic sample. (C) Schematic view of the gene structures of *ple3*, *ple4*, *ple1*, *ple5*, and *ple6*. Exon regions are shown in shaded boxes. Non-spliced introns are shown in red boxes.



(c) C1-C11 cyclization catalyzed by clade III STSs



(e) C1-C10 cyclization catalyzed by clade II_B STSs




Figure 3. (a) Schematic view of the gene structures of CpSTSs. The dark bars indicate exons and the dotted lines do introns along with intron phases. The red bars represent the computationally predicted introns that were not spliced (skipped) in cDNA from AO-TFs, of which removal resulted in functional STS expression in *E. coli* TF. Genes with the colored boxes are proposed paralogs or orthologs in the *C. pseudo-pinsitus* and *S. hirsutum* that belong to cyclization clade III (purple) or clade II_A (orange) or II_B (green). Non-functional refers to CpSTS10 lacking conserved active site motifs. (b) Proposed cyclization mechanism of terpene products catalyzed by clade II_A STSs. (c) Proposed cyclization mechanism of terpene products catalyzed by clade III STSs. (d) Schematic view of the gene structures of ShSTSs. A gene structure of functional ShSTS15 with # was reported previously (31). Genes with \$ denote the computationally predicted gene structure of STS not investigated in this work. STSs from *C. pseudo-pinsitus* are underlined in (b), (c), and (e). Stereochemistries of putative carbocation intermediates are predicted by those of cyclized products.




Figure 4. Phylogenetic tree of functionally characterized STSs except four ShSTSs (white circles) and seven PpSTSs (white circles) derived from Basidiomycetes fungi. STSs derived from *C. pseudo-pinsitus* and *S. hirsutum* are described in red and blue colors, respectively. STSs derived from Ascomycetes fungi are indicated by grey circle.



Figure 5. Schematic view of the gene structures of active and mispredicted STSs. The blue bars indicate the mispredicted sites in computationally predicted (Augustus (26)) gene structures compared to the functional gene structure shown directly above.