



HOKKAIDO UNIVERSITY

Title	Role of His63 in HutZ from <i>Vibrio cholerae</i> in the heme degradation reaction and heme binding
Author(s)	Uchida, Takeshi; Dojun, Nobuhiko; Sekine, Yukari et al.
Citation	Dalton Transactions, 48(16), 5408-5416 https://doi.org/10.1039/C9DT00926D
Issue Date	2019-04-28
Doc URL	https://hdl.handle.net/2115/77738
Type	journal article
File Information	DALTON T_supplementary information.pdf



Electronic supplementary information

Role of Hydrogen Bonding to a Heme Propionate Group in the Heme Degradation Reaction by HutZ from *Vibrio cholerae**

Takeshi Uchida,^{*1,2} Nobuhiko Dojun,² Yukari Sekine,² and Koichiro Ishimori^{1,2}

¹Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo 060-0810,
Japan

²Graduate School of Chemical Sciences and Engineering, Hokkaido University, Sapporo
060-8628, Japan

Table of Contents

Supplementary Figures

S2-S4




Fig. S1 Top view of HugZ from *Helicobacter pylori* (PDB ID code 3GAS).




Fig. S2 Regiospecificity of heme degradation by HutZ. HPLC chromatography of products from reaction of free heme (a), heme-WT (b), heme-H63L mutant (c) and heme-H63G mutant (d) with 0.1 mM H_2O_2 and 1 mM ascorbic acid.





Fig. S3 Crystal structure of HugZ from *H. pylori* (PDB 3GAS) at the heme-binding site. Residues are numbered according to the amino acid sequence of HutZ from *V. cholerae*..

(A) H63G



(B) H63F




Fig. S4 Heme titration plots of H63G and H63F mutant HutZ proteins. Heme binding curve generated from the difference absorbance of the Soret maximum by plotting ΔA_{Soret} vs. molar ratios of heme to protein, specifically, H63G (A) and H63F at pH 8.0 (B) at pH 8.0 (D). The protein was in 10 μM in 50 mM Tris-HCl, 150 mM NaCl (pH 8.0).




Fig. S5 Location of Trp109 and heme on the HugZ crystal structure (PDB 3GAS). Residues are numbered according to the amino acid sequence of HutZ from *V. cholerae*. Green and cyan show each protomer. Trp109 is present in a different protomer (green) from that (cyan) containing nearby Asp132