



HOKKAIDO UNIVERSITY

Title	Heat-stress Memory is Responsible for Acquired Thermotolerance in <i>Bangia fuscopurpurea</i>
Author(s)	Kishimoto, Ikuya; Ariga, Inori; Itabashi, Yutaka et al.
Citation	Journal of phycology, 55(5), 971-975 https://doi.org/10.1111/jpy.12895
Issue Date	2019-10
Doc URL	https://hdl.handle.net/2115/79385
Rights	This is the peer reviewed version of the following article: Heat-stress Memory is Responsible for Acquired Thermotolerance in <i>Bangia fuscopurpurea</i> , which has been published in final form at https://doi.org/10.1111/jpy.12895 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	journal article
File Information	Kishimoto et al FINAL huscu.pdf



Manuscript ID: JPY-19-058-L

HEAT-STRESS MEMORY IS RESPONSIBLE FOR ACQUIRED
THERMOTOLERANCE IN *BANGIA FUSCOPURPUREA*¹

Ikuya Kishimoto, Inori Ariga*

Graduate School of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho,
Hakodate 041-8611, Japan

Yutaka Itabashi[#] and Koji Mikami²

Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato-cho, Hakodate
041-8611, Japan

Running title: Heat-stress memory in *Bangia*

¹Received 27 March 2019, Accepted 18 June 2019.

²Corresponding author: Koji Mikami, Faculty of Fisheries Sciences, Hokkaido
University, 3-1-1 Minato-cho, Hakodate 041-8611, Japan. TEL/FAX: +81-138-40-899.
E-mail: komikami@fish.hokudai.ac.jp

*Present address: Rokkatei Confectionery Co. Ltd., 3-19 Kita 1, Nishi 24, Obihiro 080-
2496, Japan

[#] Present address: Department of Chemistry, Faculty of Science, Rikkyo University, 3-
34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan

The environmental stresses that sessile organisms experience usually fluctuate dramatically and are often recurrent. Terrestrial plants can acquire memory of exposure to sub-lethal heat stress to acquire thermotolerance and survive subsequent lethal high-temperature stress; however, little is known concerning whether seaweeds acquire thermotolerance via heat-stress memory. We have demonstrated that the red seaweed *Bangia fuscopurpurea* can indeed acquire memory of sub-lethal high-temperature stress, resulting in the acquisition of thermotolerance that protects against subsequent lethal high-temperature stress. Moreover, the maintenance of heat-stress memory was associated with a slight increase in the saturation level of membrane fatty acids. This suggests that the modification of membrane fluidity via changes in membrane fatty acid composition is involved in the establishment and maintenance of heat-stress memory in *B. fuscopurpurea*. These findings provide insights into the physiological survival and growth strategies of sessile red seaweeds to cope with recurrent changes in environmental conditions.

Key index words: *Bangia fuscopurpurea*, Fatty acid, Heat stress, Rhodophyta, Stress memory, Thermotolerance

Temperature fluctuation represents one of the most prominent environmental stresses for sessile plants (Cramer et al. 2011, Kumar et al. 2012, Pandey et al. 2017). Increases in ambient temperature can cause heat stress and affect the survival, growth, development and reproduction of plants (Barnabás et al. 2008, Zinn et al. 2010; Hartfield and Prueger 2015). In the natural environment, heat stress is often chronic or

recurring. Although the molecular mechanisms that regulate the response to a single acute heat stress have been analyzed in detail, the response to recurring stress is only beginning to be elucidated. In plants, priming by exposure to a sub-lethal heat stress promotes the acquisition of thermotolerance, which allows plants to survive a subsequent and temporally distinct second potentially lethal heat stress (Bruce et al., 2007, Bäurle 2016, Sanyal et al., 2018). These findings indicate that plants can maintain the primed state to respond strongly to a recurring stress, a phenomenon that is termed heat-stress memory (Bruce et al. 2007, Bäurle 2016, Crisp et al. 2016, Sanyal et al. 2018).

Bangiales is an industrially and economically important marine resource and a large-scale mariculture system has been established for *Pyropia yezoensis* in Japan and *P. haitanensis* and *Bangia fuscopurpurea* in China (Mumford and Miura 1988, Wang et al. 2008, Blouin et al. 2011). Because an increase in temperature affects the growth and productivity of maricultured seaweeds, much effort has been invested to establish and analyze heat stress-tolerant strains of Bangiales (Yan et al. 2010, Ding et al. 2016, Lee et al 2017, Wang et al 2018a). In addition, transcriptome analyses have generated information concerning heat-inducible and -repressible genes (Choi et al. 2013, Im et al. 2015, Sun et al. 2015, Cao et al. 2017, Wang et al. 2018a, 2018b), which has indicated that Bangiales can respond to heat stress. However, knowledge concerning the regulatory mechanisms underlying the heat-stress response, tolerance and memory in Bangiales and other seaweeds is lacking.

The effects of heat stress on survival of the red seaweed *B. fuscopurpurea* have been investigated using strains harvested from various locations (Sommerfeld and Nichols 1973, Notoya and Iijima 2003, Wang et al. 2008, Mikami and Kishimoto 2018).

Sub-lethal heat stress promotes the asexual life cycle, which releases large numbers of asexual spores to increase the number of clones (Notoya and Iijima 2003, Wang et al. 2008, Mikami and Kishimoto 2018). Because these findings suggest that *B. fuscopurpurea* responds to heat stress like other Bangiales, we addressed whether *B. fuscopurpurea* can acquire heat-stress tolerance via acquiring heat-stress memory.

We first examined whether *B. fuscopurpurea* can acquire heat-stress tolerance (see Appendix S1 for details of experimental procedure). After growth at 15°C, *B. fuscopurpurea* was primed at the sub-lethal high temperature 28°C for 1 week and was further incubated at 30, 32 or 34°C for 1 week (Fig. 1A). As reported previously (Mikami and Kishimoto 2018), the direct transfer of unprimed control samples from 15°C to 32 or 34°C for 1 week strongly reduced their viability (Figs. 1B, S1). By contrast, *B. fuscopurpurea* samples that were primed at 28°C for 1 week could survive exposure to lethal conditions for 1 week (Fig. 1B). In this case, priming for 3 days was enough to acquire the heat-stress tolerance (Fig. S2). These findings indicate that the priming of *B. fuscopurpurea* by sub-lethal high temperature promotes the acquisition of heat-stress tolerance, enabling survival of subsequent exposure to lethal high temperature as a short-term response.

To assess whether this priming in *B. fuscopurpurea* leads to heat-stress memory, we designed experiments that included the insertion of a recovery phase by returning the samples to non-stress 15°C conditions between the priming phase and triggering phase as a secondary heat-stress exposure to a lethal temperature (Fig. 1C). Importantly, the viability of samples exposed to a 1-day recovery period was equal to that of primed samples without a recovery phase (Fig 1D). Thus, we concluded that *B. fuscopurpurea* can acquire heat-stress memory and maintain it after being returned to

non-stress conditions. However, increasing the duration of the recovery phase by 1 to 5 days gradually decreased viability and 6- and 7-day recovery periods abolished heat-stress memory (Fig. 1D), indicating that increasing the length of the recovery phase reduces heat-stress memory. Thus, we conclude that heat-stress memory is detectable for only 5 days after transfer to normal growth conditions.

There is a close relationship between membrane rigidification via a reduction in fluidity and heat-stress tolerance in Poikilothermic organisms, where membrane fatty acid composition influences the fluidity of membranes (Carratù et al. 1996, Horváth et al. 1998, Murakami et al. 2000, Sangwan et al. 2002, Königshofer et al. 2008, Leach and Cowen 2014). We therefore addressed whether the acquisition of the heat-stress memory in *B. fuscopurpurea* is related to changes in the composition of membrane fatty acids (Appendix S1). Firstly, we examined whether the membrane fatty acid composition was modulated by heat stress. When *B. fuscopurpurea* was cultured at 15°C followed by 32°C for a week, the relative amounts of saturated fatty acids and monoenes increased compared to those in the 15°C controls, whereas the relative amounts of polyenes decreased (compare the white and black bars in Fig. 2A–C). Thus, heat stress induces an increased saturation level of membrane fatty acids in *B. fuscopurpurea*, which likely results in increased membrane rigidification. Secondly, we investigated the effects of priming by exposure to sub-lethal high temperatures on the composition of membrane fatty acids. Cells from samples induced to acquire heat-stress memory exhibited intermediate levels of fatty acid saturation and unsaturation compared to those in the control and non-primed 32°C-shifted samples (Fig. 2A–C; gray bars at 0 days). Thus, the modification of fatty acid composition by priming appears to be important for the acquisition and maintenance of heat-stress tolerance.

121 Lastly, we investigated the relationship between the loss of heat-stress memory (Fig.
122 1D) and changes in fatty acid composition. The relative levels of monounsaturated fatty
123 acids (monoenes) and polyunsaturated fatty acids (polyenes) in cells induced to acquire
124 heat-stress memory gradually became more similar to those observed in non-primed
125 controls (black bars), depending on the length of the recovery phase (Fig. 2B and C).
126 Thus, the loss of heat-stress memory, which resulted in more dead cells at 32°C was
127 closely associated with changes in membrane fatty acid composition. These findings
128 indicate that priming of *B. fuscopurpurea* can lead to the acquisition of heat-stress
129 memory and recovery treatments to its loss, and that both states are associated with a
130 specific membrane fatty acid composition. We therefore conclude that the maintenance
131 of heat-stress memory requires a specific membrane physical state and an appropriate
132 level of membrane fluidity.

133 Our data demonstrate that *B. fuscopurpurea* can acquire heat-stress memory as
134 a heat-stress tolerance strategy (Figs. 1 and 2), confirming that this phenomenon is
135 widely conserved among seaweeds and terrestrial plants. This represents an important
136 strategy that enables plants to adapt and survive under recurring or chronic heat stress in
137 natural environments. Although heat-stress memory involves the maintenance of a heat-
138 stress tolerance-specific composition of membrane fatty acids (Fig. 2), the regulatory
139 mechanisms that alter membrane fatty acid composition by priming are largely
140 unknown. The establishment and maintenance of heat-stress memory, which plays a
141 central role in the maintenance of thermotolerance in response to recurring heat stress in
142 terrestrial plants, is regulated epigenetically (Crisp et al. 2016, Lämke and Bäurle 2017,
143 Friedrich et al. 2019). For example, a relationship exists between the maintenance of
144 heat-stress memory and the accumulation of histone methylation markers H3K4me3

and H3K4me2 within chromatin at the promoters of genes that are highly expressed for several days during heat-stress memory (Kusch et al. 2014, Lämke et al. 2016; Liu et al. 2018). Thus, heat-stress memory is maintained by structural chromatin changes caused by H3 methylation during the memory phase. Because little is known concerning epigenetic regulation in response to abiotic stress in any Bangiales species, it is necessary to investigate whether genes associated with memory and their epigenetic control potentially regulate membrane fatty acid composition and maintain heat-stress memory in *B. fuscopurpurea*.

This work was supported in part by the KAKENHI grant (15H04539) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Barnabás, B., Jäger, K. & Fehér, A. 2008. The effect of drought and heat stress on reproductive process in cereals. *Plant Cell Environ.* 31:11–38.

Bäurle, I. 2016. Plant Heat Adaptation: priming in response to heat stress. *F1000Res.* 5:694. doi: 10.12688/f1000research.7526.1.

Blouin, N. A., Brodie, J. A., Grossman, A. C., Xu, P. & Brawley, S. H. 2011. *Porphyra*: a marine crop shaped by stress. *Trends Plant Sci.* 16:29–37.

Bruce, T. J. A., Matthes, M. C., Napier, J. A., Pickett, J. A. 2007. Stressful “memories” of plants: Evidence and possible mechanisms. *Plant Sci.* 173:603–8.

Cao, M., Wang, D., Mao, Y., Kong, F., Bi, G., Xing, Q. & Weng, Z. 2017. Integrating transcriptomics and metabolomics to characterize the regulation of EPA biosynthesis in response to Cold stress in seaweed *Bangia fuscopurpurea*. *PLoS One* 12:e0186986. doi: 10.1371/journal.pone.0186986.

169 Carratù, L., Franceschelli, S., Pardini, C. L., Kobayashi, G. S., Horvath, I., Vigh, L. &
 170 Maresca, B. 1996. Membrane lipid perturbation modifies the set point of the
 171 temperature of heat shock response in yeast. *Proc. Natl. Acad. Sci. U.S.A.* 93:3870–
 172 5.
 173 Choi, S., Hwang, M. S., Im, S., Kim, N., Jeong, W. -J., Park, E. -J & Gong, Y. -G.
 174 2013. Transcriptome sequencing and comparative analysis of the gametophyte
 175 thalli of *Pyropia tenera* under normal and high temperature conditions. *J. Appl.*
 176 *Phycol.* 25:1237–46.
 177 Cramer, G. R., Urano, K., Delrot, S., Pezzotti, M., Shinozaki, K. 2011. Effects of abiotic
 178 stress on plants: a systems biology perspective. *BMC Plant Biol.* 11:163.
 179 doi:10.1186/1471-2229-11-163
 180 Crisp, P. A., Ganguly, D., Eichten, S. R., Borevitz, J. O. & Pogson, B. J. 2016.
 181 Reconsidering plant memory: Intersections between stress recovery, RNA turnover,
 182 and epigenetics. *Sci. Adv.* 2:e1501340. doi: 10.1126/sciadv.1501340.
 183 Ding, H., Zhang, B. & Yan, X. 2016. Isolation and characterization of a heat-resistant
 184 strain with high yield of *Pyropia yezoensis* Ueda (Bangiales, Rhodophyta).
 185 *Aquacult. Fish.* 1:24–33.
 186 Friedrich, T., Faivre, L., Bäurle, I. & Schubert, D. 2019. Chromatin-based mechanisms
 187 of temperature memory in plants. *Plant Cell Environ.* 42:762–70.
 188 Hatfield, J. L. & Prueger, J. H. 2015. Temperature extremes: Effect on plant growth and
 189 development. *Weather Climate Extremes* 10:4–10.
 190 Horváth, I., Glatz, A., Varvasovszki, V., Török, Z., Páli, T., Balogh, G., Kovács, E.,
 191 Nádasdi, L., Benkő, S., Joó, F. & Vigh, L. 1998. Membrane physical state controls

the signaling mechanism of the heat shock response in *Synechocystis* PCC 6803: identification of hsp17 as a "fluidity gene". *Proc. Natl. Acad. Sci. U.S.A.* 95:3513–8.

Im, S., Choi, S., Hwang, M. S., Park, E. -J. & Choi, D. -W. 2015. *De novo* assembly of transcriptome from the gametophyte of the marine red algae *Pyropia seriata* and identification of abiotic stress response genes. *J. Appl. Phycol.* 27:1343–53.

Königshofer, H., Tromballa, H. W. & Löppert, H. G. 2008. Early events in signalling high-temperature stress in tobacco BY2 cells involve alterations in membrane fluidity and enhanced hydrogen peroxide production. *Plant Cell Environ.* 31:1771–80.

Kumar, A. A., Mishra, P., Kumari, K. & Panigrahi, K. C. 2012. Environmental stress influencing plant development and flowering. *Front. Biosci. (Schol Ed)*. 4:1315–24.

Kusch, T., Mei, A. & Nguyen, C. 2014. Histone H3 lysine 4 trimethylation regulates cotranscriptional H2A variant exchange by Tip60 complexes to maximize gene expression. *Proc. Natl. Acad. Sci. U.S.A.* 111:4850–5.

Lämke, J. & Bäurle, I. 2017. Epigenetic and chromatin-based mechanisms in environmental stress adaptation and stress memory in plants. *Genome Biol.* 18:124. doi: 10.1186/s13059-017-1263-6.

Lämke, J., Brzezinka, K., Altmann, S. & Bäurle, I. 2016. A hit-and-run heat shock factor governs sustained histone methylation and transcriptional stress memory. *EMBO J.* 35:162–75.

Leach, M. D. & Cowen, L.E. 2014. Membrane fluidity and temperature sensing are coupled via circuitry comprised of Ole1, Rsp5, and Hsf1 in *Candida albicans*. *Eukaryot. Cell* 13:1077–84.

215 Lee, H. -J., Park, E. -J. & Choi, J. -I. 2018. Isolation, Morphological characteristics and
 216 proteomic profile analysis of thermo-tolerant *Pyropia yezoensis* mutant in response
 217 to high-temperature stress. *Ocean Sci. J.* doi: org/10.1007/s12601-018-0060-9.

218 Liu, H. C., Lämke, J., Lin, S. Y., Hung, M. J., Liu, K. M., Charng, Y. Y. & Bäurle, I.
 219 2018. Distinct heat shock factors and chromatin modifications mediate the organ-
 220 autonomous transcriptional memory of heat stress. *Plant J.* 95:401–13.

221 Mikami, K., Ito, M., Taya, K., Kishimoto, I., Kobayashi, T., Itabashi, Y. & Tanaka, R.
 222 2018. Parthenosporophytes of the brown alga *Ectocarpus siliculosus* exhibit sex-
 223 dependent differences in thermotolerance as well as fatty acid and sterol
 224 composition. *Mar. Environ. Res.* 137:188–95.

225 Mikami, K. & Kishimoto, I. 2018. Temperature promoting the asexual life cycle
 226 program in *Bangia fuscopurpurea* (Bangiales, Rhodophyta) from Esashi in the
 227 Hokkaido Island, Japan. *Algal Resources* 11:25–32.

228 Mumford, T. F. & Miura, A. 1988. *Porphyra* as food: cultivation and economics. In
 229 Lembi, C.A. & Waaland, J.R. [Eds.] *Algae and Human Affairs*. Cambridge
 230 University Press, London, UK, pp. 87–117.

231 Murakami, Y., Tsuyama, M., Kobayashi, Y., Kodama, H. & Iba, K. 2000. Trienoic fatty
 232 acids and plant tolerance of high temperature. *Science* 28:476–9.

233 Notoya, M. & Iijima, N. 2003. Life history and sexuality of archeospore and apogamy
 234 of *Bangia atropurpurea* (Roth) Lyngbye (Bangiales, Rhodophyta) from Fukaura and
 235 Enoshima. *Japan. Fish. Sci.* 69:799–805,

236 Pandey, P., Irulappan, V., Bagavathiannan, M. V. & Senthil-Kumar, M. 2017. Impact of
 237 combined abiotic and biotic stresses on plant growth and avenues for crop
 238 improvement by exploiting physio-morphological traits. *Front. Plant Sci.* 8:537.

doi: 10.3389/fpls.2017.00537

- Sommerfeld, M. A. & Nichols, H. W. 1973. The life cycle of *Bangia fuscopurpurea* in culture. I. Effects of temperature and photoperiod on the morphology and reproduction of the *Bangia* phase. *J. Phycol.* 9:205–10.
- Sangwan, V., Örvar, B.L., Beyerly, J., Hirt, H. & Dhindsa, R. S. 2002. Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. *Plant J.* 31:629–38.
- Sanyal, R. P., Misra, H. S. & Saini, A. 2018. Heat-stress priming and alternative splicing-linked memory. *J. Exp. Bot.* 69:2431–4.
- Sun, P., Mao, Y., Li, G., Cao, M., Kong, F., Wang, L. & Bi, G. 2015. Comparative transcriptome profiling of *Pyropia yezoensis* (Ueda) M.S. Hwang & H.G. Choi in response to temperature stresses. *BMC Genomics* 16:463. doi: org/10.1186/s12864-015-1586-1
- Wang, W., Teng, F., Lin, Y., Ji, D., Xu, Y., Chen, C. & Xie, C. 2018a. Transcriptomic study to understand thermal adaptation in a high temperature-tolerant strain of *Pyropia haitanensis*. *PLoS One* 13:e0195842. doi: 10.1371/journal.pone.0195842.
- Wang, W., Chang, J., Zheng, H., Ji, D., Xu, Y., Chen C. & Xie, C. 2018b. Full-length transcriptome sequences obtained by a combination of sequencing platforms applied to heat shock proteins and polyunsaturated fatty acids biosynthesis in *Pyropia haitanensis*. *J. Appl. Phycol.* doi: org/10.1007/s10811-018-1624-x
- Wang, W. -J., Zhu, J. -Y., Xu, P., Xu, J. R., Lin, X. Z., Huang, C. K., Song, W. L., Peng, G., Wang, G. C. 2008. Characterization of the life history of *Bangia fuscopurpurea* (Bangiaceae, Rhodophyta) in connection with its cultivation in China. *Aquaculture* 278:101–9.

- Yan, X. -H., Lv, F., Liu, C. -J., Zheng, Y. -F. 2010. Selection and characterization of a high-temperature tolerant strain of *Porphyra haitanensis* Chang et Zheng (Bangiales, Rhodophyta). *J. Appl. Phycol.* 22:511–6.
- Zinn, K. E., Tunc-Ozdemir, M. & Harper, J. F. 2010. Temperature stress and plant sexual reproduction: uncovering the weakest links. *J. Exp. Bot.* 61:1959–68.

Supporting Information

Fig. S1. Priming-induced thermotolerance in *Bangia fuscopurpurea*. Viability of *B. fuscopurpurea* under lethal high temperatures (32 and 34°C) following prior priming at 28°C were examined. For primed experiments (right half), samples grown at 15°C were primed at 28°C for a week, then triggering was performed at 30, 32 or 34°C for a further week. For non-primed control experiments (left half), samples grown at 15°C were directly transferred to 30, 32 or 34°C and incubated for a further week. Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA as described in Mikami et al. (2018).

Fig. S2. Time course analysis for acquisition of heat-stress tolerance during the priming phase in *Bangia fuscopurpurea*. Changes in viability associated with the duration of the priming phase were examined. Algal samples grown at 15°C were separately primed at 28°C for one to seven days, then triggering was performed for all of samples at 32°C for a further week. Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA as described in Mikami et al. (2018).

Appendix S1. Details of experimental procedure.

Legends to Figures

Fig. 1. Thermotolerance is dependent on heat-stress memory in *Bangia fuscipurpurea*. (A) Schematic representation of the experimental design to assess the ability of *B. fuscipurpurea* to acquire thermotolerance. After growth at 15°C and priming at 28°C for a week, triggering was performed at 30, 32 or 34°C for a further week. The upper panel indicates the control treatment without priming. (B) Priming-induced thermotolerance; *B. fuscipurpurea* survived lethal high temperatures (32 or 34°C) following prior priming under sub-lethal conditions. (C) Schematic representation of the experimental design to assess the ability of *B. fuscipurpurea* to acquire heat-stress memory. Varying durations of the recovery phase at 15°C were inserted between priming and triggering phases. (D) Changes in viability associated with the duration of the recovery phase. Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA as described in Mikami et al. (2018).

Fig. 2. Changes in membrane fatty acid composition are associated with the acquisition and loss of heat-stress memory in *B. fuscipurpurea*. Changes in the relative amounts of saturated fatty acids (Saturates, A), monounsaturated fatty acids (Monoenes, B) and polyunsaturated fatty acids (Polyenes, C) were analyzed in control samples (white bars) or those triggered either with a lethal high temperature (32°C) without priming (black bars) or with priming and recovery treatments (gray bars). Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters

334 denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA
335 as described in Mikami et al. (2018).




Fig. 1




Fig. 2




Fig. S1 Priming-induced thermotolerance in *Bangia fuscopurpurea*.

Viability of *B. fuscopurpurea* under lethal high temperatures (32 and 34°C) following prior priming at 28°C were examined. For primed experiments (right half), samples grown at 15°C were primed at 28°C for a week, then triggering was performed at 30, 32 or 34°C for a further week. For non-primed control experiments (left half), samples grown at 15°C were directly transferred to 30, 32 or 34°C and incubated for a further week. Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA as described in Mikami et al. (2018).




Fig. S2 Time course analysis for acquisition of heat-stress tolerance during the priming phase in *Bangia fuscopurpurea*.

Changes in viability associated with the duration of the priming phase were examined. Algal samples grown at 15°C were separately primed at 28°C for one to seven days, then triggering was performed for all of samples at 32°C for a further week. Mean values $\pm$ SD per 0.1 g sample fresh weight were calculated from three independent experiments. Letters denote statistically significant differences ($p < 0.05$) as determined by one-way ANOVA as described in Mikami et al. (2018).

Appendix S1

Detailed Materials and Methods

Algal material and heat-stress treatment

Gametophytic thalli of the marine red seaweed *Bangia fuscopurpurea* were collected at Esashi, Hokkaido, Japan on 14 May 2010, and a clean single thallus of unknown sex was maintained in the laboratory via the asexual propagation producing clones as an experimental line. To prepare control and starting samples, each 0.1 g (fresh weight) of *B. fuscopurpurea* thalli was cultured in dishes (Azunoru dish $\phi 90 \times 20$ mm height, As One) containing 50 mL esterized artificial seawater at 15°C under $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ irradiation (see Mikami and Kishimoto 2018, for details). Priming of *B. fuscopurpurea* thalli was performed by cultivating 15°C-grown samples at 28°C as a sub-lethal high temperature for 1 week. To analyze thermotolerance acquisition, primed samples were exposed to 32 or 34°C as lethal triggering conditions for 1 week. To assess the acquisition of heat-stress memory, recovery phases consisting of cultivation at 15°C for one to seven days were inserted between phases of priming at 28°C and triggering at 32 or 34°C. To determine the duration required for establishment of heat-stress tolerance, samples were separately primed at 28°C for one to seven days and then all samples were transferred to 32°C and incubated for seven days. Every treatment was repeated three times. The viability of all samples was tested, and the membrane fatty acid composition was analyzed as follows.

Viability test

As described previously (Mikami and Kishimoto, 2018), 0.1 g (fresh weight) of *B.*

fuscopurpurea thalli exposed to priming, recovery and triggering conditions as described above, were visualized daily by staining with ESL medium containing 0.01% erythrosine (Wako Pure Chemical Industries, Japan). After staining for 5 min at room temperature, thalli were gently rinsed with ESL medium to remove excess erythrosine and were mounted on a slide with ESL medium. Thalli were observed and photographed using an Olympus IX73 light microscope equipped with an Olympus DP22 camera. Cells stained by the dye were defined as dead cells. Viability was calculated from the number of living and dead cells obtained using micrographs. Analysis of every treatment was repeated three times and statistical analysis was performed as described in Mikami et al. (2018).

Extraction of lipids and preparation of fatty acid methyl esters

Fresh samples of *Bangia fuscopurpurea* were immersed in boiling water for 3 min to deactivate lipid hydrolytic enzymes and were then freeze-dried and homogenized using a grinder. Lipids were extracted via the Bligh-Dyer method (Bligh and Dyer, 1959) with some modifications. In brief, 0.1 g powdered algal sample was homogenized in a mixture of 2 mL MeOH and 1 mL CHCl₃, 1 mL CHCl₃ was added and the sample was vigorously mixed. After filtration, 2 mL CHCl₃ was added to the algal residue, which was then homogenized and filtered. All the obtained solvent fractions were combined, 1 mL water was added, and the sample was mixed. The organic solvent layer was recovered by centrifugation and evaporated to dryness under reduced pressure. The resultant lipids were resuspended to a known concentration with CHCl₃/MeOH (2:1, v/v) and stored at -30°C until use.

Lipids were converted to fatty acid methyl esters by heating at 90°C for 1 h in a

solution of 1–2% H₂SO₄/MeOH (Christie and Han, 2010) and were then purified in a Pasteur pipette filled with silica gel using hexane-Et₂O (95:5, v/v) as the mobile phase. The purified fatty acid methyl esters were confirmed on a silica gel 60 F₂₅₄ aluminum TLC sheet (Merck, Darmstadt, Germany) using hexane-Et₂O (90:10, v/v) as the developing solvent (Christie and Han, 2010), and were then dissolved in 5 mg/mL hexane for gas chromatography.

GC analysis of fatty acid methyl esters

Fatty acid methyl esters were analyzed using a Shimadzu GC-14A gas chromatograph (Shimadzu, Kyoto, Japan) equipped with an Omegawax 320 column (30 m × 0.32 mm i.d., Supelco, PA, USA). Helium was used as the carrier gas at a constant flow-rate of 2.2 mL/min. The split ratio was 1:25. The column temperature was maintained at 200°C isothermally. The injector and flame-ionization detector (FID) temperatures were set at 230°C and 240°C, respectively. Peaks were monitored on a Chromatopac C-R6A and identified by comparison with the retention data of some authentic standards (GLC-462, Nu-Chek Prep., Elysian, MN), and those of known fatty acids from marine algae (Takagi et al. 1985). Analysis of samples for every treatment was repeated three times and statistical analysis was performed as described in Mikami et al. (2018).

References

- Bligh, E. G. & Dyer, W. J. 1959. A rapid method of lipid extraction and purification. *Can. J. Biochem. Physiol.* 37:911–7.
- Christie, W. W. & Han, X. 2010. Preparation of derivatives of fatty acids. *In Lipid*

Analysis, 4th ed. The Oily Press, Bridgwater, England, pp. 145–58.

Mikami, K. & Kishimoto, I. 2018. Temperature promoting the asexual life cycle program in *Bangia fuscopurpurea* (Bangiales, Rhodophyta) from Esashi in the Hokkaido Island, Japan. *Algal Resources* 11:25–32.

Takagi, T., Asahi, M. & Itabashi, Y. 1985. Fatty acid composition of twelve algae from Japanese waters. *J. Jpn. Oil Chem. Soc.* 34:1008–12.