



Title	Cp*Co-III/Chiral Carboxylic Acid-Catalyzed Enantioselective 1,4-Addition Reactions of Indoles to Maleimides
Author(s)	Kurihara, Takumaru; Kojima, Masahiro; Yoshino, Tatsuhiko et al.
Citation	Asian journal of organic chemistry, 9(3), 368-371 https://doi.org/10.1002/ajoc.201900565
Issue Date	2019-11-14
Doc URL	https://hdl.handle.net/2115/79746
Rights	This is the peer-reviewed version of the following article: T.Kurihara et al. Cp*Co-III/Chiral Carboxylic Acid - Catalyzed Enantioselective 1,4 - Addition Reactions of Indoles to Maleimides. Asian Journal of Organic Chemistry, First published 14 Nov. 2019, which has been published in final form at https://doi.org/10.1002/ajoc.201900565 article using the DOI]. This article may be used for non-commercial purposes in accordance with Wiley-VCH Terms and Conditions for Self-Archiving.
Type	journal article
File Information	WoS_91810_Yoshino.pdf



Cp*Co(III)/Chiral Carboxylic Acid-Catalyzed Enantioselective 1,4-Addition Reactions of Indoles to Maleimides

Takumaru Kurihara, Masahiro Kojima, Tatsuhiko Yoshino*, and Shigeki Matsunaga*


[a] T. Kurihara, M. Kojima, T. Yoshino, Prof. Dr. S. Matsunaga
Faculty of Pharmaceutical Sciences
Hokkaido University
Kira-ku, Sapporo 060-0812, Japan
E-mail: tyoshino@pharm.hokudai.ac.jp;
smatsuna@pharm.hokudai.ac.jp

Supporting information for this article is given via a link at the end of the document.

Abstract: We report enantioselective 1,4-addition reactions of indoles to maleimides via a C2-selective C–H activation catalyzed by an achiral Cp*Co(III)/chiral carboxylic acid system. In these reactions, a BINOL-derived chiral carboxylic acid enables the enantioselective reactions as the sole chiral source.

C–H functionalization reactions have emerged as a promising strategy for the formation of C–C or C–hetero atom bonds, and can lead to atom-^[1] and step-economical^[2] organic syntheses.^[3] Although cationic complexes of group 9 metals (Co, Rh, Ir) with a pentamethylcyclopentadienyl (Cp*) or related (Cp*) ligands can catalyze C–H functionalization reactions quite efficiently, catalytic stereocontrol is difficult to accomplish due to the lack of vacant coordination sites for additional chiral ligands.^[4] In this context, the development of well-designed chiral Cp* ligands that require no additional coordination site and elegantly enable asymmetric C–H functionalization reactions was pioneered by Cramer and co-workers^[5] followed by other research groups.^[6–9]

More recently, combinations of achiral Cp*M(III) complexes and chiral carboxylic acids (CCAs) have been exploited for enabling asymmetric C–H functionalization reactions.^[10] In Cp*M(III)-catalyzed C–H functionalization reactions, a carboxylic acid or its salt can accelerate the C–H bond activation^[11] and/or protodemetalation step;^[12] thus, the use of CCAs could potentially induce enantioselectivity in these two steps. Enantioselective C–H activation using a CCA,^[13] in which two prochiral C–H bonds of a substrate are distinguished by a CCA and one of them is cleaved selectively, has been explored by Chang,^[14] our group,^[15] and Shi^[16] (Scheme 1: Type (a)). On the other hand, enantioinduction by selective protodemetalation has been reported by Ackermann,^[17] who achieved Cp*Co(III)/CCA-catalyzed asymmetric C–H alkylations using alkenes via a selective protodemetalation (Scheme 1: Type (b)). In this case, the reversible insertion of an alkene after C–H activation generates two enantiotopic intermediates, one of which is selectively protonated by the CCA, thus leading to enantioselective alkylation.^[18] Compared to enantioselective C–H activations, enantioselective alkylations via the reversible insertion/selective protonation mechanism have received less attention, and the reactions to which it has been applied are relatively limited.^[17,19]




Scheme 1. Enantioselective C–H functionalization reactions catalyzed by achiral Cp*M(III) (M = Co, Rh, Ir) and chiral carboxylic acids (CCAs). DG = directing group.

Herein, we report our investigations into enantioselective Cp*Co(III)/CCA-catalyzed 1,4-addition reactions of indoles to maleimides via C–H activation (Scheme 1: This work). Directing group-assisted C–H activation of indoles enables site-selective C–C bond formations at the C2-position, overriding the inherent nucleophilic reactivity at the C3-position with polar electrophiles.^[20] Li, Ackermann, and co-workers have reported the racemic variants of this reaction, in which the addition of a catalytic amount of an achiral carboxylate dramatically enhanced the reactivity.^[21] We hypothesized that the rate enhancement could be due to the acceleration of the final protodemetalation step, and that the use of a CCA could enable enantioselective addition reactions.

We selected pyrimidyl-protected indole **1a** and maleimide **2a** as model substrates for the screening of several types of CCAs


Table 1. Screening CCAs and optimizing reaction conditions^[a].

entry	Co cat.	additive	CCA	% yield ^[b]	er
1	Cp*Co(CO) ₂	AgSbF ₆	4a	>95	75:25
2	Cp*Co(CO) ₂	AgSbF ₆	4b	>95	61:39
3	Cp*Co(CO) ₂	AgSbF ₆	4c	>95	69:31
4	Cp*Co(CO) ₂	AgSbF ₆	4d	63	49:51
5	Cp*Co(CO) ₂	AgSbF ₆	4e ^[c]	>95	54:46
6	Cp*Co(CO) ₂	AgSbF ₆	4f	>95	64:36
7	Cp*Co(CO) ₂	AgSbF ₆	4g	77	50:50
8 ^[d]	Cp*Co(CO) ₂	AgSbF ₆	4a	65	80:20
9 ^[d]	[Cp*Co(CH ₃ CN) ₃](SbF ₆) ₂	MS13X	4a	94	81:19



[a] Unless otherwise noted, reactions were carried out using **1a** (0.05 mmol), **2a** (0.10 mmol), Cp*Co cat. (0.0025 mmol, 5 mol %), CCA (0.005 mmol, 10 mol %), tBuOK (0.006 mmol, 12 mol %), and AgSbF₆ (0.005 mmol, 10 mol %) or MS13X (10 mg, 200 mg/mmol of **1a**) in TFE/DCM (4:1) (0.25 mL) at 35 °C for 20 h. [b] Determined by ¹H NMR analysis of the crude reaction mixture. [c] 5 mol %. [d] 10 °C, 48 h. TFE = 2,2,2-trifluoroethanol, DCM = dichloromethane.

(Table 1). Our initial trial using BINOL-derived CCA **4a** afforded the product **3aa** in excellent yield and in an enantiomeric ratio (er) of 75:25 (entry 1), supporting our hypothesis (*vide supra*). We next examined the effect of different substituents on the binaphthyl backbone on the reaction. Using a CCA with a sterically less hindered methoxy group at the 2' position (**4b**), as well as one carrying a 2-naphthyl group (**4c**) at the same location, resulted in decreased selectivity (entries 2 and 3). Unfortunately, the introduction of a substituent at the *ortho*-position of the carboxylic acid moiety (**4d**) decreased both the yield and enantioselectivity (entry 4), although a similar modification was effective in our previous work.^[14a] Dicarboxylic acid **4e**^[22] and two CCAs with non-binaphthyl backbones (**4f**, **4g**)^[15b,17b] that were used in previously reported Cp*Co(III)/CCA-catalyzed enantioselective C–H functionalization reactions did not afford better results than **4a** (entries 5–7). Based on these results, we selected **4a** as the optimal CCA and turned our attention to optimizing the reaction conditions. Carrying out the reaction at lower temperature for a longer time (10 °C, 48 h) improved the enantioselectivity to 80:20 er, but diminished the yield (entry 8). Changing the cobalt catalyst to [Cp*Co(CH₃CN)₃](SbF₆)₂/MS13X



Scheme 2. Scope and limitations of the enantioselective 1,4-addition of pyrimidylindoles **1** and maleimides **2**. Unless otherwise noted, reactions were carried out using **1** (0.20 mmol), **2** (0.40 mmol), [Cp*Co(CH₃CN)₃](SbF₆)₂ (0.01 mmol, 5 mol %), **4a** (0.02 mmol, 10 mol %), tBuOK (0.024 mmol, 12 mol %), and MS13X (40 mg, 200 mg/mmol of **1**) in TFE/DCM (4:1) (1.0 mL) at 10 °C for 72 h. [a] 20 °C.

improved the reactivity without any loss of selectivity (entry 9, 94%, 81:19 er). The MS13X molecular sieves would be able to adsorb the acetonitriles coordinating to the Co center, effectively generating an active cationic Co species.

Under the optimized reaction conditions (Table 1, entry 9), the scope and limitations of the enantioselective 1,4-addition of the pyrimidylindoles **1**^[23] to maleimides **2**^[24] were investigated (Scheme 2). Indoles bearing an electron-withdrawing group or electron-donating group at the 4-, 5-, and 6-position (**1b–1i**) provided the corresponding products (**3ba–3ia**) in 61–99% yields with 77:23–82:18 er. The absolute configuration of the major isomer of **3ea** was determined to be (S) by single-crystal X-ray diffraction analysis. Although **1j**, which contains a methyl group at the 7-position, was converted to the corresponding product (**3ja**) in high yield, the enantioselectivity was low. This was attributed to steric repulsion between the methyl group and the pyrimidyl directing group, which would distort the conformation of the metallacycle intermediate, leading to the lower selectivity. In addition to *N*-methylmaleimide **2a**, *N*-ethylmaleimide **2b** afforded the corresponding product (**3ab**) in 80% yield with 81:19 er, while *N*-benzylmaleimide **2c** exhibited lower reactivity (29% yield, 77:23 er).

A plausible catalytic cycle for this enantioselective 1,4-addition is shown in Figure 1.^[21,25] Initially, cationic $\text{Cp}^*\text{Co(III)}$ species **I** would be generated from $[\text{Cp}^*\text{Co}(\text{CH}_3\text{CN})_3](\text{SbF}_6)_2$, MS13X, **4a**, and $t\text{BuOK}$. C–H bond cleavage of indole **1** would then occur via a concerted metallation-deprotonation (CMD) mechanism^[11,26] to form a metallacycle intermediate **II**. After the coordination of maleimide **2** to **II**, which would generate intermediate **III**, the insertion of the C–C double bond would afford intermediate **IV** or **IV'**. Although this insertion step seemingly determines the stereochemistry, intermediate **III** does not contain an extra coordination site that can accommodate CCA **4a** or its anion. Therefore, it is more likely that the reaction proceeds via a reversible insertion/selective protodemetalation mechanism.^[17,19] Specifically, the insertion of the C–C double bond is expected to be reversible, which would afford the enantiomeric metallacycles **IV** and **IV'** in equilibrium, while a selective protonation of **IV** by CCA **4a** would proceed to furnish the product in an enantioselective fashion and regenerate **I**. Although we cannot completely deny a mechanism in which CCA **4a** activates maleimide **2a** as a chiral acid catalyst to promote enantioselective insertion of the C–C double bond, we consider that the reversible insertion/selective protodemetalation mechanism would be more likely because the reasonable enantioselectivity was observed in TFE, which is highly polar and would not be suitable for the recognition of substrates via hydrogen-bonding with CCAs.




Figure 1. Plausible catalytic cycle for the CCA (**4a**)-induced enantioselectivity.

In summary, we have developed an enantioselective $\text{Cp}^*\text{Co(III)}$ /CCA-catalyzed 1,4-addition reaction of indoles **1** and maleimides **2**, which proceeds via a C–H activation. In these reactions, a chiral carboxylic acid (CCA) derived from BINOL (**4a**) serves as the sole chiral source for the enantioselective addition, albeit that the observed enantioselectivity leaves room for improvement. Further investigations, including the development of new classes of CCAs, are ongoing in our group.

Acknowledgements

This work was supported in part by JSPS KAKENHI Grant Number JP15H05802 in Precisely Designed Catalysts with Customized Scaffolding, JSPS KAKENHI Grant Number JP18H04637 in Hybrid Catalysis, JSPS KAKENHI Grant Number JP17H03049 and JP19K16306.


Keywords: C–H activation • cobalt • chiral carboxylic acid • asymmetric 1,4-addition • indole

References

- [1] B. M. Trost, *Science* **1991**, *254*, 1471–1477.
- [2] P. A. Wender, B. L. Miller, *Nature* **2009**, *460*, 197–201.
- [3] Selected recent reviews on C–H functionalization reactions: a) K. Shin, H. Kim, S. Chang, *Acc. Chem. Res.* **2015**, *48*, 1040–1052; b) M. Moselage, J. Li, L. Ackermann, *ACS Catal.* **2016**, *6*, 498–525; c) T. Gensch, M. N. Hopkinson, F. Glorius, J. Wencel-Delord, *Chem. Soc. Rev.* **2016**, *45*, 2900–2936; d) J. He, M. Wasa, K. S. L. Chan, Q. Shao, J.-Q. Yu, *Chem. Rev.* **2017**, *117*, 8754–8786; e) J. R. Hummel, J. A. Boerth, J. A. Ellman, *Chem. Rev.* **2017**, *117*, 9163–9227; f) Y.-F. Yang, X. Hong, J.-Q. Yu, K. N. Houk, *Acc. Chem. Res.* **2017**, *50*, 2853–2860; g) R. R. Karimov, J. F. Hartwig, *Angew. Chem.* **2018**, *130*, 4309–4317; *Angew. Chem. Int. Ed.* **2018**, *57*, 4234–4241; h) Y. Xu, G. Dong, *Chem. Sci.* **2018**, *9*, 1424–1432; i) Y. N. Timsina, B. F. Gupton, K. C. Ellis, *ACS Catal.* **2018**, *8*, 5732–5776; j) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz, L. Ackermann, *Chem. Rev.* **2019**, *119*, 2192–2452.
- [4] Selected reviews on group 9 $\text{Cp}^*\text{M(III)}$ -catalyzed C–H functionalization reactions: a) T. Satoh, M. Miura, *Chem. Eur. J.* **2010**, *16*, 11212–11222; b) N. Kuhl, N. Schröder, F. Glorius, *Adv. Synth. Catal.* **2014**, *356*, 1443–1460; c) G. Song, X. Li, *Acc. Chem. Res.* **2015**, *48*, 1007–1020; d) S. Wang, S.-Y. Chen, X.-Q. Yu, *Chem. Commun.* **2017**, *53*, 3165–3180; e) T. Yoshino, S. Matsunaga, *Adv. Synth. Catal.* **2017**, *359*, 1245–1262; f) P. G. Chirila, C. J. Whiteoak, *Dalton Trans.* **2017**, *46*, 9721–9739; g) T. Piou, T. Rovis, *Acc. Chem. Res.* **2018**, *51*, 170–180; h) J. Park, S. Chang, *Chem. Asian J.* **2018**, *13*, 1089–1102.
- [5] For selected examples, see: a) B. Ye, N. Cramer, *Science* **2012**, *338*, 504–506; b) B. Ye, N. Cramer, *J. Am. Chem. Soc.* **2013**, *135*, 636–639; c) Y.-S. Jang, M. Dieckmann, N. Cramer, *Angew. Chem.* **2017**, *129*, 15284–15288; *Angew. Chem. Int. Ed.* **2017**, *56*, 15088–15092; d) K. Ozols, Y.-S. Jang, N. Cramer, *J. Am. Chem. Soc.* **2019**, *141*, 5675–5680; e) M. Brauns, N. Cramer, *Angew. Chem.* **2018**, *130*, 15765–15769; *Angew. Chem. Int. Ed.* **2019**, *58*, 8902–8906. For related work by other research groups, see: f) T. J. Potter, D. N. Kamber, B. Q. Mercado, J. A. Ellman, *ACS Catal.* **2017**, *7*, 150–153; g) X. Yang, G. Zheng, X. Li, *Angew. Chem.* **2019**, *131*, 328–332; *Angew. Chem. Int. Ed.* **2019**, *58*, 322–326; h) M. Tian, D. Bai, G. Zheng, J. Chang, X. Li, *J. Am. Chem. Soc.* **2019**, *141*, 9527–9532.
- [6] For other types of chiral Cp^x ligands, see: a) J. Zheng, W.-J. Cui, C. Zheng, S.-L. You, *J. Am. Chem. Soc.* **2016**, *138*, 5242–5245; b) Z.-J. Jia, C. Merten, R. Gontla, C. G. Daniluc, A. P. Antonchick, H. Waldmann, *Angew. Chem.* **2017**, *129*, 2469–2474; *Angew. Chem. Int. Ed.* **2017**, *56*, 2429–2434; c) E. A. Trifonova, N. M. Ankudinov, A. A. Mikhaylov, D. A. Chusov, Y. V. Nelyubina, D. S. Perekalin, *Angew. Chem.* **2018**, *130*, 7840–7844; *Angew. Chem. Int. Ed.* **2018**, *57*, 7714–7718; d) H. Li, R. Gontla, J. Flegel, C. Merten, S. Ziegler, A. P. Antonchick, H. Waldmann, *Angew. Chem.* **2019**, *131*, 313–317; *Angew. Chem. Int. Ed.* **2019**, *58*, 307–311; e) H. Li, X. Yan, J. Zhang, W. Guo, J. Jiang, J. Wang, *Angew. Chem.* **2019**, *131*, 6804–6808; *Angew. Chem. Int. Ed.* **2019**, *58*, 6732–6736.
- [7] A biotinylated Rh(III) complex combined with engineered streptavidins has also been used for enantioselective C–H functionalization reactions: a) T. K. Hyster, L. Knörr, T. R. Ward, T. Rovis, *Science* **2012**, *338*, 500–503; b) I. S. Hassan, A. N. Ta, M. W. Danneman, N. Semakul, M. Burns, C. H. Basch, V. N. Dippon, B. R. McNaughton, T. Rovis, *J. Am. Chem. Soc.* **2019**, *141*, 4815–4819.

- [8] Reviews on asymmetric C–H functionalization reactions with chiral Cp*M(III) catalysts: a) B. Ye, N. Cramer, *Acc. Chem. Res.* **2015**, *48*, 1308–1318; b) C. G. Newton, D. Kossler, N. Cramer, *J. Am. Chem. Soc.* **2016**, *138*, 3935–3941.
- [9] Reviews on transition metal-catalyzed asymmetric C–H functionalization reactions: a) C. G. Newton, S.-G. Wang, C. C. Oliveira, N. Cramer, *Chem. Rev.* **2017**, *117*, 8908–8976; b) T. G. Saint-Denis, R.-Y. Zhu, G. Chen, Q.-F. Wu, J.-Q. Yu, *Science*, **2018**, *359*, eaao4798; c) J. Loup, U. Dhawa, F. Pescioli, J. Wencel-Delord, L. Ackermann, *Angew. Chem.* **2019**, *131*, 12934–12949; *Angew. Chem. Int. Ed.* **2019**, *58*, 12803–12818.
- [10] For our recent account on this topic, see: T. Yoshino, S. Matsunaga, *Synlett* **2019**, *30*, 1384–1400.
- [11] D. L. Davies, S. A. Macgregor, C. L. McMullin, *Chem. Rev.* **2017**, *117*, 8649–8709.
- [12] L. Yang, B. Qian, H. Huang, *Chem. Eur. J.* **2012**, *18*, 9511–9515.
- [13] Enantioselective C–H activation has originally been achieved using Pd catalysts combined with chiral mono-protected amino acids or related ligands. For selected examples, see: a) B.-F. Shi, N. Maugel, Y.-H. Zhang, J.-Q. Yu, *Angew. Chem.* **2008**, *120*, 4960–4964; *Angew. Chem. Int. Ed.* **2008**, *47*, 4882–4886; b) F.-L. Zhang, K. Hong, T.-J. Li, H. Park, J.-Q. Yu, *Science* **2016**, *351*, 252–256; c) G. Chen, W. Gong, Z. Zhuang, M. S. Andr , Y.-Q. Chen, X. Hong, Y.-F. Yang, T. Liu, K. N. Houk, J.-Q. Yu, *Science* **2016**, *353*, 1023–1027; d) P. Jain, P. Verma, G. Xia, J.-Q. Yu, *Nat. Chem.* **2017**, *9*, 140–144; e) H. Park, P. Verma, K. Hong, J.-Q. Yu, *Nat. Chem.* **2018**, *10*, 755–762.
- [14] D. Gwon, S. Park, S. Chang, *Tetrahedron* **2015**, *71*, 4504–4511.
- [15] a) L. Lin, S. Fukagawa, D. Sekine, E. Tomita, T. Yoshino, S. Matsunaga, *Angew. Chem.* **2018**, *130*, 12224–12228; *Angew. Chem. Int. Ed.* **2018**, *57*, 12048–12052; b) S. Fukagawa, Y. Kato, R. Tanaka, M. Kojima, T. Yoshino, S. Matsunaga, *Angew. Chem.* **2019**, *131*, 1165–1169; *Angew. Chem. Int. Ed.* **2019**, *58*, 1153–1157; c) D. Sekine, K. Ikeda, S. Fukagawa, M. Kojima, T. Yoshino, S. Matsunaga, *Organometallics* **2019**, <https://doi.org/10.1021/acs.organomet.9b00407>.
- [16] Y.-H. Liu, P.-X. Li, Q.-J. Yao, Z.-Z. Zhang, D.-Y. Huang, M. D. Le, H. Song, L. Liu, B.-F. Shi, *Org. Lett.* **2019**, *21*, 1895–1899;
- [17] a) D. Zell, M. Bursch, V. M ller, S. Grimme, L. Ackermann, *Angew. Chem.* **2017**, *129*, 10514–10518; *Angew. Chem. Int. Ed.* **2017**, *56*, 10378–10382; b) F. Pescioli, U. Dhawa, J. C. A. Oliveira, R. Yin, M. John, L. Ackermann, *Angew. Chem.* **2018**, *130*, 15651–15655; *Angew. Chem. Int. Ed.* **2018**, *57*, 15425–15429.
- [18] For enantioselective C–H alkylation reactions using other metal catalysts, see a review: Z. Dong, Z. Ren, S. J. Thompson, Y. Xu, G. Dong, *Chem. Rev.* **2017**, *117*, 9333–9403.
- [19] For our recent work on Cp*Rh(III)/chiral disulfonate-catalyzed enantioselective C–H alkylation reactions, see: S. Satake, T. Kurihara, K. Nishikawa, T. Mochizuki, M. Hatano, K. Ishihara, T. Yoshino, S. Matsunaga, *Nat. Catal.* **2018**, *1*, 585–591.
- [20] For reviews on C–H activation/functionalization of indoles, see: a) A. H. Sandtorv, *Adv. Synth. Catal.* **2015**, *357*, 2403–2435; b) J. A. Leitch, Y. Bhonoah, C. G. Frost, *ACS Catal.* **2017**, *7*, 5618–5627.
- [21] Z. Zhang, S. Han, M. Tang, L. Ackermann, J. Li, *Org. Lett.* **2017**, *19*, 3315–3318.
- [22] T. Hashimoto, H. Kimura, H. Nakatsu, K. Maruoka, *J. Org. Chem.* **2011**, *76*, 6030–6037.
- [23] The pyrimidyl directing group on the nitrogen was crucial. A 2-pyridyl-protected indole afforded the racemic product, and a dimethylcarbamoyl-protected indole did not afford the desired product. The corresponding pyrrole, 5-methyl-2-(1*H*-pyrrol-1-yl)pyrimidine, also failed to give a promising result (less than 60:40 *er*).
- [24] The addition reaction to dimethyl maleate provided the product, but the enantioselectivity was low (52.5:47.5 *er*).
- [25] N. Muniraj, K. R. Prabhu, *ACS Omega* **2017**, *2*, 4470–4479.
- [26] a) D. Lapointe, K. Fagnou, *Chem. Lett.* **2010**, *39*, 1118–1126; b) L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315–1345.

Entry for the Table of Contents



The combination of an achiral Cp*Co(III) catalyst and a binaphthyl-based chiral carboxylic acid enables enantioselective 1,4-addition reactions of C–H bonds of indoles to maleimides. The chiral carboxylic acid serves as the sole chiral source to realize the enantioselective alkylation process.