




Title	Effects of biochar and litter on water relations of Japanese black pine (<i>Pinus thunbergii</i>) seedlings
Author(s)	Fujita, Saki; Watanabe, Hanami; Marozas, Vitas et al.
Citation	Journal of Forest Research, 25(2), 76-82 https://doi.org/10.1080/13416979.2020.1731071
Issue Date	2020-02-25
Doc URL	https://hdl.handle.net/2115/80488
Rights	This is an Accepted Manuscript of an article published by Taylor & Francis in Journal of Forest Research on 25 Feb 2020, available online: http://www.tandfonline.com/10.1080/13416979.2020.1731071 .
Type	journal article
File Information	Figures_JFR_revised_20200207.pdf, 図



1 **Figure 1**

2



3

4 **Figure 1**

5 Mean (±SE) of needle xylem water potential after 3 to 17 days stopping irrigation at control, biochar, litter,
6 litter + biochar treatments. Different letters indicate a statistical difference ($p < 0.05$).

7

8

9

10

11

12

13

14

15

16

Figure 2




Figure 2

Linear regression analysis of the relationship between relative fine root biomass (fine root biomass/total root biomass, %) of all treatments and xylem water potential at 17 days after ceasing irrigation. Xylem water potential and relative fine root were obtained from the same seedling ($n=4$ for each treatment).