



Title	Effects of single and repetitive valproic acid administration on the gene expression of placental transporters in pregnant rats : An analysis by gestational period.
Author(s)	Jinno, Naoki; Furugen, Ayako; Kurosawa, Yuko et al.
Citation	Reproductive toxicology (Elmsford, N.Y.), 96, 47-56 https://doi.org/10.1016/j.reprotox.2020.04.077
Issue Date	2020-05-11
Doc URL	https://hdl.handle.net/2115/81137
Rights	©2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	HUSCAP.pdf



Effects of single and repetitive valproic acid administration on the gene expression of placental transporters in pregnant rats: An analysis by gestational period.

Naoko Jinno, Ayako Furugen, Yuko Kurosawa, Yuki Kanno, Katsuya Narumi, Masaki Kobayashi*,
Ken Iseki

Laboratory of Clinical Pharmaceutics & Therapeutics, Division of Pharmasciences, Faculty of
Pharmaceutical Sciences, Hokkaido University, Kita-12-jo, Nishi-6-chome, Kita-ku, Sapporo
060-0812, Japan

*Correspondence to: Masaki Kobayashi, Ph.D. Laboratory of Clinical Pharmaceutics & Therapeutics,
Division of Pharmasciences, Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12-jo,
Nishi-6-chome, Kita-ku, Sapporo 060-0812, Japan
Phone: +81-11-706-3772; Fax: +81-11-706-3235; E-mail: masaki@pharm.hokudai.ac.jp

16 **Abbreviations**

17 ABC, ATP-binding cassette; ANOVA, analysis of variance; BCRP, breast cancer resistance protein;
18 CNT, concentrative nucleoside transporter; ENT, equilibrative nucleoside transporter; FR α , folate
19 receptor alpha; G, gestational day; GLUT, glucose transporter; HDAC, histone deacetylase; IS,
20 internal standard; LAT, L-type amino acid transporter; LC/MS/MS, liquid chromatography tandem
21 mass spectrometry; MCT, monocarboxylate transporter; MDR, multiple drug resistant; MRP,
22 multidrug resistance-associated protein; OATP, organic anion transporting polypeptide; OCT,
23 organic cation transporter; OCTN, organic cation/carnitine transporter; PCR, polymerase chain
24 reaction; SLC, solute carrier; VPA, valproic acid

25 **Abstract**

26 The use of valproic acid (VPA), an antiepileptic drug, during pregnancy, is known to increase
27 various fetal risks. Since VPA has been known to inhibit histone deacetylases (HDACs); its
28 administration could alter gene transcription levels. However, *in vivo* effects of VPA administration
29 on placental transporters have not been fully elucidated. The purpose of the present study was to
30 comprehensively evaluate the effects of single and repetitive VPA administration on the expression
31 of placental transporters and analyze them by gestational day. We investigated 18 transporters (8
32 ATP-binding cassette (ABC) and 10 solute carrier (SLC) transporters) in the placentas of pregnant
33 rats that were orally administered 400 mg/kg/day VPA for one or four days, during mid- or late
34 gestation. In the control rats, 4 ABC transporter genes (*Abcb1a*, *1b*, *Abcc2*, *Abcc4*) were upregulated,
35 3 (*Abcc3*, *Abcc5*, *Abcg2*) downregulated through gestation, whereas 1 (*Abcc1*) was not changed.
36 Regarding SLC transporters, 6 genes (*Slc7a5*, *Slc16a3*, *Slc22a3*, *Slc22a4*, *Slco2b1*, *Slco4a1*) were
37 increased, 1 (*Slc29a1*) decreased through gestation, whereas 3 (*Slc7a8*, *Slc22a5*, *Slco2a1*) showed no
38 significant change. Single VPA administration altered the expression of 9 transporters and repetitive
39 administration, 13 transporters. In particular, VPA remarkably decreased *Abcc4* and *Slc22a4* in late
40 gestation and increased *Abcc5* during mid-gestation. Our findings indicated that VPA administration
41 changed transporter expression levels in rat placenta, and suggested that sensitivity to VPA differs
42 across gestational stages.

43

44 **Key words:** Placenta; antiepileptic drug; valproic acid; ABC transporter; SLC transporter;

45 gestational period; Rat

46

47 **1. Introduction**

48 About 0.3–0.7 % of pregnant women have epilepsy (Viinikainen et al., 2006). Seizure control
49 needs to be maintained by continuous pharmacotherapy throughout pregnancy. Generally,
50 medication use during pregnancy is a fetal risk factor, needing consideration in gestational
51 pharmacotherapy.

52 Valproic acid (VPA) is a well-established and frequently used antiepileptic drug. However, it
53 increases the risk of fetal malformations, autism spectrum disorders, and cognitive defects (Baker et
54 al., 2015; Christensen et al., 2013; Jentink et al., 2010; Tomson et al., 2011). VPA treatment should
55 be avoided in women of childbearing age; however, it is prescribed to pregnant women when
56 required (Ishikawa et al., 2019; Yoshimura et al., 2018). Hence, the effect of VPA on the fetus and
57 related organs needs to be evaluated.

58 The placenta is a crucial temporary organ, which is in contact with both maternal and fetal
59 blood. ATP-binding cassette (ABC) and solute carrier (SLC) transporters play a role in nutrient,
60 metabolic waste, and xenobiotic exchange between mother and fetus (Leazer and Klaassen, 2003;
61 Staud et al., 2012). Hence, the expression levels of placental transporters are one of the factors
62 affecting the intrauterine environment and fetal growth. Since the expression of placental transporters
63 changes across gestation, their evaluation at each gestational stage is important. Additionally,
64 transporters can be affected in a sex-specific manner, in some cases (Song et al., 2017).

65 VPA has been reported to inhibit histone deacetylases (HDACs), which remove acetyl groups
66 from histones. Four classes of HDACs have been identified, and VPA acts predominantly on Class I
67 HDACs, including HDAC 1–3 and 8 (Grabiec and Potempa, 2018; Gurvich et al., 2004). Therefore,
68 VPA likely affects the expression of various genes. So far, there are several reports that have
69 investigated the effects of VPA on placental transporters. An *in vitro* study indicated that VPA
70 treatment altered the expression and function of breast cancer resistance protein (BCRP)
71 (Rubinchik-Stern et al., 2015). Moreover, we previously showed that VPA exposure induces
72 increased mRNA levels of folate receptor alpha (FR α) and proton-coupled folate transporter in
73 BeWo and JEG-3 cell lines, derived from human choriocarcinoma (Kurosawa et al., 2018).
74 Additionally, perfusion with VPA reportedly has reduced FR α and glucose transporter (GLUT) 1 in
75 *ex vivo* human placentas (Rubinchik-Stern et al., 2018; Tetro et al., 2019). However, the perfusion
76 time was short and variability among samples was high in the model. In particular, placental
77 structure, function, and transporters are dramatically altered. Therefore, an *in vivo* study is well
78 suited to sequentially investigate, throughout gestation, and assess the influence of fetal sex on
79 VPA-mediated changes. Although a previous *in vivo* study indicated that the expression of L-type
80 amino acid transporter (LAT) 1, organic anion transporting polypeptide (OATP) 4a1, and reduced
81 folate carrier were lower in placentas from VPA-treated mid-pregnant mice (Meir et al., 2016),
82 transporter expression level changes were not recorded.

83 This study aimed to comprehensively reveal the effects of both single and repetitive VPA
84 treatment on placental transporters, using pregnant rats. We analyzed the expression of 8 ABC
85 transporters, including multiple drug-resistant (MDR) 1a and 1b (*Abcb1a*, *1b*), multidrug
86 resistance-associated proteins (MRPs) 1–5 (*Abcc1–5*), and BCRP (*Abcg2*), as well as 10 SLC
87 transporters, including LAT1 and 2 (*Slc7a5* and *Slc7a8*), monocarboxylate transporter (MCT) 4
88 (*Slc16a3*), organic cation transporter (OCT) 3 (*Slc22a3*), organic cation/carnitine transporter
89 (OCTN) 1 and 2 (*Slc22a4*, 5), equilibrative nucleoside transporter (ENT) 1 (*Slc29a1*), and OATP2a1,
90 2b1, and 4a1 (*Slco2a1*, *2b1*, *4a1*), by gestational age. Furthermore, the results were secondarily
91 analyzed by fetal sex.

92 **2. Material and Methods**

93 **2.1. Chemicals and reagents.**

94 Valproic acid sodium salt (VPA Na) and valproic acid were purchased from Sigma-Aldrich
95 (St. Louis, MO, USA). Deuterium-labeled valproic acid (VPA-d₆) and 2-propyl-4-pentenoic acid
96 (4-ene-VPA) were purchased from Toronto Research Chemicals (Toronto, Canada).

97

98 **2.2. Animals.**

99 Pregnant female Wistar rats (12–13 weeks old) were provided by CLEA Japan (Tokyo,
100 Japan). The existence of a vaginal plug indicated the first day of gestation (Gestational day 0; G0).
101 G9–13 and G16–20 were considered as mid- and late gestation, respectively (Sun et al., 2015;
102 Kalisch-Smith et al., 2017). The rats were housed at 23 ± 2°C and 60 ± 10 % relative humidity under
103 a 12-h light/dark cycle, with *ad libitum* access to food and water. The experimental protocols were
104 reviewed and approved by the Hokkaido University Animal Care Committee, in accordance with the
105 National Institutes of Health guide for the care and use of laboratory animals.

106

107 **2.3. Drug administration and collection of plasma and placentas.**

108 VPA Na was dissolved in distilled water (200 mg/mL), filter-sterilized, and then orally
109 administered to pregnant rats at a daily dose of 400 mg/kg body weight, by gavage. We determined
110 the appropriate experimental dose for observing fetal effects, which does not cause maternal death,
111 based on the findings of a previous study (Vorhees, 1987). Control rats were administered the same

112 volume of distilled water. Rats of the single administration groups were treated on G12
113 (mid-gestation) or G19 (late gestation), and those in the repetitive groups were treated for 4 days on
114 G9–12 (mid-gestation) or G16–19 (late gestation). All pregnant rats were weighed before every
115 administration and euthanization.

116 Thirty minutes and 24 h post-VPA administration, blood was collected from the tail vein and
117 centrifuged at 800 g and 4°C for 15 min, to obtain plasma. Twenty-four hours post the last
118 administration, G13 or G20 rats were anesthetized with sevoflurane and euthanized by decapitation;
119 their placentas were then immediately collected, washed briefly with ice-cold PBS, and weighed.
120 Three to four placentas per dam were used for real-time PCR, and the others were frozen in liquid
121 nitrogen and stored at -80°C until further experiments.

122

123 **2.4. Real-time PCR.**

124 Total RNA was extracted from the placental homogenates using ISOGEN II (Nippon Gene,
125 Tokyo, Japan). RNA concentration was measured using Thermo Scientific™ NanoDrop 2000
126 (Thermo Fisher Scientific, Waltham, MA, USA). DNase treatment was performed using the
127 RNase-free DNase Set (QIAGEN, Venlo, Netherlands). Samples after DNase treatment were
128 reverse-transcribed using ReverTra Ace (Toyobo, Osaka, Japan) and iCycler™ (Bio-Rad
129 Laboratories, Hercules, CA), according to manufacturer's instructions. The A260/A280 and
130 A260/A230 ratios of isolated RNA were 1.95 ± 0.05 and 2.00 ± 0.10 , respectively.

131 Real-time PCR was performed using KAPA SYBR® Fast qPCR Kit (Kapa Biosystems,
132 Wilmington, MA, USA) and LightCycler® 480 System II (Roche, Basel, Switzerland), through 40
133 cycles of 95°C for 10 s, 55°C or 60°C for 20 s, and 72°C for 1 s. Primers used for the real-time PCR
134 are shown in Supplemental Table 1. Beta-actin was used as the reference gene. We determined that
135 the reference gene in the rat placenta was not influenced by gestational age or VPA administration.
136 Data were analyzed by the relative standard curve method. As a calibrator, cDNA from a G20
137 placenta (1 lot) was used. The Ct values of rat placenta samples are shown in Supplemental Table 2.

138 Fetal sex was determined by detecting the Y-chromosome-linked gene, *Sry*, which encodes a
139 sex-determining region Y protein in the placentas, in accordance with a previous study (Song et al.,
140 2017). Placentas with no *Sry* gene expression (Ct value, >35 or similar to that in negative controls)
141 were considered to be from female fetuses, whereas those with expression (Ct value, <30) were
142 considered to be from male fetuses.

143

144 **2.5. VPA and 4-ene-VPA quantification in rat plasma using liquid chromatography-tandem**
145 **mass spectrometry.**

146 For standard curve generation, blank plasma from control pregnant rats was spiked with VPA
147 and 4-ene-VPA to final concentrations of 10–500 and 0.2–100 µg/mL, respectively. VPA and
148 4-ene-VPA levels in plasma were determined as previously described (Wu and Lu, 2014; Gao et al.,
149 2011), with some modification. Briefly, 10 µL of VPA-d₆ (internal standard (IS), 100 µg/mL) and 25

150 μ L of hydrochloric acid (1 M) were added to 50 μ L of rat plasma. Next, 1 mL of toluene was added
151 to the solution and mixed for 1 min. The mixture was centrifuged at 13,000 g and 4°C for 10 min,
152 and 800 μ L of supernatant collected and distilled at 37°C under gentle nitrogen steam. The residue
153 was reconstituted in 50 μ L of mobile phase (methanol:10 mM ammonium formate solution, 80:20
154 (v/v)) and 2 μ L injected for HPLC.

155 HPLC was performed using a completely equipped Prominence 20A system (Shimadzu,
156 Kyoto, Japan). VPA and 4-ene-VPA were separated using an Inertsil ODS-3 column (2.1 $\times$ 150 mm,
157 3 μ m; GL Science Inc., Tokyo, Japan). The column temperature was maintained at 40°C and the
158 mobile phase flow rate set at 0.2 mL/min. Multiple reaction monitoring was performed using an
159 API3200TM LC-MS/MS system (Applied Biosystems, Foster City, CA). This system was operated in
160 the negative ion electrospray mode. Monitoring ions were m/z 143.1 $\rightarrow$ 143.1 for VPA, m/z 141.1 $\rightarrow$
161 141.1 for 4-ene-VPA, and m/z 149.1 $\rightarrow$ 149.1 for IS. Data were analyzed using Analyst software
162 (Applied Biosystems).

163

164 **2.6. Western blotting**

165 Whole proteins were extracted from placentas of G20 pregnant rats that were repetitively
166 administrated VPA or water (control). Placentas were homogenized in ice-cold RIPA buffer (Cell
167 Signaling Technology, Danvers, MA, USA). The lysates were kept on ice for 5 min and sonicated
168 briefly, then centrifuged at 14,000 g and 4°C for 10 min. The supernatant was used for western
169 blotting. The total protein concentration was measured using a Pierce® BCA Protein Assay Kit

170 (Thermo Fisher Scientific). Total protein was mixed with Blue Loading Buffer Pack (Cell signaling
171 Technology, Danvers, MA, USA) and denatured at 100°C for 3 min. The mixture was subjected to
172 SDS-PAGE at 20 ug protein per well and transferred onto nitrocellulose membranes (Bio-Rad
173 Laboratories). The membranes were blocked with 5% non-fat dry milk in TBS containing 0.05%
174 Tween 20 (TBST), at room temperature for 1 h, then incubated with primary antibodies diluted with
175 Can Get Signal® Solution 1 (Toyobo) overnight at 4°C. The following primary antibodies were
176 used: mouse anti-P-glycoprotein (1:200, Merck Millipore, Burlington, MA, #517310) and mouse
177 anti-actin (1:1000, Chemicon, Temecula, CA, #MAB1501) monoclonal antibodies. The bands were
178 detected using HRP-conjugated secondary antibodies (anti-mouse IgG, 1 : 4000, Southern Biotech,
179 Birmingham, AL, #1070-05), and ECL prime™ Western blotting detection reagent (GE Healthcare,
180 Buckinghamshire, England), with Image Quant LAS 4000 (GE Healthcare). The band Intensities
181 were analyzed with ImageJ analysis software (National Institutes of Health, Bethesda, MD, USA).

182

183 **2.7. Statistical analyses.**

184 Maternal body weight gain and placental weight were analyzed using Student's *t*-test for
185 comparison of the control and VPA groups. Gene expression was analyzed using a two-way analysis
186 of variance (ANOVA), with “gestational day (G)” and “VPA” or “sex” and “VPA” as factors. If an
187 interaction was present, the Tukey–Kramer test was used for multiple comparisons. Two-way
188 ANOVA and Tukey–Kramer test were performed using JMP Pro (SAS Institute, Cary, NC, USA).

189 All data are shown as mean with standard deviation (SD), and $P < 0.05$ was assessed as statistically
190 significant.

191 **3. Results**

192 **3.1. Effects of single VPA Na administration on placental transporters at G13 and G20.**

193 First, we confirmed the presence of VPA and 4-ene-VPA in the plasma from the pregnant rats
194 30 min (peak) post oral VPA Na administration. In a previous study in pregnant rats, the VPA level
195 peaked at 0.5–0.9 h post-oral VPA Na administration (Binkerd et al., 1988). The plasma levels of
196 VPA and its toxic active metabolite, 4-ene-VPA, 30 min post-administration were approximately 200
197 and 1.5 µg/mL, respectively (Table 1); this level was similar to that reported in a previous study
198 (Binkerd et al., 1988). After 24 h, their concentrations were less than or close to the lower limit of
199 quantification. There were no differences in the concentrations between mid- and late gestation.
200 Body weight gain of the pregnant rats did not differ significantly between VPA Na single
201 administration and control groups (data not shown).

202

203 **3.1.1. Alteration of placental ABC transporter expression by single VPA Na administration.**

204 The expression of 8 ABC transporter mRNAs in the placentas were assessed (Figure 1).
205 Interactions between gestational day and VPA Na administration were statistically significant for
206 rMrp1, rMrp4, and rMrp5. Multiple comparisons revealed that VPA Na decreased rMrp4 by 22 % at
207 G20, whereas increased rMrp5 by 22 % at G13 (Figure 1F and 1G). No significant difference was
208 noted in rMrp1 expression (Figure 1C). With the mRNA expression alterations associated with the
209 gestational stage, multiple comparisons showed that rMrp4 increased by 2.84-folds (Figure 1F),

210 whereas rMrp5 decreased by 0.67-folds (Figure 1G) at G20, compared with the G13 control groups.
211 The main gestational day effects were significant for rMdr1a, rMdr1b, rMrp3, and rBcrp; rMdr1a and
212 rMdr1b increased to 295 % and 262%, respectively (Figure 1A and 1B), whereas rMrp3 and rBcrp
213 decreased to 33% and 49%, respectively (Figure 1E and 1H). rMrp2 could not be evaluated because
214 it was scarcely expressed at G13 (Figure 1D).

215

216 **3.1.2. Alteration of placental SLC transporter expression by single VPA Na administration.**

217 Next, we assessed the expression of 10 SLC transporter mRNAs in the placentas (Figure 2).
218 Interactions between gestational day and VPA Na administration were statistically significant for
219 rLat1, rLat2, rOctn1, rOatp2a1, rOatp2b1, and rOatp4a1. Multiple comparisons revealed that VPA
220 Na increased rLat2 by 36 % and rOatp2a1 by 75 % at G13 (Figure 2B and 2H). In contrast, it
221 decreased rLat1 by 21 %, rOctn1 by 20 %, rOatp2b1 by 21 %, and rOatp4a1 by 18 % at G20 (Figure
222 2A, 2E, 2I, and 2J). No interactions were noted for rMct4, rOct3, and rEnt1 (Figure 2C, 2D, and 2G).
223 When compared with levels in the G13 control by multiple comparisons, rLat1, rOctn1, rOatp2b1,
224 and rOatp4a1 increased to 783%, 1660%, 485%, and 585% at G20, respectively (Figure 2A, 2E, 2I,
225 and 2J). rMct4 tended to increase to 210 %, whereas rEnt1 tended to decrease to 46 % (Figure 2C
226 and 2G).

227 **3.2. Effects of repetitive VPA Na administration on placental transporters at G13 and G20.**

228 The plasma levels of VPA and 4-ene-VPA 30 min post each administration were
229 approximately 100–300 µg/mL and 1.5–2.0 µg/mL, respectively (Table 1). After 24 h, the
230 concentrations of both VPA and 4-ene-VPA were less than or close to the lower limit of
231 quantification. There were no substantial differences between mid- and late gestation as well as
232 between single and repetitive administration. Body weight gain of VPA-administered dams was
233 approximately half of that of control dams at G20 in the late gestation group, whereas no difference
234 was noted in the mid-gestation group (Figure 3A). Placental weight was significantly decreased by
235 VPA Na at both G13 and G20 (Figure 3B).

236

237 **3.2.1. Alteration of placental ABC transporter expression by repetitive administration of VPA**
238 **Na.**

239 Interactions between gestational day and VPA Na administration were statistically significant
240 for rMdr1a, rMrp1, rMrp3, rMrp4, rMrp5, and rBcrp, among the eight ABC transporters (Figure 1).
241 Multiple comparisons revealed that VPA Na increased rMrp1 by 43 % and rMrp5 by 67 % at G13
242 (Figure 1C and 1G) and rMdr1a by 34 % at G20 (Figure 1A), whereas decreased rMrp3 by 17 % at
243 G13 and rMrp4 by 46 % at G20 (Figure 1E and 1F). With the mRNA expression alterations
244 associated with gestational stage, multiple comparisons showed that rMdr1a and rMrp4 increased to
245 277 % and 212 % (Figure 1A and 1F), whereas rMrp3, rMrp5, and rBcrp decreased to 35 %, 59 %, 16

246 and 56 %, respectively (Figure 1E, 1G, and 1H) at G20, compared with at G13. These alterations
247 were similar to those of the single administration. The main effects of VPA and gestational day on
248 rMdr1b were significant, although they did not have an interaction: rMdr1b increased to 247 %
249 through gestation and decreased by 25 % with VPA Na.

250

251 **3.2.2. Alteration of placental SLC transporter expression by repetitive VPA Na administration.**

252 Interactions between gestational day and VPA Na administration were statistically significant
253 for Lat1, rLat2, rOctn1, rOctn2, rEnt1, rOatp2a1, and rOatp4a1 (Figure 2). Multiple comparisons
254 showed that VPA Na increased rLat2 by 95 %, rOctn2 by 78 %, rEnt1 by 61 %, and rOatp2a1 by
255 131 % at G13 (Figure 2B, 2F, 2G, and 2H), whereas decreased rLat1 by 18 %, rOctn1 by 51 %, and
256 rOatp4a1 18 % at G20 (Figure 2A, 2E, and 2J). With regards to gene expression alterations during
257 gestation, multiple comparisons showed that rLat1, rOctn1, and rOatp4a1 increased by 7.6-, 17.2-,
258 and 5.7-folds, respectively (Figure 2A, 2E, and 2J). Conversely, rEnt1 expression at G20 was
259 0.49-folds lower than at G13 (Figure 2G). These alterations associated with the gestational stages
260 were the same as those of single administration.

261

262 **3.3. Analysis of fetal sexual effects on gene expression.**

263 A previous study evaluating placental response to maternal metabolic changes indicated that
264 the expression of signaling factors and nutrient transporters was altered in a sex-specific manner

265 (Song et al., 2017). We secondarily analyzed the data shown in Figures 1 and 2 by fetal sex to clarify
266 whether differences existed between male and female placentas. The analysis results of G13 and G20
267 placentas are shown in Supplemental Tables 3 and 4, respectively. Interactions between sex and VPA
268 Na were significant for rMdr1a in the G13 single administration groups and rOatp2a1 and rOatp2b1
269 in G20 repetitive administration groups. Alteration of the three transporter mRNAs is shown in
270 Supplemental Figure 1. Multiple comparisons revealed that VPA Na increased rMdr1a by 27 % and
271 decreased rOatp2a1 by 46 % in male placentas, whereas no significant changes were noted in female
272 placentas (Supplemental Figure 1A and 1B). There were no significant changes in rOatp2b1
273 expression in both male and female placentas (Supplemental Figure 1C).

274 4. Discussion

275 VPA use during pregnancy poses risks to the fetus such as malformations, autism spectrum
276 disorders, and cognitive defects (Baker et al., 2015; Christensen et al., 2013; Jentink et al., 2010;
277 Tomson et al., 2011). Therefore, evaluation of its effects on the placenta, which supports fetal
278 development and health maintenance in pregnancy, is important. In the present study, we
279 comprehensively evaluated the effects of VPA administration on the expression of rat placental
280 transporters.

281 In the study, we determined the experimental dose to observe fetal effects and not cause
282 maternal death, based on the findings of a previous study (Vorhees, 1987). In addition, previous
283 studies that have investigated the effects of prenatal VPA exposure on rat pups, chose the
284 administration dose of approximately 300–800 mg/kg (Roullet FI et al., 2013). No considerable
285 differences between VPA levels and the therapeutic VPA range, which is 42–114 µg/mL in humans
286 (Kim et al., 2011), were observed in this study. Additionally, accumulation by repetitive
287 administration and differences due to the gestational stage were not observed for VPA or 4-ene-VPA
288 (Table 1). It has been reported that 4-ene-VPA was detected in both humans and rats after
289 administration of VPA, and has been suggested to be involved in toxicity, such as hepatotoxicity
290 (Kesterson JW et al., 1984; Tennison MB et al., 1998).

291 VPA administration is known to affect fetal growth and cognitive development in rats as well
292 as humans (Ornoy, 2009; Schneider and Przewłocki, 2005). Maternal body weight loss was shown to

293 occur after the administration of 400 mg/kg VPA on G7–18 (Vorhees, 1987). Our results showed no
294 significant difference in maternal body weight (data not shown); however, body weight gain showed
295 a similar trend (Figure 3). Fetal weight also decreased in the VPA-treated group by approximately
296 1.5 g in a previous study (Vorhees, 1987). Hence, fetal growth restriction might also contribute to the
297 decreased maternal body weight gain. In humans, contradictory studies exist regarding the
298 relationship between VPA and fetal growth. A population-based cohort study indicated that VPA use
299 did not induce fetal growth restriction (Veiby et al., 2014), whereas another cohort study showed that
300 the maternal and umbilical cord levels of VPA were negatively correlated with birth length
301 (Kacirova et al., 2015). Therefore, VPA concentration in the plasma might be important for normal
302 fetal growth.

303 Mdr1a, Mdr1b, Mrp1–3, and Bcrp, which play a key role in predominantly transferring
304 xenobiotics. Mdr1a, Mdr1b, Mrp2, Mrp3 and Bcrp localize to the apical membrane of placenta.
305 Mrp1 localize to the basolateral membrane. Furthermore, Mrp1 and Mrp3 are expressed in fetal
306 vessels (Joshi et al., 2016; Ni and Mao, 2011). In this study, Mdr1a, Mdr1b, and Mrp2 were
307 increased, whereas Mrp3 and Bcrp decreased through gestation (Figure 1). It has been reported that
308 Mdr1a and Mdr1b increased with gestational stage, which was consistent with our results (Novotna
309 M et al., 2004). In regard to Mrp1-3, St-Pierre et al. showed that Mrp1 mRNA was abundantly
310 expressed in the placenta throughout gestation (St-Pierre MV et al., 2004). The group reported that
311 Mrp3 mRNA was expressed throughout gestation, whereas Mrp2 was expressed at a low level in the

312 rat placenta. Repetitive VPA administration decreased Mdr1b throughout gestation and affected
313 Mdr1a, Mrp1, and Mrp3 in a gestational age-specific manner (Figure 1). In the G20 placenta,
314 repetitive VPA administration increased Mdr1a, whereas it tended to decrease Mdr1b. Therefore, we
315 examined the expression of P-gp protein, which is encoded by *Abcb1a* and *Abcb1b*, by western
316 blotting (Supplemental Figure 2). P-gp protein tended to increase by repetitive VPA administration,
317 although the difference was not statistically significant. The result suggested that change in *Abcb1a*
318 affected the P-gp protein expression in rat placenta, and variations of gene expression by VPA can
319 alter protein levels.

320 Bcrp on the apical side and Oatp2b1 on the basolateral side collaborate to transport sulfate
321 conjugates from fetal to maternal circulation (Grube et al., 2007). As gestation progresses, these two
322 genes showed opposite changes as follows: Bcrp decreased to half the initial level (Figure 1H),
323 whereas Oatp2b1 increased by 4-folds (Figure 2I). It has been reported that Bcrp mRNA in the rat
324 placenta peaks on G15 and declines significantly to one third at term (Cygalova L et al., 2008).
325 Oatp2b1 mRNA increased throughout gestation (St-Pierre MV et al., 2004). The tendency was
326 consistent with the present results. Oatp2b1 was decreased on G20 by single VPA administration;
327 however, no significant differences were noted after repetitive treatment (Figure 2I).

328 Mrp4 on the apical side and Mrp5 on the basolateral side contribute to the transport of cyclic
329 nucleotides (Joshi et al., 2016; Wielinga et al., 2003). In this study, Mrp4 increased and Mrp5
330 decreased, following fetal development (Figure 1). Moreover, VPA remarkably decreased Mrp4 on

331 G20 (Figure 1F) and increased Mrp5 on G13 (Figure 1G). Considering these findings, VPA is
332 thought to increase cAMP efflux from the placenta to the fetus on G13, whereas decreasing it from
333 the placenta to mother on G20. In the placentas, one of the cyclic nucleotides, cAMP, is involved in
334 cell fusion and cytotrophoblast syncytialization, which is caused by an increase in Syncytin-1 via
335 cAMP/Protein kinase A signaling (Gupta et al., 2016). Hence, the altered expression of cAMP
336 transporters might affect placental cell differentiation and function.

337 Mrp4 is also involved in transporting prostaglandins (Reid et al., 2003). Additionally,
338 Oatp2a1 has been shown to transport prostaglandins (Gose et al., 2016), which was increased by
339 VPA on G13 in the present study (Figure 2H). Prostaglandin E2 is involved in cell invasion and
340 migration in the placenta (Nicola et al., 2005). Thus, VPA might alter placental kinetics of
341 prostaglandins at both G13 and G20.

342 Lat1 on the apical membrane and Lat2 on both apical and basal membranes are transporters
343 exchanging large neutral amino acids (del Amo et al., 2008; Gaccioli et al., 2015). The expression of
344 Lat1 increased remarkably as gestation progressed (Figure 2A). VPA treatment decreased Lat1 on
345 G20 (Figure 2A), whereas increased Lat2 on G13 (Figure 2B). A previous study indicated the
346 involvement of Lat1 in the transport of branched-chain amino acids in mice fetal brain, and
347 impairment of this transporter led to neurological and behavioral abnormalities associated with
348 autism spectrum disorders (Tărlungeanu et al., 2016). Since VPA induces fetal autism spectrum
349 disorders, evaluation of its effect on placental Lat1 might have important implications. Furthermore,

350 Lat1, Lat2, and Oatp4a1 transport thyroid hormones, which are necessary for fetal growth (Bernal et
351 al., 2015; Forhead and Fowden, 2014). Oatp4a1 was shown to have increased following gestation
352 and decreased by VPA on G20 (Figure 2J). The change in Oatp4a1 mRNA throughout gestation was
353 consistent with the previous results (St-Pierre MV et al., 2004).

354 Ent1 transports pyrimidine nucleosides (Nishimura et al., 2012). Ent1 decreased with
355 gestational stage and was increased by VPA on G13 (Figure 2J). The expression of other nucleoside
356 transporters, Cnt1 and Ent2, was found to be low on G20 (data not shown).

357 Mct4 mediates proton-dependent transport of monocarboxylates. Mct4 was increased by
358 gestational development. It has been reported that the MCT4 protein was strongly detected in rat
359 placentas throughout gestation (Moore et al., 2016). VPA did not affect the expression levels across
360 gestation (Figures 2C).

361 Higher levels of Oct3 in the placenta, rather than in other organs, play a role in the efflux of
362 organic cations from the fetus (Leazer and Klaassen, 2003; Sata et al., 2005). In contrast, Oct2 was
363 present at low levels in the placenta on G20 (data not shown). Oct3 was increased by gestational
364 development or VPA single administration; however, the differences were subtle (Figure 2D). It has
365 been reported that Oct3 mRNA in the rat placenta increased throughout gestation
366 (Ahmadimoghaddam et al., 2013). The tendency was consistent with our results.

367 In this study, Octn1 increased markedly following gestational development and decreased by
368 VPA administration on G20 (Figure 2E). Octn2 increased by VPA repetitive administration on G13

369 (Figure 2F). Octn1 and Octn2 play a crucial role in the uptake of L-carnitine by the placenta (Grube
370 et al., 2005; Wu et al., 2000). The defect in Octn2 was shown to induce embryonic lethality
371 (Shekhawat et al., 2018). Although the role of placental Octn1 has been poorly understood, the
372 decline in Octn1, whose substrates are similar to those of Octn2, might affect fetal growth. The
373 remarkable increase in Octn1 during pregnancy might have a critical impact; thus, further
374 investigation is needed to elucidate its role. Furthermore, previous studies showed that serum
375 carnitine levels were altered after VPA treatment (Moreno et al., 2005). Carnitine was also reported
376 to alter Octn2 expression, which might be involved in Octn2 alteration in the present study (Schürch
377 et al., 2010).

378 In the present study, variation in transporters in the placenta by VPA administration were
379 observed. As described above, repetitive VPA administration increased Mdr1a, whereas it tended to
380 decrease Mdr1b in the G20 placenta. Furthermore, VPA administration significantly decreased Mrp4
381 expression at G20. Because these are efflux transporters at the apical membrane, the changes might
382 cause variation in placental barrier function. In regard to cAMP transporters, Mrp4 and Mrp5 were
383 changed in VPA; Mrp4 decreased at G20, whereas Mrp5 increased at G13. Because cAMP signaling
384 is involved in cell fusion and syncytialization (Gupta et al., 2016), the altered expression of cAMP
385 transporters might affect placental cell differentiation and function. VPA administration affected the
386 expression SLC transporters that mediate transport of nutrition. VPA decreased Lat1, Octn1, and
387 Oatp4a1 in the G20 placenta. These are influx transporters at the apical membrane, and are important

388 for fetal development, such as amino acids, carnitine, and thyroid hormones. Additionally,
389 prostaglandin transporter Oatp2a1 was increased by VPA during G13 in the present study.
390 Prostaglandin E2 was involved in cell invasion and migration in the placenta (Nicola et al., 2005).
391 Therefore, these changes might contribute to impairment of the placental and fetal development by
392 VPA. The present results and previous knowledge suggest that the changes might be linked to
393 placental function and fetal nutrition/development. However, we have not directly evaluated the links
394 between changes in transporters and adverse effects by VPA in the present study. Future studies are
395 required to assess whether the changes of placental transporters affect placental function, fetal
396 nutrition, and development. In addition, we have not revealed changes at the protein levels, except
397 for P-gp. It is essential to evaluate protein levels for precisely justifying placental transport function.
398 Future studies should investigate the changes in proteins and function levels by VPA administration.

399 It has been reported that transporters can be affected in a sex-specific manner in some cases.
400 For instance, a maternal high-fat diet has been shown to increase GLUT3 and system A amino acid
401 transporter 2 only in the placentas from male fetuses (Song et al., 2017). Previous studies evaluating
402 the effects of VPA on the expression of placental transporters did not elucidate fetal sex differences.
403 Therefore, we secondarily analyzed by fetal sex. In this study, Mdr1a on G13 and Oatp2a1 on G20
404 were shown to have higher susceptibilities to VPA in the male placenta (Supplemental Figure 1). A
405 study showed that sex differences altered the behavior of VPA-treated rats; thus, VPA has the
406 potential to affect various aspects in a sex-specific manner (Anshu et al., 2017). However, the present

407 study had a limited number of placentas for each sex because randomized placentas were divided
408 into male or female. Thus, further investigations are needed to determine the effect of fetal sex.

409 In the present study, we used rats as the pregnant animal model to investigate the effects of
410 VPA on placental transporters because an *in vivo* study is suited to the investigation of changes
411 throughout gestation. However, species difference between human and rat placentas should be
412 considered to understand the results. First, there are structural differences in the placental barrier.
413 The human placental barrier consists of a syncytiotrophoblast layer. In rats, the placental labyrinth
414 has three trophoblastic layers. Layer II and III are thought to be the syncytium, whereas the first layer
415 that faces the maternal side does not represent a barrier (Joshi AA et al., 2016). Second, there are
416 some transporters whose localization are different between humans and rats, although the
417 transporters or orthologs we investigated in the present study have been reported to be expressed in
418 human placentas (Bleasby K et al., 2006; Gaccioli F et al., 2015; Joshi AA et al., 2016). For instance,
419 it has been reported that human MCT4 (*SLC16A3*) is expressed on the maternal side of placenta,
420 whereas rat Mct4 (*Slc16a3*) is expressed on the fetal side (Moore NP et al., 2016; Settle P et al.,
421 2004). Other transporter localizations have been reported. It has been reported that both human P-gp
422 (MDR1, *ABCB1*) and rat P-gp (Mdr1a/1b, *Abcb1a/1b*), BCRP (*ABCG2*), Bcrp (*Abcg2*), ENT1
423 (*SLC29A1*), Ent1 (*Slc29a1*), OATP4A1 (*SLCO4A1*), and Oatp4a1 (*Slco4a1*) express apical
424 membranes (Akashi T et al., 2016; Joshi AA et al., 2016; Nishikawa M et al., 2010; Nishimura T et
425 al., 2019; Sato K et al., 2003). In addition, human MRP1 (*ABCC1*) expression at the basolateral side

of the syncytiotrophoblast of human placental villi agrees with Mrp1 expression on the basolateral side of the syncytiotrophoblast in the rat labyrinth zone (St-Pierre MV et al., 2004). Human MRP2, 4 (*ABCC2*, 4) and OATP2A1 (*SLCO2A1*) have been detected in the apical membrane of the placenta (St-Pierre MV et al., 2002; Joshi AA et al., 2016), whereas their localization in the rat placenta have not been determined. Human MRP5 (*ABCC5*) have been shown to be expressed in the basolateral membrane of the syncytiotrophoblast (Joshi AA et al., 2016), whereas its localization in the rat placenta remains unclear. It has been reported that rat Octn1, 2 (*Slc22a4*, 5) is expressed in the apical membrane (St-Pierre MV., 2002). Human LAT1, 2 (*SLC7A5*, 8) have been reported to be present on the apical side of the syncytiotrophoblast, whereas LAT2 is also expressed in the basolateral membrane and in the fetal capillary endothelium (Gaccioli F et al., 2015). Lat1, 2 (*Slc7a5*, 8) is also expressed in the apical membrane in the rat placenta (Rosario FJ et al., 2011). Third, variation in placental expression of some transporters throughout gestation differs. For instance, human P-gp (*MDR1*, *ABCB1*) expression was reported to decrease with advancing gestation, whereas rat P-gp (*Mdr1a/1b*, *Abcb1a/1b*) increased (Joshi AA et al., 2016). Human placental OCT3 (*SLC22A3*) expression during the first trimester was reported to be higher than that at term, whereas rat Oct3 (*Slc22a3*) increased with advancing gestation (Ahmadimoghaddam et al., 2013). Human placental OCTN2 (*SLC22A5*) expression during the first trimester was reported to be lower than that at term (Bai et al., 2019), whereas rat Octn2 (*Slc22a5*) was not changed with advancing gestation in the present study.

445 In conclusion, we assessed the effect of VPA on the expression of rat placental transporters
446 during pregnancy and revealed that sensitivity to VPA differed following gestational development.
447 Our findings also indicated that sex differences might exist in these alterations of transporters. To the
448 best of our knowledge, this is the first study to longitudinally and systematically investigate the
449 influence of VPA on various placental transporters. As described above, prenatal exposure to VPA
450 increases various risks. Future studies are required to better understand the association of the effects
451 of VPA with fetal risks and provide an approach for relieving these risks.

452

453 **Acknowledgment**

454 **Funding**

455 This work was supported in part by a grant from the Japan Society for the Promotion of Science
456 (JSPS) KAKENHI (grant number, 18K1497208) (provided to A.F.).

457

458 **Conflicts of Interest**

459 The authors declare no conflicts of interest.

460

461 **References**

- 462 Ahmadimoghaddam D, Zemankova L, Nachtigal P, Dolezelova E, Neumanova Z, Cervený L,
463 Ceckova M, Kacerovský M, Micuda S, Staud F (2013) Organic cation transporter 3
464 (OCT3/SLC22A3) and multidrug and toxin extrusion 1 (MATE1/SLC47A1) transporter in
465 the placenta and fetal tissues: expression profile and fetus protective role at different stages of
466 gestation. *Biol Reprod* 88(3):55.
- 467 Akashi T, Nishimura T, Takaki Y, Takahashi M, Shin BC, Tomi M, Nakashima E (2016) Layer II of
468 placental syncytiotrophoblasts expresses MDR1 and BCRP at the apical membrane in rodents.
469 *Reprod Toxicol* 65:375-381.
- 470 Anshu K, Nair AK, Kumaresan UD, Kutty BM, Srinath S, and Laxmi TR (2017) Altered attentional
471 processing in male and female rats in a prenatal valproic acid exposure model of autism
472 spectrum disorder. *Autism Res* 10(12): 1929-1944.
- 473 Bai M, Zeng Q, Chen Y, Chen M, Li P, Ma Z, Sun D, Zhou H, Zheng C, Zeng S, Jiang H (2019)
474 Maternal Plasma l-Carnitine Reduction During Pregnancy Is Mainly Attributed to
475 OCTN2-Mediated Placental Uptake and Does Not Result in Maternal Hepatic Fatty Acid
476 β -Oxidation Decline. *Drug Metab Dispos* 47(6):582-591.
- 477 Baker GA, Bromley RL, Briggs M, Cheyne CP, Cohen MJ, García-Fiñana M, Gummery A, Kneen R,
478 Loring DW, Mawer G, Meador KJ, Shallcross R, and Clayton-Smith J; Liverpool and

479 Manchester Neurodevelopment Group (2015) IQ at 6 years after in utero exposure to
 480 antiepileptic drugs: a controlled cohort study. *Neurology* 84(4): 382-390.

481 Bernal J, Guadaño-Ferraz A, and Morte B (2015) Thyroid hormone transporters--functions and
 482 clinical implications. *Nat Rev Endocrinol* 11(7):406-417.

483 Binkerd PE, Rowland JM, Nau H, and Hendrickx AG (1988) Evaluation of valproic acid (VPA)
 484 developmental toxicity and pharmacokinetics in Sprague-Dawley rats. *Fundam Appl Toxicol*
 485 11(3): 485-493.

486 Bleasby K, Castle JC, Roberts CJ, Cheng C, Bailey WJ, Sina JF, Kulkarni AV, Hafey MJ, Evers R,
 487 Johnson JM, Ulrich RG, and Slatter JG (2006) Expression profiles of 50 xenobiotic
 488 transporter genes in humans and pre-clinical species: a resource for investigations into drug
 489 disposition. *Xenobiotica* 36(10-11): 963-988.

490 Christensen J, Grønborg TK, Sørensen MJ, Schendel D, Parner ET, Pedersen LH, and Vestergaard M
 491 (2013) Prenatal valproate exposure and risk of autism spectrum disorders and childhood
 492 autism. *JAMA* 309(16): 1696-1703.

493 Cygalova L, Ceckova M, Pavek P, Staud F (2008) Role of breast cancer resistance protein
 494 (Bcrp/Abcg2) in fetal protection during gestation in rat. *Toxicol Lett* 178(3):176–180.

495 del Amo EM, Urtti A, and Yliperttula M (2008) Pharmacokinetic role of L-type amino acid
 496 transporters LAT1 and LAT2. *Eur J Pharm Sci* 35(3): 161-174.

497 Forhead AJ and Fowden AL (2014) Thyroid hormones in fetal growth and prepartum maturation. *J*
498 *Endocrinol* 221(3): R87-R103.

499 Gaccioli F, Aye IL, Roos S, Lager S, Ramirez VI, Kanai Y, Powell TL, and Jansson T (2015)
500 Expression and functional characterisation of System L-amino acid transporters in the human
501 term placenta. *Reprod Biol Endocrinol* 13: 57.

502 Gao S, Miao H, Tao X, Jiang B, Xiao Y, Cai F, Yun Y, Li J, and Chen W (2011) LC-MS/MS
503 method for simultaneous determination of valproic acid and major metabolites in human
504 plasma. *J Chromatogr B Analyt Technol Biomed Life Sci* 879(21): 1939-1944.

505 Gose T, Nakanishi T, Kamo S, Shimada H, Otake K, and Tamai I (2016) Prostaglandin transporter
506 (OATP2A1/SLCO2A1) contributes to local disposition of eicosapentaenoic acid-derived
507 PGE3. *Prostaglandins Other Lipid Mediat* 122: 10-17.

508 Grabiec AM and Potempa J (2018) Epigenetic regulation in bacterial infections: targeting histone
509 deacetylases. *Crit Rev Microbiol* 44(3): 336-350.

510 Grube M, Meyer Zu Schwabedissen H, Draber K, Präger D, Möritz KU, Linnemann K, Fusch C,
511 Jedlitschky G, and Kroemer HK (2005) Expression, localization, and function of the carnitine
512 transporter octn2 (slc22a5) in human placenta. *Drug Metab Dispos* 33(1): 31-37.

513 Grube M, Reuther S, Meyer Zu Schwabedissen H, Köck K, Draber K, Ritter CA, Fusch C,
514 Jedlitschky G, and Kroemer HK (2007) Organic anion transporting polypeptide 2B1 and

515 breast cancer resistance protein interact in the transepithelial transport of steroid sulfates in
516 human placenta. *Drug Metab Dispos* 35(1): 30-35.

517 Gupta SK, Malhotra SS, Malik A, Verma S, and Chaudhary P (2016) Cell signaling pathways
518 involved during invasion and syncytialization of trophoblast cells. *Am J Reprod Immunol*
519 75(3): 361-371.

520 Gurvich N, Tsygankova OM, Meinkoth JL, and Klein PS (2004) Histone deacetylase is a target of
521 valproic acid-mediated cellular differentiation. *Cancer Res* 64(3): 1079-1086.

522 Ishikawa T, Obara T, Jin K, Nishigori H, Miyakoda K, Suzuka M, Ikeda-Sakai Y, Akazawa M,
523 Nakasato N, Yaegashi N, Kuriyama S, and Mano N (2019) Examination of the prescription of
524 antiepileptic drugs to prenatal and postpartum women in Japan from a health administrative
525 database. *Pharmacoepidemiol Drug Saf* 28(6): 804-811.

526 Jentink J, Loane MA, Dolk H, Barisic I, Garne E, Morris JK, and de Jong-van den Berg LT;
527 EUROCAT Antiepileptic Study Working Group (2010) Valproic acid monotherapy in
528 pregnancy and major congenital malformations. *N Engl J Med* 362(23): 2185-2193.

529 Joshi AA, Vaidya SS, St-Pierre MV, Mikheev AM, Desino KE, Nyandege AN, Audus KL, Unadkat
530 JD, and Gerk PM (2016) Placental ABC transporters: Biological impact and pharmaceutical
531 significance. *Pharm Res* 33(12): 2847-2878.

532 Kacirova I, Grundmann M, and Brozmanova H (2015) Serum levels of valproic acid during delivery
 533 in mothers and in umbilical cord—correlation with birth length and weight. *Biomed Pap Med*
 534 *Fac Univ Palacky Olomouc Czech Repub* 159(4): 569-575.

535 Kalisch-Smith JI, Simmons DG, Dickinson H, and Moritz KM (2017) Review: Sexual dimorphism
 536 in the formation, function and adaptation of the placenta. *Placenta* 54: 10-16.

537 Kim KB, Seo KA, Kim SE, Bae SK, Kim DH, and Shin JG (2011) Simple and accurate quantitative
 538 analysis of ten antiepileptic drugs in human plasma by liquid chromatography/tandem mass
 539 spectrometry. *J Pharm Biomed Anal* 56(4): 771-777.

540 Kesterson JW, Granneman GR, and Machinist JM (1984) The hepatotoxicity of valproic acid and its
 541 metabolites in rats. I. Toxicologic, biochemical and histopathologic studies. *Hepatology*
 542 4(6):1143-1152.

543 Kurosawa Y, Furugen A, Nishimura A, Narumi K, Kobayashi M, and Iseki K (2018) Evaluation of
 544 the effects of antiepileptic drugs on folic acid uptake by human placental choriocarcinoma
 545 cells. *Toxicol In Vitro* 48, 104-110.

546 Leazer TM and Klaassen CD (2003) The presence of xenobiotic transporters in rat placenta. *Drug*
 547 *Metab Dispos* 31(2): 153-167.

548 Meir M, Bishara A, Mann A, Udi S, Portnoy E, Shmuel M, and Eyal S (2016) Effects of valproic
 549 acid on the placental barrier in the pregnant mouse: Optical imaging and transporter
 550 expression studies. *Epilepsia* 57(6): e108-112.

551 Moore NP, Picut CA, and Charlap JH (2016) Localisation of Lactate Transporters in Rat and Rabbit
552 Placentae. *Int J Cell Biol* 2016:2084252.

553 Moreno FA, Macey H, and Schreiber B (2005) Carnitine levels in valproic acid-treated psychiatric
554 patients: A cross-sectional study. *J Clin Psychiatry* 66(5): 555-558.

555 Nicola C, Timoshenko AV, Dixon SJ, Lala PK, and Chakraborty C (2005) EP1 receptor-mediated
556 migration of the first trimester human extravillous trophoblast: the role of intracellular
557 calcium and calpain. *J Clin Endocrinol Metab* 90(8):4736-4746.

558 Nishikawa M, Iwano H, Yanagisawa R, Koike N, Inoue H, Yokota H (2010) Placental transfer of
559 conjugated bisphenol A and subsequent reactivation in the rat fetus. *Environ Health Perspect*
560 118(9):1196-1203.

561 Nishimura T, Chishu T, Tomi M, Nakamura R, Sato K, Kose N, Sai Y, and Nakashima E (2012)
562 Mechanism of nucleoside uptake in rat placenta and induction of placental CNT2 in
563 experimental diabetes. *Drug Metab Pharmacokinet* 27(4): 439-446.

564 Nishimura T, Sano Y, Takahashi Y, Noguchi S, Uchida Y, Takagi A, Tanaka T, Katakura S,
565 Nakashima E, Tachikawa M, Maruyama T, Terasaki T, Tomi M (2019) Quantification of
566 ENT1 and ENT2 Proteins at the Placental Barrier and Contribution of These Transporters to
567 Ribavirin Uptake. *J Pharm Sci* 108(12):3917-3922.

568 Ni Z and Mao Q (2011) ATP-binding cassette efflux transporters in human placenta. *Curr Pharm*
569 *Biotechnol* 12(4): 674-685.

570 Novotna M, Libra A, Kopecky M, Pavek P, Fendrich Z, Semecky V, and Staud F (2004)
 571 P-glycoprotein expression and distribution in the rat placenta during pregnancy. *Reprod*
 572 *Toxicol* 18(6): 785-792.

573 Ornoy A (2009) Valproic acid in pregnancy: how much are we endangering the embryo and fetus?
 574 *Reprod Toxicol* 28(1): 1-10.

575 Reid G, Wielinga P, Zelcer N, van der Heijden I, Kuil A, de Haas M, Wijnholds J, and Borst P
 576 (2003) The human multidrug resistance protein MRP4 functions as a prostaglandin efflux
 577 transporter and is inhibited by nonsteroidal antiinflammatory drugs. *Proc Natl Acad Sci U S A*
 578 100(16): 9244-9249.

579 Rosario FJ, Jansson N, Kanai Y, Prasad PD, Powell TL, Jansson T (2011) Maternal protein
 580 restriction in the rat inhibits placental insulin, mTOR, and STAT3 signaling and
 581 down-regulates placental amino acid transporters. *Endocrinology* 152(3):1119-29.

582 Rouillet FI, Lai JK, Foster JA (2013) In utero exposure to valproic acid and autism--a current review
 583 of clinical and animal studies. *Neurotoxicol Teratol* 36:47-56.

584 Rubinchik-Stern M, Shmuel M, Bar J, Kovo M, and Eyal S (2018) Adverse placental effects of
 585 valproic acid: Studies in perfused human placentas. *Epilepsia* 59(5): 993-1003.

586 Rubinchik-Stern M, Shmuel M, and Eyal S (2015) Antiepileptic drugs alter the expression of
 587 placental carriers: An in vitro study in a human placental cell line. *Epilepsia* 56(7):
 588 1023-1032.

589 Sata R, Ohtani H, Tsujimoto M, Murakami H, Koyabu N, Nakamura T, Uchiumi T, Kuwano M,
 590 Nagata H, Tsukimori K, Nakano H, and Sawada Y (2005) Functional analysis of organic
 591 cation transporter 3 expressed in human placenta. *J Pharmacol Exp Ther* 315(2): 888-895.
 592 Sato K, Sugawara J, Sato T, Mizutamari H, Suzuki T, Ito A, Mikkaichi T, Onogawa T, Tanemoto M,
 593 Unno M, Abe T, Okamura K (2003) Expression of organic anion transporting polypeptide E
 594 (OATP-E) in human placenta. *Placenta* 24(2-3):144-8.
 595 Schneider T and Przewłocki R (2005) Behavioral alterations in rats prenatally exposed to valproic
 596 acid: animal model of autism. *Neuropsychopharmacology* (1): 80-89.
 597 Schürch R, Todesco L, Novakova K, Mevissen M, Stieger B, and Krähenbühl S (2010) The plasma
 598 carnitine concentration regulates renal OCTN2 expression and carnitine transport in rats. *Eur*
 599 *J Pharmacol* 635(1-3): 171-176.
 600 Settle P, Mynett K, Speake P, Champion E, Doughty IM, Sibley CP, D'Souza SW, and Glazier J
 601 (2004) Polarized lactate transporter activity and expression in the syncytiotrophoblast of the
 602 term human placenta. *Placenta* 25(6):496-504.
 603 Shekhawat PS, Sonne S, Matern D, and Ganapathy V (2018) Embryonic lethality in mice due to
 604 carnitine transporter OCTN2 defect and placental carnitine deficiency. *Placenta* 69: 71-73.
 605 Song L, Sun B, Boersma GJ, Cordner ZA, Yan J, Moran TH, and Tamashiro K (2017) Prenatal
 606 high-fat diet alters placental morphology, nutrient transporter expression, and mtorc1
 607 signaling in rat. *Obesity (Silver Spring)* 25(5): 909-919.

608 Staud F, Cervený L, and Cecková M (2012) Pharmacotherapy in pregnancy; effect of ABC and SLC
609 transporters on drug transport across the placenta and fetal drug exposure. *J Drug Target*
610 20(9): 736-763.

611 St-Pierre MV, Hagenbuch B, Ugele B, Meier PJ, Stallmach T (2002) Characterization of an organic
612 anion-transporting polypeptide (OATP-B) in human placenta. *J Clin Endocrinol Metab*
613 87(4):1856-63.

614 St-Pierre MV, Stallmach T, Freimoser Grundschober A, Dufour JF, Serrano MA, Marin JJ,
615 Sugiyama Y, Meier PJ (2004) Temporal expression profiles of organic anion transport
616 proteins in placenta and fetal liver of the rat. *Am J Physiol Regul Integr Comp Physiol*
617 287(6):R1505-1516.

618 St-Pierre MV, Ugele B, Gambling L, Shiverick KT (2002) Mechanisms of drug transfer across the
619 human placenta-a workshop report. *Placenta* 23 Suppl A:S159-164

620 Sun B, Fu A, Wang R, and Zhang Y (2015) Influence of carbon dioxide pneumoperitoneum on the
621 growth hormone-insulin-like growth factor I axis in mid- and late-pregnancy rats. *J Obstet*
622 *Gynaecol Res* 41(9): 1394-1398.

623 Tărlungeanu DC, Deliu E, Dotter CP, Kara M, Janiesch PC, Scalise M, Galluccio M, Tesulov M,
624 Morelli E, Sonmez FM, Bilguvar K, Ohgaki R, Kanai Y, Johansen A, Esharif S, Ben-Omran
625 T, Topcu M, Schlessinger A, Indiveri C, Duncan KE, Caglayan AO, Gunel M, Gleeson JG,

626 and Novarino G (2016) Impaired amino acid transport at the blood brain barrier is a cause of
 627 autism spectrum disorder. *Cell* 167(6): 1481-1494.e18.

628 Tennison MB, Miles MV, Pollack GM, Thorn MD, and Dupuis RE (1988) Valproate metabolites and
 629 hepatotoxicity in an epileptic population. *Epilepsia* 29(5):543-547.

630 Tetro N, Imbar T, Wohl D, Eisenberg I, Yagel S, Shmuel M, and Eyal S (2019) The effects of
 631 valproic acid on early pregnancy human placentas: Pilot ex vivo analysis in cultured placental
 632 villi. *Epilepsia* 60(5): e47-e51.

633 Tomson T, Battino D, Bonizzoni E, Craig J, Lindhout D, Sabers A, Perucca E, and Vajda F; EURAP
 634 study group (2011) Dose-dependent risk of malformations with antiepileptic drugs: an
 635 analysis of data from the EURAP epilepsy and pregnancy registry. *Lancet Neurol* 10(7):
 636 609-617.

637 Veiby G, Daltveit AK, Engelsen BA, and Gilhus NE (2014) Fetal growth restriction and birth defects
 638 with newer and older antiepileptic drugs during pregnancy. *J Neurol* 261(3): 579-588.

639 Viinikainen K, Heinonen S, Eriksson K, and Kälviäinen R (2006) Community-based, prospective,
 640 controlled study of obstetric and neonatal outcome of 179 pregnancies in women with
 641 epilepsy. *Epilepsia* 47(1): 186-192.

642 Vorhees CV (1987) Teratogenicity and developmental toxicity of valproic acid in rats. *Teratology*
 643 35(2): 195-202.

644 Wielinga PR, van der Heijden I, Reid G, Beijnen JH, Wijnholds J, and Borst P (2003)
 645 Characterization of the MRP4- and MRP5-mediated transport of cyclic nucleotides from
 646 intact cells. *J Biol Chem* 278(20): 17664-17671.

647 Wu CY and Lu CY (2014) Derivatization oriented strategy for enhanced detection of valproic acid
 648 and its metabolites in human plasma and detection of valproic acid induced reactive oxygen
 649 species associated protein modifications by mass spectrometry. *J Chromatogr A* 1374: 14-22.

650 Wu X, George RL, Huang W, Wang H, Conway SJ, Leibach FH, and Ganapathy V (2000) Structural
 651 and functional characteristics and tissue distribution pattern of rat OCTN1, an organic cation
 652 transporter, cloned from placenta. *Biochim Biophys Acta* 1466(1-2): 315-327.

653 Yoshimura K, Hashimoto T, Sato Y, Sato A, Takeuchi T, Watanabe H, Terao T, Nakazato M, and
 654 Iyo M (2018) Survey of anticonvulsant drugs and lithium prescription in women of
 655 childbearing age in Japan using a public National Insurance Claims Database. *Clinical*
 656 *Neuropsychopharmacology and Therapeutics* 9: 20-29.
 657

658 **Figure legends**

659 **Figure 1** Effects of single and repetitive VPA Na administration on the expression of rMdr1a (A),
660 rMdr1b (B), rMrp1 (C), rMrp2 (D), rMrp3 (E), rMrp4 (F), rMrp5 (G), and rBcrp (H) mRNAs in rat
661 placentas. Pregnant rats were orally administered 400 mg/kg VPA Na (black) or distilled water as a
662 control (white). Rats in the single administration groups were treated during G12 (mid-gestation) or
663 G19 (late gestation), and those in the repetitive groups were treated for 4 d during G9–12
664 (mid-gestation) or G16–19 (late gestation). Expression of transporters was assessed by real-time PCR.
665 Each column represents the mean with SD (Single administration group: n = 9–12 placentas from
666 three dams; Repetitive administration group: n = 10–12 placentas from three dams). *: $P < 0.05$
667 when compared to the G20 control. **: $P < 0.01$ when compared to the G20 control. †: $P < 0.05$
668 when compared to the G13 control. ††: $P < 0.01$ when compared to the G13 control. ND: not
669 determined because of low expression. G: gestational day. Gene expression was analyzed using an
670 ANOVA, with “G” and “VPA” as factors. If an interaction was present, the Tukey–Kramer test was
671 used for multiple comparisons.

672

673 **Figure 2** Effects of single and repetitive VPA Na administration on the expression of rLat1 (A),
674 rLat2 (B), rMct4 (C), rOct3 (D), rOctn1 (E), rOctn2 (F), rEnt1(G), rOatp2a1 (H), rOatp2b1 (I), and
675 rOatp4a1 (J) mRNAs in rat placentas. Pregnant rats were orally administered 400 mg/kg VPA Na
676 (black) or distilled water as a control (white). Rats of the single administration groups were treated

during G12 (mid-gestation) or G19 (late gestation), and those in the repetitive groups were treated for 4 d during G9–12 (mid-gestation) or G16–19 (late gestation). Expression of transporters was assessed by real-time PCR. Each column represents the mean with SD (Single administration group: n = 9–12 placentas from three dams; Repetitive administration group: n = 10–12 placentas from three dams). *: $P < 0.05$ when compared to the G20 control. **: $P < 0.01$ when compared to the G20 control. ††: $P < 0.01$ when compared to the G13 control. G: gestational day. Gene expression was analyzed using an ANOVA, with “G” and “VPA” as factors. If an interaction was present, the Tukey–Kramer test was used for multiple comparisons.

Figure 3 Body weight gain (A) and placental weight (B) of pregnant rats orally administered 400 mg/kg VPA Na or distilled water as control for 4 days on G9–12 or G16–19, respectively. (A) Maternal body weight gain was calculated by subtracting the body weight on the first day of administration from those of each gestational day. Each point represents the mean with SD (n = 3 dams). *: $P < 0.05$ when compared to the control. (B) Each column represents the mean with SD (n = 40 placentas for G13 control, 34 for G13 VPA, 32 for G20 control, 33 for G20 VPA placentas from three dams. **: $P < 0.01$ when compared to each control. G: gestational day. Student’s t-test was used for comparison of the control and VPA groups.

696 **Supplemental Figure 1** Interactive effects of VPA Na administration and fetal sex on rMdr1a
697 expression in G13 rat placentas (A), and those of rOatp2a1 (B) and rOatp2b1 (C) in G20 rat
698 placentas. Pregnant rats were orally administered 400 mg/kg/day VPA Na (black) or distilled water
699 as a control (white) on G12 (A) or G16–19 (B, C). Expression of transporters was assessed by
700 real-time PCR. Each column represents the mean with SD ($n = 3\text{--}7$ placentas from two or three
701 dams). *: $P < 0.05$ when compared to the male control. **: $P < 0.01$ when compared to the male
702 control. G: gestational day. Gene expression was analyzed using a two-way analysis of ANOVA,
703 with “sex” and “VPA” as factors. If an interaction was present, the Tukey–Kramer test was used for
704 multiple comparisons.

705
706 **Supplemental Figure 2** Effects of repetitive VPA Na administration on P-gp protein expression in
707 rat placentas (G20). Pregnant rats were orally administered 400 mg/kg/day VPA Na (black) or
708 distilled water as a control (white), for 4 d. Whole proteins were extracted from placentas of G20
709 pregnant rats with repetitive administrated of VPA or water (control). Expression of P-gp was
710 assessed by western blotting. Fetal sex was not determined for the protein samples. (A) Data shown
711 are typical results of three independent experiments. (B) Each column represents the mean with SD
712 ($n = 12$ placentas from three dams).

713 **Table 1** Concentration of VPA and 4-ene-VPA in the plasma of pregnant rats orally administered 400 mg/kg VPA Na.

			Mid-gestation				Late gestation					
			VPA		4-ene-VPA		VPA		4-ene-VPA			
			(µg/mL)		(µg/mL)		(µg/mL)		(µg/mL)			
			Mean	SD	Mean	SD	Mean	SD	Mean	SD		
Single administration	30 min	G12	211	28.9	1.49	0.288	G19	175	57.2	1.50	0.165	
	24 h	G13	a		a		G20	a		0.827	0.543	
Repetitive administration	Day 1	30 min	G9	159	38.8	1.62	0.0858	G16	241	4.93	1.41	0.230
	Day 2	24 h	G10	a		a		G17	a		a	
		30 min		160	14.5	2.02	0.680		261	157	1.36	0.361
	Day 3	24 h	G11	a		a		G18	a		a	
		30 min		286	151	1.61	0.205		200	30.1	1.76	0.301
	Day 4	24 h	G12	a		a		G19	a		a	
		30 min		98.8	67.0	2.08	0.544		146	102	1.95	0.151
	Day 5	24 h	G13	a		a		G20	a		0.437	0.308

714
715 Concentration of VPA and 4-ene-VPA was quantified by LC/MS/MS. Data are shown as the mean with SD (n = 3 dams). a: less than the lower
716 limit of quantification (LLOQ). LLOQs of VPA and 4-ene-VPA were 10 µg/mL and 0.2 µg/mL, respectively.

722 **Supplemental Table 1** Primer sequences for real-time PCR.

723

Name	Primer sequence (5' to 3')	Product size (bp)
rBeta-actin (<i>Actb</i>)	Forward : CTATCGGCAATGAGCGGTTC Reverse : GAGGTCTTTACGGATGTCAACG	134
rBcrp (<i>Abcg2</i>)	Forward : GTTTGGACTCAAGCACAGCA Reverse : TGAGTTTCCCAGAAGCCAGT	150
rMrp1 (<i>Abcc1</i>)	Forward : CGAATGTCCTCTGAGATGGAGAC Reverse : CTCTACACGGCCTGAATGGG	138
rMrp2 (<i>Abcc2</i>)	Forward : TGATCGGTTTCGTGAAGAGCT Reverse : ACGCACATTCCCAACACAAA	139
rMrp3 (<i>Abcc3</i>)	Forward : CGTTCCGATTCACTTCTTC Reverse : TCTGGGCAAGGATTTGTGTC	124
rMrp4 (<i>Abcc4</i>)	Forward : GGACACTGAACTAGCAGAATC Reverse : TGTATTAACCTCGTCAGTTCTCG	150
rMrp5 (<i>Abcc5</i>)	Forward : CCACCATCCATGCCTATAACAA Reverse : CCCCCTGGTGGTGTATCAG	158
rMdr1a (<i>Abcb1a</i>)	Forward : GCAGGTTGGCTGGACAGATT Reverse : GGAGCGCAATTCCATGGATA	70
rMdr1b (<i>Abcb1b</i>)	Forward : CTGCTATCATCCACGGAACC Reverse : GCTGACGGTCTGTGTACTGTTG	140
rLat1 (<i>Slc7a5</i>)	Forward : CCTACGGAGGATGGAACCTATCTGA Reverse : TGGGCAAGGAGATGATGATG	93
rLat2 (<i>Slc7a8</i>)	Forward : TCCACATTTGGTGGAGTCAA Reverse : TGGATCATGGCTAACACGCT	101
rOct3 (<i>Slc22a3</i>)	Forward : TATGCAGCGGACAGATACGG Reverse : AAAATTTCGGTGCAAACGCCA	96
rOctn1 (<i>Slc22a4</i>)	Forward : TGATGTCTGTGATGCTGTGG Reverse : ATATATCTCCGAGGCAGGGTTC	172
rOctn2 (<i>Slc22a5</i>)	Forward : CGATCCCAGTGAGTTACAAGAC Reverse : GAGAAAGTCCGAAGTAGCCC	149
rOatp2a1 (<i>Slco2a1</i>)	Forward : GCCAGATACCCACAAGGAGA Reverse : GATGGCGAATAGGATGGAGA	166
rOatp2b1 (<i>Slco2b1</i>)	Forward : ACGACTTTGCCCACCATAGC Reverse : CCACGTAAAGGCGTAGCATGA	117
rOatp4a1 (<i>Slco4a1</i>)	Forward : AGAACGTCAAGTCGAGCTATTCTG Reverse : GGCCCACTTCTGTGTAAACATTT	123
rEnt1 (<i>Slc29a1</i>)	Forward : GGCCTGTGCAGTTGTCATTC Reverse : CCTCCTCTTGGCTCCTCTCC	153
rMct4 (<i>Slc16a3</i>)	Forward : GGGTCATCACTGGCTTGGGT Reverse : GGAACACGGGACTGCCTGC	123
rSry	Forward : AAGCCTTACAGAAGCCGAAAAA Reverse : TGTGGCACTTTAACCCTTCGA	120

724 **Supplemental Table 2** Ct value of rat placental samples.

Gene		Ct value	
		Single administration (G20 Control)	
		Mean	SD
ABC transporters	rBcrp	29.5	0.6
	rMdr1a	27.8	0.2
	rMdr1b	25.1	0.6
	rMrp1	27.2	0.5
	rMrp2	30.8	1.3
	rMrp3	28.7	0.4
	rMrp4	28.4	0.3
	rMrp5	27.2	0.3
SLC transporters	rLat1	24.6	0.4
	rLat2	30.6	0.6
	rOctn1	26.6	0.3
	rOctn2	27.2	0.3
	rOct3	26.2	0.3
	rOatp2a1	24.7	0.4
	rOatp2b1	29.6	0.5
	rOatp4a1	23.9	0.3
	rEnt1	29.3	0.6
	rMct4	23.1	0.2

725
 726 Data are shown as the mean with SD from placentas of G20 control rats of single administration
 727 groups (n = 9–12 placentas from three dams). ND: not determined because of low expression. G:
 728 gestational day. For measurements of each gene expression, 50-fold (rMrp2, rMrp3 G20, rMrp5, and
 729 rOatp2b1), 100-fold (rBcrp, rMdr1a, rMdr1b, rMrp1, rMrp4, rOctn1, rOctn2, rOct3, rOatp2a1,
 730 rOatp4a1, and rMct4) 200-fold (rMrp3 G13, rLat2), or 500-fold (rLat1 and rEnt1) diluted samples
 731 were amplified by real-time PCR.

732
 733

734 **Supplemental Table 3** Analysis of fetal sexual effects on gene expression (G13 rat placentas).

Gene	Single administration									Repetitive administration								
	Target gene / beta-actin mRNA (relative to male control)									Target gene / beta-actin mRNA (relative to male control)								
	Male				Female				P value	Male				Female				Interaction
	Control		VPA		Control		VPA			Control		VPA		Control		VPA		
Mean	SD	Mean	SD	Mean	SD	Mean	SD	Interaction	Mean	SD	Mean	SD	Mean	SD	Mean	SD	(P value)	
rBcrp	1.000	0.218	0.950	0.205	0.947	0.265	0.768	0.111	0.4797	1.000	0.215	0.804	0.211	0.984	0.233	0.818	0.062	0.8587
rMdr1a	1.000	0.081	1.273	0.203	1.001	0.136	1.007	0.093	0.0361	1.000	0.137	0.990	0.200	0.957	0.194	0.931	0.175	0.9103
rMdr1b	1.000	0.264	1.020	0.202	1.064	0.368	1.079	0.345	0.9871	1.000	0.199	0.624	0.112	0.981	0.170	0.907	0.203	0.0549
rMrp1	1.000	0.124	1.118	0.143	0.940	0.115	1.104	0.031	0.6424	1.000	0.189	1.555	0.330	1.059	0.080	1.361	0.259	0.2172
rMrp2	ND		ND		ND		ND			ND		ND		ND		ND		
rMrp3	1.000	0.112	1.252	0.319	1.068	0.220	1.042	0.150	0.1536	1.000	0.129	0.842	0.073	1.001	0.099	0.797	0.054	0.4180
rMrp4	1.000	0.103	1.024	0.210	0.925	0.128	0.993	0.110	0.7236	1.000	0.145	1.118	0.218	0.872	0.447	1.076	0.117	0.5272
rMrp5	1.000	0.207	1.274	0.249	1.034	0.313	1.215	0.170	0.6605	1.000	0.311	1.751	0.670	0.913	0.104	1.433	0.590	0.5697
rLat1	1.000	0.090	0.919	0.123	1.070	0.146	0.966	0.192	0.8769	1.000	0.092	0.893	0.186	1.221	0.336	1.150	0.425	0.8733
rLat2	1.000	0.240	1.221	0.349	0.927	0.268	1.405	0.294	0.3096	1.000	0.275	2.198	0.920	1.047	0.301	1.826	0.924	0.5018
rOcn1	1.000	0.143	1.268	0.246	1.238	0.090	1.660	0.489	0.5035	1.000	0.235	1.175	0.365	0.926	0.078	1.035	0.297	0.7831
rOcn2	1.000	0.309	1.093	0.471	0.951	0.342	1.204	0.185	0.5959	1.000	0.339	1.794	0.753	0.979	0.326	1.726	0.241	0.9065
rOct3	1.000	0.155	1.382	0.281	1.079	0.209	1.106	0.292	0.0935	1.000	0.100	0.855	0.352	0.985	0.161	1.141	0.216	0.1289
rOatp2a1	1.000	0.215	1.856	0.743	1.039	0.197	1.721	0.287	0.6349	1.000	0.218	2.534	1.525	0.931	0.115	1.893	0.783	0.4455
rOatp2b1	1.000	0.151	1.200	0.262	1.047	0.267	1.273	0.406	0.9088	1.000	0.157	1.290	0.274	1.010	0.118	1.071	0.582	0.4055
rOatp4a1	1.000	0.104	1.091	0.162	1.096	0.143	0.988	0.227	0.1613	1.000	0.132	1.180	0.222	0.976	0.112	1.226	0.341	0.7034
rEnt1	1.000	0.221	1.204	0.478	1.031	0.448	1.358	0.168	0.7005	1.000	0.278	1.664	0.612	0.965	0.228	1.490	0.439	0.6974
rMct4	1.000	0.108	1.237	0.173	1.067	0.166	1.095	0.133	0.1174	1.000	0.035	0.862	0.155	0.990	0.051	1.003	0.103	0.0810

735

736 Data are shown as the mean with SD (n = 5–7 placentas from three dams). ND: not determined because of low expression. The data shown in
737 Figures 1 and 2 of G13 rat placentas were analyzed by fetal sex. Gene expression was analyzed using a two-way ANOVA, with “sex” and “VPA”
738 as factors. If *p* value was < 0.05, an interaction was considered to be present.

739 **Supplemental Table 4** Analysis of fetal sexual effects on gene expression (G20 rat placentas).


Gene	Single administration									Repetitive administration								
	Target gene / beta-actin mRNA (relative to male control)									Target gene / beta-actin mRNA (relative to male control)								
	Male				Female				Interaction (<i>P</i> value)	Male				Female				Interaction (<i>P</i> value)
	Control		VPA		Control		VPA			Control		VPA		Control		VPA		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD	Mean	SD	
rBcrp	1.000	0.144	1.180	0.601	0.906	0.294	0.844	0.086	0.4835	1.000	0.290	1.386	0.451	0.924	0.187	0.756	0.117	0.0701
rMdr1a	1.000	0.091	0.985	0.148	0.885	0.070	0.949	0.177	0.5206	1.000	0.282	1.319	0.180	0.957	0.271	1.247	0.135	0.9003
rMdr1b	1.000	0.169	0.950	0.257	1.248	0.486	1.059	0.158	0.6200	1.000	0.121	0.832	0.134	1.274	0.203	1.026	0.022	0.5633
rMrp1	1.000	0.160	0.942	0.190	1.044	0.112	0.957	0.155	0.8428	1.000	0.190	0.764	0.171	0.893	0.133	0.896	0.003	0.1023
rMrp2	1.000	0.399	0.761	0.568	0.794	0.612	0.528	0.333	0.9536	1.000	0.933	1.254	0.923	0.693	0.241	0.455	0.186	0.4522
rMrp3	1.000	0.161	1.218	0.402	1.225	0.393	1.405	0.219	0.8993	1.000	0.158	0.894	0.261	1.017	0.206	0.951	0.231	0.8409
rMrp4	1.000	0.095	0.742	0.180	1.203	0.330	1.030	0.222	0.6864	1.000	0.186	0.570	0.222	1.118	0.226	0.601	0.113	0.6502
rMrp5	1.000	0.088	0.914	0.162	0.966	0.169	0.873	0.106	0.9622	1.000	0.219	1.181	0.266	1.022	0.231	0.971	0.135	0.2840
rLat1	1.000	0.107	0.801	0.044	1.046	0.157	0.812	0.105	0.7224	1.000	0.164	0.745	0.103	0.851	0.090	0.770	0.083	0.1100
rLat2	1.000	0.137	0.859	0.104	0.967	0.171	0.858	0.229	0.8340	1.000	0.116	0.831	0.213	0.845	0.100	0.943	0.132	0.0702
rOctn1	1.000	0.111	0.771	0.116	0.927	0.143	0.761	0.185	0.6353	1.000	0.131	0.479	0.074	1.086	0.155	0.601	0.041	0.7378
rOctn2	1.000	0.107	0.996	0.145	0.871	0.205	0.854	0.194	0.9376	1.000	0.083	0.911	0.200	1.118	0.156	1.277	0.081	0.0916
rOct3	1.000	0.083	1.032	0.166	1.052	0.134	1.189	0.099	0.3865	1.000	0.398	1.448	0.256	1.282	0.228	1.621	0.577	0.7230
rOatp2a1	1.000	0.183	0.861	0.159	1.074	0.264	0.779	0.408	0.5357	1.000	0.101	0.541	0.070	1.061	0.149	0.862	0.256	0.0494
rOatp2b1	1.000	0.118	0.819	0.151	1.003	0.156	0.755	0.129	0.6052	1.000	0.184	1.262	0.156	1.054	0.208	0.928	0.049	0.0240
rOatp4a1	1.000	0.181	0.812	0.099	0.968	0.151	0.792	0.180	0.9302	1.000	0.240	0.772	0.109	0.973	0.078	0.883	0.049	0.2805
rEnt1	1.000	0.324	0.879	0.172	1.191	0.091	1.063	0.136	0.9637	1.000	0.113	0.657	0.127	0.828	0.248	0.684	0.184	0.2387
rMct4	1.000	0.064	0.868	0.167	0.887	0.181	0.926	0.481	0.5004	1.000	0.204	1.146	0.183	1.123	0.181	1.178	0.093	0.5819

740


741 Data are shown as the mean with SD (n = 3–7 placentas from one to three dams). The data shown in Figures 1 and 2 of G20 rat placentas were
742 analyzed by fetal sex. Gene expression was analyzed using a two-way ANOVA, with “sex” and “VPA” as factors. If *p* value was < 0.05, an
743 interaction was considered to be present.

Figure 1


(A) rMdr1a



(B) rMdr1b



(C) rMrp1



(D) rMrp2






Figure 1


(E) rMrp3



(F) rMrp4



(G) rMrp5



(H) rBcrp




Figure 2






Figure 2


(E) rOcn1



(F) rOcn2



(G) rEnt1



(H) rOatp2a1





Figure 2

(I) rOatp2b1



(J) rOatp4a1






Figure 3

A




B




Supplemental Figure 1


A rMdr1a
(G13, single)
Sex: $P = 0.0369$
VPA: $P = 0.0299$
Interaction: $P = 0.0361$



B rOatp2a1
(G20, repetitive)
Sex: $P = 0.0062$
VPA: $P < 0.0001$
Interaction: $P = 0.0494$




C rOatp2b1
(G20, repetitive)
Sex: $P = 0.0923$
VPA: $P = 0.3992$
Interaction: $P = 0.0240$



Supplemental Figure 2

A



B

