



Title	The Merger of Photoredox and Cobalt Catalysis
Author(s)	Kojima, Masahiro; Matsunaga, Shigeki
Citation	Trends in Chemistry, 2(5), 410-426 https://doi.org/10.1016/j.trechm.2020.01.004
Issue Date	2020-05
Doc URL	https://hdl.handle.net/2115/81686
Type	journal article
File Information	WoS_94704_Kojima.pdf



The Merger of Photoredox and Cobalt Catalysis

Masahiro Kojima* and Shigeki Matsunaga*

Faculty of Pharmaceutical Sciences, Hokkaido University, Kita-12 Nishi-6, Kita-ku, Sapporo, 060-0812, Japan

*Correspondence: m-kojima@pharm.hokudai.ac.jp (M.Kojima);
smatsuna@pharm.hokudai.ac.jp (S. Matsunaga).

Abstract

In the last decade, synthetic chemists have discovered the outstanding generality and potential of visible-light-driven photoredox catalysis, which converts visible light into chemical energy and realizes numerous transformations of small molecules. The current state-of-the-art strategy in photoredox catalysis, combining photoredox and transition-metal catalysis, has received considerable attention in organometallic chemistry. In parallel to the rapid development of nickel/photoredox and copper/photoredox catalysis, cobalt/photoredox catalysis has emerged as a distinct new option in this area. This short review covers the general strategy, characteristics compared to other metallaphotoredox systems, and future perspectives of cobalt/photoredox catalysis. To the best of our knowledge, this is the first review encompassing the general combination of cobalt catalysis and photoredox catalysis in synthetic chemistry.

Keywords: photoredox catalysis; cobalt; dehydrogenation; dual catalysis; metallaphotoredox catalysis

Merger of 3d-metal catalysis and photoredox catalysis

Photochemistry historically represents a vital part of synthetic chemistry. Although direct excitation of organic compounds by UV light has long been mainstream in synthetic photochemistry [1], visible-light-driven photoreactions have recently started to flourish as a safer and more user-friendly alternative. This renaissance in photochemistry relies on the emergence of photoredox catalysts [2-7]. In general, a photoredox catalyst absorbs visible light and reaches its **excited state** (see Glossary) with a lifetime that is typically > 100 ns. The photocatalyst in its excited state serves as a one-electron oxidant or reductant that enables the transfer of one electron between the catalyst and the organic substrates. Thus, photoredox catalysts are suitable for generating radicals from stable organic molecules and have contributed to recent development of radical-mediated transformations [8-10]. Following its success in radical-mediated transformations, the application of a photoredox phenomenon in transition-metal-catalyzed reactions, the so-called metallaphotoredox catalysis [11-15], has attracted the interest of researchers in both photochemistry and organometallic chemistry.

A pioneering contribution by Sanford and colleagues employed an organopalladium [16] complex generated *in situ*, that allows the formation of C–C bonds *via* organometallic complexes that reside in unusually high **oxidation states**. This pioneering work has quickly been extended by a study into other metals as potential partners in metallaphotoredox chemistry, and this merger has been established for 3d, 4d, and 5d metal catalysts. Among these, 3d metals such as cobalt, nickel, and copper are particularly appealing because they are more earth abundant than to 4d or 5d metals. Moreover, several mechanistic studies have indicated the involvement of one-electron redox

processes in the thermal catalytic cycle [17-19], as well as the feasibility of controlling their **redox potential** by a judicious choice of ancillary ligands. These aspects promise potential for their compatibility with photoredox catalysis, which operates under mild conditions, provides facile access to radical intermediates, and is characterized by a rich choice of redox potentials for one-electron transfers (Figure 1A).

Indeed, nickel/photoredox [20-22] and copper/photoredox [23, 24] cooperative systems have evolved rapidly, showcasing their potential in challenging C–C- [25-29] or C–heteroatom- [30-35] bond-forming reactions as well as highly enantioselective transformations (Figure 1B, right) [36-38]. Since the primary discoveries on metallaphotoredox catalysis was analogous to cross-coupling-type reactions previously catalyzed by palladium (Suzuki–Miyaura, Buchwald–Hartwig, etc.), studies on nickel or copper (which are established alternatives for palladium catalysts in thermal cross-coupling reactions) have dominated the area, while the potential of other first row transition metals including cobalt was largely unexplored. On the other hand, the merger of photoredox catalysis with cobalt catalysis has a long history in inorganic chemistry (Figure 1B, left). The reduction of protons into molecular hydrogen is an indispensable transformation in **artificial photosynthesis** and, in that context, the combination of visible-light-driven photocatalysts and cobalt complexes has been intensively studied [39]. In addition, early contributions on **biomimetic chemistry** regarding vitamin B12 and its model compounds under photochemical conditions [40, 41] should also be regarded as pioneering work in cobalt/photoredox cooperative catalysis. Despite these precedents and recent developments in cobalt/photoredox cooperative catalysis, to the best of our knowledge there is still no review that focuses on this topic from the

perspective of synthetic chemistry. This review aims to cover recent examples (typically those reported between 2014 and 2019) on the merger of cobalt and photoredox catalysis, focusing on (1) dehydrogenative transformations (cross-coupling hydrogen evolution and acceptorless dehydrogenation), and (2) C–C- or C–heteroatom-bond-forming reactions in which organocobalt intermediates presumably play a vital role (e.g., hydrofunctionalization of alkenes, cycloaddition of alkynes, allylation/deallylation, and reactions using stoichiometric oxidant or reductant). Despite the diversity of these precedents, they all share the general mechanistic strategies depicted in Figure 1C: The reduction of the cobalt catalyst by the reduced photocatalyst is responsible for low-valent cobalt or cobalt-hydride-mediated transformations. On the other hand, one-electron oxidation of a cobalt complex would provide a high-valent cobalt complex that could subsequently mediate an oxidative transformation.

Cross-coupling hydrogen evolution (CCHE)

The oxidative functionalization of C–H bonds to C–C or C–heteroatom bonds represents a desirable method for streamlining multistep syntheses of complex molecules, as it avoids the pre-functionalization steps that are usually associated with the preparation of reaction partners in classical cross-coupling reactions [42, 43]. Early examples of such transformations are based on the use of equimolar amounts of chemical oxidants, which is not ideal with respect to sustainability, safety, and functional-group tolerance. On the other hand, oxidative couplings that occur under the concomitant evolution of hydrogen gas (cross-coupling hydrogen evolution; CCHE) have emerged as a more advanced technology in this context [44-48]. While the proof-of-concept study on CCHE employed a heterogeneous graphene-supported ruthenium oxide nanocomposite for the hydrogen

evolution [49], it was quickly demonstrated that cobaloxime derivatives are able to play the same role under homogeneous conditions (*vide infra*).

An early example of the intermolecular formation of C–C bonds by CCHE was reported by Wu and colleagues who introduced a combination of the organophotocatalyst eosin Y and Co(dmgh)₂Cl₂ for the noble-metal-free CCHE between tetrahydroisoquinolines and indoles (Figure 2A) [50]. In their proposed catalytic cycle, photoexcited eosin Y is responsible for the one-electron oxidation of amines, while the second oxidation of the α -aminoalkyl radical intermediate is mediated by the cobaloxime cocatalyst. This successive two-electron oxidation leads to the formation of an iminium intermediate, which leads to the formation of a new C–C bond upon reaction with a nucleophile. Finally, molecular hydrogen is generated by the two-electron reduction of cobaloxime followed by protonolysis. Through the similar photochemical oxidation of substrates, dehydrogenative functionalization of amino acids [51] and isochromanones [52] were also possible. Furthermore, Luo and Wu and colleagues reported an asymmetric variant of the dehydrogenative coupling assisted by a chiral amine catalyst [53].

Cobaloxime/photoredox cooperative catalysis has been proven to be generally applicable to synthetically valuable C–C bond-forming reactions including synthesis of heterocycles [54], oxidative Heck [55, 56], and Minisci-type [57, 58] reactions. Notably, the catalytic dehydrogenative formation of C–C bonds was also extended to provide access to more complex molecular scaffolds: Lei and colleagues reported the dehydrogenative coupling between styrene derivatives and alkynes, which affords multisubstituted naphthalene rings in a single step (Figure 2B) [59]. The high oxidation potential of an excited Acr⁺-

Mes organophotocatalyst enables the oxidation of styrene derivatives. Then, the addition of alkynes to the resulting radical cation is responsible for the formation of the first C–C bond. The second C–C bond formation occurs by an intramolecular Friedel–Crafts-type reaction, followed by an aromatization through the release of two protons and an electron. This aromatization is facilitated by the simultaneous formation of cobalt hydride, whose protonation results in the release of hydrogen gas and the regeneration of the cobaloxime catalyst. Consecutive formations of C–C and C–N bonds were also demonstrated in the similar reaction design for the preparation of 3,4-dihydroisoquinoline derivatives [60] or isoquinoline derivatives [61].

Since CCHE realizes oxidative transformations in the absence of chemical oxidants, its advantage is evident for the construction of C–heteroatom bonds, for which the starting materials or products are susceptible to undesirable oxidation. Lei and colleagues have demonstrated the intramolecular C–H thiolation of aromatic thioamides using a combined cobaloxime/photoredox catalytic system (Figure 2C) [62]. The products were obtained in lower yield when chemical oxidants were used, presumably because thioamides readily decompose by these reagents. From a mechanistic perspective, the one-electron oxidation of a thioamide by the excited $\text{Ru}(\text{bpy})_3^{2+}$ photoredox catalyst provides access to a thiyl radical, which undergoes addition to an aromatic ring. The oxidation of the resulting radical intermediate by the reduced cobaloxime affords the benzothiazole product, while hydrogen gas is released by protonolysis of the cobalt hydride. The authors proposed that precise tuning of the proton transfer steps by basic additives contributes to high catalytic turnovers.

Wu and Tung and colleagues reported another impressive example of CCHE, i.e., the C–H amination and hydroxylation of benzene (Figure 2D) [63]. The UV-light-driven, highly oxidizing photocatalyst QuCN^+ oxidizes benzene to afford a benzene radical cation. The addition of a heteroatom nucleophile to the intermediate and release of a proton delivers a cyclohexadienyl radical intermediate, which is presumably oxidized by the reduced cobaloxime to afford the products with a new C–N or C–O bond. Protonation of the resulting cobalt hydride regenerates the cobaloxime cocatalyst under concomitant release of hydrogen gas. It is proposed that overoxidation of the products is inhibited by a rapid electron back transfer between the product and the photocatalyst. Other challenging dehydrogenative bond-forming reactions (e.g., C–N [64-66]; C–O [64, 67-69]; C–P [70-72]; C–S [62, 73]; and C–Si [74]) by CCHE have also been reported, providing further evidence for the versatility of this strategy to construct C–heteroatom bonds.

Acceptorless dehydrogenation

Desaturation of a C–C or C–heteroatom single bond to a double bond is a fundamental process for the preparation of fine chemicals and other industrial applications. Among these transformations, acceptorless dehydrogenation is considered to be ideal because chemical oxidants are not required, and useful hydrogen gas is the sole byproduct [48]. However, catalytic acceptorless dehydrogenation under thermal conditions generally requires high reaction temperatures because the release of hydrogen gas together with the desaturation is thermodynamically unfavorable in terms of enthalpy.

In order to realize the acceptorless dehydrogenation of alkanes under milder conditions, Sorensen and colleagues have employed a cooperative base-metal catalytic system that consists of a tetra-*n*-butylammonium decatungstate photocatalyst and cobaloxime (Figure

3A) [75]. Notably, the acceptorless dehydrogenation of unactivated alkanes proceeds at room temperature under irradiation with UV light. The authors proposed that the excited decatungstate photocatalyst abstracts a hydrogen atom from the alkane, which affords a radical intermediate and the partially hydrogenated photocatalyst. Then, a second hydrogen-atom transfers from the radical intermediate to the cobaloxime affords an alkene and cobalt hydride. The release of hydrogen gas by the reaction between the reduced photocatalyst and the cobalt hydride regenerates the two base-metal catalysts. The same cooperative base-metal catalysts are also applicable to the dehydroformylation of aliphatic aldehydes (Figure 3B) [76]. In this case, the photochemical cleavage of the C–H bond is followed by a decarbonylation, which leads to an sp^3 -hybridized carbon-centered radical. This radical intermediate engages with the cobaloxime catalyst to afford an alkene and molecular hydrogen.

Li and colleagues have demonstrated that the merger of a photoredox catalyst and a cobaloxime catalyst is a viable strategy for the acceptorless dehydrogenation of partially saturated *N*-heterocycles (Figure 3C) [77]. The catalytic dehydrogenation provides access to valuable *N*-heteroarenes such as quinolines, quinoxalines, indoles, and thiazoles, and thus represents broad substrate generality due to exceptionally mild reaction conditions (under irradiation with visible light at room temperature). The photoredox catalyst $Ru(bpy)_3^{2+}$ facilitates the one-electron oxidation of saturated amines, which allows the generation of an α -aminoalkyl radical intermediate. Subsequent electron transfer from the radical intermediate to the photochemically reduced cobaloxime affords cobalt hydride. The ensuing protonolysis of cobalt hydride results in the formation of molecular hydrogen and the regeneration of the cobaloxime catalyst. This chemistry was later proven to be

compatible with aqueous reaction media [78]. It should also be noted here that similar cooperative catalysis can facilitate the dehydrogenative coupling of two *N*-heterocycles by finely tuning the reaction conditions [79]. In addition to C–N bond, N–N bond was able to be desaturated to afford azobenzene derivatives by cobaloxime/photoredox system [80].

An interesting extension of the catalytic acceptorless dehydrogenation is the decarboxylative desaturation of carboxylic acids. According to the report by Ritter and colleagues, the combination of the photocatalyst Ir[dF(CF₃)ppy]₂(dtbpy)PF₆ and a cobaloxime cocatalyst provided a general system that converts aliphatic carboxylic acids into alkenes (Figure 3D) [81]. In the proposed reaction mechanism, single-electron oxidation of a carboxylate by the photoredox catalyst affords an alkyl radical via decarboxylation. Engagement of the radical with the reduced cobaloxime mediates the formation of an alkene, which is followed by the release of hydrogen gas from cobalt hydride. The net transformation can be accomplished under stoichiometric additive-free conditions, which is appealing for the conversion of abundant fatty acids into industrially valuable α -olefins. Tunge and Cartwright have reported a relevant dehydrogenation of amino acids into enamides using a combination of cobaloxime and an organophotocatalyst, thus demonstrating the generality of such combinations for acceptorless dehydrogenations [82]. Larionov and colleagues have reported a catalytic decarboxylative dehydrogenation under enzyme-compatible conditions, which allowed the direct transformation of esters into alkenes [83].

Hydrofunctionalization of alkenes

Considering that cobalt hydride species derived from cobaloximes can be readily generated and are compatible with a series of reaction conditions (*vide supra*), it is reasonable to assume that cobalt-hydride-mediated transformations should be possible by using cobalt/photoredox catalysis. Considering that cobalt hydride is responsible for the Drago–Mukaiyama hydration and related hydrofunctionalizations of alkenes [84], the cobalt/photoredox approach found applications in the hydrofunctionalization of alkenes using photochemically generated cobalt hydride species.

Rovis and Thullen have reported the hydroaminoalkylation of conjugated dienes using a dual catalyst system consisting of a cobalt diphosphine complex and the photocatalyst $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbpy})\text{PF}_6$ (Figure 4A) [85]. From a mechanistic perspective, the one-electron reduction of the Co^{II} complex generates Co^{I} *in situ*, which is subsequently protonated to afford the cobalt hydride. Its insertion into conjugated dienes generates π -allylcobalt species. On the other hand, amines are oxidized by the excited photocatalyst to furnish nucleophilic α -aminoalkyl radicals. The photoredox-mediated reduction of the organocobalt complex, followed by an addition of the α -aminoalkyl radical affords a new C–C bond and regenerates the low-valent cobalt complex. Conjugated dienes with electron-withdrawing groups generally furnish the hydroalkylated products in good yield, and the authors unambiguously confirmed that cobalt catalysis is indispensable to achieve high product yields. This system is also applicable to the transformation of aliphatic amines or simple conjugated dienes without functional groups. The transformation also allows the construction of quaternary carbon atoms, which might be facilitated by sterically less hindered nature of the α -aminoalkyl radicals.

Another application for photochemically-generated cobalt hydrides has been disclosed by König and colleagues, who reported the isomerization of terminal alkenes into internal alkenes using a combination of cobalt acetylacetonate and 4CzIPN (Figure 4B) [86]. The excited organophotocatalyst was quenched using a Hantzsch ester, and the electron transfer from the reduced photocatalyst generates a Co^{I} complex. A subsequent protonation affords a cobalt hydride, which serves as a catalyst for the isomerization of alkenes [87]. In the proposed mechanism, the insertion of cobalt hydride into terminal alkenes affords an alkylcobalt intermediate. A subsequent β -hydride elimination, which might be accelerated by visible light, affords thermodynamically more favorable internal alkenes and regenerates the cobalt hydride. The combined catalytic system was applied to the isomerization of hydrocarbons and natural-product derivatives, affording the internal alkenes in high yield. It is worthwhile noting that sequential (or “chain-walking”) isomerization is also possible with this system.

The cobalt-catalyzed hydrofunctionalization was successfully applied to C–C bond forming reaction by Maji and coworkers who reported reductive hydroalkenylation of electron deficient alkenes (Figure 4C) [88]. In their mechanistic hypothesis, cobalt hydride which was obtained by protonation of Co^{I} inserts to electron deficient alkene. Insertion of an alkyne is followed by the release of the hydroalkenylated product and reductive regeneration of the Co^{I} catalyst.

Collins and colleagues proposed distinct access to cobalt hydride and application to hydroalkynylation of alkynes [89]. This report represents prospects of the cobalt/photoredox system for hydrofunctionalization of alkynes.

Cycloaddition of alkynes

The cobalt-catalyzed cycloaddition of alkynes has been studied mainly relying on the well-defined low-valent cobalt catalysts [90-92]. Inspired by these precedents, the application of photochemically generated low-valent cobalt complexes for the cycloaddition of alkynes has attracted attention in the merger of cobalt and photoredox catalysis. A pioneering contribution has been made by Rovis and Ruhl, who have employed a combination of $\text{CoBr}_2(\text{PCy}_3)_3$ and $\text{Ir}[\text{dF}(\text{CF}_3)\text{ppy}]_2(\text{dtbpy})\text{PF}_6$ for the [2+2+2] cyclotrimerization of diynes and alkynes (Figure 5A) [93]. The catalytic system contains catalytic amounts of an organic amine, which most likely serves as a reductant for the initial generation of a low-valent cobalt catalyst. In the proposed catalytic cycle, a low-valent cobalt(0) complex, generated by a photoredox-mediated reaction, enables the oxidative cyclization of alkynes. The scope of this cyclotrimerization includes aliphatic alkynes, diynes with different tethers, and internal alkynes. The authors insightfully applied the catalytic system to the photolithographic modification of a polymer surface, given that the light-driven reaction provides spatial control over the reaction. This “light-gated” control might be due to sophisticated electron-transfer steps between the organocobalt intermediate and the photocatalyst. This light-gated strategy was also effective in controlling the molecular weight in polymer syntheses, meaning that temporal control of the reaction is possible [94]. Interestingly, a relevant [2+2+2] cycloaddition

was later achieved in the absence of a photoredox catalyst [95]. In that case, the organocobalt intermediate absorbs visible light and is involved in electron-transfer steps.

Zhao and Wu and colleagues have demonstrated another application of photoredox-induced low-valent cobalt catalysis, in which the cycloaddition of alkynes and carbon dioxide affords oxygen-containing heterocycles (Figure 5B) [96]. Although poorly functional-group-tolerant organometallic reductants are often required to incorporate carbon dioxide into organic molecules, an organic amine can be used here as a mild stoichiometric reductant. The amine is oxidized by the excited photocatalyst Ir(ppy)₂(dtbpy)PF₆, and an electron transfer from the reduced photocatalyst to a cobalt diphosphine complex affords a low-valent Co^I complex. This cobalt complex mediates the oxidative cycloaddition of alkynes and carbon dioxide. A subsequent migration of the acetyl group and a transmetallation with zinc bromide releases a tricyclic intermediate that eventually cyclizes to afford γ -hydroxybutenolides, and the low-valent cobalt complex is regenerated by a photoredox-mediated reduction. The scope of the reaction includes multisubstituted five-membered heterocycles with cyclopropyl groups, electron-rich or -deficient arenes, and sterically hindered acyl groups. The same catalytic system mediates the cycloaddition with other substituted alkynes to afford coumarin derivatives and substituted cinnamic acid derivatives. These results indicate that photochemically induced low-valent cobalt catalysis exhibits reasonable functional-group tolerance and can be considered as a useful option in synthetic chemistry.

Allylation and deallylation

Transition-metal-catalyzed allylic substitution reactions constitute an intensively investigated area in synthetic organometallic chemistry. Among all conceivable metal catalysts, noble group 9 metals such as rhodium [97] and iridium [98-100] represent a privileged class of catalysts due to their intrinsic tendency to afford synthetically versatile branched products. On the other hand, cobalt-mediated allylic substitution reactions have received less attention, despite the abundance of cobalt salts in the earth's crust and the intrinsic similarity between cobalt, rhodium, and iridium. Active catalysts for allylic substitution reactions are often low-valent group 9 metals such as Rh^{I} or Ir^{I} , and therefore, their low-valent congener Co^{I} should serve as a potent catalyst for allylation reactions, as suggested by recent thermal reactions [101, 102].

Gryko and coworkers have reported an insightful combination of a titanium oxide photocatalyst and vitamin B12 for the deallylation of phenol derivatives (Figure 6A) [103]. The authors proposed that a low-valent cobalt complex, generated *in situ* by electron transfer from the titanium photocatalyst to vitamin B12, serves as the active species in the catalytic cycle. The resulting low-valent cobalamine is highly nucleophilic, and its addition to the substrate results in the formation of an allylcobalt intermediate and the release of the deallylated arenes. A subsequent protonation and reduction of the allylcobalamine regenerates the low-valent cobalamine catalyst. A subsequent treatment with hydrochloric acid is desirable as enol ethers are the major byproducts in this reaction, which might be derived from the isomerization of the double bond of the allyl group. The deallylation proceeds in high yield for sterically hindered 2,4,6-methylphenol or phenols with electron-withdrawing groups and a deallylation to afford an aliphatic alcohol is also possible. Unfortunately, the transfer of the allyl group to other nucleophiles was not

accomplished in this catalytic system, possibly due to the limited electrophilicity of the allylcobalamine intermediate.

More recently, Kojima and Matsunaga and colleagues have reported the organophotoredox-promoted regioselective allylation of active methylene nucleophiles (Figure 6B) [104]. The authors employed a Co^{II} diphosphine complex as the catalyst precursor and generated a low-valent cobalt complex *in situ* by electron transfer from the 4CzIPN organophotocatalyst in the presence of an organic amine. The oxidative addition of allylic electrophiles to the generated Co^{I} complex affords a π -allylcobalt intermediate. A nucleophilic addition of a soft carbon nucleophile to the electrophilic π -allylcobalt species induces the formation of a C–C bond and regenerates the Co^{I} catalyst. The reaction afforded branched allylic substitution products in high regioselectivity, suggesting a similarity between the cobalt-catalyzed method and rhodium- or iridium-catalyzed allylation reactions. In addition, the catalytic allylation was successfully expanded to the regioselective substitution of an allyl sulfone. Considering that substitution of allyl sulfones is challenging for rhodium- or iridium-based catalysts, this report is the first example of the superiority of cobalt catalysts compared to established Rh- and Ir-based catalytic systems.

Reactions using stoichiometric oxidant or reductant

Metallaphotoredox catalysis enables the use of environmentally benign oxidant instead of stoichiometric transition metals such as copper or silver salts. Rueping and coworkers elegantly demonstrated the photoredox-enabled aerobic oxidative C–H functionalization with rhodium catalysis [105]. Inspired by the precedent, Sundararaju and colleagues

reported a C–H activation/oxidative cycloaddition of alkynes using a merger of Co^{II} catalysis and eosin Y under aerobic conditions (Figure 7A) [106]. It is proposed that after the irreversible, turnover limiting C–H bond cleavage, successive coordination of an alkyne, migratory insertion, and reductive elimination affords the product along with the release of Co^{I} catalyst. Finally, the photoredox-mediated electron transfer regenerates Co^{III} using molecular oxygen as an oxidant. Terminal alkenes are also able to involve in the relevant oxidative C–H alkylation while the turnover limiting step in this reaction was not the C–H bond cleavage [107]. These combinations represent rare examples of cobalt/photoredox catalysis in the presence of an oxidant, and serve as a starting point for the combination of high-valent cobalt-catalyzed C–H bond functionalizations with photoredox catalysis [108].

Cobalt/photoredox dual system allows the use of milder and more tractable reagents for net reductive transformations. Lloret-Fillol and coworkers reported reduction of ketones and aldehydes catalyzed by dual cobalt catalysis and photocatalysis using triethylamine or *N,N*-diisopropylethylamine as a reductant (Scheme 7B) [109]. Interestingly, photochemically generated Co^{II} hydride reduces the carbonyl compounds either via stepwise single electron transfer/hydrogen atom transfer mechanism or via hydride transfer mechanism. The excited copper-based photoredox catalyst is reductively quenched by amines and transfers an electron to the polydentate cobalt catalyst.

Casitas and Lloret-Fillol and colleagues applied virtually the same combination of cobalt and photoredox catalysis for reduction of unactivated alkyl chlorides (Figure 7C) [110]. This transformation clearly demonstrates the advantage of metallaphotoredox chemistry

because unactivated alkyl chlorides are known to be inert toward conventional photoredox catalysis. In this case a respective polydentate nickel complex was studied in detail, and the mechanism of cobalt/photoredox catalysis for dechlorination is proposed in Figure 7C assuming that the two metal catalysts should work in similar mechanisms. Oxidative addition of an alkyl chloride to low valent metal complex affords alkyl metal intermediate. On the other hand, photoredox catalyst is reduced by amines assisted by visible light. Single electron transfer from the reduced photoredox catalyst to the alkyl metal intermediate prompts homolysis of the metal-carbon bond, and an alkyl radical is formed and the low valent metal complex is regenerated. The alkyl radical undergoes intramolecular addition to the tethered alkene and the cyclized product is obtained.

Concluding remarks

Although the merger of cobalt catalysis and photoredox catalysis has previously been a focus in inorganic chemistry, it is now clear that cobalt/photoredox cooperative catalysis also constitutes an innovative technology in synthetic chemistry. While nickel/photoredox or copper/photoredox system are useful for cross-coupling-type reactions which were previously catalyzed by palladium, cobalt/photoredox catalysis provides new options to dehydrogenative transformations and reactions which were efficiently catalyzed by noble group 9 metals such as branch-selective allylic alkylation. Since cobalt can exist in a variety of oxidation states (e.g., 0, I, II, and III) and engage both in one-electron transfers and two-electron mediated bond forming processes, its combination with photoredox catalysis validates numerous reaction pathways that were not accessible in thermal transformations. Rational control of the reaction pathways is thus essential for productive transformations, and more mechanistic studies as well as the variety of established ligands

and photoredox catalysts would help us establish practical reaction conditions. In addition, employment of other photocatalysts [111-113] or combination with hydrogen atom transfer catalyst [114-117] are promising strategies for functionalization of less reactive small molecules, and should expand the scope of cobalt/photoredox catalysis. Since several more tasks (enantioselective reactions, comparison with thermal reactions, merger with Co-catalyzed C-H functionalizations [118-120] and mechanistic understanding; see Outstanding Questions for details) remain to be addressed, cobalt/photoredox dual catalysis continues to deserve intense attention in synthetic community as a worthwhile challenge in metallaphotoredox chemistry.

Acknowledgements

This work was supported in part by JSPS KAKENHI grant JP15H05802 within the remit of Precisely Designed Catalysts with Customized Scaffolding, as well as JSPS KAKENHI grants JP17H03049 and JP18H06097.

Glossary

Artificial photosynthesis: an artificial photochemical process which enables synthesis of an energy carrying molecule (for example, molecular hydrogen) from an abundant molecule (for example, water) toward green production of chemical fuel.

Biomimetic chemistry: an area of chemistry which aims to mimic highly efficient chemical phenomena in biological system by rational design and application of artificial molecules or chemical methods

Excited state: a higher energy state of a molecule or a system in quantum chemistry compared to its original, thermally stable state (or “ground state”), typically accessed by an absorption of a photon

Oxidation state: a formal degree of oxidation of a specific atom in a molecule calculated by its formal charge and covalent bonds, which is informative for simplified interpretation of electrophilicity and potential reactivity of the atom

Redox potential: a measure in electrochemistry which quantitatively describes the voltage required for electron transfer from the specific molecule to an electrode: comparison of the values between different chemical species helps predict the feasibility of an electron transfer between the two

Disclaimer Statement

The authors declare no competing financial interest.

1. Kärkäs, M.D. *et al.* (2016) Photochemical Approaches to Complex Chemotypes: Applications in Natural Product Synthesis. *Chem. Rev.* 116, 9683-9747.
2. Prier, C.K. *et al.* (2013) Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* 113, 5322-5363.
3. Schultz, D.M. and Yoon, T.P. (2014) Solar Synthesis: Prospects in Visible Light Photocatalysis. *Science* 343, 1239176.
4. Ravelli, D. *et al.* (2016) Carbon–Carbon Bond Forming Reactions via Photogenerated Intermediates. *Chem. Rev.* 116, 9850-9913.
5. Romero, N.A. and Nicewicz, D.A. (2016) Organic Photoredox Catalysis. *Chem. Rev.* 116, 10075-10166.
6. Marzo, L. *et al.* (2018) Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis? *Angew. Chem. Int. Ed.* 57, 10034-10072.
7. McAtee, R.C. *et al.* (2019) Illuminating Photoredox Catalysis. *Trends Chem.* 1, 111-125.
8. Studer, A. and Curran, D.P. (2016) Catalysis of Radical Reactions: A Radical Chemistry Perspective. *Angew. Chem. Int. Ed.* 55, 58-102.
9. Yan, M. *et al.* (2016) Radicals: Reactive Intermediates with Translational Potential. *J. Am. Chem. Soc.* 138, 12692-12714.
10. Yi, H. *et al.* (2017) Recent Advances in Radical C–H Activation/Radical Cross-Coupling. *Chem. Rev.* 117, 9016-9085.
11. Skubi, K.L. *et al.* (2016) Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* 116, 10035-10074.
12. Fabry, D.C. and Rueping, M. (2016) Merging Visible Light Photoredox Catalysis with Metal Catalyzed C–H Activations: On the Role of Oxygen and Superoxide Ions as Oxidants. *Acc. Chem. Res.* 49, 1969-1979.
13. Twilton, J. *et al.* (2017) The merger of transition metal and photocatalysis. *Nat. Rev. Chem.* 1, 0052.
14. Hopkinson, M.N. *et al.* (2016) Merging Visible Light Photoredox and Gold Catalysis. *Acc. Chem. Res.* 49, 2261-2272.
15. De Abreu, M. *et al.* Synergistic Photoredox/Transition-Metal Catalysis for Carbon–Carbon Bond Formation Reactions. *Eur. J. Org. Chem.* Early View. DOI: 10.1002/ejoc.201901146.
16. Kalyani, D. (2011) Room-Temperature C–H Arylation: Merger of Pd-Catalyzed C–H Functionalization and Visible-Light Photocatalysis. *J. Am. Chem. Soc.* 133, 18566-18569.

17. Tsou, T.T. and Kochi, J.K. (1979) Mechanism of biaryl synthesis with nickel complexes. *J. Am. Chem. Soc.* 101, 7547-7560.
18. Jones, G.O. (2010) Computational Explorations of Mechanisms and Ligand-Directed Selectivities of Copper-Catalyzed Ullmann-Type Reactions. *J. Am. Chem. Soc.* 132, 6205-6213.
19. Schley, N.D. and Fu, G.C. (2014) Nickel-Catalyzed Negishi Arylations of Propargylic Bromides: A Mechanistic Investigation. *J. Am. Chem. Soc.* 136, 16588-16593.
20. Gui, Y.-Y. (2016) Photoredox sheds new light on nickel catalysis: from carbon-carbon to carbon-heteroatom bond formation. *Org. Chem. Front.* 3, 522-526.
21. Tellis, J.C. *et al.* (2016) Single-Electron Transmetalation via Photoredox/Nickel Dual Catalysis: Unlocking a New Paradigm for sp^3 - sp^2 Cross-Coupling. *Acc. Chem. Res.* 49, 1429-1439.
22. Milligan, J.A. *et al.* (2019) Alkyl Carbon-Carbon Bond Formation by Nickel/Photoredox Cross-Coupling. *Angew. Chem. Int. Ed.* 58, 6152-6163.
23. Hossain, A. *et al.* (2019) Copper's rapid ascent in visible-light photoredox catalysis. *Science* 364, eaav9713.
24. Reiser, O. (2016) Shining Light on Copper: Unique Opportunities for Visible-Light-Catalyzed Atom Transfer Radical Addition Reactions and Related Processes. *Acc. Chem. Res.* 49, 1990-1996.
25. Tellis, J.C. *et al.* (2014) Single-electron transmetalation in organoboron cross-coupling by photoredox/nickel dual catalysis. *Science* 345, 433-436.
26. Zuo, Z. *et al.* (2014) Merging photoredox with nickel catalysis: Coupling of α -carboxyl sp^3 -carbons with aryl halides. *Science* 345, 437-440.
27. Le, C. *et al.* (2018) A radical approach to the copper oxidative addition problem: Trifluoromethylation of bromoarenes. *Science* 360, 1010-1014.
28. Ye, Y. and Sanford, M.S. (2012) Merging Visible-Light Photocatalysis and Transition-Metal Catalysis in the Copper-Catalyzed Trifluoromethylation of Boronic Acids with CF_3I . *J. Am. Chem. Soc.* 134, 9034-9037.
29. Kautzky, J.A. *et al.* (2018) Decarboxylative Trifluoromethylation of Aliphatic Carboxylic Acids. *J. Am. Chem. Soc.* 140, 6522-6526.
30. Terrett, J.A. *et al.* (2015) Switching on elusive organometallic mechanisms with photoredox catalysis. *Nature* 524, 330.
31. Liang, Y. *et al.* (2018). Decarboxylative sp^3 C-N coupling via dual copper and photoredox catalysis. *Nature* 559, 83-88.

32. Ahn, J.M. *et al.* (2017) Design of a Photoredox Catalyst that Enables the Direct Synthesis of Carbamate-Protected Primary Amines via Photoinduced, Copper-Catalyzed N-Alkylation Reactions of Unactivated Secondary Halides. *J. Am. Chem. Soc.* 139, 18101-18106.
33. Mao, R. *et al.* (2018) Decarboxylative C(sp³)-N cross-coupling via synergetic photoredox and copper catalysis. *Nat. Catal.* 1, 120-126.
34. Yoo, W.-J. *et al.* (2015) Visible-Light-Mediated Chan-Lam Coupling Reactions of Aryl Boronic Acids and Aniline Derivatives. *Angew. Chem. Int. Ed.* 54, 6587-6590.
35. Griffin, J.D. *et al.* (2017) Reversing the Regioselectivity of Halofunctionalization Reactions through Cooperative Photoredox and Copper Catalysis. *Angew. Chem. Int. Ed.* 56, 2097-2100.
36. Kainz, Q.M. *et al.* (2016) Asymmetric copper-catalyzed C-N cross-couplings induced by visible light. *Science* 351, 681-684.
37. Li, Y. *et al.* (2018) Copper(II)-Catalyzed Asymmetric Photoredox Reactions: Enantioselective Alkylation of Imines Driven by Visible Light. *J. Am. Chem. Soc.* 140, 15850-15858.
38. Wang, D. *et al.* (2017) Enantioselective Decarboxylative Cyanation Employing Cooperative Photoredox Catalysis and Copper Catalysis. *J. Am. Chem. Soc.* 139, 15632-15635.
39. Eckenhoff, W.T. *et al.* (2013) Cobalt complexes as artificial hydrogenases for the reductive side of water splitting. *Biochim. Biophys. Acta Bioenergetics* 1827, 958-973.
40. Giedyk, M. *et al.* (2015) Vitamin B12 catalysed reactions. *Chem. Soc. Rev.* 44, 3391-3404.
41. Tahara, K. *et al.* (2018) Learning from B12 enzymes: biomimetic and bioinspired catalysts for eco-friendly organic synthesis. *Beilstein J. Org. Chem.* 14, 2553-2567.
42. Girard, S.A. *et al.* (2014) The Cross-Dehydrogenative Coupling of C-H Bonds: A Versatile Strategy for C-C Bond Formations. *Angew. Chem. Int. Ed.* 53, 74-100.
43. Liu, C. *et al.* (2011) Bond Formations between Two Nucleophiles: Transition Metal Catalyzed Oxidative Cross-Coupling Reactions. *Chem. Rev.* 111, 1780-1824.
44. He, K.-H. and Li, Y. (2014) Oxidant-Free Dehydrogenative Coupling Reactions via Hydrogen Evolution. *ChemSusChem* 7, 2788-2790.

45. Tang, S. *et al.* (2018) Oxidative R(1)–H/R(2)–H Cross-Coupling with Hydrogen Evolution. *J. Am. Chem. Soc.* 140, 13128-13135.
46. Sauermann, N. *et al.* (2018) Electrocatalytic C–H Activation. *ACS Catal.* 8, 7086-7103.
47. Wang, H. *et al.* (2019) Recent Advances in Oxidative R1–H/R2–H Cross-Coupling with Hydrogen Evolution via Photo-/Electrochemistry. *Chem. Rev.* 119, 6769-6787.
48. Cartwright, K.C. *et al.* Cobaloxime-Catalyzed Hydrogen Evolution in Photoredox-Facilitated Small-Molecule Functionalization. *Eur. J. Org. Chem.* Early View. DOI: 10.1002/ejoc.201901170.
49. Meng, Q.-Y. *et al.* (2013) A Cascade Cross-Coupling Hydrogen Evolution Reaction by Visible Light Catalysis. *J. Am. Chem. Soc.* 135, 19052-19055.
50. Zhong, J.-J. *et al.* (2014) Cross-Coupling Hydrogen Evolution Reaction in Homogeneous Solution without Noble Metals. *Org. Lett.* 16, 1988-1991.
51. Gao, X.-W. *et al.* (2015) Visible Light Catalysis Assisted Site-Specific Functionalization of Amino Acid Derivatives by C–H Bond Activation without Oxidant: Cross-Coupling Hydrogen Evolution Reaction. *ACS Catal.* 5, 2391-2396.
52. Xiang, M. *et al.* (2015) Activation of C–H Bonds through Oxidant-Free Photoredox Catalysis: Cross-Coupling Hydrogen-Evolution Transformation of Isochromans and β -Keto Esters. *Chem. – Eur. J.* 21, 18080-18084.
53. Yang, Q. *et al.* (2017) Visible-Light-Promoted Asymmetric Cross-Dehydrogenative Coupling of Tertiary Amines to Ketones by Synergistic Multiple Catalysis. *Angew. Chem. Int. Ed.* 56, 3694-3698.
54. Wu, C.-J. *et al.* (2016) An Oxidant-Free Strategy for Indole Synthesis via Intramolecular C–C Bond Construction under Visible Light Irradiation: Cross-Coupling Hydrogen Evolution Reaction. *ACS Catal.* 6, 4635-4639.
55. Hu, X. *et al.* (2018) Photoinduced oxidative activation of electron-rich arenes: alkenylation with H₂ evolution under external oxidant-free conditions. *Chem. Sci.* 9, 1521-1526.
56. Cao, H. *et al.* (2018) Photo-induced Decarboxylative Heck-Type Coupling of Unactivated Aliphatic Acids and Terminal Alkenes in the Absence of Sacrificial Hydrogen Acceptors. *J. Am. Chem. Soc.* 140, 16360-16367.
57. Koeller, J. *et al.* (2019) Visible-Light-Induced Decarboxylative C–H Adamantylation of Azoles at Ambient Temperature. *Synthesis* 51, 1284-1292.

58. Tian, W.-F. *et al.* (2019) Visible-Light Photoredox-Catalyzed Decarboxylative Alkylation of Heteroarenes Using Carboxylic Acids with Hydrogen Release. *Org. Lett.* 21, 6930-6935.
59. Zhang, G. *et al.* (2018) Oxidative [4+2] annulation of styrenes with alkynes under external-oxidant-free conditions. *Nat. Commun.* 9, 1225.
60. Hu, X. *et al.* (2018) Selective Oxidative [4+2] Imine/Alkene Annulation with H₂ Liberation Induced by Photo-Oxidation. *Angew. Chem. Int. Ed.* 57, 1286-1290.
61. Tian, W.-F. *et al.* (2018) Visible-Light Photoredox-Catalyzed Iminyl Radical Formation by N–H Cleavage with Hydrogen Release and Its Application in Synthesis of Isoquinolines. *Org. Lett.* 20, 1421-1425.
62. Zhang, G. *et al.* (2015) External Oxidant-Free Oxidative Cross-Coupling: A Photoredox Cobalt-Catalyzed Aromatic C–H Thiolation for Constructing C–S Bonds. *J. Am. Chem. Soc.* 137, 9273-9280.
63. Zheng, Y.-W. *et al.* (2016) Photocatalytic Hydrogen-Evolution Cross-Couplings: Benzene C–H Amination and Hydroxylation. *J. Am. Chem. Soc.* 138, 10080-10083.
64. Yi, H. *et al.* (2017) Photocatalytic Dehydrogenative Cross-Coupling of Alkenes with Alcohols or Azoles without External Oxidant. *Angew. Chem. Int. Ed.* 56, 1120-1124.
65. Niu, L. *et al.* (2017) Photo-induced oxidant-free oxidative C–H/N–H cross-coupling between arenes and azoles. *Nat. Commun.* 8, 14226.
66. Chen, H. *et al.* (2018) External Oxidant-Free Regioselective Cross Dehydrogenative Coupling of 2-Arylimidazoheterocycles and Azoles with H₂ Evolution via Photoredox Catalysis. *Adv. Synth. Catal.* 360, 3220-3227.
67. Zhang, G. *et al.* (2016) Anti-Markovnikov Oxidation of β -Alkyl Styrenes with H₂O as the Terminal Oxidant. *J. Am. Chem. Soc.* 138, 12037-12040.
68. Shao, A. *et al.* (2018) External Oxidant-Free Dehydrogenative Lactonization of 2-Arylbenzoic Acids via Visible-Light Photocatalysis. *J. Org. Chem.* 83, 3582-3589.
69. Zhang, M. *et al.* (2018) Photoredox and cobalt co-catalyzed C(sp²)–H functionalization/C–O bond formation for synthesis of lactones under oxidant- and acceptor-free conditions. *Org. Chem. Front.* 5, 749-752.
70. Niu, L. *et al.* (2017) Visible-Light-Induced External Oxidant-Free Oxidative Phosphonylation of C(sp²)–H Bonds. *ACS Catal.* 7, 7412-7416.

71. Niu, L. *et al.* (2018) Visible light-mediated oxidative C(sp³)–H phosphonylation for α -aminophosphonates under oxidant-free conditions. *Chem. Commun.* 54, 1659-1662.
72. Luo, K. *et al.* (2016) Cross-Coupling Hydrogen Evolution by Visible Light Photocatalysis Toward C(sp²)–P Formation: Metal-Free C–H Functionalization of Thiazole Derivatives with Diarylphosphine Oxides. *Org. Lett.* 18, 452-455.
73. Zhang, G. *et al.* (2016) Visible-light induced oxidant-free oxidative cross-coupling for constructing allylic sulfones from olefins and sulfinic acids. *Chem. Commun.* 52, 10407-10410.
74. Yu, W.-L. *et al.* (2019) Dehydrogenative Silylation of Alkenes for the Synthesis of Substituted Allylsilanes by Photoredox, Hydrogen-Atom Transfer, and Cobalt Catalysis. *Angew. Chem. Int. Ed.* 58, 10941-10945.
75. West, J.G. *et al.* (2015) Acceptorless dehydrogenation of small molecules through cooperative base metal catalysis. *Nat. Commun.* 6, 10093.
76. Abrams, D.J. *et al.* (2017) Toward a mild dehydroformylation using base-metal catalysis. *Chem. Sci.* 8, 1954-1959.
77. He, K.-H. *et al.* (2017) Acceptorless Dehydrogenation of N-Heterocycles by Merging Visible-Light Photoredox Catalysis and Cobalt Catalysis. *Angew. Chem. Int. Ed.* 56, 3080-3084.
78. Sahoo, M.K. and Balaraman, E. (2019) Room temperature catalytic dehydrogenation of cyclic amines with the liberation of H₂ using water as a solvent. *Green Chem.* 21, 2119-2128.
79. Jia, Z. *et al.* (2019) Photoredox Mediated Acceptorless Dehydrogenative Coupling of Saturated N-Heterocycles. *ACS Catal.* 9, 3589-3594.
80. Sahoo, M.K. *et al.* (2018) Photocatalysis Enabling Acceptorless Dehydrogenation of Diaryl Hydrazines at Room Temperature. *ACS Catal.* 8, 7727-7733.
81. Sun, X. *et al.* (2018) Catalytic dehydrogenative decarboxyolefination of carboxylic acids. *Nat. Chem.* 10, 1229-1233.
82. Cartwright, K.C. and Tunge, J.A. (2018) Decarboxylative Elimination of N-Acyl Amino Acids via Photoredox/Cobalt Dual Catalysis. *ACS Catal.* 8, 11801-11806.
83. Nguyen, V.T. *et al.* (2019). Alkene Synthesis by Photocatalytic Chemoenzymatically Compatible Dehydrodecarboxylation of Carboxylic Acids and Biomass. *ACS Catal.* 9, 9485-9498.
84. Green, S.A. *et al.* (2018) The High Chemofidelity of Metal-Catalyzed Hydrogen Atom Transfer. *Acc. Chem. Res.* 51, 2628-2640.

85. Thullen, S.M. and Rovis, T. (2017) A Mild Hydroaminoalkylation of Conjugated Dienes Using a Unified Cobalt and Photoredox Catalytic System. *J. Am. Chem. Soc.* 139, 15504-15508.
86. Meng, Q.-Y. *et al.* (2019) Controllable Isomerization of Alkenes by Dual Visible-Light-Cobalt Catalysis. *Angew. Chem. Int. Ed.* 58, 5723-5728.
87. Crossley, S.W.M. *et al.* (2014) Simple, Chemoselective, Catalytic Olefin Isomerization. *J. Am. Chem. Soc.* 136, 16788-16791.
88. Rai, P. *et al.* (2019) Photoredox/Cobalt Dual Catalysis for Visible-Light-Mediated Alkene–Alkyne Coupling. *Org. Lett.* 21, 3755-3759.
89. Grenier-Petel, J.-C. and Collins, S.K. (2019) Photochemical Cobalt-Catalyzed Hydroalkynylation To Form 1,3-Enynes. *ACS Catal.* 9, 3213-3218.
90. Vollhardt, K.P.C. (1984) Cobalt-Mediated [2 + 2 + 2]-Cycloadditions: A Maturing Synthetic Strategy [New Synthetic Methods (43)]. *Angew. Chem. Int. Ed. Engl.* 23, 539-556.
91. Gandon, V. *et al.* (2006) Recent progress in cobalt-mediated [2 + 2 + 2] cycloaddition reactions. *Chem. Commun.* 2209-2217.
92. Gandeepan, P. and Cheng, C.-H. (2015) Cobalt Catalysis Involving π Components in Organic Synthesis. *Acc. Chem. Res.* 48, 1194-1206.
93. Ruhl, K.E. and Rovis, T. (2016) Visible Light-Gated Cobalt Catalysis for a Spatially and Temporally Resolved [2+2+2] Cycloaddition. *J. Am. Chem. Soc.* 138, 15527-15530.
94. Ravetz, B.D. *et al.* (2018) External Regulation of Cobalt-Catalyzed Cycloaddition Polymerization with Visible Light. *ACS Catal.* 8, 5323-5327.
95. Ravetz, B.D. *et al.* (2018) Photoinduced Ligand-to-Metal Charge Transfer Enables Photocatalyst-Independent Light-Gated Activation of Co(II). *ACS Catal.* 9, 200-204.
96. Hou, J. *et al.* (2018) Visible-Light-Driven Alkyne Hydro-/Carboxylation Using CO₂ via Iridium/Cobalt Dual Catalysis for Divergent Heterocycle Synthesis. *J. Am. Chem. Soc.* 140, 5257-5263.
97. Turnbull, B.W.H. and Evans, P.A. (2018) Asymmetric Rhodium-Catalyzed Allylic Substitution Reactions: Discovery, Development and Applications to Target-Directed Synthesis. *J. Org. Chem.* 83, 11463-11479.
98. Hartwig, J.F. and Stanley, L.M. (2010) Mechanistically Driven Development of Iridium Catalysts for Asymmetric Allylic Substitution. *Acc. Chem. Res.* 43, 1461-1475.

99. Lafrance, M. *et al.* (2012) Direct, Enantioselective Iridium-Catalyzed Allylic Amination of Racemic Allylic Alcohols. *Angew. Chem. Int. Ed.* 51, 3470-3473.
100. Cheng, Q. *et al.* (2019) Iridium-Catalyzed Asymmetric Allylic Substitution Reactions. *Chem. Rev.* 119, 1855-1969.
101. Sun, M. *et al.* (2019). Construction of Vicinal Quaternary Carbon Centers via Cobalt-Catalyzed Asymmetric Reverse Prenylation. *Org. Lett.* 21, 1278-1282.
102. Ghorai, S. *et al.* (2019). Cobalt-Catalyzed Regio- and Enantioselective Allylic Amination. *J. Am. Chem. Soc.* 141, 11430-11434.
103. Giedyk, M. *et al.* (2017) Photoinduced Vitamin B12-Catalysis for Deprotection of (Allyloxy)arenes. *Org. Lett.* 19, 2670-2673.
104. Takizawa, K. *et al.* (2019) Cobalt-Catalyzed Allylic Alkylation Enabled by Organophotoredox Catalysis. *Angew. Chem. Int. Ed.* 58, 9199-9203.
105. Fabry, D.C. *et al.* (2014) Combining Rhodium and Photoredox Catalysis for C–H Functionalizations of Arenes: Oxidative Heck Reactions with Visible Light. *Angew. Chem. Int. Ed.* 53, 10228-10231.
106. Kalsi, D. *et al.* (2018) Room-Temperature C–H Bond Functionalization by Merging Cobalt and Photoredox Catalysis. *ACS Catal.* 8, 8115-8120.
107. Kalsi, D. *et al.* (2019). C–H and N–H bond annulation of aryl amides with unactivated olefins by merging cobalt(III) and photoredox catalysis. *Chem. Commun.* 55, 11626-11629.
108. Dwivedi, V. *et al.* (2019) Electrochemical-/Photoredox Aspects of Transition Metal-Catalyzed Directed C–H Bond Activation. *ChemCatChem* 11, 5160-5187.
109. Call, A. *et al.* (2017) Dual cobalt–copper light-driven catalytic reduction of aldehydes and aromatic ketones in aqueous media. *Chem. Sci.* 8, 4739-4749.
110. Claros, M. *et al.* (2019) Reductive Cyclization of Unactivated Alkyl Chlorides with Tethered Alkenes under Visible-Light Photoredox Catalysis. *Angew. Chem. Int. Ed.* 58, 4869-4874.
111. Perry, I.B. *et al.* (2018) Direct arylation of strong aliphatic C–H bonds. *Nature* 560, 70-75.
112. Dewanji, A. *et al.* (2019) The Dual Role of Benzophenone in Visible-Light/Nickel Photoredox-Catalyzed C–H Arylations: Hydrogen-Atom Transfer and Energy Transfer. *Angew. Chem. Int. Ed.* 58, 3566-3570.
113. Shen, Y. *et al.* (2018) sp³ C–H Arylation and Alkylation Enabled by the Synergy of Triplet Excited Ketones and Nickel Catalysts. *J. Am. Chem. Soc.* 140, 12200-12209.

114. Shaw, M.H. *et al.* (2016) Native functionality in triple catalytic cross-coupling: sp^3 C–H bonds as latent nucleophiles. *Science* 352, 1304-1308.
115. Shields, B.J. and Doyle, A.G. (2016) Direct $\text{C}(\text{sp}^3)$ –H Cross Coupling Enabled by Catalytic Generation of Chlorine Radicals. *J. Am. Chem. Soc.* 138, 12719-12722.
116. Heitz, D.R. *et al.* (2016) Photochemical Nickel-Catalyzed C–H Arylation: Synthetic Scope and Mechanistic Investigations. *J. Am. Chem. Soc.* 138, 12715-12718.
117. Kato, S. *et al.* (2017) Hybrid Catalysis Enabling Room-Temperature Hydrogen Gas Release from N-Heterocycles and Tetrahydronaphthalenes. *J. Am. Chem. Soc.* 139, 2204-2207.
118. Gao, K. and Yoshikai, N. (2014) Low-Valent Cobalt Catalysis: New Opportunities for C–H Functionalization. *Acc. Chem. Res.* 47, 1208-1219.
119. Moselage, M. *et al.* (2016). Cobalt-Catalyzed C–H Activation. *ACS Catal.* 6, 498-525.
120. Yoshino, T. and Matsunaga, S. (2017) (Pentamethylcyclopentadienyl)cobalt(III)-Catalyzed C–H Bond Functionalization: From Discovery to Unique Reactivity and Selectivity. *Adv. Synth. Catal.* 359, 1245-1262.

Figure Captions.

Figure 1. Merger of 3d-metal Catalysis and Photoredox Catalysis. (A) 3d-metal catalysts and photoredox catalysts are compatible in nature. (B) Cobalt/photoredox cooperative catalysis is an emerging technology in organic synthesis. (C) Representative mechanisms in cobalt/photoredox catalysis. Abbreviations: PC, photoredox catalyst; A, electron acceptor; D, electron donor.

Figure 2. Cobalt/Photoredox-catalyzed Cross-coupling Hydrogen Evolution (CCHE). (A) A representative CCHE between amines and nucleophiles. (B) CCHE for the construction of naphthalenes *via* the formation of two C–C bonds. (C) CCHE enables an oxidative transformation of thioamides. (D) Direct formation of phenols from benzene by CCHE. Abbreviations: LED, light-emitting diode; PMP, 4-methoxyphenyl.

Figure 3. Cobalt/Photoredox-Catalyzed Acceptorless Dehydrogenation. (A) Dehydrogenation of unreactive alkanes. (B) Dehydroformylation of aldehydes *via* a dehydrogenation. (C) Dehydrogenation of *N*-heterocycles provides *N*-heteroarenes. (D) Application of the catalytic system to the conversion of carboxylic acids into alkenes. Abbreviations: Py, pyridine; M.S., molecular sieves; DME, 1,2-dimethoxyethane.

Figure 4. Hydrofunctionalization of Alkenes by Cobalt/Photoredox Cooperative Catalysis. (A) Hydroaminoalkylation of dienes by merging cobalt catalysis and radical chemistry. (B) A photochemically generated cobalt hydride mediates the catalytic isomerization of alkenes. Abbreviations: acac, acetylacetonate; DMF, *N,N*-dimethylformamide.


Figure 5. Cycloaddition of Alkynes mediated by Cobalt/Photoredox Dual Catalysis. (A) Visible-light-gated [2+2+2] cycloaddition. (B) Low-valent cobalt catalysis enables the capture of CO₂ by cycloaddition. Abbreviations: CFL, compact fluorescent lightbulb; Cy, cyclohexyl; Ts, *p*-toluenesulfonyl.

Figure 6. Allylation and deallylation by Cobalt/Photoredox Dual Catalysis. (A) Photochemically activated Vitamin B12 mediates the deallylation of phenols. (B) Low-valent cobalt-catalyzed allylation exhibits high regioselectivity. Abbreviations: B, branched isomer; L: linear isomer; dppp, 1,3-diphenylphosphinopropane.


Figure 7. Reactions using stoichiometric oxidant or reductant. (A) Photoredox-enabled aerobic oxidative C–H bond alkenylation/cyclization. (B) Cobalt-catalyzed reduction of carbonyls *via* net hydride transfer. (C) Reductive dechlorination/cyclization by low-valent cobalt catalyst. Abbreviations: TFE, 2,2,2-trifluoroethanol; E, ethoxycarbonyl.

Figures.

(A) Merger of 3d-metal catalysis and photoredox catalysis



(B) Background of the merger of Co catalysis and photoredox catalysis



(C) General mechanisms














Figure 1.

(A) CCHE between tetrahydroisoquinolines and indoles**(B) CCHE for successive formation of two C-C bonds****(C) Intramolecular C-S bond formation by CCHE****(D) C-H hydroxylation of benzene****Figure 2.**


(A) Acceptorless dehydrogenation of alkanes



(B) Dehydrogenative dehydroformylation of aldehydes



(C) Acceptorless dehydrogenation of *N*-heterocycles



(D) Dehydrogenative decarboxyolefination of carboxylic acids






Figure 3.

(A) Hydroaminoalkylation of dienes



(B) Isomerization of alkenes



(C) Hydroalkenylation of alkenes








Figure 4.

(A) [2+2+2] Cycloaddition



(B) Cycloaddition with CO₂








Figure 5.

(A) Deallylation of phenol derivatives



(B) Regioselective allylic alkylation






Figure 6.

(A) Oxidative C-H functionalization using O₂



(B) Reduction of ketones and aldehydes using Et₃N



(C) Reductive cyclization of alkyl chlorides using Et₃N





Figure 7.