



HOKKAIDO UNIVERSITY

Title	Long-term prognostic factors of patients with hepatocellular carcinoma who survive over 10 years after hepatectomy
Author(s)	Shimada, Shingo; Kamiyama, Toshiya; Orimo, Tatsuya et al.
Citation	Journal of Surgical Oncology, 121(8), 1209-1217 https://doi.org/10.1002/jso.25910
Issue Date	2020-06-15
Doc URL	https://hdl.handle.net/2115/81916
Rights	This is the peer reviewed version of the following article: Shimada, S, Kamiyama, T, Orimo, T, et al. Long-term prognostic factors of patients with hepatocellular carcinoma who survive over 10 years after hepatectomy. J Surg Oncol. 2020; 121: 1209-1217, which has been published in final form at https://doi.org/10.1002/jso.25910 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Type	journal article
File Information	J Surg Oncol 121_1209.pdf



Original article

Long-term prognostic factors of patients with hepatocellular carcinoma who survive over ten years after hepatectomy.

Shingo Shimada, MD, PhD,¹ Toshiya Kamiyama, MD, PhD, FACS,¹ Tatsuya Orimo, MD, PhD,¹
Akihisa Nagatsu, MD,¹ Yoh Asahi, MD,¹ Yuzuru Sakamoto, MD,¹ Hirofumi Kamachi, MD, PhD,¹ and
Akinobu Taketomi, MD, PhD, FACS.¹

Affiliation and address of the authors: ¹Department of Gastroenterological Surgery I, Hokkaido
University Graduate School of Medicine, Sapporo, Japan
North 15-West 7, Kita-Ku, Sapporo, Hokkaido 060-8638, Japan

Corresponding author: Shingo Shimada

Department of Gastroenterological Surgery I, Hokkaido University Graduate School of Medicine,
Sapporo, Japan North 15-West 7, Kita-Ku, Sapporo, Hokkaido 060-8638, Japan

Tel.: +81-11-7065927; Fax: +81-11-7177515

E-mail: shingoshimada1979@true.ocn.ne.jp

1 Short title

2 Long-term favorable factors of HCC

3

4 Authorship

5 SS and TK conceived the study concept and design, were involved with patient care and drafted the
6 manuscript and literature review. TO, AN, YA, YS, HK, and AT were involved with the formation of
7 the study concept and design, patient care and drafting of the manuscript and literature review.

8 All authors have read and approved the final version of the manuscript.

9

10 Grant support

11 We have no grant support for this study.

12

13 Competing interests

14 The authors declare no competing interests in relation to this study.

15

16 Synopsis

17 Important factors for long-term prognoses in HCC patients were solitary tumor, small tumors, and no
18 advanced fibrosis. Even patients who relapse might survive long term if they have late or solitary
19 intrahepatic recurrence, nonsevere cirrhosis, and curative treatment at recurrence.

20

1 Data Availability Statement

2 The data that support the findings of this study are available on request from the corresponding author.

3 The data are not publicly available due to privacy or ethical restrictions.

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

Abstract

Background and objectives: The aim was to evaluate long-term prognostic factors in hepatocellular carcinoma (HCC) patients who survived over ten years after hepatectomy and compare prognostic factors between patients with recurrence who died and survived 10 years after initial hepatectomy.

Methods: We analyzed the HCC patients without recurrence over 10 years after hepatectomy (n=35), those with recurrence who survived over 10 years (n=48), and those who died within 10 years (n=132).

Results: The rate of recurrence was 16.3 %, ten-year overall survival (OS) rate was 38.6 %, and the ten-year recurrence-free survival (RFS) rate was 16.7 %. Nonviral, solitary tumor, well differentiation, and without severe fibrosis were independent favorable factors for long-term RFS. High cholinesterase levels, small tumor and without portal vein invasion were independent favorable factors for long-term survival among patients with recurrence. Long-term survivors with recurrence showed significantly low early recurrence, extrahepatic recurrence, multiple intrahepatic recurrence.

Conclusion: Important factors for long-term prognoses in HCC patients were solitary tumor, small tumors, and no advanced fibrosis. A treatment for nonviral hepatitis is needed to achieve long-term RFS. Even patients who relapse might survive long term if they have late or solitary intrahepatic recurrence, nonsevere cirrhosis, and curative treatment at recurrence.

Keywords: long-term prognoses; hepatectomy; hepatocellular carcinoma; nonviral hepatitis; solitary intrahepatic recurrence

Text

INTRODUCTION

It is estimated that 18.1 million new cancer cases and 9.6 million cancer-related deaths occurred in 2018 around the world [1]. Liver cancer is the seventh most frequent cancer, with an estimated 841,080 cases per year, and the second leading cause of cancer-related death; liver cancer is responsible for approximately 781,631 deaths per year [1]. Hepatocellular carcinoma (HCC) shows high recurrence rates, ranging from 70-75 % within five years after hepatectomy [2, 3]. Thus, these patients have many opportunities for the treatment of recurrence. According to the report of the Liver Cancer Study Group of Japan, the 3-year, 5-year, and 10-year overall survival (OS) rates of patients with HCC who undergo hepatectomy are 72.3 %, 56.8 % and 32.0 %, respectively [4]. Survival rates decrease over time. Several prognostic factors for the survival and recurrence of patients with resected HCC have been reported and include tumor size, tumor number, tumor marker, vascular invasion, and liver function [5-7]. However, reports on the prognostic factors of long-term survivors with resected HCC are rare. Therefore, in this study, we evaluated long-term prognostic factors in patients with HCC who survived over ten years after hepatectomy. We also compared prognostic factors between patients who died and survived 10 years after initial hepatectomy among those with recurrence.

METHODS

Patients and methods

Between January 2000 and November 2007, 339 consecutive patients with HCC underwent primary hepatectomy at the Gastroenterological Surgery I Unit of the Hokkaido University Hospital in Sapporo, Japan. We investigated 215 patients with confirmed outcomes ten years after hepatectomy, excluding deaths caused by other diseases. The hepatitis B virus (HBV) group included patients with HBV infection alone, the hepatitis C virus (HCV) group included patients with HCV infection alone, and the non-B non-C (NBNC) group included patients without HBV and HCV infection.

We divided the patients into three groups as follows: patients without recurrence over 10 years after hepatectomy (recurrence free: Group RF; n=35, 35/215, 16.3 %), patients with recurrence who survived over 10 years after hepatectomy (recurrence survive: Group RS; n=48, 48/215, 22.3 %), and patients who died within 10 years after hepatectomy excluding death caused by other diseases (recurrence dead: Group RD; n=132, 132/215, 61.4 %). First, we evaluated long-term survival and recurrence factors at primary hepatectomy and prognostic factors for survival among patients with recurrence. Second, we evaluated the timing of recurrence and treatment after recurrence.

This study was approved by the Hokkaido University Hospital Voluntary Clinical Study Committee (approval 018-0303) and was performed in accordance with the Helsinki Declaration guidelines.

Hepatectomy

The type of surgical procedure was usually determined based on the patients' liver function

reserve, i.e., according to the results of the indocyanine green retention test at 15 min (ICGR15) [8]. In principle, anatomical resection was performed for patients with an ICGR15 result less than 25 %. All surgeries showed that the surgical margins were histologically or macroscopically free. Fibrosis was defined as f3, bridging fibrosis or f4; cirrhosis was defined according to the general rules for the clinical and pathological study of primary liver cancer set by the Liver Cancer Study Group of Japan [4]. Postoperative morbidity was assessed using the validated Clavien–Dindo classification system [9]. Serious complications were categorized as grades III–V and defined as morbidity requiring surgical or radiological intervention.

Follow-up after hepatectomy

Follow-up studies after liver resection were conducted at 3-month intervals and included physical, serological (liver function, serum alpha-fetoprotein [AFP], and serum protein induced by vitamin K absence-II [PIVKA-II] tests), and radiological examinations (ultrasound sonography [US] and contrast-enhanced computed tomography [CT] scans or contrast-enhanced magnetic resonance imaging [MRI]).

Statistical analyses

Univariate analyses were performed using the chi-square test for noncontinuous variables. Logistic regression model analyses were performed for multivariate analyses. OS and recurrence-free survival (RFS) were analyzed by using the Kaplan-Meier method with the log-rank test or the Cox proportional hazards model. JMP Pro 14.0.0 for Windows (SAS Institute, Cary, NC) was used for the statistical analyses.

RESULTS

Prognoses of the study cohort

In this cohort of 215 study patients, the median survival time (MST) and ten-year OS rate were 69 months and 38.6 %, respectively. The ten-year RFS rate was 16.7 % (Figure 1).

Long-term prognostic factors for recurrence

Table 1a presents the clinicopathological factors related to long-term RFS between Groups RF and RS+RD. The univariate analysis showed a significantly higher proportion of females, not NBNC, platelets $\geq 80,000/\text{mm}^3$, albumin (Alb) ≥ 4.0 g/dl, total bilirubin (T-bil) < 1.0 mg/dl, ICGR15 < 15 %, AFP < 20 ng/ml, tumor size < 4.5 cm, solitary tumor, well differentiation, patients without portal vein invasion (PVI), without f3/f4, anatomical resection, and blood loss at surgery < 200 ml in Group RF than in Group RS+RD.

The multivariate analysis indicated that not NBNC, solitary tumor, well differentiation, and without f3/f4 were independent favorable factors for long-term RFS (Table 1b).

Long-term prognostic factors for survival

Table 2a presents the clinicopathological factors related to long-term OS between Groups RF+RS and RD. The univariate analysis showed a significantly higher proportion of age ≥ 60 years, Alb ≥ 4.0 g/dl, cholinesterase (ChE) ≥ 200 IU/L, ICGR15 < 15 %, AFP < 20 ng/ml, PIVKA-II < 100 mAU/ml, tumor size < 4.5 cm, solitary tumor, well differentiation, without PVI and hepatic vein invasion (HVI), without f3/f4, and blood loss at surgery < 200 ml in Group RF+RS than in Group RD.

The multivariate analysis indicated that AFP < 20 ng/ml, tumor size < 4.5 cm, solitary tumor, well differentiation, and without f3/f4 were independent favorable factors for long-term OS (Table 2b).

Long-term prognostic factors for survival among patients with recurrence

Table 3a presents the clinicopathological factors related to survival between Groups RS and RD. The univariate analysis showed a significantly higher proportion of age $\geq$ 60 years, Alb $\geq$ 4.0 g/dl, ChE $\geq$ 200 IU/L, AFP < 20 ng/ml, PIVKA-II < 100 mAU/ml, tumor size < 4.5 cm, solitary tumor, well differentiation, without PVI and HVI, without f3/f4, and blood loss at surgery < 200 ml in Group RS than in Group RD.

The multivariate analysis indicated that ChE $\geq$ 200 IU/L, tumor size < 4.5 cm, and without PVI were independent favorable factors for survival among patients with recurrence (Table 3b).

Timing of recurrence, recurrence site and number, and treatment after recurrence

Regarding the timing of recurrence, the rates of recurrence within 2 years and 6 months after hepatectomy in Group RS were 42 % and 6 %, respectively; those in Group RD were 85 % and 36 %, respectively. The median recurrence time after hepatectomy in Groups RS and RD was 32 (2-115) months and 8 (0.4-68) months, respectively ($p < 0.01$). Group RD had significantly earlier recurrence (Figure 2).

Regarding the initial recurrence sites, extrahepatic recurrence was significantly higher in Group RD than in Group RS (65.2 % vs. 18.7 %) ($p < 0.01$) (Table 4). In addition, multiple recurrence was a significantly unfavorable factor among patients with liver recurrence alone. Table 4 also shows the treatment methods after recurrence. Rehepatectomy, liver transplantation or ablation was

performed more often in Group RS than in Group RD. Moreover, the number of patients treated with systemic chemotherapy, including molecular target drugs or radiotherapy, was significantly lower in Group RS than in Group RD.

DISCUSSION

The present study indicated that not NBNC, solitary tumor, well differentiation, and no underlying high-grade liver fibrosis were important factors for long-term survival and no recurrence over ten years. The group of patients who died within ten years after hepatectomy had significantly more individuals with early and extrahepatic recurrence than those who survived with recurrence. Multiple recurrence was a significantly unfavorable factor among patients with liver recurrence alone. The patients who survived with recurrence more frequently underwent hepatectomy, liver transplantation or ablation than patients who died within ten years.

Few previous studies have focused on the prognoses of long-term survivors with HCC for more than 10 years worldwide [10-22]. This study showed that the ten-year OS rate was 38.6 %, which is consistent with previous reports, with OS rates ranging from 21.8-32.0 % [4,15,16,19,22]. The rate of no recurrence was 16.3 %, and the rate of recurrence within ten years after hepatectomy was 83.7 %. These results are also consistent with those of other reports, with rates ranging from 8.7-30.6 % [15,19,22]. Eguchi et al. reported that the ten-year RFS rate after hepatectomy for HCC was 22.4 % and that solitary tumor, tumor size less than 5 cm, simple nodular type, and low tumor markers were favorable factors for long-term RFS [18]. Kim et al. reported that the ten-year RFS rate

after hepatectomy for HCC was 24.1 % and that HBe-Ag positivity, satellite nodules, vascular invasion, and high ICGR15 values were unfavorable factors for long-term RFS [19]. Another report showed that ten-year RFS rates after hepatectomy for HCC were 9.9-11.7 % [11,16]. In 2005, Shimada et al. reported that HCV-Ab positivity was an unfavorable factor for long-term RFS [15]. In contrast, NBNC was an unfavorable factor for long-term RFS in this study. This discrepancy might be explained by the following: in recent years, HCV has been controlled with antiviral drugs, and it was reported that the sustained virologic response (SVR) rate was 95.2 % with HCV, 95.9 % with Child-Pugh class A and 88.3 % with Child-Pugh class B [23]. However, no treatment has been established for NBNC. Several studies have claimed that underlying liver fibrosis and cirrhosis might predispose patients to multicentric hepatocarcinogenesis [24,25], and the risk of HCC was 24 times higher in patients with severe cirrhosis than in patients with normal or mild fibrosis [26]. Although no significance was observed in the multivariate analysis, female sex was significantly favorable for RFS in the univariate analysis. Some studies have reported that a potential role of androgen is associated with liver carcinogenesis [27, 28]. Regarding long-term RFS, tumor size was not an independent factor, but solitary tumor was a strongly favorable factor in the multivariate analysis. It has been reported that solitary large HCC has better molecular characteristics than nodular HCC; these characteristics are derived from differences in gene expression, e.g., cathepsin L, integrin alpha6, and RhoC [29].

On the other hand, our results showed that low AFP, small tumor size, solitary tumor, well differentiation, and without f3/f4 were independent favorable factors for long-term OS. These results

are consistent with those of previous reports [15,16,19,21,22]. Regarding long-term prognostic factors among patients with HCC who experienced recurrence within ten years after hepatectomy, small tumor size, high ChE and without PVI were independent favorable factors for long-term survival. The patients who survived more than ten years after hepatectomy showed significantly less early recurrence and less extrahepatic recurrence. Large HCC is a risk factor for PVI [30]. PVI is also known as a risk factor for the early recurrence of HCC [31].

Serum ChE is known as a biomarker of liver cirrhosis [32]. No underlying severe liver cirrhosis is very important for the treatment of secondary liver recurrence to achieve curative treatment. Because improved liver function, i.e., ChE, was one of the favorable prognostic factors for HCC [33] and, in this study, the absence of high-grade liver fibrosis was an important factor for long-term survival and no recurrence over ten years, the improvement in fibrosis for the treatment of viral hepatitis and nonalcoholic steatohepatitis (NASH) was necessary for long-term survival regardless of recurrence.

In this study, early recurrence is a strong unfavorable factor. Zheng et al. reported that 6 % of long-term survivors experienced recurrence within 2 years, while 77 % of short-term survivors experienced recurrence [22]. Poon et al. reported that early recurrence (within 1 year) arises mainly from intrahepatic metastases, which tend to be multifocal, and that late recurrence (after 1 year) is more likely to be related to multicentric occurrence in origin [34]. Consistent with these results, intrahepatic recurrence in non-long-term survivors might originate from not only multicentric occurrence but also intrahepatic secondary metastases. This study showed that multiple recurrence

was a significantly unfavorable factor among patients with liver recurrence alone. Multiple intrahepatic recurrence is associated with intrahepatic metastases [35].

Radical treatments, such as rehepatectomy, liver transplantation or ablation, are often performed in long-term survivors [20,21]. Minagawa et al. reported that repeat hepatic resection was very useful for improving prognoses, especially in patients with single HCC at primary resection, negative portal invasion at repeat resection, and a disease-free interval of more than 1 year after initial hepatectomy [36]. Belghiti et al. reported the usefulness of salvage liver transplantation, and the results of salvage liver transplantation were almost the same as those of first liver transplantation [37]. In addition, it was reported that though hepatic resection was the most effective treatment for recurrent HCC, ablation could be a safe and effective option for the majority of patients with recurrence who were unsuitable for surgery in a systematic review of 19 studies [38].

In this study, the long-term survival of patients with extrahepatic recurrence was insufficient. However, Nakagawa et al. reported that patients who had a disease-free interval of more than 1 year and AFP $\geq$ 500 ng/ml after pulmonary metastasectomy of HCC showed 3- and 5-year survival rates of 89 % and 74 %, respectively [39]. Thus, we might try aggressive surgical management for suitable patients.

Conclusions

The important factors for long-term prognoses of more than ten years in patients with HCC were solitary tumor, small tumors, and no advanced fibrosis. In addition, the development of a

treatment for nonviral hepatitis is needed to achieve long-term RFS. Even patients who experience relapse might be long-term survivors if they have late or solitary intrahepatic recurrence, nonsevere cirrhosis, and curative treatment at recurrence.

Acknowledgments

The authors would like to thank the staff of the Gastroenterological Surgery I Unit of the Hokkaido University Graduate School of Medicine for their kind cooperation.

Abbreviations list

HCC: Hepatocellular carcinoma; HBs-Ag: HBs-antigen; HCV-Ab: HCV-antibody; ICGR15: indocyanine green retention test at 15 min; AFP: Alpha-fetoprotein; PIVKA-II: Protein induced by vitamin K absence-II; US: Ultrasound sonography; CT: Computed tomography; MRI: Magnetic resonance imaging; OS: Overall survival; RFS: Relapse-free survival; MST: Median survival time; NBNC: non-B non-C; PVI: Portal vein invasion; TACE: Transarterial chemoembolization; PT: Prothrombin time; HVI: Hepatic vein invasion; ChE: Cholinesterase.

References

1. Bray F, Ferlay J, Soerjomataram I, et al.: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394-424.
2. Marubashi S, Gotoh K, Akita H, et al.: Analysis of recurrence patterns after anatomical or non-anatomical resection for hepatocellular carcinoma. *Ann Surg Oncol* 2015;22:2243-2252.
3. Kluger MD, Salceda JA, Laurent A, et al.: Liver resection for hepatocellular carcinoma in 313 Western patients: tumor biology and underlying liver rather than tumor size drive prognosis. *J Hepatol* 2015;62:1131-1140.
4. Liver cancer study group of Japan (eds).: General rules for the clinical and pathological study of primary liver cancer. The 6th edition. Kanehara & CO., LTD., Tokyo. 2015.
5. Kamiyama T, Orimo T, Shimada S, et al.: Survival outcomes of hepatectomy for stage B Hepatocellular carcinoma in the BCLC classification. *World J Surg Oncol* 2017;15:156.
6. Wakayama K, Kamiyama T, Shimada S, et al.: Huge hepatocellular carcinoma greater than 10 cm in diameter worsens prognosis by causing distant recurrence after curative resection. *J Surg Oncol* 2017;115:324-329.
7. Matsushima H, Takami Y, Ryu T, et al.: Prognosis of hepatocellular carcinoma patients who achieved long-term recurrence-free survival after curative therapy: impact of the ALBI grade. *J Gastrointest Surg* 2018;22:1230-1238.
8. Kamiyama T, Nakanishi K, Yokoo H, et al.: Perioperative management of hepatic resection toward

- 1 zero mortality and morbidity: analysis of 793 consecutive cases in a single institution. *J Am Coll*
2 *Surg* 2010;211:443-449.
- 3 9. Clavien PA, Barkun J, de Oliveira ML, et al.: The Clavien-Dindo classification of surgical
4 complications: five-year experience. *Ann Surg* 2009;250:187-196.
- 5 10. Tsunoda T, Segawa T, Eto T, et al.: Long-term survivors after hepatectomy for hepatocellular
6 carcinoma. *J Gastroenterol Hepatol* 1990;5:595-600.
- 7 11. Shirabe K, Shimada M, Kajiyama K, et al.: Clinicopathologic features of patients with
8 hepatocellular carcinoma surviving > 10 years after hepatic resection. *Cancer* 1998;83:2312-2316.
- 9 12. Zhou XD, Tang ZY, Yang BH, et al.: Experience of 1000 patients who underwent hepatectomy
10 for small hepatocellular carcinoma. *Cancer* 2001;91:1479-1486.
- 11 13. Poon RT, Ng IO, Fan ST, et al.: Clinicopathological features of long-term survivors and disease-
12 free survivors after resection of hepatocellular carcinoma: a study of a prospective cohort. *J Clin*
13 *Oncol* 2001;19:3037-3044.
- 14 14. Bilimoria MM, Lauwers GY, Doherty DA, et al.: Underlying liver disease, not tumor factors,
15 predicts long-term survival after resection of hepatocellular carcinoma. *Arch Surg* 2001;136:528-535.
- 16 15. Shimada K, Sano T, Sakamoto Y, et al.: A long-term follow up and management study of
17 hepatocellular carcinoma patients surviving for 10 years or longer after curative hepatectomy. *Cancer*
18 2005;104:1939-1947.
- 19 16. Fukuda S, Itamoto T, Amano H, et al.: Clinicopathologic features of hepatocellular carcinoma
20 patients with compensated cirrhosis surviving more than 10 years after curative hepatectomy. *World*

- 1 J Surg 2007;31:345-352.
- 2 17. Zhou XD, Tang ZY, Ma ZC, et al.: Twenty-year survivors after resection for hepatocellular
3 carcinoma - analysis of 53 cases. J Cancer Res Clin Oncol 2009;135:1067-1072.
- 4 18. Eguchi S, Kanematsu T, Arii S, et al.: Recurrence-free survival more than 10 years after liver
5 resection for hepatocellular carcinoma. Br J Surg 2011;98:552-557.
- 6 19. Kim SH, Choi SB, Lee JG, et al.: Prognostic factors and 10-year survival in patients with
7 hepatocellular carcinoma after curative hepatectomy. J Gastrointest Surg 2011;15:598-607.
- 8 20. Wu KT, Wang CC, Lu LG, et al.: Hepatocellular carcinoma: clinical study of long-term survival
9 and choice of treatment modalities. World J Gastroenterol 2013;19:3649-3657.
- 10 21. Franssen B, Jibara G, Tabrizian P, et al.: Actual 10-year survival following hepatectomy for
11 hepatocellular carcinoma. HPB (Oxford) 2014;16:830-835.
- 12 22. Zheng J, Kuk D, Gonen M, et al.: Actual 10-year survivors after resection of hepatocellular
13 carcinoma. Ann Surg Oncol 2017;24:1358-1366.
- 14 23. Calvaruso V, Cabibbo G, Cacciola I, et al.: Incidence of hepatocellular carcinoma in patients with
15 HCV-associated cirrhosis treated with Direct-acting antiviral agents. Gastroenterology
16 2018;155:411-421.
- 17 24. Yamamoto J, Kosuge T, Takayama T, et al.: Recurrence of hepatocellular carcinoma after surgery.
18 Br J Surg 1996;83:1219-1222.
- 19 25. Ko S, Kanehiro H, Hisanaga M, et al.: Liver fibrosis increases the risk of intrahepatic recurrence
20 after hepatectomy for hepatocellular carcinoma. Br J Surg 2002;89:57-62.

- 1 26. Yoshida H, Shiratori Y, Moriyama M, et al.: Interferon therapy reduces the risk for hepatocellular
2 carcinoma: national surveillance program of cirrhotic and noncirrhotic patients with chronic
3 hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy.
4 Ann Intern Med 1999;131:174-181.
- 5 27. Nagasue N, Yu L, Yukaya H, et al.: Androgen and oestrogen receptors in hepatocellular
6 carcinoma and surrounding liver parenchyma: impact on intrahepatic recurrence after hepatic
7 resection. Br J Surg 1995;82:542-547.
- 8 28. De Maria N, Manno M, Villa E.: Sex hormones and liver cancer. Mol Cell Endocrinol
9 2002;193:59-63.
- 10 29. Yang LY, Wang W, Peng JX,et al.: Differentially expressed genes between solitary large
11 hepatocellular carcinoma and nodular hepatocellular carcinoma. World J Gastroenterol
12 2004;10:3569-3573.
- 13 30. Shimada S, Kamiyama T, Yokoo H, et al.: Clinicopathological characteristics of hepatocellular
14 carcinoma with microscopic portal venous invasion and the role of anatomical liver resection in
15 these cases. World J Surg 2017;41:2087-2094.
- 16 31. Shirabe K, Kanematsu T, Matsumata T,et al.: Factors linked to early recurrence of small
17 hepatocellular carcinoma after hepatectomy: univariate and multivariate analyses. Hepatology
18 1991;14:802-805.
- 19 32. Ramachandran J, Sajith KG, Priya S, et al.: Serum cholinesterase is an excellent biomarker of
20 liver cirrhosis. Trop gastroenterol 2014;35:15-20.

33. Kogo M, Kano A, Kiuchi Y, et al.: Prognostic index for survival in patients after treatment for primary hepatocellular carcinoma. *Dig Dis Sci* 2007;52:2444-2451.
34. Poon RT, Fan ST, Ng IO, et al.: Different risk factors and prognosis for early and late intrahepatic recurrence after resection of hepatocellular carcinoma. *Cancer* 2000;89:500-507.
35. Hao S, Fan P, Chen S, et al.: Distinct recurrence risk factors for intrahepatic metastasis and multicenter occurrence after surgery in patients with hepatocellular carcinoma. *J Gastrointest Surg* 2017; 21(2):312-320.
36. Minagawa M, Makuuchi M, Takayama T, et al.: Selection criteria for repeat hepatectomy in patients with recurrent hepatocellular carcinoma. *Ann Surg* 2003;238:703-710.
37. Belghiti J, Cortes A, Abdalla EK, et al.: Resection prior to liver transplantation for hepatocellular carcinoma. *Ann Surg* 2003;238:885-893.
38. Thomasset SC, Dennison AR, Garcea G.: Ablation for recurrent hepatocellular carcinoma: a systematic review of clinical efficacy and prognostic factors. *World J Surg* 2015;39:1150-1160.
39. Nakagawa T, Kamiyama T, Nakanishi K, et al.: Pulmonary resection for metastases from hepatocellular carcinoma: factors influencing prognosis. *J Thorac Cardiovasc Surg* 2006;131:1248-1254.

1 **Figure legends**

2 **Fig. 1**

3 Overall survival (OS) and Relapse-free survival (RFS) of this cohort.

4

5 **Fig. 2**

6 Recurrence rates and time after hepatectomy in the RS and RD groups.

7

Table 1. Clinicopathological characteristics of RF and RS+RD

(a) Clinicopathological characteristics by univariate analyses

Characteristics		RF	RS+RD	<i>p</i>
		(n = 35)	(n = 180)	
Epidemiology				
Age	< 60	16 (46 %)	86 (48 %)	0.82
	≥ 60	19 (54 %)	94 (52 %)	
Sex	male	25 (71 %)	154 (86 %)	0.04
	Female	10 (29 %)	26 (14 %)	
HBs-Ag	positive	20 (57 %)	84 (47 %)	0.25
	negative	15 (43 %)	96 (53 %)	
HCV-Ab	positive	12 (34 %)	48 (27 %)	0.35
	negative	23 (66 %)	132 (73 %)	
NBNC	yes	3 (9 %)	48 (27 %)	0.02
	No	32 (91 %)	132 (73 %)	
Biochemical Factors				
Platelets	< 80,000 /mm ³	0 (0 %)	19 (11 %)	0.04
	≥ 80,000 /mm ³	35 (100 %)	161 (89 %)	
Albumin	< 4.0 g/dl	8 (23 %)	77 (43 %)	0.02
	≥ 4.0 g/dl	27 (77 %)	103 (57 %)	
Total bilirubin	≥ 1.0 mg/dl	3 (9 %)	48 (27 %)	0.02
	< 1.0 mg/dl	32 (91 %)	132 (73 %)	
PT	< 80 %	5 (14 %)	48 (27 %)	0.11
	≥ 80 %	30 (86 %)	132 (73 %)	
ChE	< 200 IU/L	7 (20 %)	62 (34 %)	0.09
	≥ 200 IU/L	28 (80 %)	118 (66 %)	
ICGR15	≥ 15 %	8 (23 %)	91 (51 %)	<0.01
	< 15 %	27 (77 %)	89 (49 %)	
AFP	≥ 20 ng/ml	13 (37 %)	101 (56 %)	0.03
	< 20 ng/ml	22 (63 %)	79 (44 %)	
PIVKA-II	≥ 100 mAU/ml	17 (49 %)	105 (58 %)	0.28
	< 100 mAU/ml	18 (51 %)	75 (42 %)	
Tumor Factors				
Tumor size	≥ 4.5 cm	13 (37 %)	100 (56 %)	0.04
	< 4.5 cm	22 (63 %)	80 (44 %)	

Tumor number	multiple	4 (11 %)	82 (46 %)	<0.01
	solitary	31 (89 %)	98 (54 %)	
Macroscopic type	simple nodular	20 (57 %)	87 (48 %)	0.34
	others	15 (43 %)	93 (52 %)	
Differentiation	well	5 (14 %)	9 (5 %)	0.04
	others	30 (86 %)	171 (95 %)	
PVI	yes	5 (14 %)	56 (31 %)	0.04
	no	30 (86 %)	124 (69 %)	
HVI	yes	1 (3 %)	22 (12 %)	0.10
	no	34 (97 %)	158 (88 %)	
Underlying Liver				
Fibrosis	f3/4	9 (26 %)	107 (59 %)	<0.01
	f0-2	26 (74 %)	73 (41 %)	
Surgical Factors				
Anatomical resection	yes	31 (89 %)	131 (73 %)	0.04
	no	4 (11 %)	49 (27 %)	
Operative time	≥ 180 min	32 (91 %)	173 (96 %)	0.22
	< 180 min	3 (9 %)	7 (4 %)	
Blood loss	≥ 200 ml	21 (60 %)	139 (77 %)	0.03
	< 200 ml	14 (40 %)	41 (23 %)	

RF, Recurrence-free group; RS, Recurrence-survival group; RD, Recurrence-death group; HBs-Ag, HBs-antigen; HCV-Ab, HCV-antibody; NBNC, without HBV and HCV; PT, prothrombin time; ChE, cholinesterase; ICGR15, indocyanine green retention rate at 15 min; AFP, alpha-fetoprotein; PIVKA-II, protein induced by vitamin K absence-II; PVI, portal venous invasion; HVI, hepatic venous invasion; f3, bridging fibrosis; f4, cirrhosis.

Table 1. Clinicopathological characteristics of RF and RS+RD

(b) Clinicopathological characteristics by multivariate analyses

Characteristics	Odds ratio	95% CI	<i>P</i>
Sex Female	2.3688	0.6920-8.1081	0.16
Not NBNC	5.8812	1.6237-21.3019	<0.01
Platelets $\geq 80,000$ /mm ³	2.0707	0.1204-35.6142	0.61
Albumin ≥ 4.0 g/dl	1.7456	0.5993-5.0845	0.30
Total bilirubin < 1.0 mg/dl	3.6286	0.7536-17.4732	0.10
ICGR15 < 15 %	2.0019	0.6500-6.1657	0.22
AFP < 20 ng/ml	1.7616	0.6688-4.6403	0.25
Tumor size < 4.5 cm	1.0021	0.3457-2.9047	0.99
Solitary tumor	11.7775	2.6214-52.9132	<0.01
Well differentiation	18.6768	2.7686-125.9936	<0.01
PVI negative	1.4960	0.4180-5.3539	0.53
Fibrosis f3/4 negative	7.8509	2.7037-22.7969	<0.01
Anatomical resection	2.1250	0.4769-9.4690	0.32
Blood loss < 200 ml	1.6918	0.5858-4.8863	0.33

NBNC, without HBV and HCV; ICGR15, indocyanine green retention rate at 15 min; AFP, alpha-fetoprotein; PVI, portal venous invasion; f3, bridging fibrosis; f4, cirrhosis.

Table 2. Clinicopathological characteristics of RF+RS and RD

(a) Clinicopathological characteristics by univariate analyses

Characteristics		RF+RS	RD	<i>p</i>
		(n = 83)	(n = 132)	
Epidemiology				
Age	< 60	31 (37 %)	71 (54 %)	0.01
	≥ 60	52 (63 %)	61 (46 %)	
Sex	male	65 (78 %)	114 (86 %)	0.12
	Female	18 (22 %)	18 (14 %)	
HBs-Ag	positive	46 (55 %)	58 (44 %)	0.10
	negative	37 (45 %)	74 (56 %)	
HCV-Ab	positive	22 (27 %)	38 (29 %)	0.71
	negative	61 (73 %)	94 (71 %)	
NBNC	yes	15 (18 %)	36 (27 %)	0.12
	No	68 (82 %)	96 (73 %)	
Biochemical Factors				
Platelets	< 80,000 /mm ³	4 (5 %)	13 (10 %)	0.18
	≥ 80,000 /mm ³	79 (95 %)	119 (90 %)	
Albumin	< 4.0 g/dl	22 (27 %)	63 (48 %)	<0.01
	≥ 4.0 g/dl	61 (73 %)	69 (52 %)	
Total bilirubin	≥ 1.0 mg/dl	15 (18 %)	36 (27 %)	0.12
	< 1.0 mg/dl	68 (82 %)	96 (73 %)	
PT	< 80 %	19 (23 %)	34 (26 %)	0.63
	≥ 80 %	64 (77 %)	98 (74 %)	
ChE	< 200 IU/L	17 (20 %)	52 (39 %)	<0.01
	≥ 200 IU/L	66 (80 %)	80 (61 %)	
ICGR15	≥ 15 %	30 (36 %)	69 (52 %)	0.02
	< 15 %	53 (64 %)	63 (48 %)	
AFP	≥ 20 ng/ml	32 (39 %)	78 (59 %)	<0.01
	< 20 ng/ml	51 (61 %)	54 (41 %)	
PIVKA-II	≥ 100 mAU/ml	38 (46 %)	84 (64 %)	0.01
	< 100 mAU/ml	45 (54 %)	48 (36 %)	
Tumor Factors				
Tumor size	≥ 4.5 cm	25 (30 %)	86 (65 %)	<0.01
	< 4.5 cm	58 (70 %)	46 (35 %)	

Tumor number	multiple	19 (23 %)	67 (51 %)	<0.01
	solitary	64 (77 %)	65 (59 %)	
Macroscopic type	simple nodular	47 (57 %)	60 (45 %)	0.11
	others	36 (43 %)	72 (55 %)	
Differentiation	well	10 (12 %)	4 (3 %)	<0.01
	others	73 (88 %)	128 (97 %)	
PVI	yes	12 (14 %)	49 (37 %)	<0.01
	no	71 (86 %)	83 (63 %)	
HVI	yes	3 (4 %)	20 (15 %)	<0.01
	no	80 (96 %)	112 (85 %)	
Underlying Liver				
Fibrosis	f3/4	36 (43 %)	79 (60 %)	0.01
	f0-2	47 (57 %)	53 (40 %)	
Surgical Factors				
Anatomical resection	yes	68 (82 %)	94 (71 %)	0.07
	no	15 (18 %)	38 (29 %)	
Operative time	≥ 180 min	78 (94 %)	127 (96 %)	0.44
	< 180 min	5 (6 %)	5 (4 %)	
Blood loss	≥ 200 ml	52 (63 %)	108 (82 %)	<0.01
	< 200 ml	31 (37 %)	24 (18 %)	

RF, Recurrence-free group; RS, Recurrence-survival group; RD, Recurrence-death group; HBs-Ag, HBs-antigen; HCV-Ab, HCV-antibody; NBNC, without HBV and HCV; PT, prothrombin time; ChE, cholinesterase; ICGR15, indocyanine green retention rate at 15 min; AFP, alpha-fetoprotein; PIVKA-II, protein induced by vitamin K absence-II; PVI, portal venous invasion; HVI, hepatic venous invasion; f3, bridging fibrosis; f4, cirrhosis.

Table 2. Clinicopathological characteristics of RF+RS and RD

(b) Clinicopathological characteristics by multivariate analyses

Characteristics	Odds ratio	95% CI	<i>p</i>
Age $\geq$ 60	0.5016	0.2453-1.0255	0.05
ChE $\geq$ 200 IU/L	1.2705	0.5748-2.8083	0.55
Albumin $\geq$ 4.0 g/dl	1.1253	0.5261-2.4071	0.76
ICGR15 < 15 %	1.6285	0.7970-3.3276	0.18
AFP < 20 ng/ml	2.0013	1.0142-3.9493	0.04
PIVKA-II < 100 mAU/ml	0.8602	0.3933-1.8813	0.70
Tumor size < 4.5 cm	4.1825	1.8065-9.6837	<0.01
Solitary tumor	2.6579	1.2656-5.5820	<0.01
Well differentiation	4.7835	1.0637-21.5107	0.04
PVI negative	1.3650	0.5722-3.2562	0.48
HVI negative	2.3858	0.5700-9.9853	0.23
Fibrosis f3/f4	3.5454	1.5677-8.0179	<0.01
Blood loss < 200 ml	1.6677	0.7614-3.6526	0.20

ChE, cholinesterase; Alb, albumin; ICGR15, indocyanine green retention rate at 15 min; AFP, alpha-fetoprotein; PIVKA-II, protein induced by vitamin K absence-II; PVI, portal venous invasion; HVI, hepatic venous invasion; f3, bridging fibrosis; f4, cirrhosis.

Table 3. Clinicopathological characteristics of RS and RD

(a) Clinicopathological characteristics by univariate analyses

Characteristics		RS	RD	p
		(n = 48)	(n = 132)	
Epidemiology				
Age	< 60	15 (31 %)	71 (54 %)	<0.01
	≥ 60	33 (69 %)	61 (46 %)	
Sex	male	40 (83 %)	114 (86 %)	0.60
	female	8 (17 %)	18 (14 %)	
HBs-Ag	positive	26 (54 %)	58 (44 %)	0.22
	negative	22 (46 %)	74 (56 %)	
HCV-Ab	positive	10 (21 %)	38 (29 %)	0.28
	negative	38 (79 %)	94 (71 %)	
NBNC	yes	12 (25 %)	36 (27 %)	0.76
	No	36 (75 %)	96 (73 %)	
Biochemical Factors				
Platelets	< 80,000 /mm ³	4 (8 %)	13 (10 %)	0.75
	≥ 80,000 /mm ³	44 (92 %)	119 (90 %)	
Albumin	< 4.0 g/dl	14 (29 %)	63 (48 %)	0.02
	≥ 4.0 g/dl	34 (71 %)	69 (52 %)	
Total bilirubin	≥ 1.0 mg/dl	12 (25 %)	36 (27 %)	0.76
	< 1.0 mg/dl	36 (75 %)	96 (73 %)	
PT	< 80 %	14 (29 %)	34 (26 %)	0.64
	≥ 80 %	34 (71 %)	98 (74 %)	
ChE	< 200 IU/L	10 (21 %)	52 (39 %)	0.02
	≥ 200 IU/L	38 (79 %)	80 (61 %)	
ICGR15	≥ 15 %	22 (46 %)	69 (52 %)	0.44
	< 15 %	26 (54 %)	63 (48 %)	
AFP	≥ 20 ng/ml	19 (40 %)	78 (59 %)	0.02
	< 20 ng/ml	29 (60 %)	54 (41 %)	
PIVKA-II	≥ 100 mAU/ml	21 (44 %)	84 (64 %)	0.01
	< 100 mAU/ml	27 (56 %)	48 (36 %)	
Tumor Factors				
Tumor size	≥ 4.5 cm	12 (25 %)	86 (65 %)	<0.01
	< 4.5 cm	36 (75 %)	46 (35 %)	

Tumor number	multiple	15 (31 %)	67 (51 %)	0.02
	solitary	33 (69 %)	65 (49 %)	
Macroscopic type	simple nodular	27 (56 %)	60 (45 %)	0.20
	others	21 (44 %)	72 (55 %)	
Differentiation	well	5 (10 %)	4 (3 %)	0.04
	others	43 (90 %)	12 (97 %)	
PVI	yes	7 (15 %)	49 (37 %)	<0.01
	no	41 (85 %)	83 (63 %)	
HVI	yes	2 (4 %)	20 (15 %)	0.04
	no	46 (96 %)	112 (85 %)	
Underlying Liver				
Fibrosis	f3/4	28 (58 %)	79 (60 %)	0.85
	f0-2	20 (42 %)	53 (40 %)	
Surgical Factors				
Anatomical resection	yes	37 (77 %)	94 (71 %)	0.43
	no	11 (23 %)	38 (29 %)	
Operative time	≥ 180 min	46 (96 %)	127 (96 %)	0.90
	< 180 min	2 (4 %)	5 (4 %)	
Blood loss	≥ 200 ml	31 (65 %)	108 (82 %)	0.01
	< 200 ml	17 (35 %)	24 (18 %)	

RS, Recurrence-survival group; RD, Recurrence-death group; HBs-Ag, HBs-antigen; HCV-Ab, HCV-antibody; NBNC, without HBV and HCV; PT, prothrombin time; ChE, cholinesterase; ICGR15, indocyanine green retention rate at 15 min; AFP, alpha-fetoprotein; PIVKA-II, protein induced by vitamin K absence-II; PVI, portal venous invasion; HVI, hepatic venous invasion; f3, bridging fibrosis; f4, cirrhosis.

Table 3. Clinicopathological characteristics of RS and RD
(b) Clinicopathological characteristics by multivariate analyses

Characteristics	Odds ratio	95% CI	<i>p</i>
Age $\geq$ 60	0.5548	0.2555-1.2046	0.13
ChE $\geq$ 200 IU/L	3.3088	1.3460-8.1335	<0.01
Albumin $\geq$ 4.0 g/dl	1.1073	0.4585-2.6743	0.82
AFP < 20 ng/ml	2.0606	0.9039-4.6972	0.08
PIVKA-II < 100 mAU/ml	0.7171	0.2841-1.8103	0.48
Tumor size < 4.5 cm	3.1530	1.2952-7.6754	0.01
Solitary tumor	1.7216	0.7371-4.0207	0.20
Well differentiation	3.0987	0.5785-16.5992	0.18
PVI negative	3.6316	1.0789-12.2237	0.03
HVI negative	1.8456	0.3314-10.2790	0.48
Blood loss < 200 ml	1.4836	0.6035-3.6473	0.39

ChE, cholinesterase; AFP, alpha-fetoprotein; PIVKA-II, protein induced by vitamin K absence-II; PVI, portal venous invasion; HVI, hepatic venous invasion.

Table 4. Initial recurrence patterns and treatment after recurrence

	RS (n = 48)	RD (n = 132)	<i>p</i>
Recurrence site			
Liver only	39 (81 %)	46 (35 %)	<0.01
Extrahepatic	9 (19 %)	86 (65 %)	
Tumor number of liver recurrence alone	(n=39)	(n=46)	
Multiple	9 (23 %)	31 (67 %)	<0.01
Treatment after recurrence			
Re-hepatectomy	23 (48 %)	20 (15 %)	<0.01
Liver transplantation	5 (10 %)	3 (2 %)	0.01
Ablations	29 (60 %)	30 (23 %)	<0.01
TACE	27 (56 %)	88 (67 %)	0.19
Resection of metastases	7 (15 %)	23 (17 %)	0.79
Systemic chemotherapy including molecular target drug	9 (19 %)	56 (42 %)	<0.01
Radiotherapy	1 (2 %)	36 (27 %)	<0.01

RS, Recurrence-survival group; RD, Recurrence-death group; TACE, transarterial chemoembolization.

