HOKKAIDO UNIVERSITY

Title	Association between oral, social, and physical frailty in community -dwelling older adults
Author(s)	Hironaka, Sanae; Kugimiya, Yoshihiro; Watanabe, Yutaka et al.
Citation	Archives of gerontology and geriatrics, 89, 104105 https://doi.org/10.1016/j.archger.2020.104105
Issue Date	2020-07
Doc URL	https://hdl.handle.net/2115/82114
Rights	©2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Arch Gerontol Geriatr_2020_89_104105.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Association between oral, social, and physical frailty in community-dwelling older adults

Sanae Hironaka[a,b], Yoshihiro Kugimiya[c], Yutaka Watanabe[b,d], Keiko Motokawa[b], Hirohiko Hirano[b,e], Hisashi Kawai[f], Takeshi Kera[f,g], Motonaga Kojima[h], Yoshinori Fujiwara[i], Kazushige Ihara[j], Hunkyung Kim[b], Shuichi Obuchi[f], Yasuaki Kakinoki[a]

[a]Department of Special Needs and Geriatric Dentistry, Kyushu Dental University, Kitakyushu, Japan

[b]Research Team for Promoting Independence and Mental Health, Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

[c]Department of Removable Prosthodontics and Gerodontology, Tokyo Dental College, Tokyo, Japan

[d]Gerodontology, Department of Oral Health Science, Faculty of Dental Medicine, Hokkaido University, Sapporo, Japan

[e]Department of Oral Surgery and Dentistry, Tokyo Metropolitan Geriatric Hospital, Tokyo, Japan

[f]Research Team for Human Care, Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

[g]Department of Physical Therapy, Takasaki University of Health and Welfare, Takasaki, Japan

[h]Department of Physical Therapy, University of Tokyo Health Sciences, Tokyo, Japan

[i]Research Team for Social Participation and Community Health, Tokyo Metropolitan Institute of Gerontology, Tokyo, Japan

[j]Department of Social Medicine, Hirosaki University Graduate School of Medicine, Hirosaki City, Aomori, Japan

Corresponding author:

Yutaka Watanabe DDS, PhD

Associate Professor

Gerodontology, Department of Oral Health Science, Faculty of Dental Medicine,

Hokkaido University

Nishi-7, Kita-13, Kita-ku, Sapporo, Hokkaido 060-8586, Japan.

Phone & Fax: 81-11-706-4582

Email: ywata@den.hokudai.ac.jp

Research Team for Promoting Independence and Mental Health, Tokyo Metropolitan Institute of Gerontology

35-2 Sakae-cho, Itabashi-ku, Tokyo 173-0015, Japan.

Abstract

Objectives Oral frailty (OF) has been shown to be a predictor of disability. Therefore, it is important to be able to identify factors associated with OF in order to prevent long-term dependence. The purpose of this study was to clarify the relationships between OF, social frailty (SF), and physical frailty (PF) in community-dwelling older adults, with the overarching aim of informing the future development of effective measures to prevent frailty.

Methods Oral, physical, and social function, nutritional and psychological status, and medical history were examined in 682 community-dwelling individuals (267 men, 415 women) aged ≥ 65 years. Ordinal logistic regression analysis with SF and PF as independent variables was performed with pass analysis to determine the relationship between the different types of frailty.

Results Logistic regression analysis revealed significant associations between OF and decline in social function, physical function, and nutritional status, and an increase in the number of medications used. Path analysis showed that SF was directly related to OF and that OF and SF were directly related to PF.

Conclusions These findings suggest that a decline in social function may directly influence a decline in oral and physical function. The results of this study provide initial evidence, that may guide the future development of measures that aim to prevent and manage OF.

Key words: oral frailty, social frailty, physical frailty, path analysis, oral function, polypharmacy

Abbreviations list

GFI, goodness of fit index

IADL, instrumental activities of daily living

JST-IC, Japan Science and Technology Agency Index of Competence

LTCI, Japanese public long-term care insurance system

MMSE, Mini-Mental State Examination

MNA-SF, Mini Nutritional Assessment-Short Form

NFI, normed fit index

ODK, oral diadochokinesis

OF, oral frailty

PF, physical frailty

RMSEA, root mean square error of approximation

SF, social frailty

1. Introduction

Frailty is an important consideration in the development of strategies that aim to optimize a healthy life expectancy in the super-aged population of Japan. Frailty is defined as "a clinical state in which there is an increase in an individual's risk of dependence and/or mortality when exposed to a stressor that can be reduced by taking appropriate measures" (Morley et al., 2013). Risk factors for frailty include chronic disease (Danon-Hersch, Rodondi, Spagnoli, & Santos-Eggimann, 2012; Denfeld et al., 2017), poor nutritional status (Soysal et al., 2019), and polypharmacy (Saum et al., 2017). There is also evidence suggesting a relationship between frailty, number of teeth present, and oral function (Hakeem, Bernabé, & Sabbah, 2019; Tanaka et al., 2018).

Previous studies assessed oral function with only a single measure (e.g., tongue pressure or occlusal force) (Nakagawa et al., 2017; Tsuga, Maruyama, Yoshikawa, Yoshida, & Akagawa, 2011). Minakuchi et al. (2018) suggested that a multidimensional approach is necessary to evaluate oral function, which includes mastication, swallowing, and phonation. No recent studies have demonstrated that a combination of oral functional impairments can predict a decline in general function and disability.

Tanaka et al. (2018) coined the term "oral frailty" (OF) to recognize the multidimensional nature of poor oral health status. In their longitudinal study on community-dwelling older adults, they found that OF was a predictor of physical frailty (PF), dependency, and mortality.

Establishing management for OF appears to be important for the prevention of dependency. The relation factor of OF requires clarification before obtaining the basic information needed to establish appropriate management for OF. Moreover, there is an association between a decline in social function and decreases in subjective masticatory performance, number of teeth present, and tongue pressure (Mikami et al., 2019; Nagayoshi et al., 2017; Takeuchi, Aida, Kondo, & Osaka, 2013). Disability in social role functioning and intellectual activity were also identified as predictors of low performance in instrumental activities of daily living (IADL) (Fujiwara et al., 2003). This is of concern given that poor social relationships are recognized as a risk factor for mortality (Holt-Lunstad, Smith, & Layton, 2010). Makizako et al. (2015b) devised the term "social frailty" (SF) in a longitudinal study, that found a higher risk of disability in individuals with multiple indicators of social decline (e.g., living alone and decreased frequency of going out).

Although two longitudinal studies in community-dwelling older adults have confirmed that both, OF (Tanaka et al., 2018) and SF (Makizako et al., 2015b) negatively impact health status, the relationship between OF and SF remains unclear. In the meantime, the effect seems to be limited in the alone measure for frailty, which is a multifaceted concept. If the relationship between OF, SF, and PF, which is a predictor of dependency, is clarified, more multifaceted and effective correspondence may be possible.

We hypothesized that there would be associations between OF, SF, and PF. Therefore, in this study, we investigated the relationships between OF, SF, and PF in community-dwelling older adults using the definitions established for each type of frailty and a comprehensive range of variables, including nutritional status, sex, and medical history.

2. Materials and Methods

2.1 *Study Design*

This study had a cross-sectional design, and included subjects' data collected from comprehensive health checkups conducted annually by the Tokyo Metropolitan Institute of Gerontology (Tokyo, Japan) as a part of the Otasha Kenshin study initiated in 2011. The subjects in these health checkups were selected from a total of 7,162 older adults (aged 65–84 years) residing in nine towns in the Itabashi ward of Tokyo in 2011. Invitations for health checkups were sent to 6,699 older people, excluding those who were residing in long-term care institutions and those who had been subjects in previous research. In 2012, invitations for health checkups were sent to 7,015 older adults, excluding those who had reached 65 years of age or did not reside at the same address (according to the inhabitant registry) in the preceding fiscal year. Subsequent invitations for health checkups were sent in 2013–2018 to new study subjects, who had reached 65 years of age and to older adults who had participated in previous health checkups. In 2018, invitations were mailed to a total of 2,440 older adults, including 1,463 who had participated in previous health checkups (in 2011–2017) and 977 new study subjects, who had reached 65 years of age.

We excluded individuals receiving long-term care via the Japanese public long-term care insurance (LTCI) system and those with cognitive impairment (as indicated by a Mini-Mental State Examination [MMSE] score ≤ 18) to ensure the reliability of the subjective assessments, which included a self-administered questionnaire (Tanaka et al., 2018). We also excluded individuals with missing cognitive or oral function assessment data. All study investigators received training, that included measurement methods (using standardized evaluation criteria) and the use of evaluation instruments. The study protocol was approved by the Ethics Committee of the Tokyo Metropolitan Institute of Gerontology (Approval No. 48 in 2011; Approval No. 16 in 2018) and conducted in accordance with the Strengthening the Reporting of Observational Studies in Epidemiology statement.

2.2 *Measurements*

2.2.1 *Basic characteristics*

Information on sex, age, years of education, smoking status, and alcohol consumption was collected using a self-administered questionnaire. We used the Japan Science and Technology Agency Index of Competence (JST-IC) score to assess the ability to perform IADL (Iwasa et al., 2018).

2.2.2 *Nutritional status*

Any one of three registered dietitians evaluated the nutritional status of each subject. Body mass index was calculated from height and weight using the standard formula. Limb muscle mass was measured by the bioelectrical impedance method using an InBody 770 system (Biospace, Seoul, Korea). Skeletal muscle mass index was calculated by dividing the sum of limb muscle mass by the square of height in meters (Murakami et al., 2015). Nutritional status was assessed using the Mini Nutritional Assessment-Short Form (MNA-SF), on which a score of ≤ 11 is considered to indicate a risk of malnutrition (Soysal et al., 2019).

2.2.3 *Physical function*

Any one of four specialist researchers assessed physical function. Grip strength was measured using a Smedley-type hand dynamometer (Yagami, Nagoya, Japan) and gait speed by the 5-m walk test (Kera et al., 2018). Researchers asked subjects to stand and grip the hand dynamometer as tightly as possible using their preferred hand. Tests were carried out twice, and the higher value was selected. To evaluate subjects' gait, their usual walking speed was measured on a 5-m course, with 3-m acceleration and deceleration courses placed at each end of the courses. Researchers asked subjects to walk at their usual speed.

2.2.4 *Cognitive psychological function*

Any one of three nurses or two psychologists assessed cognitive function using the Japanese version of the MMSE (Folstein, Folstein, & McHugh, 1975; Sakuma et al., 2017). MMSE scores ≥ 19 were classified into three groups (≤ 24, moderate cognitive impairment; ≤ 27, mild cognitive impairment; and ≥ 28, preserved cognitive function) (Kaufer et al., 2008; Sakuma et al., 2017). Following the method devised by Fujisawa et al. (2005), the tendency for depression was investigated using a self-administered questionnaire containing the following five items: "Is your life pretty full? (last 2 weeks)" (yes); "Do you still enjoy doing the things you used to do? (last 2 weeks)" (yes); "Do you think it is too much trouble to do the things you used to do? (last 2 weeks)" (yes); "Do you feel that you are a useful person who is needed by others?" (yes); and "Do you feel tired without any specific reason? (last 2 weeks)" (yes). Subjects who met two or more criteria were considered to be depressed (Ihara et al., 2016).

2.2.5 *Medical history and number of medications*

Either of two nurses recorded the medical history and number of medications taken. Information was sought about the following diseases, each of which has been associated with PF: hypertension (Danon-Hersch et al., 2012), heart disease (Denfeld et al., 2017), diabetes mellitus (Danon-Hersch et al., 2012), hyperlipidemia (Ramsay et al., 2015), osteoporosis (Li et al., 2017), chronic obstructive pulmonary disease (Lee, Auyeung, Leung, Kwok, & Woo, 2014), osteoarthritis (Castell et al., 2015), spinal stenosis (Yagi et al., 2018), malignant neoplasm (Calado et al., 2016), anemia (Danon-Hersch, Rodondi, Spagnoli, & Santos-Eggimann, 2016), stroke (Calado, Ferriolli, Moriguti, Martinez, & Lima, 2016), chronic kidney disease (Walker et al., 2013), and fracture after 60 years of age (Kojima, 2016). Subjects were asked how many medications they were taking (0, 1, 2, 3, 4, or ≥ 5). The use of five or more medications related to PF was defined as polypharmacy (Saum et al., 2017).

2.2.6 *Oral function*

Any one of four dentists or four dental hygienists assessed oral function. The parameters measured were the number of teeth present, masticatory performance, tongue pressure, and oral diadochokinesis (ODK), which represents the dexterity of the tongue. The number of teeth present was defined as the number of

erupted teeth in the oral cavity, excluding retained roots and teeth with severe periodontitis. Masticatory performance was assessed using color-changeable chewing gum (Masticatory Performance Evaluating Gum Xylitol; Lotte Co., Ltd., Tokyo, Japan). The a* value for the gums after 1 minute of mastication was measured in three places using a color-difference meter (CR-20 Color Reader; Konica Minolta, Tokyo, Japan), with the mean value defined as the masticatory performance (Kugimiya et al., 2020). Tongue pressure was assessed using a JMS tongue pressure device (JMS Co., Ltd., Hiroshima, Japan). The tongue pressure probe was pinched between the tongue and palate, and tongue pressure was applied with maximal voluntary muscular effort for approximately 7 seconds. Three measurements were recorded, and the mean value was defined as the tongue pressure (Tsuga et al., 2011). ODK was assessed using an oral functions measurement device (Kenkokun Handy; Takei Scientific Instruments Co., Ltd., Niigata, Japan). The study subjects were asked to repeat each given syllable /ta/ sequentially as fast as possible for 5 seconds. The number of pronunciations per second was recorded (Watanabe et al., 2017).

2.3 *Definitions of frailty*

2.3.1 *Oral frailty*

Referencing the report by Tanaka et al. (2018), the cut-off values for the individual components were defined as follows: number of teeth present (< 20); masticatory performance (men, < 14.2; women, < 10.8); tongue pressure (men, < 27.4 kPa; women, < 26.5 kPa); and ODK (men, < 5.2 times/second; women, < 5.4 times/second). As a subjective evaluation, subjects who responded “yes” to "Do you have any difficulties eating tough foods compared to 6 months ago?" were considered to have decreased masticatory performance, and those who responded “yes” to "Have you choked on your tea or soup recently?" were considered to have decreased swallowing function. OF was evaluated according to the responses to six objective and subjective items, as previously described (Tanaka et al., 2018). Individuals who did not respond positively to any relevant items were regarded as robust. Those who responded positively to 1–2 items and ≥ 3 items were categorized as having as oral prefrailty (preOF) and OF, respectively.

2.3.2 *Social frailty*

SF was evaluated as in the report by Makizako et al. (2015b). Cut-offs were set for each of the five items as follows: “Do you go out less frequently compared with last year?” (yes), “Do you sometimes visit your friends?” (no), “Do you feel you are helpful to friends or family?” (no), “Do you live alone?” (yes), and “Do you talk with someone every day?” (no). Individuals who did not have any relevant items were regarded as robust. Those who had 1 item and ≥ 2 items were categorized as having social prefrailty (preSF) and SF, respectively.

2.3.3 *Physical frailty*

As in the report by Makizako, Shimada, Doi, Tsutsumimoto, and Suzuki (2015a), PF was assessed for each of the five categories with the following cut-off values: (1) unintentional weight loss, “Have you lost 2 kg or more in the past 6 months?” (yes); (2) exhaustion, “In the last 2 weeks, have you felt tired for no reason?”

(yes); (3) low activity, “Do you engage in moderate levels of physical exercise or sports aimed at health?” and “Do you engage in low levels of physical exercise aimed at health?” (subjects who answered “no” to both of these questions were classified as having low activity); (4) weakness, decreased grip strength (men, < 26.0 kg, women, < 18.0 kg), and (5) slowness, decreased gait (< 1.0 m/s). Individuals who did not have any relevant items were regarded as robust. Those who had 1–2 items and ≥ 3 items were categorized as having physical prefrailty (prePF) and PF, respectively.

2.4 *Statistical analysis*

Sex-related differences were analyzed using the Mann-Whitney *U* test for continuous variables and the chi-square test for categorical variables. Continuous variables were Bonferroni-corrected after the Jonckheere-Terpstra trend test to examine trends in the robust OF, preOF, and OF groups. Categorical variables were analyzed using the chi-square test. Ordinal logistic regression analyses were then performed, treating the three OF groups as objective variables to identify factors associated with OF. Currently, there are few reports on factors associated with OF, but there is a report on the association between OF and PF (Tanaka et al., 2018). Therefore, the factors related to PF were treated as independent variables. The number of medications was analyzed as a continuous variable with a cut-off value of five. Next, path analysis of the results of the ordinal logistic regression analysis was performed in order to examine the relationship between OF, SF, and PF. The fit of the models in the path analysis was evaluated using the goodness of fit index (GFI), normed fit index (NFI), and root mean square error of approximation (RMSEA). All statistical analyses were performed using SPSS Statistics version 23 and SPSS Amos version 23 (IBM Corp., Armonk, NY, USA) with a significance level of less than 5%.

3. Results

3.1 *Subject characteristics*

Figure 1 shows a flow chart outlining the process used to select subjects for this study. Invitations for health checks were mailed to a total of 2,440 older adults, of whom 769 (31.5%) eventually participated. Individuals with an MMSE score ≤ 18 (n=6, 0.2%), those receiving LTCI support (n=16, 0.7%), and those with missing cognitive or oral function data (n=65, 2.7%) were excluded, leaving 682 (27.9%) subjects (267 men, 415 women; mean age 73.3 ± 6.6 years) for inclusion in the analyses. The OF, PF, and SF rates are compared according to sex in Table 1. There was no significant sex-related difference in the OF or PF rate; however, SF was found to be more common in men. Table 2 shows the results for the study endpoints according to OF status. Three hundred and eighty (55.7%) of the 682 subjects had preOF and 65 (9.5%) had OF. Subjects with OF were older, were less independent in IADL, had lower physical and cognitive function, and tended to be depressed. Appendices A1-A3 show the respective correspondence rate for each item potentially associated with OF, SF, and PF.




Figure 1. Flow chart demonstrating the process used to select the study subjects.

LTCI, Japanese Public Long-Term Care Insurance System; MMSE, Mini-Mental State Examination.

Table 1. Characteristics of the study population

Variable	Overall (n = 682)		Men (n = 267)		Women (n = 415)		*P*-value
	Mean ± SD / n (%)	Median, [Q1, Q3]	Mean ± SD / n (%)	Median, [Q1, Q3]	Mean ± SD / n (%)	Median [Q1, Q3]	
Age, years	73.3 ± 6.6	72.0 [68.0, 78.0]	72.8 ± 6.6	71.0 [67.0, 78.0]	73.6 ± 6.5	73.0 [68.0, 79.0]	0.101
Education, years	13.4 ± 2.8	12.0 [12.0, 16.0]	14.2 ± 2.8	16.0 [12.0, 16.0]	12.9 ± 2.7	12.0 [12.0, 14.0]	<0.001
Current or past smoker, n (%)	248 (36.4)		183 (68.5)		65 (15.7)		<0.001
Drinks alcohol, n (%)	321 (47.1)		182 (68.2)		139 (33.5)		<0.001
JST-IC score	12.1 ± 2.8	12.0 [11.0, 14.0]	12.0 ± 2.9	12.0 [10.0, 14.0]	12.2 ± 2.7	13.0 [11.0, 14.0]	0.222
BMI	22.8 ± 3.3	22.7 [20.6, 24.7]	23.6 ± 3.2	23.3 [21.7, 25.4]	22.3 ± 3.4	22.1 [20.1, 24.2]	<0.001
SMI	6.5 ± 1.0	6.3 [5.7, 7.3]	7.5 ± 0.7	7.5 [7.0, 7.9]	5.9 ± 0.7	5.8 [5.4, 6.3]	<0.001
MNA-SF score	12.8 ± 1.3	13.0 [12.0, 14.0]	13.0 ± 1.3	13.0 [12.0, 14.0]	12.6 ± 1.3	13.0 [12.0, 14.0]	<0.001
≤ 11, n (%)	126 (18.5)		39 (14.6)		87 (21.0)		0.037
Handgrip strength, kg	27.9 ± 8.5	26.0 [22.0, 34.0]	35.5 ± 6.9	36.0 [31.0, 40.0]	22.9 ± 5.1	23.0 [19.0, 26.0]	<0.001

Gait speed, m/s	1.4 ± 0.3	1.4 [1.2, 1.6]	1.4 ± 0.3	1.4 [1.2, 1.6]	1.4 ± 0.3	1.4 [1.2, 1.6]	0.643
MMSE score	28.6 ± 1.6	29.0 [28.0, 30.0]	28.4 ± 1.7	29.0 [28.0, 30.0]	28.7 ± 1.5	29.0 [28.0, 30.0]	0.028
≤ 23, n (%)	10 (1.5)		5 (1.9)		5 (1.2)		
24−27, n (%)	126 (18.5)		57 (21.3)		69 (16.6)		0.218
≥ 28, n (%)	546 (80.1)		205 (76.8)		341 (82.2)		
Depression, n (%)	153 (22.4)		51 (19.1)		102 (24.6)		0.094
Oral function							
Present teeth, n	21.6 ± 7.9	24.0 [18.0, 27.0]	22.5 ± 7.3	25.0 [20.0, 27.0]	21.1 ± 8.2	24.0 [18.0, 27.0]	0.046
Masticatory performance	22.0 ± 4.9	22.8 [19.6, 25.4]	23.3 ± 4.2	23.9 [21.1, 26.3]	21.1 ± 5.1	22.16 [18.8, 24.8]	<0.001
Tongue pressure, kPa	33.0 ± 7.5	32.9 [28.1, 37.8]	33.7 ± 8.4	33.3 [27.8, 38.9]	32.6 ± 6.9	32.6 [28.3, 36.9]	0.109
ODK, times/s	6.4 ± 0.7	6.4 [6.0, 6.8]	6.4 ± 0.7	6.4 [6.0, 6.8]	6.4 ± 0.7	6.4 [6.0, 6.8]	0.853
Subjective difficulty in eating	147 (21.6)		42 (15.7)		105 (25.3)		0.003

Subjective difficulty in swallowing	182	(26.7)	67	(25.1)	115	(25.7)	0.451
Medical history							
Hypertension, n (%)	282	(41.3)	129	(48.3)	153	(36.9)	0.003
Heart disease, n (%)	109	[16.0)	51	(19.1)	58	(14.0)	0.075
Diabetes mellitus, n (%)	74	(10.9)	38	(14.2)	36	(8.7)	0.023
Hyperlipidemia, n (%)	256	(37.5)	90	(33.7)	166	(40.0)	0.098
Osteoporosis, n (%)	113	(16.6)	11	(4.1)	102	(24.6)	<0.001
Chronic obstructive pulmonary disease, n (%)	3	(0.4)	3	(1.1)	0	(0.0)	0.030
Osteoarthritis, n (%)	98	(14.4)	19	(7.1)	79	(19.0)	<0.001
Spinal stenosis, n (%)	68	(10.0)	25	(9.4)	43	(10.4)	0.671

Malignant neoplasm, n (%)	98	(14.4)	35	(13.1)	63	(15.2)	0.451
Anemia, n (%)	20	(2.9)	8	(3.0)	12	(2.9)	0.937
Stroke, n (%)	40	(5.9)	25	(9.4)	15	(3.6)	0.002
Chronic kidney disease, n (%)	3	(0.4)	1	(0.4)	2	(0.5)	0.836
Fracture, n (%)	122	(17.9)	31	(11.6)	91	(21.9)	0.001
Medications ≥ 5, n (%)	174	(25.5)	81	(30.3)	93	(22.4)	0.020
Oral frailty							
Robust, n (%)	237	(34.8)	100	(37.5)	137	(33.0)	
Prefrailty, n (%)	380	(55.7)	147	(55.1)	233	(56.1)	0.238
Frailty, n (%)	65	(9.5)	20	(7.5)	45	(10.8)	
Social frailty							
Robust, n (%)	287	(42.1)	123	(46.1)	164	(39.5)	0.001
Prefrailty, n (%)	237	(34.8)	71	(26.6)	166	(40.0)	

Frailty, n (%)	158	(23.2)	73	(27.3)	85	(20.5)	
Physical frailty							
Robust, n (%)	346	(50.7)	136	(50.9)	210	(50.6)	
Prefrailty, n (%)	312	(45.7)	122	(45.7)	190	(45.8)	0.984
Frailty, n (%)	24	(3.5)	9	(3.4)	15	(3.6)	

Categorical variables are shown as the number (percentage) and were analyzed using the chi-square test. Continuous variables are expressed as the mean and standard deviation and were analyzed using the Mann-Whitney *U* test. A *P*-value of < 0.05 was considered statistically significant. The number of subjects taking more than five medications was not documented; therefore, the number of subjects on polypharmacy are shown as a proportion. BMI, body mass index; JST-IC, Japan Science and Technology Agency Index of Competence; MMSE, Mini-Mental State Examination; MNA-SF, Mini Nutritional Assessment-Short Form; ODK, oral diadochokinesis; Q1, first quartile; Q3, third quartile; SD, standard deviation; SMI, skeletal muscle mass index

.

Table 2. Evaluation of items potentially affecting oral frailty status

Variable	Robust (n = 237)		preOF (n = 380)		OF (n = 65)		*P*-value
	Mean ± SD n (%)	Median [Q1, Q3]	Mean ± SD n (%)	Median [Q1, Q3]	Mean ± SD n (%)	Median [Q1, Q3]	
Age, years	71.1 ± 5.7	69.0 [67.0, 75.0]	74.0 ± 6.6	74.0 [68.0, 79.0]	77.1 ± 6.5	78.0 [73.0, 81.0]	<0.001
Men, n (%)	100 (42.2)		147 (38.7)		20 (30.8)		0.238
Education, years	13.9 ± 2.9	14.0 [12.0, 16.0]	13.2 ± 2.7	12.0 [12.0, 16.0]	12.5 ± 2.8	12.0 [10.5, 14.0]	<0.001
Current or past smoker, n (%)	72 (30.4)		150 (39.5)		26 (40.0)		0.060
Drinks alcohol, n (%)	123 (51.9)		182 (47.9)		16 (24.6)		<0.001
JST-IC score	12.8 ± 2.3	13.0 [11.0, 15.0]	12.0 ± 2.8	12.0 [10.0, 14.0]	10.6 ± 3.4	11.0 [7.5, 13.0]	<0.001
BMI	23.1 ± 3.4	23.0 [21.0, 24.9]	22.7 ± 3.3	22.6 [20.5, 24.5]	22.4 ± 3.4	22.3 [20.0, 24.7]	0.049
SMI	6.7 ± 1.1	6.6 [5.8, 7.5]	6.5 ± 1.0	6.3 [5.7, 7.3]	6.1 ± 0.9	5.9 [5.4, 6.6]	<0.001
MNA-SF score	13.0 ± 1.2	13.0 [12.0, 14.0]	12.7 ± 1.3	13.0 [12.0, 14.0]	12.4 ± 1.5	13.0 [11.0, 14.0]	0.002
≤ 11, n (%)	32 (13.5)		75 (19.7)		19 (29.2)		0.010

Handgrip strength, kg	29.7	±	8.6	28.0 [24.0, 37.0]	27.5	±	8.3	26.0 [22.0, 33.0]	23.5	±	7.4	24.0 [17.0, 27.0]	<0.001
Gait speed, m/s	1.4	±	0.2	1.5 [1.3, 1.6]	1.4	±	0.3	1.4 [1.2, 1.5]	1.2	±	0.3	1.3 [1.1, 1.4]	<0.001
MMSE score	28.9	±	1.4	29.0 [28.0, 30.0]	28.5	±	1.7	29.0 [28.0, 30.0]	28.0	±	1.9	29.0 [27.0, 29.0]	<0.001
≤23, n (%)	1		(0.4)		6		(1.6)		3		(4.6)		
24−27, n (%)	33		(13.9)		76		(20.0)		17		(26.2)		0.009
≥ 28, n (%)	203		(85.7)		298		(78.4)		45		(69.2)		
Depression, n (%)	30		(12.7)		95)		[25.0)		(28)		[43.1)		<0.001
Oral functions													
Present teeth, n	26.1	±	2.6	26.0 [24.0, 28.0]	20.5	±	8.1	23.0 [16.0, 27.0]	11.4	±	8.2	13.0 [4.0, 18.0]	—
Masticatory performance	23.8	±	3.0	24.2 [21.9, 26.0]	21.9	±	4.3	22.6 [19.2, 25.1]	15.4	±	7.4	16.8 [10.2, 21.6]	—
Tongue pressure, kPa	35.3	±	5.8	34.1 [31.1, 38.3]	32.3	±	7.7	32.1 [26.6, 37.6)	28.5	±	9.1	26.2 [21.4, 36.1]	—
ODK, times/s	6.6	±	0.6	6.6 [6.2, 7.0]	6.4	±	0.7	6.4 [6.0, 6.8]	5.9	±	1.0	6.2 [5.2, 6.6]	—
Subjective difficulty in eating	0		(0.0)		92		(24.2)		55		(84.6)		—

Subjective difficulty in swallowing	0	(0.0)	144	(37.9)	38	(58.5)	—
Medical history							
Hypertension, n (%)	92	(38.8)	160	(42.1)	30	(46.2)	0.513
Heart disease, n (%)	31	(13.1)	63	(16.6)	15	(23.1)	0.134
Diabetes mellitus, n (%)	21	(8.9)	43	(11.3)	10	(15.4)	0.296
Hyperlipidemia, n (%)	87	(36.7)	145	(38.2)	24	(36.9)	0.931
Osteoporosis, n (%)	26	(11.0)	74	(19.5)	13	(20.0)	0.016
Chronic obstructive pulmonary disease, n (%)	0	(0.0)	2	(0.5)	1	(1.5)	0.234
Osteoarthritis, n (%)	30	(12.7)	61	(16.1)	7	(10.8)	0.346
Spinal stenosis, n (%)	22	(9.3)	41	(10.8)	5	(7.7)	0.675

Malignant neoplasm, n (%)	22	(9.3)	70	(18.4)	6	(9.2)	0.003
Anemia, n (%)	5	(2.1)	10	(2.6)	5	(7.7)	0.053
Stroke, n (%)	8	(3.4)	21	(5.5)	11	(16.9)	<0.001
Chronic kidney diseases, n (%)	1	(0.4)	1	(0.3)	1	(1.5)	0.356
Fracture, n (%)	35	(14.8)	71	(18.7)	16	(24.6)	0.154
Medications ≥ 5, n (%)	36	(15.2)	111	(29.2)	27	(41.5)	<0.001
Social frailty							
Robust, n (%)	118	(49.8)	155	(40.8)	14	(21.5)	
Prefrailty, n (%)	84	(35.4)	131	(34.5)	22	(33.8)	<0.001
Frailty, n (%)	35	(14.8)	94	(24.7)	29	(44.6)	
Physical frailty							
Robust, n (%)	151	(63.7)	178	(46.8)	17	(26.2)	
Prefrailty, n (%)	82	(34.6)	188	(49.5)	42	(64.6)	<0.001

Frailty, n (%)	4	(1.7)		14	(3.7)		6	(9.2)

Categorical variables are shown as the number (percentage) and were analyzed using the chi-square test. A *P*-value < 0.05 was considered statistically significant. Continuous variables were analyzed with the Bonferroni-corrected Jonckheere-Terpstra test. A *P*-value < 0.006 was considered statistically significant. The Jonckheere-Terpstra test was not used because it was clear that oral functions decreased with OF. The number of subjects taking more than five medications was not documented; Therefore, the number of subjects on polypharmacy are shown as a proportion. BMI, body mass index; JST-IC, Japan Science and Technology Agency Index of Competence; MMSE, Mini-Mental State Examination; MNA-SF, Mini Nutritional Assessment-Short Form; ODK, oral diadochokinesis; Q1, first quartile; Q3, third quartile; SD, standard deviation; SMI, skeletal muscle mass index.

3.2 *Ordinal logistic regression analysis*

The results of the ordinal logistic regression analysis, which was performed to investigate the factors associated with OF, are shown in Table 3. There were significant associations with age (odds ratio [OR] 1.075, 95% confidence interval [CI] 1.045 – 1.104), MNA-SF score (OR 0.837, 95% CI 0.735 – 0.953), stroke (OR 2.423, 95% CI 1.197 – 4.899), number of medications used (OR 1.127, 95% CI 1.010 – 1.259), SF (OR 1.680, 95% CI 1.080 – 2.612), and prePF (OR 1.726, 95% CI 0.202 – 2.479).

Table 3. Ordinal logistic regression analyses

Variable	OR	95% CI	*P*-value
Age	1.075	1.045 – 1.104	< 0.001
Sex (0: men, 1: women)	0.797	0.555 – 1.145	0.219
Education	0.948	0.892 – 1.007	0.082
MNA-SF scores	0.837	0.735 – 0.953	0.007
MMSE scores	0.987	0.888 – 1.097	0.810
Depression (0: no, 1: yes)	1.508	0.970 – 2.347	0.068
Hypertension (0: no, 1: yes)	0.780	0.539 – 1.129	0.188
Heart disease (0: no, 1: yes)	1.021	0.653 – 1.597	0.927
Diabetes mellitus (0: no, 1: yes)	1.213	0.716 – 2.054	0.474
Hyperlipidemia (0: no, 1: yes)	0.868	0.614 – 1.228	0.425
Osteoporosis (0: no, 1: yes)	0.929	0.581 – 1.486	0.758
Osteoarthritis (0: no, 1: yes)	0.976	0.617 – 1.543	0.918
Spinal stenosis (0: no, 1: yes)	0.929	0.542 – 1.590	0.788
Malignant neoplasm (0: no, 1: yes)	1.408	0.899 – 2.208	0.136
Anemia (0: no, 1: yes)	1.451	0.572 – 3.680	0.433

Stroke (0: no, 1: yes)		2.423	1.197 – 4.899	0.014
Fracture (0: no, 1: yes)		1.010	0.662 – 1.540	0.963
Number of medications		1.127	1.010 – 1.259	0.033
SF	Robust	1.000（Reference）		
	preSF	1.138	0.790 – 1.639	0.488
	SF	1.680	1.080 – 2.612	0.021
PF	Robust	1.000（Reference）		
	prePF	1.726	1.202 – 2.479	0.003
	PF	1.347	0.546 – 3.327	0.519

MNA-SF and MMSE were analyzed as continuous variables. Age, MNA-SF, stroke, number of medications, SF, and prePF were significantly associated with OF. CI, confidence interval; MMSE, Mini-Mental State Examination; MNA-SF, Mini Nutritional Assessment-Short Form; OF, oral frailty; OR, odds ratio; PF, physical frailty; SF, social frailty

3.3 *Path analysis*

Path analysis is a statistical method used to analyze relationships between multiple factors. Figure 2 shows the path diagram derived by path analysis in this study. The best fit was selected among the path diagrams created by referring to the results of the ordinal logistic regression analysis. The pathway shows that the starting factor affected the stabbing factor. The diagram also shows that SF was directly related to OF and that the path coefficient from SF to OF was 0.141. Furthermore, both OF and SF were directly related to PF. The path coefficient from OF to PF was 0.155 and that from SF to PF was 0.206. All pathways were statistically significant. The GFI was 0.982, the NFI was 0.961, and the RMSEA was 0.026.




Figure 2. A pathway diagram demonstrating the relationship between OF and SF. MNA-SF (Mini Nutritional Assessment-Short Form), MMSE (Mini-Mental State Examination), OF (oral frailty), SF (social frailty), PF (physical frailty), and number of medications used were analyzed as categorical variables. Sex (0, male; 1, female), medications (0, ≤ 4; 1, ≥ 5), OF, SF, PF (0, robust; 1, prefrailty; 2, frailty), MNA-SF (0, ≥ 12; 1, ≤ 11), MMSE (0, ≥ 28; 1, 24 – 27; 2, ≤ 23). OF was directly related to PF, and SF was in the background of OF. SF was directly associated with PF, and SF was associated with PF via OF.

4. Discussion

Longitudinal studies have shown that OF, an indicator of decline in oral function, and SF, an indicator of decline in social function, are both predictors of dependency (Makizako et al., 2015b; Tanaka et al., 2018). A decline in oral and social function appears connected to the deterioration of an individual's general health status. However, previous studies have not clearly defined the relationships between OF, SF, and PF. If a relationship between the three frailties could be clarified, a more multifaceted and effective method for preventing dependency could be established. The results of the present study suggest that SF directly relates to OF and that both, OF and SF are directly related to PF.

The rate of PF in this study was 3.5%, which was lower than the prevalence of 7.4% reported by a meta-analysis of Japanese community-dwelling older adults aged 65 years or older (Kojima et al., 2017). In the present study, the mean JST-IC score for IADL was 12.1 ± 2.8, which is higher than the mean value of 9.5 previously reported for individuals aged 65 years or older (mean, 74.0 ± 2.8 years) in other regions of Japan (Iwasa et al., 2018). This study found that the ability to perform IADL was maintained; therefore, it is reasonable to assume that the subjects in this study maintained good physical function and that the rate of

PF was low. Furthermore, this study found that deterioration from robust oral function to OF tended to be more common in subjects who were older, less educated, had poor nutritional status, had poorer physical and cognitive function, tended to be depressed, and were on polypharmacy. These findings are similar to those reported by Tanaka et al. (2018).

In accordance with previous studies, this investigation defined three stages of OF, SF, and PF, comprising robust (no functional decline), prefrailty (mild functional decline), and frailty (accumulation of several types of functional decline) (Makizako et al., 2015a; Makizako et al., 2015b; Tanaka et al., 2018). These definitions reflect the association of a more severe stage with a greater risk of dependency.

Ordinal logistic regression analysis using the factors associated with PF as independent variables identified the factors associated with OF. Using path analysis, we examined how the factors associated with OF in the ordinal logistic regression analysis were associated with OF, SF, PF, and cognitive decline (Feng et al., 2017), which is an indicator of the psychological aspects of frailty. This study included a path analysis that could infer relationships between different types of frailty and clarify the factors associated with OF.

The logistic regression analysis showed significant associations of OF with age, MNA-SF scores, history of stroke, number of medications used, SF, and prePF.

The findings of the present study regarding the association between multidimensional aspects of oral and social function are consistent with those in several previous reports (Mikami et al., 2019; Nagayoshi et al., 2017; Nakanishi et al., 1999; Takeuchi et al., 2013).

There were reports of an association between decline in oral function and decline in physical function (Tanaka et al., 2018; Watanabe et al., 2017). Although the present study found no association between OF and PF, there was an association between prePF and OF. It was inferred that the lack of association between OF and PF in this study was due to the low PF rate of 3.5% (n = 24). A relationship between OF and the MNA-SF score was identified; this is consistent with the findings of previous studies on the association between a decline in oral function and decreased nutritional status (Kwon et al., 2017; Kwon, Suzuki, Kumagai, Shinkai, & Yukawa, 2006; Sura, Madhavan, Carnaby, & Crary, 2017).

Older adults often have multiple comorbidities and are likely to be on polypharmacy (Sirois et al., 2019). Xerostomia is one of the adverse effects of several medications commonly used in older adults and is associated with caries, deterioration of masticatory performance, and dysphagia (Barbe, 2018). The risk of hazardous events caused by an increase in the number of medications may be associated with a decline in oral function; however, this relationship is unclear, because no information regarding the adverse effects of medications was collected in this study. The mechanism of declining oral function and number of medications used or medication-related adverse events warrants further investigation in future studies.

Stroke has a variety of sequelae, including both, motor and sensory impairments (Winstein et al., 2016), and is responsible for considerable disability (Naruse, Sakai, Matsumoto, & Nagata, 2015). Patients requiring nursing care were excluded from this study, suggesting that the functional impairment attributable to stroke was mild. However, we demonstrated an association between stroke and OF; therefore, it may be

that various symptoms resulting from a history of stroke were related to a decline in oral function, even if they were mild.

The results of path analysis showed that SF was directly related to OF. Kobayashi, Wardle, & von Wagner (2015) reported an association between decline in social function and decreased health literacy. Preservation of health literacy is useful for health maintenance in daily life; however, a decline in health literacy leads to worsening health care and ability to self-manage disease (Sørensen et al., 2012). Oral diseases, such as caries and periodontal disease, are associated with lifestyle habits, including diet and hygiene (Petersen, 2008). Therefore, a lowering of general health literacy, which stems from a decline in social function, seems to be connected with a lowering of oral health literacy. Decreased oral health literacy can lead to poor oral hygiene, tooth loss, and even a decline in oral function. Furthermore, the opportunities for conversation are likely to decrease alongside a decline in social function. Decreased speaking may lead to reduced tongue pressure as a result of a decrease in the tongue movements involved in speech intelligibility. Furthermore, tongue pressure is associated with masticatory performance and speed of tongue movement (Kugimiya et al., 2019; Sagawa et al., 2019). These reports suggest that decreased opportunities for conversation lead to decreased tongue pressure, decreased masticatory performance, and slowing of tongue movement. Therefore, the results of this path analysis, which demonstrate the possibility of SF in the background of OF, seem to be appropriate. This study found that SF was directly associated with PF, which is in line with previous research (Makizako et al., 2019). Shimada et al. (2010) reported that loss of social roles and difficulty in responding to environmental changes reduces the frequency of activity by narrowing the range of behaviors in older adults. Narrowing of the range and frequency of activity decreases the amount of physical activity, and this is linked to PF. This study found that SF was not only directly related to PF, but also indirectly via OF. Moreover, a decline in oral function was identified to be one of the factors through which a decline in social function was connected to a decline in physical function. The process by which this occurs may involve not only a decrease in physical activity, but also a decline in oral function due to a reduction in the amount of interaction with others and decreased opportunities for conversation. The decline in social function may lead to a lowering of oral health literacy via a weakening of relationships with others, and in turn, cause a decline in oral function. Therefore, strategies to prevent or manage a decline in oral function should also focus on maintaining and improving social function. This will involve securing opportunities to engage with others, with the aim of maintaining oral health literacy.

Tanaka et al. (2018) reported that age and polypharmacy were directly related to OF. Barbe (2018) also reported an association of medication-induced xerostomia with caries, deterioration of masticatory performance, and dysphagia. Comorbidity and the long-term morbidity associated with several chronic diseases are also associated with frailty (Castrejón-Pérez, Gutiérrez-Robledo, Cesari, & Pérez-Zepeda, 2017). An accumulation of adverse events associated with chronic disease and polypharmacy and may be related to not only deterioration of general health status but also a decline in oral function.

This study also found a direct association between the MNA-SF score and OF. There are reports of an association between masticatory performance and vitamin and mineral deficiency (Kwon et al., 2017),

deterioration in subjective masticatory performance with a decline in diversity in the diet (Kwon et al., 2006), and dysphagia with malnutrition (Sura et al., 2017). A decline in oral function, such as masticatory performance and swallowing, may negatively impact the intake of nutrients. The results of this study are consistent with these reports. It is clear that the risk of malnutrition was associated with a transition from robust oral function to OF. Therefore, management of OF may prevent undernutrition.

The factors found to be directly related to OF in this study support the findings of previous reports. Furthermore, MNA-SF was directly associated with SF, suggesting that OF may lead to SF via malnutrition. The interaction between decreased social function and oral function may indeed be bidirectional, and should be investigated in future longitudinal and interventional studies.

Age, the MNA-SF score, and polypharmacy were directly related to SF. Makizako et al. (2015b) reported an association between age and SF and Tani et al. (2015) identified an association between unhealthy dietary behaviors and eating or living alone. Furthermore, given that use of five or more medications was identified as a risk factor for falls (Kojima et al., 2012) and that falls are associated with social isolation (Pohl, Cochrane, Schepp, & Woods, 2018), the relationship between polypharmacy and SF is likely to be consistent. The relationship between SF and its background factors is in line with previous reports, and the results in this regard also support the validity of this study.

Polypharmacy and the MMSE score were directly related to PF in this investigation; this is consistent with previous investigations that identified associations between polypharmacy and PF (Saum et al., 2017) and between cognitive decline and PF (Feng et al., 2017). Furthermore, the factors identified to be in the background of PF in this study are consistent with previous reports. Age, the MNA-SF score, and polypharmacy were directly related to the MMSE score, supporting previous reports, which identified an association between cognitive decline and aging (Sakuma et al., 2017), malnutrition (Ozawa et al., 2013), and polypharmacy (Niikawa et al., 2017). This study also identified that aging was directly related to polypharmacy, an association that has also been reported before (Saum et al., 2017).

The path model used in this investigation had a high level of fit, and all pathways showed a relationship between OF, SF, and PF, as in previous research. The congruence between these results and those in previous reports confirm the generalizability of these findings.

There are several limitations to this study. First, the associations between decline in oral function and cognitive decline (Kugimiya et al., 2019; Watanabe et al., 2018), social relationships and risk of dementia (Kuiper et al., 2015), and dietary patterns and risk of dementia (Ozawa et al., 2013) have already been reported. Although the path analysis showed a relationship between the MNA-SF and MMSE scores, there was no significant pathway between OF or SF and the MMSE score. The path analysis also indicated a circular relationship between OF, SF, and the MNA-SF score, suggesting that a deterioration in nutritional status was involved in the process which decreased oral and social function lead to cognitive decline. OF, SF, malnutrition, and cognitive decline are associated with PF and dependency (Feng et al., 2017; Makizako et al., 2015b; Soysal et al., 2019; Tanaka et al., 2018). The fact that the relationships between these four factors could be inferred in this study is one of the important grounds for investigating effective measures

for the multifaceted concept of frailty in the future. These relationships could not be determined in this study because of its cross-sectional design; therefore, longitudinal and intervention studies are needed. Second, although Makizako et al. (2015a) have reported an association between age and PF, there was no significant pathway between these two variables in this study, possibly because the older subjects had already participated in previous health checkups and maintained their level of physical function over the course of the study. Third, although Soysal et al. (2019) reported an association between malnutrition and PF, we could not identify a significant pathway between the MNA-SF score and PF in this study. The MNA-SF score was indirectly associated with PF via SF, and the MMSE score but was not directly associated with PF. However, there seemed to be a connection between malnutrition and a decline in physical function via declines in social and cognitive function. Therefore, it will be necessary to consider decreased social function and cognition when the relationships between malnutrition and decline in physical function are examined in the future. Fourth, this study only included older individuals who were able to independently attend for assessment and excluded those with severe deterioration of physical and cognitive function; this could have introduced a degree of selection bias. Therefore, the results of this study cannot be generalized to persons with severe psychosomatic impairment. Finally, the cross-sectional nature of the study meant that no causal relationships could be identified; however, the path model used in the study had a high level of fit. All OF, SF, and PF pathways were significant, as in previous reports, and are generalizable to older independent adult populations. This study suggests that the relationship between OF, SF, and PF would lead to a general deterioration in health. This may provide an important perspective for future research on frailty.

The logistic regression analysis showed that the factors associated with OF were a decline in social function, a decline in physical function, decreased nutritional status, and an increased number of medications used. These are known frailty-related factors (Makizako et al., 2015b; Saum et al., 2017; Soysal et al., 2019). It is important to detect OF at an early stage and use a multifaceted approach to maintain general health. The results of the path analysis highlight the possibility that SF is connected to OF. Therefore, the development of comprehensive and effective methods to manage declines in oral function will also need to consider evaluation of social function and its role in the maintenance of oral health literacy.

5. Conclusion

In this study, OF, SF, and PF were found to be mutually related in community-dwelling older adults. SF was directly related to OF, and OF in turn was directly related to PF. A decline in social function may contribute to decreased oral and physical function. This implies that optimal management of OF will also require evaluation and management of SF. These associations will be important considerations when devising appropriate measures for management of OF.

Funding

This work was supported by JSPS KAKENHI (grant numbers: 16K01853 and 18K18462), 2018 Health and Labor Sciences Research Grants (46–35), and 2018 Research Funding for Longevity Sciences (28–30

and 29–42) from the National Center for Geriatrics and Gerontology (NCGG), Japan.

Declarations of interest

None.

Acknowledgments

We express our gratitude to the local authorities responsible for the Itabashi Ward for their support and thank everyone who participated in the study. We would also like to thank the members of the Otasha Kenshin study staff for their cooperation.

References

Barbe, A. G. (2018). Medication-induced xerostomia and hyposalivation in the elderly: Culprits, complications, and management. *Drugs & Aging*, *35*, 877–885. https://doi.org/10.1007/s40266-018-0588-5.

Calado, L. B., Ferriolli, E., Moriguti, J. C., Martinez, E. Z., & Lima, N. K. (2016). Frailty syndrome in an independent urban population in Brazil (FIBRA study): A cross-sectional populational study. *Sao Paulo Medical Journal, 134*, 385–392. https://doi.org/10.1590/1516-3180.2016.0078180516.

Castell, M. V., van der Pas, S., Otero, A., Siviero, P., Dennison, E., Denkinger, M., et al. (2015). Osteoarthritis and frailty in elderly individuals across six European countries: Results from the European Project on OSteoArthritis (EPOSA). *BMC Musculoskeletal Disorders*, *16*, 359. https://doi.org/10.1186/s12891-015-0807-8.

Castrejón-Pérez, R. C., Gutiérrez-Robledo, L. M, Cesari, M., & Pérez-Zepeda, M. U. (2017). Diabetes mellitus, hypertension and frailty: A population-based, cross-sectional study of Mexican older adults. *Geriatrics & Gerontology International*, *17*, 925–930. https://doi.org/10.1111/ggi.12805.

Danon-Hersch, N., Rodondi, N., Spagnoli, J., & Santos-Eggimann, B. (2012). Prefrailty and chronic morbidity in the youngest old: An insight from the Lausanne Cohort Lc65+. *Journal of the American Geriatrics Society*, *60*, 1687–1694. https://doi.org/10.1111/j.1532-5415.2012.04113.x.

Denfeld, Q. E., Winters-Stone, K., Mudd, J. O., Gelow, J. M., Kurdi S., & Lee, C. S. (2017). The prevalence of frailty in heart failure: a systematic review. *International Journal of Cardiology*, *236*, 283–289. https://doi.org/10.1016/j.ijcard.2017.01.153.

Feng, L., Nyunt, M. S., Gao, Q., Feng, L., Lee, T. S., Tsoi, T., et al. (2017). Physical frailty, cognitive impairment, and the risk of neurocognitive disorder in the Singapore longitudinal ageing studies. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences, 72*, 369–375. https://doi.org/10.1093/gerona/glw050.

Fhon, J. R., Rodrigues, R. A., Neira, W. F., Huayta, V. M., & Robazzi, M. L. (2016). Fall and its association with the frailty syndrome in the elderly: Systematic review with meta-analysis. *University of São*

Paulo Nursing School Journal, *50*, 1005–1013. https://doi.org/10.1590/S0080-623420160000700018.

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, *12*, 189–198. https://doi.org/10.1016/0022-3956(75)90026-6.

Fujisawa, D., Tanaka, E., Sakamoto, S., Neichi, K., Nakagawa, A., & Ono, Y. (2005). The development of a brief screening instrument for depression and suicidal ideation for elderly: The Depression and Suicide Screen. *Psychiatry and Clinical Neurosciences*, *59*, 634–638. https://doi.org/ 10.1111/j.1440-1819.2005.01429.x.

Fujiwara, Y., Shinkai, S., Kumagai, S., Amano, H., Yoshida, Y., Yoshida, H., et al. (2003). Longitudinal changes in higher-level functional capacity of an older population living in a Japanese urban community. *Archives of Gerontology and Geriatrics*, *36*, 141–153. https://doi.org/10.1016/s0167-4943(02)00081-x.

Hakeem, F. F., Bernabé, E., & Sabbah, W. (2019). Association between oral health and frailty: A systematic review of longitudinal studies. *Gerodontology*, *36*, 205–215. https://doi.org/10.1111/ger.12406.

Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine*, *7*, e1000316. https://doi.org/10.1371/journal.pmed.1000316.

Ihara, K., Yoshida, H., Jones, P. B., Hashizume, M., Suzuki, Y., Ishijima, H., et al. (2016). Serum BDNF levels before and after the development of mood disorders: A case–control study in a population cohort. *Translational Psychiatry*, *6*, e782. https://doi.org/10.1038/tp.2016.47.

Iwasa, H., Masui, Y., Inagaki, H., Yoshida, Y., Shimada, H., Otsuka, R., et al. (2018). Assessing competence at a higher level among older adults: Development of the Japan Science and Technology Agency Index of Competence (JST-IC). *Aging Clinical and Experimental Research*, *30*, 383–393. https://doi.org/10.1007/s40520-017-0786-8.

Kaufer, D. I., Williams, C. S., Braaten, A. J., Gill, K., Zimmerman, S., & Sloane, P. D. (2008). Cognitive screening for dementia and mild cognitive impairment in assisted living: comparison of 3 tests. *Journal of the American Directors Association*, *9*, 586–593. https://doi.org/10.1016/j.jamda.2008.05.006.

Kera, T., Kawai, H., Hirano, H., Kojima, M., Watanabe, Y., Fujiwara, Y., et al. (2018). Comparison of body composition and physical and cognitive function between older Japanese adults with no diabetes, prediabetes and diabetes: A cross-sectional study in community-dwelling Japanese older people. *Geriatrics & Gerontology International*, *18*, 1031–1037. https://doi.org/10.1111/ggi.13301.

Kobayashi, L. C., Wardle, J., & von Wagner, C. (2015). Internet use, social engagement and health literacy decline during ageing in a longitudinal cohort of older English adults. *Journal of Epidemiology and Community Health*, *69*, 278–283. https://doi.org/10.1136/jech-2014-204733.

Kojima, T., Akishita, M., Nakamura, T., Nomura, K., Ogawa, S., Iijima, K., et al. (2012). Polypharmacy as a risk for fall occurrence in geriatric outpatients. *Geriatrics & Gerontology International*, *12*, 425–430. https://doi.org/10.1111/j.1447-0594.2011.00783.x.

Kojima, G. (2016). Frailty as a predictor of fractures among community-dwelling older people: A systematic review and meta-analysis. *Bone*, *90*, 116–122. https://doi.org/10.1016/j.bone.2016.06.009.

Kojima, G., Iliffe, S., Taniguchi, Y., Shimada, H., Rakugi, H., & Walters, K. (2017). Prevalence of frailty in Japan: A systematic review and meta-analysis. *Journal of Epidemiology*, *27*, 347–353. https://doi.org/10.1016/j.je.2016.09.008.

Kugimiya, Y., Ueda, T., Watanabe, Y., Takano, T., Edahiro, A., Awata, S., et al. (2019). Relationship between mild cognitive decline and oral motor functions in metropolitan community-dwelling older Japanese: The Takashimadaira Study. *Archives of Gerontology and Geriatrics*, *81*, 53–58. https://doi.org/10.1016/j.archger.2018.11.008.

Kugimiya, Y., Watanabe Y., Igarashi, K., Hoshino, D., Motokawa, K., Edahiro, A., et al. (2020). Factors associated with masticatory performance in community-dwelling older adults: A cross-sectional study. *Journal of the American Dental Association. 151*, 118–126. https://doi.org/10.1016/j.adaj.2019.10.003

Kuiper, J. S., Zuidersma, M., Oude Voshaar, R. C., Zuidema, S. U., van den Heuvel, E. R., Stolk, R. P., et al. (2015). Social relationships and risk of dementia: A systematic review and meta-analysis of longitudinal cohort studies. *Ageing Research Reviews*, *22*, 39–57. https://doi.org/10.1016/j.arr.2015.04.006.

Kwon, J., Suzuki, T., Kumagai, S., Shinkai, S., & Yukawa, H. (2006). Risk factors for dietary variety decline among Japanese elderly in a rural community: A 8-year follow-up study from TMIG-LISA. *European Journal of Clinical Nutrition*, *30*, 305–311. https://doi.org/10.1038/sj.ejcn.1602314.

Kwon, S. H., Park, H. R., Lee, Y. M., Kwon, S. Y., Kim, O. S., Kim, H. Y., & Lim, Y. S. (2017). Difference in food and nutrient intakes in Korean elderly people according to chewing difficulty: Using data from the Korea National Health and Nutrition Examination Survey 2013 (6th). *Nutrition Research and Practice*, *11*, 139–146. https://doi.org/10.4162/nrp.2017.11.2.139.

Lee, J. S., Auyeung, T. W., Leung, J., Kwok, T., & Woo, J. (2014). Transitions in frailty states among community-living older adults and their associated factors. *Journal of the American Directors Association*, *15*, 281–286. https://doi.org/10.1016/j.jamda.2013.12.002.

Li, G., Thabane, L., Papaioannou, A., Ioannidis, G., Levine, M. A., & Adachi, J. D. (2017). An overview of osteoporosis and frailty in the elderly. *BMC Musculoskeletal Disorders*, 18, 46. https://doi.org/10.1186/s12891-017-1403-x.

Makizako, H., Shimada, H., Doi, T., Tsutsumimoto, K., & Suzuki, T. (2015a). Impact of physical frailty on disability in community-dwelling older adults: a prospective cohort study. *BMJ Open*, *5*, e008462. https://doi.org/10.1136/bmjopen-2015-008462.

Makizako, H., Shimada, H., Tsutsumimoto, K., Lee, S., Doi, T., Nakakubo, S., et al. (2015b). Social frailty in community-dwelling older adults as a risk factor for disability. *Journal of the American Directors Association*, *16*, 1003.e7–11. https://doi.org/10.1016/j.jamda.2015.08.023.

Makizako, H., Kubozono, T., Kiyama, R., Takenaka, T., Kuwahata, S., Tabira, T., et al. (2019). Associations of social frailty with loss of muscle mass and muscle weakness among community-dwelling older adults. *Geriatrics & Gerontology International*, *19*, 76–80. https://doi.org/10.1111/ggi.13571.

Mikami, Y., Watanabe, Y., Motokawa, K., Shirobe, M., Motohashi, Y., Edahiro, A., et al. (2019). Association between decrease in frequency of going out and oral function in older adults living in major urban areas. *Geriatrics & Gerontology International*, *19*, 792–797. https://doi.org/10.1111/ggi.13715.

Minakuchi, S., Tsuga, K., Ikebe, K., Ueda, T., Tamura, F., Nagao, K, et al. (2018). Oral hypofunction in the population: Position paper of the Japanese Society of Gerodontology in 2016. *Gerodontology*, *35*, 317–324. https://doi.org/10.1111/ger.12347.

Morley, J. E., Vellas, B., van Kan, G. A., Anker, S. D., Bauer, J. M., Bernabei, R., et al. (2013). Frailty consensus: A call to action. *Journal of the American Directors Association*. *14*, 392–397. https://doi.org/10.1016/j.jamda.2013.03.022.

Murakami, M., Hirano, H., Watanabe, Y., Sakai, K., Kim, H., & Katakura, A. (2015). Relationship between chewing ability and sarcopenia in Japanese community-dwelling older adults. *Geriatrics & Gerontology International*, *15*, 1007–1012. https://doi.org/10.1111/ggi.12399.

Nagayoshi, M., Higashi, M., Takamura, N., Tamai, M., Koyamatsu, J., Yamanashi, H., et al. (2017). Social networks, leisure activities and maximum tongue pressure: Cross-sectional associations in the Nagasaki Islands Study. *BMJ Open*, *7*, e014878. https://doi.org/10.1136/bmjopen-2016-014878.

Nakagawa, K., Matsuo, K., Takagi, D., Morita, Y., Ooka, T., Hironaka, S., & Mukai, Y. (2017). Effects of gum chewing exercises on saliva secretion and occlusal force in community-dwelling elderly individuals: A pilot study. *Geriatrics & Gerontology International*, *17*, 48–53. https://doi.org/10.1111/ggi.12670.

Nakanishi, N., Hino, Y., Ida, O., Fukuda, H., Shinsho, F., & Tatara, K. (1999). Associations between self-assessed masticatory disability and health of community-residing elderly people. *Community Dentistry and Oral Epidemiology*, *27*, 366–371. https://doi.org/10.1111/j.1600-0528.1999.tb02033.x.

Naruse, T., Sakai, M., Matsumoto, H., & Nagata, S. (2015). Diseases that precede disability among latter-stage elderly individuals in Japan. *Bioscience Trends*, *9*, 270–274.https://doi.org/10.5582/bst.2015.01059.

Niikawa, H., Okamura, T., Ito, K., Ura, C., Miyamae, F., Sakuma, N., et al. (2017). Association between polypharmacy and cognitive impairment in an elderly Japanese population residing in an urban community. *Geriatrics & Gerontology International*, *17*, 1286–1293. https://doi.org/10.1111/ggi.12862.

Ozawa, M., Ninomiya, T., Ohara, T., Doi, Y., Uchida, K., Shirota, T., et al. (2013). Dietary patterns and risk of dementia in an elderly Japanese population: The Hisayama Study. *American Journal of Clinical Nutrition*, *97*, 1076–1082. https://doi.org/10.3945/ajcn.112.045575.

Petersen, P. E. (2008). World Health Organization global policy for improvement of oral health – World Health Assembly 2007. *International Dental Journal*, *58*, 115–121. https://doi.org/10.1111/j.1875-595x.2008.tb00185.x.

Pohl, J. S., Cochrane, B. B., Schepp, K. G., Woods, N. F. (2018). Falls and social isolation of older adults in the National Health and Aging Trends Study. *Research in Gerontological Nursing*, *11*, 61–70. https://doi.org/10.3928/19404921-20180216-02.

Ramsay, S. E., Arianayagam, D. S., Whincup, P. H., Lennon, L. T., Cryer, J., Papacosta, A. O., et al. (2015). Cardiovascular risk profile and frailty in a population-based study of older British men. *Heart*, 101, 616–622. https://doi.org/10.1136/heartjnl-2014-306472.

Sagawa, K., Furuya, H., Ohara, Y., Yoshida, M., Hirano, H., Iijima, K., & Kikutani, T. (2019). Tongue function is important for masticatory performance in the healthy elderly: A cross-sectional survey of community-dwelling elderly. *Journal of Prosthodontic Research*, *63*, 31–34. https://doi.org/10.1016/j.jpor.2018.03.006.

Sakuma, N., Ura, C., Miyamae, F., Inagaki, H., Ito, K., Niikawa, H., et al. (2017). Distribution of Mini-Mental State Examination scores among urban community-dwelling older adults in Japan. *International Journal of Geriatric Psychiatry*, *32*, 718–725. https://doi.org/10.1002/gps.4513.

Saum, K. U., Schöttker, B., Meid, A. D., Holleczek, B., Haefeli, W. E., Hauer, K., & Brenner, H. (2017). Is polypharmacy associated with frailty in older people? Results From the ESTHER Cohort Study. *Journal of the American Geriatrics Society*, *65*, e27–e32. https://doi.org/10.1111/jgs.14718.

Shimada, H., Ishizaki, T., Kato, M., Morimoto, A., Tamate, A., Uchiyama, Y., & Yasumura, S. (2010). How often and how far do frail elderly people need to go outdoors to maintain functional capacity? *Archives of Gerontology and Geriatrics*, *50*, 140–146. https://doi.org/10.1016/j.archger.2009.02.015.

Sirois, C., Domingues, N. S., Laroche, M. L., Zongo, A., Lunghi, C., Guénette, L., et al. (2019). Polypharmacy definitions for multimorbid older adults need stronger foundations to guide research, clinical practice and public health. *Pharmacy (Basel)*, *29*, 7(3). pii: E126. https://doi.org/10.3390/pharmacy7030126.

Sørensen, K., Van den Broucke, S., Fullam, J., Doyle, G., Pelikan, J., Slonska, Z., & Brand, H; (HLS-EU) Consortium Health Literacy Project European. (2012). Consortium Health Literacy Project European. Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health*, *12*, 80. https://doi.org/10.1186/1471-2458-12-80.

Soysal, P., Veronese, N., Arik, F., Kalan, U., Smith, L., & Isik, A. T. (2019). Mini Nutritional Assessment Scale-Short Form can be useful for frailty screening in older adults. *Clinical Interventions in Aging*, *14*, 693–699. https://doi.org/10.2147/CIA.S196770.

Sura, L., Madhavan, A., Carnaby, G., & Crary, M. A. (2012). Dysphagia in the elderly: Management and nutritional considerations. *Clinical Interventions in Aging*, *7*, 287–298. https://doi.org/10.2147/CIA.S23404.

Takeuchi, K., Aida, J., Kondo, K., & Osaka, K. (2013). Social participation and dental health status among older Japanese adults: A population-based cross-sectional study. *PLoS One*, *8*, e61741. https://doi.org/10.1371/journal.pone.0061741.

Tanaka, T., Takahashi, K., Hirano, H., Kikutani, T., Watanabe, Y., Ohara, Y., et al. (2018). Oral frailty as a risk factor for physical frailty and mortality in community-dwelling elderly. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, *73*, 1661–1667.https://doi.org/10.1093/gerona/glx225.

Tani, Y., Kondo, N., Takagi, D., Saito, M., Hikichi, H., Ojima, T., et al. (2015). Combined effects of eating alone and living alone on unhealthy dietary behaviors, obesity and underweight in older Japanese adults: Results of the JAGES. *Appetite*, *95*, 1–8. https://doi.org/10.1016/j.appet.2015.06.005.

Tsuga, K., Maruyama, M., Yoshikawa, M., Yoshida, M., & Akagawa, Y. (2011). Manometric evaluation of oral function with a hand-held balloon probe. *Journal of Oral Rehabilitation*, *38*, 680–685. https://doi.org/10.1111/j.1365-2842.2011.02202.x.

Walker, S. R., Gill, K., Macdonald, K., Komenda, P., Rigatto, C., Sood, M. M., et al. (2013). Association of frailty and physical function in patients with non-dialysis CKD: A systematic review. *BMC Nephrology*, *14*, 228. https://doi.org/10.1186/1471-2369-14-228.

Watanabe, Y., Hirano, H., Arai, H., Morishita, S., Ohara, Y., Edahiro, A., et al. (2017). Relationship between frailty and oral function in community-dwelling elderly adults. *Journal of the American Geriatrics Society, 65*, 66–76. https://doi.org/10.1111/jgs.14355.

Watanabe, Y., Arai, H., Hirano, H., Morishita, S., Ohara, Y., Edahiro, A., et al. (2018). Oral function as an indexing parameter for mild cognitive impairment in older adults. *Geriatrics & Gerontology International, 18*, 790–798. https://doi.org/10.1111/ggi.13259.

Winstein, C. J., Stein, J., Arena, R., Bates, B., Cherney, L. R., Cramer, S. C., et al. (2016). Guidelines for adult stroke rehabilitation and recovery: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, *47*, e98–e169. https://doi.org/10.1161/STR.0000000000000098.

Yagi, M., Fujita, N., Okada, E., Tsuji, O., Nagoshi, N., Tsuji, T., et al. (2018). Impact of frailty and comorbidities on surgical outcomes and complications in adult spinal disorders. *Spine*, *43*, 1259–1267. https://doi.org/10.1097/BRS.0000000000002596.

Appendices

Table A1. Correspondence rate for each item potentially associated with oral frailty

	Overall	Men	Women
Oral frailty	n (%)	n (%)	n (%)
Number of present teeth	190 (27.9)	64 (24.0)	126 (30.4)
Masticatory performance	27 (4.0)	7 (2.6)	20 (4.8)
Tongue pressure	141 (20.7)	62 (23.2)	79 (19.0)
ODK /ta/	39 (5.7)	9 (3.4)	30 (7.2)
Subjective difficulty in eating (yes)	147 (21.6)	42 (15.7)	105 (25.3)
Subjective difficulty in swallowing (yes)	182 (26.7)	67 (25.1)	115 (27.7)

ODK, oral diadochokinesis

Table A2. Correspondence rate for each item potentially associated with social frailty

Social frailty	Overall n (%)	Men n (%)	Women n (%)
Living alone (yes)	185 (27.1)	42 (15.7)	143 (34.5)
Going out less frequently compared with last year (yes)	127 (18.6)	50 (18.7)	77 (18.6)
Visits friends sometimes (no)	211 (30.9)	108 (40.4)	103 (24.8)
Feels helpful to friends or family (no)	57 (8.4)	36 (13.5)	21 (5.1)
Talks with someone every day (no)	65 (9.5)	27 (10.1)	38 (9.2)

Table A3. Correspondence rate for each item potentially associated with physical frailty

	Overall	Men	Women
Physical frailty	n (%)	n (%)	n (%)
Unintentional weight loss	83 (12.2)	47 (17.6)	36 (8.7)
Exhaustion	171 (25.1)	59 (22.1)	112 (27.0)
Low activity	102 (15.0)	39 (14.6)	63 (15.2)
Weakness	85 (12.5)	25 (9.4)	60 (14.5)
Slowness	47 (6.9)	17 (6.4)	30 (7.2)