HOKKAIDO UNIVERSITY

Title	Landing instructions focused on pelvic and trunk lateral tilt decrease the knee abduction moment during a single-leg drop vertical jump
Author(s)	Chijimatsu, Masato; Ishida, Tomoya; Yamanaka, Masanori et al.
Citation	Physical Therapy in Sport, 46, 226-233 https://doi.org/10.1016/j.ptsp.2020.09.010
Issue Date	2020-11
Doc URL	https://hdl.handle.net/2115/82776
Rights	© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Rights(URL)	https://creativecommons.org/licenses/by-nc-nd/4.0/
Type	journal article
File Information	Manuscript_Chijimatsu20210126_HUSCAP.pdf

Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

Landing Instructions Focused on Pelvic and Trunk Lateral Tilt Decrease the Knee Abduction Moment during a Single-leg Drop Vertical Jump

Authors: Masato Chijimatsu [a, b], Tomoya Ishida [a, *], Masanori Yamanaka [c], Shohei Taniguchi [a], Ryo Ueno [d], Ryohei Ikuta [e], Mina Samukawa [a], Takumi Ino [f], Satoshi Kasahara [a], Harukazu Tohyama [a]

Affiliations: [a]Faculty of Health Sciences, Hokkaido University, Sapporo, Hokkaido, Japan; [b]Department of Physical Therapy, Matsuda Orthopedic Memorial Hospital, Sapporo, Hokkaido, Japan; [c]Faculty of Health Sciences, Hokkaido Chitose College of Rehabilitation, Chitose, Hokkaido, Japan; [d] Department of Sport Science, University of Innsbruck, Innsbruck, Austria; [e]Hachioji Sports Orthopaedic Clinic, Hachioji, Tokyo, Japan; [f] Faculty of Health Sciences, Hokkaido University of Science, Sapporo, Hokkaido, Japan

Corresponding author

- Tomoya Ishida, PhD
- Faculty of Health Sciences, Hokkaido University, West 5, North 12, Kitaku, Sapporo, Hokkaido 060-0812, Japan
- Phone: +81 11-706-3531
- Email: t.ishida@hs.hokudai.ac.jp

Declarations:

- **Conflict of interest**: None declared.
- **Ethical approval**: This study was approved by the Institutional Review Board of Hokkaido University (approval number: 17-87).
- **Funding**: None declared.
- **Informed consent**: Written informed consent was obtained from all subjects.

Acknowledgments:

The authors would like to acknowledge our subjects for their willingness to participate in the study. We thank American Journal Experts (https://www.aje.com/) for editing a draft of this manuscript.

ABSTRACT

Objectives: To investigate the effects of pelvic and trunk lateral tilt-focused landing instructions on the knee abduction moment during the single-leg drop vertical jump task.

Design: Descriptive laboratory study.

Setting: Motion analysis laboratory.

Participants: Fifteen young, healthy female participants.

Main Outcome Measures: The participants performed 15 single-leg drop vertical jumps. Landing instructions with self-video recordings were provided so that the participants' pelvis and trunk remained horizontal in the frontal plane. Pelvic, trunk and knee kinematics and kinetics were evaluated using a three-dimensional motion analysis system before and after the landing instructions.

Results: The peak knee abduction moment significantly decreased postinstruction (preinstruction 22.6 ± 15.3 Nm, postinstruction 17.9 ± 15.4 Nm, $P = 0.004$), as did pelvic and trunk lateral tilt ($P < 0.01$). The knee abduction and internal rotation angles at initial contact significantly decreased postinstruction ($P = 0.037$, $P = 0.007$), with no significant change in the peak knee abduction and internal rotation angles from pre- to postinstruction.

Conclusions: Landing instructions focused on pelvic and trunk lateral tilt are effective in

decreasing the knee abduction moment during the single-leg drop vertical jump. Pelvic and trunk lateral tilt should be controlled to decrease the knee abduction moment during single-leg landing.

KEYWORDS

Anterior cruciate ligament, Injury prevention, Core, Single-leg landing

1. INTRODUCTION

Anterior cruciate ligament (ACL) injury is one of the most serious sport injuries. It takes 6 to 12 months to return to sport activities after ACL reconstruction, and some athletes who undergo ACL reconstruction give up returning to sports activities (Ardern, Webster, Taylor, & Feller, 2011). Approximately 70% of ACL injuries are observed in noncontact situations, such as jump landing, cutting and pivoting (Agel, Arendt, & Bershadsky, 2005; Boden, Dean, Feagin, & Garrett, 2000). The incidence of ACL injuries is 2 to 8 times higher in female athletes than in male athletes (Agel et al., 2005; Olsen, Myklebust, Engebretsen, & Bahr, 2004). Although ACL injury prevention programs that target female athletes have demonstrated a preventive effect (Kiani, Hellquist, Ahlqvist,

Gedeborg, Michaëlsson, & Byberg, 2010; LaBella, Huxford, Grissom, Kim, Peng, & Christoffel, 2011), the incidence of ACL injuries in female athletes remains high (Agel, Rockwood, & Klossner, 2016).

The current ACL injury prevention programs consist of different exercise types but the landing stabilization exercise to ensure proper landing technique and alignment is most effective (Petushek, Sugimoto, Stoolmiller, Smith, & Myer, 2019). Therefore, it is necessary to reveal landing instructions to modify poor landing technique for ACL injury prevention.

Large knee abduction and internal rotation in the small knee flexion during landing have been considered as biomechanical risk factors for ACL injuries and these factors should be improved for ACL injury prevention (Hewett et al., 2005; Kiapour et al., 2016; Koga et al., 2010). In particular, the large knee abduction moment during landing has been considered a major biomechanical risk factor for noncontact ACL injuries in female athletes (Hewett et al., 2005; Levine et al., 2013). Some studies performed with cadaveric landing simulations have suggested that knee abduction moment is one of the major contributing factors to ACL injuries (Bates et al., 2019; Navacchia, Bates, Schilaty, Krych, & Hewett, 2019; Ueno, Navacchia, Bates, Schilaty, Krych, & Hewett, 2020). Therefore,

the knee abduction moment during landing should be decreased by preventive training to prevent ACL injuries (Sugimoto, Myer, Foss, & Hewett, 2015).

Landing instructions are commonly used to improve landing biomechanics. Verbal instruction, video feedback, and the combination of verbal instruction and video feedback are used to provide the landing instructions (Dallinga, Benjaminse, Gokeler, Cortes, Otten, & Lemmink, 2017; Milner, Fairbrother, Srivatsan, & Zhang, 2012; Oñate, Guskiewicz, Marshall, Giuliani, Yu, & Garrett, 2005). Verbal instruction or video feedback alone did not decrease the knee abduction moment, during double-leg landing tasks (Dallinga et al., 2017; Milner et al., 2012; Oñate et al., 2005), while the landing instructions using the combination of verbal instruction and video feedback immediately decreased the knee abduction moment during double-leg landing tasks (Herman et al., 2009). Many previous studies have examined landing instructions during double-leg landing, but few have examined landing instructions during single-leg landing, which is one of the situations in which ACL injuries most frequently occur (Krosshaug et al., 2007; Olsen et al., 2004). Landing instructions increase knee flexion angle and decrease ground reaction force during single-leg landing (Cowling, Steele, & McNair, 2003; Elias, Hammill, & Mizner, 2015). However, none have reported the effects of landing instructions on the knee

abduction moment during single-leg landing. Previous studies have reported that the knee abduction moment is larger during single-leg landing than during double-leg landing (Wang, 2011; Yeow, Lee, & Goh, 2011). In addition, the knee abduction moment during single-leg landing is not correlated with the knee abduction moment during double-leg landing (Taylor, Ford, Nguyen, & Shultz, 2016). Therefore, it is unknown whether landing instructions decrease the knee abduction moment during single-leg landing.

The trunk motions in the frontal plane have been considered to influence knee abduction moment. A previous study showed that there is a significant correlation between the peak knee abduction moment and trunk lateral tilt angle at initial contact (IC) during single-leg landing (Dempsey, Elliott, Munro, Steele, & Lloyd, 2012). Other studies have reported that a larger trunk lateral tilt toward the stance leg increases the peak knee abduction moment during side-step cutting (Dempsey, Lloyd, Elliott, Steele, Munro, & Russo, 2007; Jamison, Pan, & Chaudhari, 2012). In addition, the increase in intended trunk lateral tilt or rotation during landing increased the knee abduction and internal rotation angles and moments (Critchley et al., 2020; Hinshaw, Davis, Layer, Wilson, Zhu, & Dai, 2019; Saito, Okada, Sasaki, & Wakasa, 2020). However, the effects of the landing instruction to decrease trunk lateral motion on knee kinematics were unclear. Pelvic lateral

120 tilt has also been shown to influence the frontal-plane knee moment during single-leg
121 standing and gait (Dunphy, Casey, Lomond, & Rutherford, 2016; Takacs & Hunt, 2012).
122 However, pelvic lateral tilt and its effect on knee abduction moment in jump-landing
123 training have not been studied (Dempsey, Elliott, Munro, Steele, Lloyd, 2014; Neilson,
124 Ward, Hume, Lewis, & McDaid, 2019). According to previous research, controlling
125 pelvic lateral tilt as well as trunk lateral tilt can decrease the knee abduction moment
126 during single-leg landing.

127 The primary purpose of the present study was to investigate the effects of landing
128 instructions focused on pelvic and trunk lateral tilt on the knee abduction moment during
129 single-leg landing in female participants. In addition, the second purpose was to
130 investigate the effects of the landing instructions on the knee abduction angle, flexion and
131 internal rotation angles and moments during single-leg landing because these variables
132 are also considered as biomechanical risk factors for ACL injuries (Hewett et al., 2005;
133 Kiapour et al., 2016; Koga et al., 2010) . The hypotheses were that the pelvic and trunk
134 lateral tilt angles decrease after the landing instructions are provided and that,
135 consequently, the knee abduction moment also decreases.

136

2. METHODS

2.1. Participants

Fifteen female participants (mean ± SD: age 20.7 ± 0.7 years; height 160.8 ± 5.2 cm; weight 54.7 ± 5.9 kg) participated in this study. A priori power analysis in a pilot study using 6 participants showed that 15 participants were required to achieve a statistical power of 0.8 with an alpha level of 0.05 and an effect size of 0.8. All participants had experience with regular sports activities (4 basketball, 2 volleyball, 2 badminton, 4 tennis, 2 track and field, 1 dancing) that were defined as sports activities for 30 minutes a day at least 3 times per week (Schmitz, Kulas, Perrin, Riemann, & Shultz, 2007). Exclusion criteria were a history of musculoskeletal injuries within the last 6 months, surgeries or fractures of the lower extremities or trunk. None of the participants met any of the exclusion criteria. All participants read and signed informed consent forms prior to their inclusion in this study. This research was approved by the institutional review board of our university (approval number: 17-87).

2.2. Procedures

The experimental procedure is displayed in Fig. 1. The participants performed a warm-

154 up on a stationary bicycle for 5 minutes. Next, data were collected during a static standing
155 trial for each participant. A single-leg drop vertical jump (SDVJ) was used for the practice
156 session as well as for the before-after landing instructions (Fig. 2). The SDVJ was selected
157 because the combination of the trunk lateral tilt angle and knee abduction angle displayed
158 during this task has been reported to be a significant predictor of noncontact knee injuries,
159 including ACL tears (Dingenen et al., 2015). All participants were barefoot to exclude the
160 effects of shoes on lower extremity kinematics and kinetics (Hong, Yoon, Kim, & Shin,
161 2014). The dominant leg (the side used for kicking a ball), which was the right leg in all
162 participants, was then assessed. Each participant stood on a 30-cm-tall box on their
163 dominant leg, then dropped from the box, and landed with their dominant leg on a force
164 plate. They then performed a maximum vertical jump immediately after landing. The
165 participants were asked to look forward and to keep their elbows and hands up to ear level
166 to avoid marker occlusion throughout the SDVJ task (Dingenen et al., 2015; Ishida et al.,
167 2015). Trials in which the nondominant leg touched the ground or the participant lost
168 balance during the test were defined as failed trials and were excluded from the analysis.
169 The landing instruction protocol and the number of practice trials were determined on
170 the basis of a previous study (Oñate et al., 2005). The participants were allowed to rest

171 between each trial and session, as needed. After a static standing trial was collected, the
172 preinstruction test consisting of 3 SDVJ trials was conducted (Fig. 1). During the
173 preinstruction test, the trials were recorded using a digital video camera (DCR-TRV900
174 NTSC, SONY, Tokyo, Japan) placed in a frontal plane view of the landing zone. After the
175 preinstruction test, the participants were instructed to land “keeping the trunk vertical and
176 pelvis horizontal without tilting in the frontal plane” by checking their preinstruction test
177 videos. The video of each trial was shown twice - the first time at normal speed and the
178 second one at a slower speed. Then 3 practice sessions were provided with each session
179 including 5 SDVJ trials. The trials were recorded using the digital video camera during
180 practice sessions, as in the preinstruction test. Between the sessions, the same “keep the
181 trunk vertical and pelvis horizontal without tilting in the frontal plane” landing
182 instructions were provided while showing the videos of their previous sessions. Landing
183 instructions were not individualized to each participant. After the practice sessions
184 followed by the landing instructions, postinstruction tests consisting of 3 SDVJ trials were
185 conducted.
186



Fig. 1. Experimental procedure

SDVJ indicates a single-leg drop vertical jump task.



Fig. 2. Single-leg drop vertical jump (SDVJ) task

Participants dropped from a 30-cm-tall box with one leg and then performed a maximum vertical jump on the same leg.

2.3. Data collection

All data were collected with Cortex 5.0.1 (Motion Analysis Corporation, Santa Rosa, CA, USA), a motion analysis system with seven high-speed cameras (Hawk cameras; Motion Analysis Corporation), and a force plate (Type 9286, Kistler AG, Winterthur, Switzerland). The sampling rates were set to be 200 Hz for the marker coordinate data and 1,000 Hz for the force plate data. Retroreflective markers were placed on the sacrum, right iliac crest, and both acromions, anterosuperior iliac spines (ASIS), greater trochanters, medial and lateral femoral condyles, medial and lateral malleoli, heels, and second and fifth metatarsal heads, and cluster markers were placed on the thigh and shank of the dominant leg (Ishida et al., 2015).

2.4. Data analysis

The marker coordinate data and force plate data were low-pass filtered using a zero-lag fourth-order Butterworth filter at 12 Hz. The knee joint angles and moments were calculated with SIMM 6.0.2 software (MusculoGraphics, Santa Rosa, CA, USA) using inverse kinematics and inverse dynamics, respectively, in a rigid-body dynamics pipeline. Each participant's anthropometric model was created based on the static trial and the pre-

212 measured foot length and width. The segment inertia properties were adjusted based on
213 each participant's body mass and segment lengths (de Leva, 1996). The segment
214 coordinate system was located such that the pelvis was at the midpoint of the right and
215 left ASIS markers, the femur was at the center of the femoral head, the tibia was at the
216 midpoint between the femoral medial and lateral condyles, and the tibia coordinate system
217 was fixed in the tibia (Delp, 1990). The pelvic and trunk lateral tilt were calculated as the
218 frontal plane projection angle with respect to the global coordinate system by using
219 MATLAB (MathWorks, Natick, Massachusetts, USA). The trunk lateral tilt angle was
220 defined as the angle between the line connecting the left shoulder marker and the left ASIS
221 marker and a vertical line, and positive values corresponded to tilting toward the support-
222 leg side (DiCesare, Bates, Myer, & Hewett, 2014). The pelvic lateral tilt angle was defined
223 as the angle between the line connecting both ASIS markers and a horizontal line, and
224 positive values corresponded to pelvic elevation on the contralateral-leg side (Fig. 3). All
225 angles measured during the static standing trial were set to 0°.
226



Fig. 3. The trunk and pelvic lateral tilt angles. (a) The trunk lateral tilt angle was defined as the angle between the line connecting the left shoulder marker and the left ASIS marker and a vertical line. (b) The pelvic lateral tilt angle was defined as the angle between the line connecting the right and left ASIS markers and a horizontal line.

IC was defined as when the vertical ground reaction force first exceeded 10 N, and toe-off (TO) was defined as when the vertical ground reaction force first fell below 10 N after IC (Ford, Myer, & Hewett, 2007). The landing phase was defined as the period from IC to peak knee flexion during the first landing. The peak knee abduction, flexion and internal rotation moments and peak ground reaction force during the landing phase were computed. The pelvic and trunk lateral tilt and knee abduction, flexion and internal rotation angles were calculated at IC and when the peak angle was attained during the landing phase. In addition, the jump height and the stance time were calculated to assess the potential effect of the landing instructions on performance (Dai, Garrett, Gross, Padua, Queen, & Yu, 2015). The jump height was calculated by subtracting the height of the sacrum marker in the static standing trial from the maximal height of the sacrum marker after TO. The stance time was calculated as the duration of stance phase from IC to TO. The mean of three trials for each of the pre- and postinstruction tests was used for statistical analysis.

2.5. Statistical analysis

Paired t-tests were performed to determine the effects of instructions on the kinematic,

kinetic and jump performance data. Pearson's correlation coefficients were used to determine the association between the change in peak knee abduction moment from pre- to postinstruction and the changes in pelvic and trunk lateral tilt angle from pre- to postinstruction. All statistical analyses were performed using IBM SPSS Statistics, version 22 (IBM, Armonk, NY, USA). The level of significance was set to be $P < 0.05$. In addition, effect sizes were calculated for all outcomes with Cohen's *d* using G*Power 3.1. (Institute of Experimental Psychology, Heinrich Heine University, Dusseldorf, Germany). The effect sizes were interpreted as follows: 0.80 indicated a large effect, 0.50 indicated a medium effect, and 0.20 indicated a small effect (Cohen, 1988).

3. RESULTS

All participants presented the pelvic and trunk lateral tilt angle at IC and peak during the preinstruction test. The landing instructions significantly decreased the pelvic lateral tilt angle at IC, with a large effect size ($P = 0.002$, $d = 1.00$, Fig. 4a). The peak pelvic lateral tilt angle also decreased from pre- to postinstruction, with a large effect size ($P = 0.001$, $d = 1.06$, Fig. 4a). After receiving the instructions, 13/15 (87%) of the participants had a decreased pelvic lateral tilt angle at IC and a decreased peak pelvic lateral tilt angle.

Additionally, the landing instructions significantly decreased the trunk lateral tilt angle at IC, with a large effect size ($P < 0.001$, $d = 1.49$, Fig. 4b). The peak trunk lateral tilt angle also decreased from pre- to postinstruction, with a large effect size ($P < 0.001$, $d = 1.66$, Fig. 4b). After receiving the instructions, 15/15 (100%) of the participants demonstrated decreases in the trunk lateral tilt angle at IC and the peak trunk lateral tilt angle.



Fig. 4. The pelvic (a) and trunk (b) lateral tilt angles recorded before and after the landing instructions focused on pelvic and trunk lateral tilt. The error bars indicate standard deviations. IC: initial contact

278 All participants demonstrated the knee abduction moment during the preinstruction
279 test. The peak knee abduction moment significantly decreased from pre- to postinstruction,
280 with a large effect size (P = 0.004, d = 1.31, Fig. 5, Table 1). After receiving the
281 instructions, 14/15 (93%) of the participants demonstrated decreases in the peak knee
282 abduction moment. There were no significant correlations between the change in the peak
283 knee abduction moment from pre- to postinstruction and the changes in the pelvic and
284 trunk lateral tilt angles at IC from pre- to postinstruction (pelvic tilt: P = 0.564, R = -
285 0.162; trunk tilt: P = 0.253, R = -0.315). Additionally, there were no significant
286 correlations between the change in the peak knee abduction moment from pre- to
287 postinstruction and the changes in the peak pelvic and trunk lateral tilt angles from pre-
288 to postinstruction (pelvic tilt: P = 0.857, R = -0.051; trunk tilt: P = 0.401, R = -0.234).
289 There were no significant differences in the peak knee flexion and internal rotation
290 moments and the peak vertical ground reaction force (Table 1).
291



Fig. 5. The peak knee abduction moment recorded before and after the landing instructions focused on pelvic and trunk lateral tilt. The error bars indicate standard deviations.

297 **Table 1**. Comparison of external knee joint moments and vertical ground reaction force
298 between the tests performed before and after landing instructions focused on pelvic and
299 trunk lateral tilt.

	preinstruction	postinstruction	*P* value
Peak external moment (Nm)			
knee flexion moment	142.9 ± 38.4	143.7 ± 30.3	0.815
knee abduction moment	22.6 ± 15.3	17.9 ± 15.4	**0.004**
knee internal rotation moment	27.3 ± 18.7	28.8 ± 19.7	0.468
Peak vertical ground reaction force (N)	2356.6 ± 400.3	2337.9 ± 448.5	0.402

300 The data are presented as the mean ± SD.
301

The knee abduction and internal rotation angles at IC significantly decreased postinstruction, with medium and large effect sizes ($P = 0.037$, $d = 0.78$; $P = 0.007$, $d = 0.82$), while there were no significant differences in the peak knee abduction and internal rotation angles (Table 2). For the knee flexion, there were no significant changes from pre- to postinstruction (Table 2). The jump height significantly decreased from pre- to postinstruction, with a large effect size ($P = 0.004$, $d = 0.88$), while the stance time did not change (Table 3).

310 **Table 2**. Comparison of knee kinematics between the tests performed before and after

311 landing instructions focused on pelvic and trunk lateral tilt.

	preinstruction	postinstruction	*P* value
Knee flexion angle (deg.)			
at IC	17.3 ± 5.4	16.8 ± 5.5	0.641
peak angle	60.6 ± 6.6	61.9 ± 5.5	0.423
Knee abduction angle (deg.)			
at IC	0.9 ± 2.1	0.3 ± 2.0	**0.037**
peak angle	9.2 ± 5.9	8.5 ± 5.9	0.163
Knee internal rotation angle (deg.)			
at IC	1.1 ± 6.7	-0.3 ± 7.1	**0.007**
peak angle	11.4 ± 7.6	11.0 ± 7.5	0.362

312 IC: initial contact

313 The data are presented as the mean ± SD.

314

315 **Table 3**. Comparison of the jump performance between the tests performed before and

316 after landing instructions focused on pelvic and trunk lateral tilt.

	preinstruction	postinstruction	*P* value
Jump height (mm)	196.9 ± 20.0	182.5 ± 22.5	**0.004**
Stance time (ms)	318.4 ± 40.0	323.9 ± 39.3	0.370

317 The data are presented as the mean ± SD.

318

4. DISCUSSION

The primary purpose of this study was to determine the effects of landing instructions to keep the pelvis horizontal and the trunk vertical in the frontal plane on the knee abduction moment during the SDVJ task. The most important findings were that the landing instructions significantly decreased the peak knee abduction moment as well as the pelvic and trunk lateral tilt angles during the SDVJ. However, the knee flexion angle and vertical ground reaction force did not change from pre- to postinstruction. These findings supported our hypothesis that the knee abduction moment also decreases by decreasing the pelvic and trunk lateral tilt angles after the landing instructions.

In the present study, the expected changes in the frontal-plane pelvic and trunk kinematics were observed postinstruction. Therefore, the protocol for the landing instructions and the content of the landing instructions (i.e., to keep the trunk vertical and pelvis horizontal without tilting them in the frontal plane) is valid. Providing landing instructions and short practice sessions is an effective method for reducing pelvic and trunk lateral tilt during the SDVJ.

The peak knee abduction moment significantly decreased postinstruction compared with preinstruction. This is the first study to report that landing instructions decrease the

peak knee abduction moment during the single-leg landing task. This finding is consistent with those from previous studies that have reported that the increase in trunk lateral tilt or rotation increase the knee abduction moment during landing (Critchley et al., 2020; Dempsey et al., 2012). A previous study on cutting maneuvers reported results that are consistent with those in the present study (Dempsey, Lloyd, Elliott, Steele, & Munro, 2009). It has been suggested that the knee abduction moment during single-leg stance is affected by the center of mass (COM) position relative to the knee (Celebrini, Eng, Miller, Ekegren, Johnston, & MacIntyre, 2012). When the pelvic and trunk lateral tilt angles decrease during single-leg support, the COM moves toward the medial side of the stance leg (Takacs & Hunt, 2012). Then, the moment arm for knee abduction is shortened. This concept is assumed to be the mechanism by which knee abduction moment decreased in the present study. The present findings suggested that the landing instructions focused on pelvic and trunk lateral tilt decreased the peak knee abduction moment by decreasing the pelvic and trunk lateral tilt angles.

It was expected that the larger the decreases in the pelvic and trunk lateral tilt angles were, the larger the decrease in the peak knee abduction moment from pre- to postinstruction. However, there were no significant correlations between the change in

353 the peak knee abduction moment and the changes in the pelvic and trunk lateral tilt angles
354 in the present study. All participants demonstrated decreases in the trunk lateral tilt angle,
355 but some did not demonstrate decreases in the peak knee abduction moment or pelvic
356 lateral tilt angle. In this study, the landing instructions decreased the peak knee abduction
357 moment by decreasing the pelvic and trunk lateral tilt angles, but the magnitude of the
358 effects of decreased pelvic and trunk lateral tilt angle may have differed among
359 participants. Therefore, the linear relationships between the change in the peak knee
360 abduction moment and the changes in the pelvic and trunk lateral tilt angles may not have
361 been detected. The effects of other factors that were correlated with the knee abduction
362 moment during single-leg landing (e.g., trunk rotation, knee flexion and rotation and foot
363 rotation) should be investigated in the future (Dempsey et al., 2012).

364 In a previous study, a 6-week landing training program did not decrease the peak knee
365 abduction moment during the single-leg landing with ball catching (Dempsey et al., 2014).
366 In that study, although the landing instructions were focused on increasing knee flexion,
367 reducing trunk lateral tilt and reducing trunk rotation, a significant change was found only
368 in knee flexion, which increased. Therefore, it may have been difficult for participants to
369 focus on knee and trunk motions while catching a ball. The present study showed that

simple instructions to keep the pelvis and trunk positioned horizontally decreased the peak knee abduction moment. Future studies should be conducted to investigate the effects of simple landing instructions focused on pelvic and trunk lateral tilt on the knee abduction moment during single-leg landings in game-like situations.

Decreases in knee abduction moment are very important for ACL injury prevention. In a cadaver study, combined external loads of knee abduction moment, internal tibial rotation moment, and anterior tibial shear force under axial impact produced clinically relevant ACL injuries, and only knee abduction moment significantly contributed to peak ACL strain at injuries (Levine et al., 2013). Excessive knee abduction moment is considered a contributing factor of ACL injuries. (Bates et al., 2019; Navacchia et al., 2019; Ueno et al., 2020). Therefore, landing instructions focused on pelvic lateral tilt as well as trunk lateral tilt should be given to decrease the knee abduction moment during single-leg landing.

A previous study reported that the trunk lateral tilt angle during the SDVJ task was one of the significant predictors of noncontact knee injuries, including ACL tears (Dingenen et al., 2015). A video analysis study of ACL injuries during single-leg movements showed that the trunk lateral tilt angles of ACL-injured female athletes were

larger than those of control female athletes (Hewett, Torg, & Boden, 2009). There was a significant association between a large pelvic lateral tilt angle and ACL injury risk in female athletes (Leppänen et al., 2020). The present study showed that landing instructions are effective in reducing pelvic and trunk lateral tilt. Therefore, landing instructions focused on pelvic lateral tilt as well as trunk lateral trunk lean should be incorporated into ACL injury prevention training programs. In prevention programs that have recently been shown to be successful, landing instructions about controlling not only the knee position but also the trunk position have been given (Omi et al., 2018).

The findings regarding the knee abduction angle at IC after landing instructions are consistent with those from previous studies that have reported that the increase in intended trunk lateral tilt or rotation increases the knee abduction angle at IC or at the early landing phase (Critchley et al., 2020; Hinshaw et al., 2019). As a result of the decrease in the knee abduction moment, the knee abduction angle was also expected to decrease. However, the results of this study did not reveal a change in the peak knee abduction angle, unlike the findings of a previous study, which reported that the increase in intended trunk lateral tilt increases the peak knee abduction angle (Saito et al., 2020). The peak knee abduction angle may be increased by factors other than the external knee abduction moment. A

previous study suggested that knee abduction motion during the double-leg drop vertical jump is high to prepare the individual for the subsequent jump after landing (Ishida et al., 2018). An alternative intervention other than providing landing instructions focused on pelvic and trunk lateral tilt should be developed to decrease the knee abduction angle during the SDVJ.

The findings regarding the decreased knee internal rotation angle at IC after the landing instructions in the present study supported the results from previous studies on the effects of intended trunk lateral tilt (Hinshaw et al., 2019). The peak knee internal rotation angle and moment did not change from pre- to postinstruction in the present study. The increase in intended trunk rotation increased the peak knee internal rotation angle and moment (Critchley et al., 2020). Adding landing instructions focused on pelvic and trunk rotation may help to decrease the internal rotation angle and moment during landing.

The results in this study showed that the knee flexion angle and moment and vertical ground reaction force did not change from pre- to postinstruction. In previous studies, landing instructions provided to increase knee flexion and to decrease the impact of landing have demonstrated the intended effects during single-leg landing (Cowling et al., 2003; Elias et al., 2015). The landing instructions were “land with your knee bent” or

421 “land as softly and quietly as possible” in the previous studies. The landing instructions
422 focused only on the pelvic and trunk lateral tilt did not change the knee flexion angle and
423 moment or the vertical ground reaction force in this study. The instructions that address
424 modifications for pelvic and trunk lateral tilt, knee flexion and landing impact may
425 improve both the knee abduction moment and vertical ground reaction force. However,
426 the combined effects of these instructions have not been reported, and future research is
427 warranted.

428 The results of this study showed that the stance time did not change, while the jump
429 height significantly decreased after receiving the instructions. A previous study also
430 reported performance deterioration with the immediate modification of landing
431 techniques, as found in the present study (Dai et al., 2015). In previous studies, 6-week
432 ACL injury prevention programs improved ACL injury biomechanical risks and
433 performance at the same time (Chappell & Limpisvasti, 2008; Myer, Ford, Palumbo, &
434 Hewett, 2005). A certain period of training may be needed to improve landing techniques
435 while maintaining jump performance.

436 This study had some limitations. First, the present study tested only the immediate
437 effect of landing instructions. Therefore, additional studies are needed to assess the

retention and transfer of these effects to actual sports activities. Second, the present study included only female participants. A previous study showed that the trunk lateral tilt angle of male participants was smaller than that of female participants during the single-leg squat task (Nakagawa, Moriya, Maciel, & Serrão, 2012). The landing instructions used in this study may be less effective in male participants than in female participants.

5. CONCLUSION

The present study showed that landing instructions focused on pelvic and trunk lateral tilt decrease the peak knee abduction moment as well as the pelvic and trunk lateral tilt angles during single-leg landing. These findings indicate that the control of the pelvis and trunk should be emphasized to decrease the knee abduction moment during single-leg landing.

REFERENCES

Agel, J., Arendt, E. A., & Bershadsky, B. (2005). Anterior cruciate ligament injury in national collegiate athletic association basketball and soccer. *The American Journal of Sports Medicine*, *33*(4), 524–531. https://doi.org/10.1177/0363546504269937

455 Agel, J., Rockwood, T., & Klossner, D. (2016). Collegiate ACL injury rates across 15
456 sports: National collegiate athletic association injury surveillance system data update
457 (2004-2005 through 2012-2013). *Clinical Journal of Sport Medicine*, *26*(6), 518–
458 523. https://doi.org/10.1097/JSM.0000000000000290
459 Ardern, C. L., Webster, K. E., Taylor, N. F., & Feller, J. A. (2011). Return to sport
460 following anterior cruciate ligament reconstruction surgery: A systematic review
461 and meta-analysis of the state of play. *British Journal of Sports Medicine*, *45*(7),
462 596–606. https://doi.org/10.1136/bjsm.2010.076364
463 Bates, N. A., Mejia Jaramillo, M. C., Vargas, M., McPherson, A. L., Schilaty, N. D.,
464 Nagelli, C. V., Krych, A. J., & Hewett, T. E. (2019). External loads associated with
465 anterior cruciate ligament injuries increase the correlation between tibial slope and
466 ligament strain during in vitro simulations of in vivo landings. *Clinical*
467 *Biomechanics*, *61*, 84–94. https://doi.org/10.1016/j.clinbiomech.2018.11.010
468 Boden, B., Dean, G. S., Feagin, J. A., Jr, & Garrett, W. (2000). Mechanisms of anterior
469 cruciate ligament injury. *Orthopedics*, 23(6), 573-578.
470 Celebrini, R. G., Eng, J. J., Miller, W. C., Ekegren, C. L., Johnston, J. D., & MacIntyre,
471 D. L. (2012). The effect of a novel movement strategy in decreasing ACL risk

472 factors in female adolescent soccer players. *Journal of Strength Condition Research*,
473 *26*(12), 3406–3417. https://doi.org/10.1519/JSC.0b013e3182472fef
474 Chappell, J. D., & Limpisvasti, O. (2008). Effect of a neuromuscular training program
475 on the kinetics and kinematics of jumping tasks. *The American Journal of Sports*
476 *Medicine*, *36*(6), 1081–1086. https://doi.org/10.1177/0363546508314425
477 Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. (2nd ed.).
478 Hillsdale, New Jersey: Lawrence Erlbaum Associates.
479 https://doi.org/10.4324/9780203771587
480 Cowling, E. J., Steele, J. R., & McNair, P. J. (2003). Effect of verbal instructions on
481 muscle activity and risk of injury to the anterior cruciate ligament during landing.
482 *British Journal of Sports Medicine*, *37*(2), 126–130.
483 https://doi.org/10.1136/bjsm.37.2.126
484 Critchley, M. L., Davis, D. J., Keener, M. M., Layer, J. S., Wilson, M. A., Zhu, Q., &
485 Dai, B. (2020). The effects of mid-flight whole-body and trunk rotation on landing
486 mechanics: implications for anterior cruciate ligament injuries. *Sports Biomechanics*,
487 *19*(4), 421–437. https://doi.org/10.1080/14763141.2019.1595704
488 Dai, B., Garrett, W. E., Gross, M. T., Padua, D. A., Queen, R. M., & Yu, B. (2015). The

effects of 2 landing techniques on knee kinematics, kinetics, and performance during stop-jump and side-cutting tasks. *The American Journal of Sports Medicine*, *43*(2), 466–474. https://doi.org/10.1177/0363546514555322

Dallinga, J., Benjaminse, A., Gokeler, A., Cortes, N., Otten, E., & Lemmink, K. (2017). Innovative video feedback on jump landing improves landing technique in males. *International Journal of Sports Medicine*, *38*(2), 150–158. https://doi.org/10.1055/s-0042-106298

de Leva, P. (1996). Adjustments to Zatsiorsky-Seluyanov's segment inertia parameters. *Journal of Biomechanics*, *29*(9), 1223–1230. https://doi.org/ 10.1016/0021-9290(95)00178-6

Delp, S. L. (1990). *Surgery simulation: A computer graphics system to analyze and design musculoskeletal reconstructions of the lower limb.* Ph.D. thesis, Department of Mechanical Engineering, Stanford University.

Dempsey, A. R., Elliott, B. C., Munro, B. J., Steele, J. R., & Lloyd, D. G. (2012). Whole body kinematics and knee moments that occur during an overhead catch and landing task in sport. *Clinical Biomechanics*, *27*(5), 466–474. https://doi.org/10.1016/j.clinbiomech.2011.12.001

506 Dempsey, A. R., Elliott, B. C., Munro, B. J., Steele, J. R., & Lloyd, D. G. (2014). Can
507 technique modification training reduce knee moments in a landing task? *Journal of*
508 *Applied Biomechanics*, *30*(2), 231–236. https://doi.org/10.1123/jab.2013-0021
509 Dempsey, A. R., Lloyd, D. G., Elliott, B. C., Steele, J. R., & Munro, B. J. (2009).
510 Changing sidestep cutting technique reduces knee valgus loading. *The American*
511 *Journal of Sports Medicine*, *37*(11), 2194–2200.
512 https://doi.org/10.1177/0363546509334373
513 Dempsey, A. R., Lloyd, D. G., Elliott, B. C., Steele, J. R., Munro, B. J., & Russo, K. A.
514 (2007). The effect of technique change on knee loads during sidestep cutting.
515 *Medicine and Science in Sports and Exercise*, *39*(10), 1765–1773.
516 https://doi.org/10.1249/mss.0b013e31812f56d1
517 DiCesare, C. A., Bates, N. A., Myer, G. D., & Hewett, T. E. (2014). The validity of 2-
518 dimensional measurement of trunk angle during dynamic tasks. *International*
519 *Journal of Sports Physical Therapy*, *9*(4), 420–427.
520 Dingenen, B., Malfait, B., Nijs, S., Peers, K. H. E., Vereecken, S., Verschueren, S. M. P.,
521 & Staes, F. F. (2015). Can two-dimensional video analysis during single-leg drop
522 vertical jumps help identify non-contact knee injury risk? A one-year prospective

study. *Clinical Biomechanics*, *30*(8), 781–787. https://doi.org/10.1016/j.clinbiomech.2015.06.013

Dunphy, C., Casey, S., Lomond, A., & Rutherford, D. (2016). Contralateral pelvic drop during gait increases knee adduction moments of asymptomatic individuals. *Human Movement Science*, *49*, 27–35. https://doi.org/10.1016/j.humov.2016.05.008

Elias, A. R. C., Hammill, C. D., & Mizner, R. L. (2015). Changes in quadriceps and hamstring cocontraction following landing instruction in patients with anterior cruciate ligament reconstruction. *Journal of Orthopaedic & Sports Physical Therapy*, *45*(4), 273–280. https://doi.org/10.2519/jospt.2015.5335

Ford, K. R., Myer, G. D., & Hewett, T. E. (2007). Reliability of landing 3D motion analysis: Implications for longitudinal analyses. *Medicine and Science in Sports and Exercise*, *39*(11), 2021–2028. https://doi.org/10.1249/mss.0b013e318149332d

Herman, D. C., Oñate, J. A., Weinhold, P. S., Guskiewicz, K. M., Garrett, W. E., Yu, B., & Padua, D. A. (2009). The effects of feedback with and without strength training on lower extremity biomechanics. *The American Journal of Sports Medicine*, *37*(7), 1301–1308. https://doi.org/10.1177/0363546509332253

Hewett, T. E., Myer, G. D., Ford, K. R., Heidt, R. S., Jr, Colosimo, A. J., Mclean, S. G.,

van den Bogert, A. J., Paterno, M.V., & Succop, P. (2005). Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: A prospective study. *The American Journal of Sports Medicine*, *33*(4), 492–501. https://doi.org/10.1177/0363546504269591

Hewett, T. E., Torg, J. S., & Boden, B. P. (2009). Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: Lateral trunk and knee abduction motion are combined components of the injury mechanism. *British Journal of Sports Medicine*, *43*(6), 417–422. https://doi.org/10.1136/bjsm.2009.059162

Hinshaw, T. J., Davis, D. J., Layer, J. S., Wilson, M. A., Zhu, Q., & Dai, B. (2019). Mid-flight lateral trunk bending increased ipsilateral leg loading during landing: a center of mass analysis. *Journal of Sports Sciences*, *37*(4), 414–423. https://doi.org/10.1080/02640414.2018.1504616

Hong, Y. G., Yoon, Y. J., Kim, P., & Shin, C. S. (2014). The kinematic/kinetic differences of the knee and ankle joint during single-leg landing between shod and barefoot condition. *International Journal of Precision Engineering and Manufacturing*, *15*(10), 2193–2197. https://doi.org/10.1007/s12541-014-0581-9

557 Ishida, T., Koshino, Y., Yamanaka, M., Ueno, R., Taniguchi, S., Samukawa, M., Saito,
558 H., Matsumoto, H., Aoki, Y., & Tohyama, H. (2018). The effects of a subsequent
559 jump on the knee abduction angle during the early landing phase. *BMC*
560 *Musculoskeletal Disorders*, *19*(1), 379. https://doi.org/10.1186/s12891-018-2291-4
561 Ishida, T., Yamanaka, M., Takeda, N., Homan, K., Koshino, Y., Kobayashi, T.,
562 Matsumoto, H., & Aoki, Y. (2015). The effect of changing toe direction on knee
563 kinematics during drop vertical jump: A possible risk factor for anterior cruciate
564 ligament injury. *Knee Surgery, Sports Traumatology, Arthroscopy*, *23*(4), 1004–
565 1009. https://doi.org/10.1007/s00167-013-2815-2
566 Jamison, S. T., Pan, X., & Chaudhari, A. M. W. (2012). Knee moments during run-to-cut
567 maneuvers are associated with lateral trunk positioning. *Journal of Biomechanics*,
568 *45*(11), 1881–1885. https://doi.org/10.1016/j.jbiomech.2012.05.031
569 Kiani, A., Hellquist, E., Ahlqvist, K., Gedeborg, R., Michaëlsson, K., & Byberg, L.
570 (2010). Prevention of soccer-related knee injuries in teenaged girls. *Archives of*
571 *Internal Medicine*, *170*(1), 43–49. https://doi.org/10.1001/archinternmed.2009.289
572 Kiapour, A. M., Demetropoulos, C. K., Kiapour, A., Quatman, C. E., Wordeman, S. C.,
573 Goel, V. K., & Hewett, T. E. (2016). Strain response of the anterior cruciate ligament

574 to uniplanar and multiplanar loads during simulated landings. *The American Journal*
575 *of Sports Medicine*, *44*(8), 2087–2096. https://doi.org/10.1177/0363546516640499
576 Koga, H., Nakamae, A., Shima, Y., Iwasa, J., Myklebust, G., Engebretsen, L., Bahr, R.,
577 Krosshaug, T. (2010). Mechanisms for noncontact anterior cruciate ligament
578 injuries: Knee joint kinematics in 10 injury situations from female team handball
579 and basketball. *The American Journal of Sports Medicine*, *38*(11), 2218–2225.
580 https://doi.org/10.1177/0363546510373570
581 Krosshaug, T., Nakamae, A., Boden, B. P., Engebretsen, L., Smith, G., Slauterbeck, J. R.,
582 Hewett, T.E., & Bahr, R. (2007). Mechanisms of anterior cruciate ligament injury in
583 basketball: Video analysis of 39 cases. *The American Journal of Sports Medicine*,
584 *35*(3), 359–367. https://doi.org/10.1177/0363546506293899
585 LaBella, C. R., Huxford, M. R., Grissom, J., Kim, K. Y., Peng, J., & Christoffel, K. K.
586 (2011). Effect of neuromuscular warm-up on injuries in female soccer and basketball
587 athletes in urban public high schools: Cluster randomized controlled trial. *Archives*
588 *of Pediatrics and Adolescent Medicine*, *165*(11), 1033–1040.
589 https://doi.org/10.1001/archpediatrics.2011.168
590 Leppänen, M., Rossi, M. T., Parkkari, J., Heinonen, A., Äyrämö, S., Krosshaug, T.,

591 Vasankari, T., Kannus, P., & Pasanen, K. (2020). Altered hip control during a
592 standing knee-lift test is associated with increased risk of knee injuries.
593 *Scandinavian Journal of Medicine and Science in Sports*, *30*(5), 922-931.
594 https://doi.org/10.1111/sms.13626
595 Levine, J. W., Kiapour, A. M., Quatman, C. E., Wordeman, S. C., Goel, V. K., Hewett,
596 T. E., & Demetropoulos, C. K. (2013). Clinically relevant injury patterns after an
597 anterior cruciate ligament injury provide insight into injury mechanisms. *The*
598 *American Journal of Sports Medicine*, *41*(2), 385–395.
599 https://doi.org/10.1177/0363546512465167
600 Milner, C. E., Fairbrother, J. T., Srivatsan, A., & Zhang, S. (2012). Simple verbal
601 instruction improves knee biomechanics during landing in female athletes. *Knee*,
602 *19*(4), 399–403. https://doi.org/10.1016/j.knee.2011.05.005
603 Myer, G. D., Ford, K. R., Palumbo, J. P., & Hewett, T. E. (2005). Neuromuscular training
604 improves performance and lower-extremity biomechanics in female athletes.
605 *Journal of Strength and Conditioning Research*, *19*(1), 51–60.
606 https://doi.org/10.1519/13643.1
607 Nakagawa, T. H., Moriya, E. T. U., Maciel, C. D., & SerrãO, F. V. (2012). Trunk, pelvis,

608 hip, and knee kinematics, hip strength, and gluteal muscle activation during a single-
609 leg squat in males and females with and without patellofemoral pain syndrome.
610 *Journal of Orthopaedic & Sports Physical Therapy*, *42*(6), 491–501.
611 https://doi.org/10.2519/jospt.2012.3987
612 Navacchia, A., Bates, N. A., Schilaty, N. D., Krych, A. J., & Hewett, T. E. (2019). Knee
613 abduction and internal rotation moments increase ACL force during landing through
614 the posterior slope of the tibia. *Journal of Orthopaedic Research*, *37*(8), 1730-1742.
615 https://doi.org/10.1002/jor.24313
616 Neilson, V., Ward, S., Hume, P., Lewis, G., & McDaid, A. (2019, September 1). Effects
617 of augmented feedback on training jump landing tasks for ACL injury prevention:
618 A systematic review and meta-analysis. *Physical Therapy in Sport*, *39*, 126–135.
619 https://doi.org/10.1016/j.ptsp.2019.07.004
620 Olsen, O. E., Myklebust, G., Engebretsen, L., & Bahr, R. (2004). Injury mechanisms for
621 anterior cruciate ligament injuries in team handball. *The American Journal of Sports*
622 *Medicine*, *32*(4), 1002–1012. https://doi.org/10.1177/0363546503261724
623 Omi, Y., Sugimoto, D., Kuriyama, S., Kurihara, T., Miyamoto, K., Yun, S., Kawashima,
624 T., & Hirose, N. (2018). Effect of hip-focused injury prevention training for anterior

cruciate ligament injury reduction in female basketball players: A 12-year prospective intervention study. *The American Journal of Sports Medicine*, *46*(4), 852–861. https://doi.org/10.1177/0363546517749474

Oñate, J. A., Guskiewicz, K. M., Marshall, S. W., Giuliani, C., Yu, B., & Garrett, W. E. (2005). Instruction of jump-landing technique using videotape feedback: Altering lower extremity motion patterns. *The American Journal of Sports Medicine*, *33*(6), 831–842. https://doi.org/10.1177/0363546504271499

Petushek, E. J., Sugimoto, D., Stoolmiller, M., Smith, G., & Myer, G. D. (2019). Evidence-based best-practice guidelines for preventing anterior cruciate ligament injuries in young female athletes: A systematic review and meta-analysis. *The American Journal of Sports Medicine*, *47*(7), 1744–1753. https://doi.org/10.1177/0363546518782460.

Saito, A., Okada, K., Sasaki, M., & Wakasa, M. (2020). Influence of the trunk position on knee kinematics during the single-leg landing: implications for injury prevention. *Sports Biomechanics*, Online ahead of print, 1-14. https://doi.org/10.1080/14763141.2019.1691642

Schmitz, R. J., Kulas, A. S., Perrin, D. H., Riemann, B. L., & Shultz, S. J. (2007). Sex

differences in lower extremity biomechanics during single leg landings. *Clinical Biomechanics*, *22*(6), 681–688. https://doi.org/10.1016/j.clinbiomech.2007.03.001

Sugimoto, D., Myer, G. D., Foss, K. D. B., & Hewett, T. E. (2015). Specific exercise effects of preventive neuromuscular training intervention on anterior cruciate ligament injury risk reduction in young females: Meta-analysis and subgroup analysis. *British Journal of Sports Medicine*, *49*(5), 282–289. https://doi.org/10.1136/bjsports-2014-093461

Takacs, J., & Hunt, M. A. (2012). The effect of contralateral pelvic drop and trunk lean on frontal plane knee biomechanics during single limb standing. *Journal of Biomechanics*, *45*(16), 2791–2796. https://doi.org/10.1016/j.jbiomech.2012.08.041

Taylor, J. B., Ford, K. R., Nguyen, A.-D., & Shultz, S. J. (2016). Biomechanical comparison of single- and double-leg jump landings in the sagittal and frontal plane. *Orthopaedic Journal of Sports Medicine*, *4*(6), 2325967116655158. https://doi.org/10.1177/2325967116655158

Ueno, R., Navacchia, A., Bates, N. A., Schilaty, N. D., Krych, A. J., & Hewett, T. E. (2020). Analysis of internal knee forces allows for the prediction of rupture events in a clinically relevant model of anterior cruciate ligament injuries. *Orthopaedic*

659 *Journal of Sports Medicine*, *8*(1), 2325967119893758.
660 https://doi.org/10.1177/2325967119893758

661 Wang, L. I. (2011). The lower extremity biomechanics of single- and double-leg stop-
662 jump tasks. *Journal of Sports Science and Medicine*, *10*(1), 151–156.

663 Yeow, C. H., Lee, P. V. S., & Goh, J. C. H. (2011). An investigation of lower extremity
664 energy dissipation strategies during single-leg and double-leg landing based on
665 sagittal and frontal plane biomechanics. *Human Movement Science*, *30*(3), 624–635.
666 https://doi.org/10.1016/j.humov.2010.11.010

667